1 **Classification**: Biological Sciences: Microbiology
2
3 **Title**: Ligand-Mediated Biofilm Formation via Enhanced Physical Interaction Between a
4 Diguanylate Cyclase and Its Receptor
5

6 **Authors**: David Giacalone[1], T. Jarrod Smith[1], Alan J. Collins[1], Holger Sondermann[3], Lori J. Koziol[2],
7 and George A. O'Toole[1]
8
Author Affiliations:
[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth,
Hanover, NH 03755
[2]Department of Biology, New England College, Henniker, NH 03242
[3]Department of Molecular Medicine, College of Veterinary Medicine, Cornell University, Ithaca,
NY 14853
15
16 **Corresponding author**: Geoirge A. O'Toole; address: Geisel School of Medicine at Dartmouth,
17 Dept. of Microbiology and Immunology, Remson 202, 66 North College St., Hanover, NH 03755;
18 phone: 603-650-1248; email: georgeo@Dartmouth.edu

19
Key words: biofilm, c-di-GMP, receptor, *Pseudomonas fluorescens*, CACHE domain, signaling

Abstract. The second messenger, cyclic dimeric GMP (c-di-GMP) regulates biofilm formation
and surface attachment for many bacteria. Biofilm formation of *Pseudomonas fluorescens* Pf0-
1 is controlled by the transfer of c-di-GMP to the inner-membrane protein, LapD. LapD-bound
c-di-GMP in turn inhibits proteolytic cleavage of the adhesin LapA by the periplasmic protease
LapG, which allows localization of LapA to the cell surface and thereby promotes biofilm
formation. LapD is central to a complex network of c-di-GMP-mediated biofilm formation. In
this study, we examine how signaling specificity of c-di-GMP by a DGC is achieved by small
ligand sensing through characterization of the <u>ca</u>lcium <u>ch</u>annel <u>che</u>motaxis (CACHE) receptor of
the DGC, GcbC. We provide evidence that biofilm formation is enhanced by the
environmentally relevant organic acid citrate, in a GcbC-dependent manner through enhanced
LapA localization to the cell surface. In the presence of citrate, the GcbC shows enhanced
physical interaction with LapD. GcbC also shows increased c-di-GMP production when LapD is
present, and this enhanced GcbC-LapD interaction by citrate further drives c-di-GMP synthesis.
Given that LapD can interact with a dozen different DGCs of *P. fluorescens*, many of which have
ligand-binding domains, the ligand-mediated enhanced signaling of LapD-GcbC described here
is likely a conserved mechanism of signaling in this network.

Significance Statement. Biofilm formation is a highly regulated event controlled by c-di-GMP
signaling. In many bacteria, dozens of DGCs produce this dinucleotide signal, however it is
unclear how undesired cross-talk is mitigated in the context of this soluble signal, and how c-di-
GMP signaling is regulated in the context of environmental inputs. Here we show a ligand-
mediated mechanism of signaling specificity whereby citrate enhances catalysis of the c-di-GMP
synthesizing diguanylate cyclase (DGC) via increased physical interaction with its cognate
receptor. We envision a scenario wherein a "cloud" of weakly interacting DGCs can increase
specific interaction with their receptor in response to appropriate environmental signals,
concomitantly boosting c-di-GMP production, ligand-specific signaling and biofilm formation.

Text

For most bacteria, surface attachment and biofilm formation is a highly regulated event (1, 2).
The bacterial intracellular second messenger, cyclic dimeric GMP (c-di-GMP) regulates biofilm
formation by regulating a diversity of biofilm-relevant outputs, including surface attachment
(3), flagellar motility (4), extracellular polysaccharide production (5, 6), adhesin localization (7),
and transcriptional control of pathways important for attachment (8). An important research
theme has been the focus on c-di-GMP signaling specificity in the context of microbes that can
have >50 proteins that make, degrade and bind this second messenger.

Biofilm formation by *Pseudomonas fluorescens* Pf0-1 occurs when the adhesin LapA localizes to
the cell surface (9). LapA is maintained on the cell surface when the inner membrane protein

LapD binds c-di-GMP, which in turn inhibits cleavage of LapA by the periplasmic protease LapG
(10, 11). The c-di-GMP-bound LapD sequesters LapG, thus this protease is not available to
target the N-terminal cleavage site of LapA (11). One example of how c-di-GMP is specifically
transferred to the LapD receptor is by physical interaction with a diguanylate cyclase (DGC)
(12). The DGC called GcbC has been shown to physically interact with LapD, a c-di-GMP
receptor located in the inner membrane, utilizing a surface exposed α-helix of the GGDEF
domain on GcbC and the surface exposed α-helix of the EAL domain of LapD (12). This direct
interaction model was proposed as one means to confer signaling specificity.

GcbC is an inner membrane protein that contains a calcium channel chemotaxis receptor
(CACHE) located N-terminal to its GGDEF domain. CACHE domains are responsible for small
ligand sensing (13). Many signal transduction proteins, including DGCs and histidine kinases,
contain CACHE domains (14). GcbC, along with five other DGCs encoded on the *P. fluorescens*
Pf0-1 genome (Pfl01_2295, Pfl01_2297, Pfl01_1336, Pfl01_3550, and Pfl01_3800), contains a
CACHE domain located N-terminally to the GGDEF domain, suggesting that these six DGCs are
capable of sensing and responding to small ligands. The predicted domain organization of
GcbC, Pfl01_2295, and Pfl01_2297 are shown in Figure 1A.

In this study, we analyze the CACHE domain of GcbC and identify the environmentally relevant
organic acid citrate as a ligand for this DGC. We show citrate-enhanced physical interaction
with GcbC and LapD, and that this enhanced interaction promotes increased c-di-GMP synthesis
by GcbC, thereby promoting biofilm formation. We propose that ligand-mediated
enhancement of DGC interactions with a receptor could serve as a general mechanism to
confer specificity to this complex signaling network.

Results

Citrate-mediated Biofilm Enhancement is Dependent on GcbC Activity. We identified a CACHE
domain in the periplasmic loop of GcbC (Figure 1A). To identify small molecules that the CACHE
domain of GcbC may bind, we performed a CLUSTAL alignment of the amino acid sequence of
CACHE domains of known structure (15). GcbC showed the highest amino acid similarity to
rpHK1S-Z16 (PDB ID: 3LIF) of *Rhodopseudomonas palustris* with 31% identity (Fig. S1). When
crystallized, the CACHE domain of rpHK1S-Z16 was bound to citrate and methyl-2,4-
pentanediol, two small ligands recruited from the crystallization cocktail (15).

In a previous study, citrate was shown to act as a calcium chelator thus inhibiting proteolytic
cleavage of LapA by LapG, and thereby stimulating biofilm formation (16). However, in a *lapG*
mutant, citrate was capable of further stimulating biofilm levels (16) thus suggesting there was
a second role for citrate in promoting biofilm formation, a conclusion supported by the putative
small molecule-binding sites present in GcbC. To investigate this point, we tested the ability of
citrate to promote biofilm formation by *P. fluorescens* Pf0-1. Citrate-mediated enhancement of
biofilm formation observed for the wild-type (WT) strain was abolished in the *gcbC* mutant (Fig.

1B), suggesting that citrate-mediated enhancement of biofilm formation is dependent on the
presence of GcbC. We selected Pfl01_2295 and Pfl01_2297 to serve as controls to determine if
other CACHE domains also respond to citrate to promote biofilm formation. We used a low
biofilm forming *P. fluorescens* Pf0-1 strain, lacking four DGCs, referred to as Δ4DGC, which
shows minimal biofilm formation; the four DGCs deleted in the Δ4DGC strain, identified
previously, are *gcbA*, *gcbB*, *gcbC*, and *wspR* (17). Of these three CACHE domain-containing
DGCs expressed in the Δ4DGC mutant background (GcbC, Pfl01_2295, and Pfl01_2297), only
the strain expressing GcbC responded to citrate with increased biofilm formation (Fig. 1C).
GcbC synthesizes c-di-GMP via the GGDEF motif and mutation of this motif to GGAAF eliminates
synthesis of c-di-GMP but this mutant variant is stably expressed (17). When GcbC-GGAAF was
expressed in the Δ4DGC mutant background, biofilm formation was abolished and notably,
citrate-mediated biofilm formation was also abolished (Fig. 1C).

Importantly, citrate enhanced biofilm formation is dependent on LapA (Fig. S2), indicating that
citrate acts via the known LapD-LapG-LapA pathway. We used *P. fluorescens* strains containing
a HA-tagged LapA variant to detect the amount of LapA at the cell surface as a function of the
presence of citrate, as reported (7,10). In a WT *P. fluorescens* strain, citrate caused a 159%
increase in LapA pixel density, which suggests a higher abundance of LapA at the cell surface in
the presence of citrate (Fig. 1D). Only when GcbC was present and catalytically active did
citrate cause an increase in cell surface-associated LapA (Fig. 1D). Taken together, these data
show that citrate-mediated stimulation of biofilm formation in WT *P. fluorescens* Pf0-1 requires
the active diguanylate cyclase GcbC, and is associated with enhanced cell surface LapA.

The Putative Ligand Binding Site of the CACHE Domain is Important for GcbC-mediated
Biofilm Formation. CACHE domains are ubiquitous, extracellular, ligand-binding domains (13,
14). In a previous study, the RXYF motif was found to be the most conserved feature among the
characterized CACHE domains (15). The study found that the X residue is either negatively
charged or polar, and the tyrosine residue of the RXYF motif points towards the ligand-binding
site (15). A mutation of the RXYF motif of the CACHE domain of KinD, a histidine kinase in
Bacillus subtilis, caused this microbe to lose its ability to respond to root exudates, forming
reduced biofilm levels on tomato roots compared to a WT strain. (18). Thus, we predicted that
a mutation within the RXYF motif of GcbC (Fig. S1) would result in impairment of citrate-
dependent biofilm formation when expressed in a Δ4DGC mutant background. We mutated
the tyrosine residue of the RXYF motif to a phenylalanine (Y141F) in a HA-tagged variant of
GcbC (17) and expressed GcbC-Y141F-HA in the Δ4DGC mutant. Mutation of the RXYF motif
resulted in the inability of GcbC to promote biofilm formation or respond to citrate (Fig. 2A).
We assessed the stability of the GcbC-Y141F variant, which showed a 70% reduction in level
compared to WT GcbC (Fig. 2B).

We sought to identify the putative site where citrate might bind to GcbC using the known
structures of CACHE domains (Fig. 2C, left; template PDB ID 3LIB.). Like in the other CACHE
domains, the tyrosine residue of the RXYF motif of GcbC was predicted to point towards the
ligand-binding site (Fig. 2C, Fig, S3). Furthermore, based on the CACHE domain model of GcbC,
the amino acids R139, R162, and R172 were predicted to shape the predicted ligand-binding
site (Fig. 2C, right). Based on the model, we predict that three arginine residues can coordinate
the three carboxylic acid groups of citrate (Fig. 2C, left). R139 is also part of the RXYF motif.
Each of the arginine residues forming the putative ligand site were mutated and introduced
into the Δ4DGC mutant background. Mutating R172 (R172A and R172E) resulted in instability of
GcbC (Table S1), however GcbC-R139A, GcbC-R139E, and GcbC-R162A were detected by
Western blot, with GcbC-R139E variant present at ~WT levels (Fig. 2D). The strain carrying
GcbC-R139E variant did not show a significant enhancement of biofilm formation in the
presence of citrate (Fig. 2E). Together, our data indicate that the RXYF motif and a putative
citrate-binding arginine triad are critical for GcbC-dependent, citrate-mediated enhancement of
biofilm formation. We further expanded our search for important conserved residues within
the CACHE domain that were predicted based on alignments with other CACHE domain
proteins, however the other twenty-seven mutant proteins we constructed were unstable
(Table S1).

Citrate-mediated Interaction of GcbC with LapD Enhances Synthesis of c-di-GMP. In our
published model, GcbC mediates biofilm formation by transferring c-di-GMP to LapD through
physical interaction of the $\alpha5^{GGDEF}$ helix of GcbC with the $\alpha2^{EAL}$ helix of LapD (12). We asked
whether citrate might exert its effect of stimulating biofilm formation via stabilization of the
LapD-GcbC signaling complex. To test if citrate bolstered GcbC-LapD interaction, we exploited
the bacterial two-hybrid system used to initially demonstrate interaction between these
proteins. We did observe a modest but significant enhancement of LapD-GcbC interaction in
the presence of citrate. No such enhancement was observed for the control interactions: LapD-
Pfl01_2295, or GcbC with two other LapD-like dual domain proteins (Fig. 3A). Citrate also did
not enhance GcbC-GcbC dimerization (Fig. 3A), which is perhaps expected, as dimerization is
required for diguanylate cyclase activity (19, 20). Further, the catalytically inactive variant of
GcbC still showed citrate-stimulated interaction (Fig. 3B).

We further explored whether citrate could also enhance the diguanylate cyclase activity of
GcbC. We used the bacterial two-hybrid (B2H) plasmids and strains to express GcbC and LapD
outside of their native context, and to better focus on how the interaction of these two
proteins might specifically impact GcbC's activity. The activity of GcbC was assessed by
measuring the level of c-di-GMP extracted from the indicated strains. The level of c-di-GMP
measured in the *E. coli* strain carrying the vector controls was <2 pmol c-di-GMP/mg dry
weight. This low background of c-di-GMP provided a useful tool to measure differences in c-di-
GMP levels derived from GcbC in the presence and absence of citrate. GcbC alone did not

synthesize detectable levels of c-di-GMP and citrate did not promote c-di-GMP production (Fig.
3C). Co-expression of GcbC with LapD resulted in ~ 20 pmol c-di-GMP/mg dry weight, and the
level of c-di-GMP was significantly increased upon addition of citrate (Fig. 3C). The increase in c-
di-GMP required catalytically active GcbC, and was specific to GcbC interacting with LapD (Fig.
3C). Pfl01_0192 is a dual domain protein and was shown to interact weakly with GcbC (Fig. 3A),
but background levels of c-di-GMP were detected when GcbC and Pfl01_0192 were co-
expressed +/- citrate (Fig. 3C).

Given that stability of GcbC-R139E was equivalent to WT GcbC levels (Fig. 2D), we tested for
interaction of GcbC-R139E with LapD. Mutation of the arginine residue in the RXYF motif did
not affect basal GcbC-LapD interaction. However, citrate-enhanced interaction of GcbC-R139E-
LapD interaction was significantly reduced (but not eliminated) compared to citrate-enhanced
WT GcbC-LapD interaction (Fig. 3D). Thus, our data indicate that LapD-GcbC interaction
enhances c-di-GMP production, and the addition of citrate stimulates both interation of these
proteins and c-di-GMP synthesis, likely via the CACHE domain of GcbC.

The CACHE Domain Participates in GcbC-LapD Interaction. Thus far, we showed the CACHE
domain to be important for citrate-mediated biofilm formation via increasing GcbC-LapD
interaction and enhance GcbC activity. In a previous study (12), the four point mutations,
E477A, Q478A, F481A, K485A (Quad Alanine) were introduced to the surface-exposed $\alpha5^{GGDEF}$
helix of GcbC and expressed in a Δ4DGC mutant background, which resulted in reduced biofilm
formation of *P. fluorescens* due to reduced interaction with LapD (12). However, the GcbC-
Quad Alanine mutant, expressed in the Δ4DGC mutant background still showed a significant
citrate-mediated enhancement of biofilm formation (Fig. S4A), thus suggesting the possibility of
a second interface of GcbC-LapD interaction that is enhanced by citrate.

We next assessed whether the CACHE domain was required for citrate-enhanced interaction
between GcbC and LapD. The two transmembrane domains plus the periplasmic portion of
GcbC showed only a modest level of dimerization with full length GcbC, which was not
stimulated by citrate (Fig. S4B). Importantly, the periplasmic domain of GcbC interacted with
LapD at a level similar to full length GcbC, but citrate-enhanced interaction was abolished (Fig.
S4B). Together, these data indicated that CACHE domain is required but not sufficient for
citrate-mediated, enhanced interaction of LapD and GcbC; however, this domain appears
responsible for basal LapD-GcbC interaction.

LapD Interacts With Numerous DGCs. We next explored if the mechanism we defined for GcbC-
LapD interactions might apply to any of the other 20 DGCs in *P. fluorescens*. We found that
LapD is a central hub of DGC interaction; LapD interacts with a dozen different DGCs in a
pairwise bacterial two-hybrid assay (Fig. 4A). Included among the twelve DGCs that interact

with LapD are Pfl01_2295 and Pfl01_2297, and the putative SadC homolog, Pfl01_4451 (Fig.

4A). SadC is a DGC identified for its role in the early stages of biofilm development in *P.*

aeruginosa (21), and the Δ*sadC* mutant of *P. aeruginosa* shows approximately a 50% reduction

of global, cellular c-di-GMP levels compared to WT PA14 (22). We have not yet identified

ligands that enhance interactions and/or activity of these other DGCs.

Discussion. A key open question relating to c-di-GMP signaling is understanding how specificity

of an output is mediated in the context of up to dozens of enzymes or receptors making,

breaking and binding this dinucleotide. Here, our data suggest that an extracellular ligand can

modulate the activity of a DGC and do so via interaction with its cognate receptor. Our data are

consistent with the model that citrate binding to the CACHE domain of GcbC enhances

interaction with LapD. We propose the increased interaction between GcbC and LapD to have

two important consequences: stabilization of a complex that allows direct transfer of the GcbC-

generated c-di-GMP signal to the LapD receptor, and equally importantly, activation of the DGC

activity of GcbC. Indeed, when expressed in a heterologous system, GcbC showed almost no

capacity to synthesize c-di-GMP; it is only when co-expressed with LapD that a significant

increase in c-di-GMP synthesis above background was detected. An additional boost in cyclic

nucleotide production was measured when citrate was added to the medium in a strain co-

expressing GcbC and LapD, indicating that the enhanced GcbC-LapD interaction also enhanced

DGC activity. Furthermore, recent work from the Sondermann lab (23) proposed a model

where GcbC and LapD, along with LapG, form a large signaling complex that facilitates LapD-

GcbC interactions (Fig. 4B). Thus, our data supports a model for a ligand-based mechanism to

enhance signaling specificity by c-di-GMP in the context of a large signaling complex.

We have identified citrate as a potential ligand that binds to GcbC via its CACHE domain. There

is an abundance of CACHE domains present on signal transduction proteins (14), including

proteins involved in c-di-GMP signaling, and yet, the role of CACHE domains fused to DGCs is

poorly understood. Our data show that citrate is a potential ligand for the CACHE domain

associated with GcbC. *P. fluorescens*, a plant symbiont, forms biofilms on tomato roots and the

root exudates contain organic acids including citrate (24, 25, 26). It is possible, however, that

GcbC senses other organic acids. For example, in *P. syringe* pv. *actinidiae*, the CACHE domain

PscD (PDB ID: 5G4Z) binds glycolate, acetate, propionate, and pyruvate (27). It is also possible

that different ligands sensed via the CACHE domain of GcbC can dictate which proteins interact

with this DGC (Fig. 4A). Together, our data indicate that extracellular ligands, via their ability to

modulate protein-protein interactions and/or DGC activity, can modulate c-di-GMP signaling

specificity. Given the large number of ligand-binding domains associated with c-di-GMP-

metabolizing proteins, the data presented here could represent a general means of regulating

c-di-GMP-controlled outputs by enhancing the signaling between c-di-GMP-metabolizing

enzymes and their effectors.

Enzymes that make c-di-GMP are part of a complex network, with LapD as a central receptor for
DGCs to interact with (Fig. 4A). It is highly unlikely that the twelve DGCs that make up this DGC
network all interact with LapD without regulation to control signaling specificity. A more likely
scenario could be that ligands are used as a signal to regulate signaling of a DGC to interact with
LapD and outcompete with one another. It is also possible that other ligand-mediated
mechanisms exist, such as a ligand-enhanced homodimerization of a DGC. Further
characterizing and identifying what small molecules DGCs respond to could provide insight into
how c-di-GMP signaling specificity is regulated within this complex signaling network. For
example, we envision a scenario wherein a "cloud" of weakly interacting DGCs can increase
specific interaction with their receptor in response to appropriate environmental signals,
concomitantly boosting c-di-GMP production, ligand-specific signaling and biofilm formation.
Our work provides insight of a ligand-based mechanism of how signaling specificity occurs
among a complex network of enzymes and receptors that make, break, and bind c-d-GMP.

Material and Methods

Strains and Media. Bacterial strains used in this study are listed in Table S2, and were cultured
and maintained in lysogeny broth (LB) or on 1.5% agar LB plates. *P. fluorescens* was grown at
30°C and *E. coli* was grown at 37°C. *E. coli* S17-1-λ-pir was used for maintenance and transfer of
plasmids. *Saccharomyces cerevisiae* strain InvSc1 was used for plasmid modification as
described previously (28, 29). K10T-1 medium was prepared as described previously (30).
Sodium citrate was added to 1.5% agar LB plates and K10T-1 media to a final concentration of
13.6 mM (0.4% wt/vol) for all the experiments described. The following antibiotics were used
as indicated: gentamycin (15μg/mL for *E. coli*, 30μg/mL for *P. fluorescens*), kanamycin (50μg/mL
for *E. coli*), and carbenicillin (50μg/mL for *E. coli*).

Biofilm Assay. Biofilm assays were performed as described previously (1). *P. fluorescens* Pf0-1
strains were incubated in K10T-1 medium with and without 0.4% sodium citrate, as indicated,
for 6 hours at 30°C. Biofilms were stained with 0.1% crystal violet, washed with water and then
solubilized with a 45% methanol, 45% dH$_2$O, and 10% glacial acetic acid solution. The optical
density (OD) of the solubilized crystal violet solution was measured at 550 nm to determine the
amount of biofilm formed.

Dot Blot LapA Localization Assay. Localization of LapA to the cells surface was measured using
a HA-tagged LapA variant integrated into the chromosome of *P. fluorescens* as described
previously (7, 10, 31). Overnight LB-grown cultures were subcultured in 5mL K10T-1 medium in
the presence and absence of 0.4% sodium citrate for 6 hours at 30°C. Samples were normalized
by the lowest OD$_{600}$ value.

Bacterial Two-Hybrid Assay. Bacterial two-hybrid assays were performed using *E. coli* BTH101
cells based on a previously described system (32). Briefly, ~100ng of each bacterial two hybrid

plasmid was cotransformed into *E. coli* BTH101 by electroporation. *E. coli* BTH101 cells were
incubated on LB agar supplemented with 50μg/mL kanamycin, 50μg/mL carbenicillin, and
0.5mM isopropyl β-D-1-thiogalactopyranoside (IPTG) for 24 hours at 30°C. At 24 hours, either β-
galactosidase or c-di-GMP levels were quantified as described below. β-galactosidase assays
were performed as exactly described previously (12) to quantify the extent of protein-protein
interaction. β-galactosidase levels are presented in Miller Units.

c-di-GMP Quantification Assay. c-di-GMP was extracted from *E. coli* BTH101 cells after
incubation on LB agar plates at 30°C for 24 hours. The cells were scraped from the plate surface
with 1 mL of dH_2O, then pelleted and resuspended in 0.250 mL nucleotide extraction buffer
(40% methanol, 40% acetonitrile, 20% dH_2O, and 0.1N formic acid), followed by incubation at -
20°C for 1 hour. Cells were pelleted again and the reaction was neutralized by transfer of 0.2
ml nucleotide extract to 8μl of 15% NH_4CO_3. Nucleotide extracts were vacuum-dried and
resuspended in 0.2 mL HPLC grade H_2O. c-di-GMP concentration was analyzed by liquid
chromatography-mass spectrometry and compared to a standard curve of known c-di-GMP
concentration, as reported (12). The moles of c-di-GMP were normalized to the dry weight of
the cell pellet from which the nucleotides were extracted.

Statistical Analysis. Student's t-test was used to test for statistical significance. $P < 0.05$; *, $P <$
0.01; **, $P < 0.001$; ***

Acknowledgement. We thank members of the lab for helpful discussions.

Literature Cited.

1. O'Toole GA, Kolter R (1998) Initiation of biofilm formation in *Pseudomonas fluorescens*
   WCS365 proceeds via multiple, convergent signaling pathways: A genetic analysis. *Mol*
   Microbiol 28: 449-461.
2. Pruss BM, Besemann C, Denton A, Wolfe AJ (2006) A complex transcription network
   controls the early stages of biofilm development by *Escherichia coli*. *J Bacteriol* 188(11):
   3731-3739.
3. Römling U, Gomelsky M, Galperin MY (2005) c-di-GMP: The dawning of a novel bacterial
   signaling system. *Mol Microbiol* 57(3): 629-639.
4. Baker AE, Diepold A, Kuchma SL, Scott JE, Ha DG, Orazi G, Armitage JP, O'Toole GA
   (2016) A PilZ domain protein FlgZ mediates cyclic-di-GMP-dependent swarming motility
   control in *Pseudomonas aeruginosa*. *J Bacteriol* 198(13): 1837-1846.
5. Ross P, Weinhouse H, Aloni Y, Michaeli D, Weinberger-Ohana P, Mayer R, Braun S, de
   Vroom E, van der Marel GA, van Boom JH, Benziman M (1987) Regulation of cellulose
   synthesis in *Acetobacter xylinum* by cyclic diguanlyic acid. *Nature* 325(6101): 279-281.

6. Merrit JH, Brothers KM, Kuchma SL, O'Toole HA (2007) SadC reciprocally influences biofilm formation and swarming motility via modulation of exopolysaccharide production and flagellar function. *J Bacteriol* 189(22): 8154-8164.

7. Monds RD, Newell PD, Gross RH, O'Toole GA (2007) Phosphate-dependent modulation of c-di-GMP levels regulates *Pseudomonas fluorescens* Pf0-1 biofilm formation by controlling secretion of the adhesin LapA. *Mol Microbiol* 63(3): 659-679.

8. Martínez-Gil M, Ramos-González MI, Espinosa-Urgel M (2014) Roles of cyclic di-GMP and the Gac system in transcriptional control of the genes coding for *Pseudomonas putida* adhesins LapA and LapF. *J Bacteriol* 196(8): 1484-1495.

9. Hinsa SM, Espinosa-Urgel M, Ramos JL, O'Toole HA (2003) Transition from reversible to irreversible attachment during biofilm formation by *Pseudomonas fluorescens* WCS365 requires an ABC transporter and a large secreted protein. *Mol Microbiol* 49: 905-918.

10. Newell PD, Monds RD, O'Toole GA (2009) LapD is a bis-(3'-5')-cyclic dimeric GMP-binding protein that regulates surface attachment by *Pseudomonas fluorescens* Pf0-1. *Proc Natl Acad Sci USA* 106(9): 3461-3466.

11. Navarro MV, Newell PD, Krasteva PV, Chatterjee D, Madden DR, O'Toole GA, Sondermann HA (2011) Structural basis for c-di-GMP-mediated inside-out signaling controlling periplasmic proteolysis. *PLoS Biol* 9(2): e1000588. doi: 10.1371.

12. Dahlstrom KM, Giglio KM, Collins AJ, Sondermann H, O'Toole GA (2015) Contribution of physical interactions to signaling specificity between a diguanylate cyclase and its effector. *mBio* 6(6):e01978-15. doi:10.1128/mBio.01978-15.

13. Anantharamn V, Aravind L (2000) Cache – a signaling domain common to animal Ca(2+)-channel subunits and a class of prokaryotic chemotaxis receptors. *Trends Biochem Sci* 25(11): 535-537.

14. Upadhyay AA, Fleetwood AD, Adebali O, Finn RD, Zhulin IB (2016) Cache domains that are homologous to, but different from PAS domains comprise the largest superfamily of extracellular sensors in prokaryotes. *PLoS Comput Biol* 12(4): e1004862. doi: 10.1371/journal.pcbi.1004862.

15. Zhang Z, Hendrickson WA (2010) Structural characterization of the predominant family of histidine kinase sensor domains. *J Mol Biol* 400(3): 335-353.

16. Boyd CD, Chatterjee D, Sondermann H, O'Tool GA (2012) LapG, required for modulating biofilm formation by *Pseudomonas fluorescens* Pf0-1, is a calcium-dependent protease. *J Bacteriol* 194(16): 4406-4414.

17. Newell PD, Yoshioka S, Hvorecny KL, Monds RD, O'Toole GA (2011) Systematic analysis of diguanylate cyclases that promote biofilm formation by *Pseudomonas fluorescens* Pf0-1. *J Bacteriol* 193(18): 4685-4698.

18. Chen Y, Cao S, Chai Y, Clardy J, Kolter R, Guo JH, Losick R (2012) A *Bacillus subtilis* sensor kinase involved in triggering biofilm formation on the roots of tomato plants. *Mol Microbiol* 85(3): 418-430.

19. Chan C, Paul R, Samoray D, Amiot NC, Giese B, Jenal U, Schirmer T (2004) Structural basis of activity and allosteric control of diguanylate cyclase. *Proc Natl Acad Sci USA* 101(49): 17084-17089.

20. Ryjenkov DA, Tarutina M, Moskvin OV, Gomelsky M (2005) Cyclic diguanylate is a ubiquitous signaling molecule in bacteria: insights into biochemistry of the GGDEF protein domain. *J Bacteriol* 187(5): 1792-1798.

21. Merritt JH, Brothers KM, Kuchma SL, O'Toole GA (2007) SadC reciprocally influences biofilm formation and swarming motility via modulation of exopolysaccharide production and flagellar function. *J Bacteriol* 189(22): 8154-8164.

22. Merritt JH, Ha DG, Cowles KN, Lu W, Morales DK, Rabinowitz J, Gitai Z, O'Toole GA (2010) Specific control of *Pseudomonas aeruginosa* surface-associated behaviors by two c-di-GMP diguanylate cyclases. *Mbio* 1(4): e00183-10. doi:10.1128/mBio. 00183-10.

23. Cooley RB, O'Donnell JP, Sondermann H (2016) Coincidence detection and bi-directional transmembrane signaling control a bacterial second messenger receptor. *Elife* 5: e21848 doi: 10.7554/eLife.21848.

24. Haas D, Défago G (2005) Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat Rev Microbiol* 3(4): 307-319.

25. Jones DL (1998) Organic acids in the rhizosphere – a critical review. *Plant Soil* 205: 25-44.

26. de Weert S, Vermeiren H, Mulders IH, Kuiper I, Hendrickx N, Bloemberg GV, Vanderleyden J, De Mot R, Lugtenberg BJ (2002) Flagella-driven chemotaxis towards exudate components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. *Mol Plant Microbe Interact* 15(11): 1173-1180.

27. Brewster JL, McKellar JL, Finn TJ, Newman J, Peats TS, Gerth ML (2016) Structural basis for ligand recognition by a Cache chemosensory domain that mediates carboxylate sensing in *Pseudomonas syringae*. *Sci Rep* 6:35198. doi: 10.1038/srep35198.

28. Shanks RMA, Calazza NC, Hinsa SM, Toutain CM, O'Toole GA (2006) *Saccharomyces cerevisiae*-based molecular tool kit for manipulation of gram-negative bacterial genes. *Appl Environ Microbiol* 72(7): 5027-5036.

29. Bascom-Slack CA, Dawson D (1998) A physical assay for detection of early meiotic recombination intermediates in *Saccharomyces cerevisiae*. *Mol Gen Genet* 258: 512-520.

30. Monds RD, Newell PD, Schwartzman JA, O'Toole GA (2006) Conservation of the Pho regulon in *Pseudomonas fluorescens* Pf0-1. *Appl Environ Microbiol* 72(3): 1910-1924.

31. Newell PD, Boyd CD, Sodermann H, O'Toole GA (2011) A c-di-GMP effector system control cell adhesion by inside-out signaling and surface protein cleavage. *PloS Biol* 9(2): e1000587. doi: 10.1371/journal.pbio.1000587.

32. Karimova G, Pidoux J, Ullmann A, Ladant D (1998) A bacterial two-hybrid system based on a reconstituted signal transduction pathway. *Proc Natl Acad Sci USA* 95: 5752-5756.

33. Shultz J, Milpetz F, Bork P, Ponting CP (1998) SMART, a simple modular architecture research tool: Identification of signaling domains. *Proc Natl Acad Sci USA* 95: 5857-5864.

399   34. Ulrich LE, Zhulin IB (2010) The MiST2 database: a comprehensive genomics resource on
400   microbial signal transduction. *Nucleic Acids Res* 38: D401-407 doi:10.1093/nar/gkp940.
401   35. Simon R, Preifer U, Puhler A (1983) A broad host range mobilization system for *in vivo*
402   genetic engineering: transposon mutagenesis in gram negative bacteria. *Nat Biotechnol*
403   1: 784-791.

  Figures

Fig. 1. Citrate stimulates biofilm formation via GcbC. (A) Predicted domain organization of
GcbC, Pfl01_2295, and Pfl01_2297 as predicted by SMART (33). The domains are indicated in
each block. TM = transmembrane domain. Analysis of Pfl01_2295 by MiST2 (34) predicts a
second transmembrane domain. (B) Biofilm formation by WT *P. fluorescens* and the Δ*gcbC*
mutant in the presence and absence of 0.4% citrate (n = 3; $\pm$ SD). (C) Biofilm formation by the
indicated strains +/- citrate. In this panel, the Δ4DGC mutant background is used, with the WT
GcbC and the catalytically inactive variant (GGAAF) introduced into this mutant on plasmids.
Pfl01_2297 and Pfl01_2295 are two other CACHE-domain containing diguanylate cyclases in *P.*
fluorescens that serve as controls. (D) Quantification of cell-surface levels of LapA in the
presence and absence of 0.4% citrate. Representative blots are shown. For panels B, C, and D,
the values shown are an average of 3 replicates ($\pm$ SD); horizontal black bars indicate a P value
of either <0.05 (*), <0.01 (**), or <0.001 (***) with a student's t-test compared to the condition
without citrate comparing each strain without versus with citrate.

Fig. 2. Effects of CACHE domain mutations on biofilm formation. (A) Quantitation of the effect
of CACHE domain mutant, GcbC Y141F, on biofilm formation in the presence and absence of
citrate. (B) Western blot to assess the relative stability of GcbC-HA and GcbC-Y141F-HA in the
presence and absence of citrate. (C) Homology model of the CACHE domain of GcbC based on
rpHK1S-Z16 (PDB ID: 3LIF) and vpHK1S-Z8 (PDB ID: 3LID) (see Figure S3) to identify the putative
ligand-binding site. (D) Western blot assessment of the relative stability of GcbC-R139A-HA,
GcbC-R139E-HA and GcbC-R162A-HA. (E) Quantitative analysis of biofilm formation by strains
expressing the indicated CACHE domain mutants in the presence and absence of citrate. For
panels A and E, data shown is the average of three replicates ($\pm$ SD); horizontal black bars
indicate a P value of either <0.05 (*) or <0.001 (***) with a student's t-test comparing each
strain without versus with citrate.

Fig. 3. Citrate-mediated biofilm enhancement. (A) The effect of citrate on interaction with the indicated proteins is compared by the B2H assay. Pfl01_2295 = CACHE domain-containing DGC; Pfl01_0192 = GGDEF and EAL domain-containing protein; Pfl01_2920 = phosphodiesterase. Briefly, the proteins of interest are fused to two halves of the catalytic domain of an adenylate cyclase, T25 and T18, from *Bordetella pertussis* are transformed into *E. coli* BTH101 cells (32). Plasmids pKT25 and pUT18C are the vector-only controls. If the two proteins interact, the adenylate cyclase activity is reconstituted, promoting cAMP synthesis, which in turn activates transcription of *lacZ*. The β-galactosidase activity is assessed using X-gal as a substrate, which serves as a read out for the degree of interaction (32). Citrate did not enhance interaction of the vectors or of the positive control (GCN4 leucine zipper protein). Citrate only enhanced the interaction between LapD and GcbC. (B) The effect of catalytic activity of GcbC on interaction with LapD were compared in the presence and absence of 0.4% citrate. (C) c-di-GMP production in *E. coli* BTH101 was determined for strains carrying the indicated proteins. Only co-expression of LapD-GcbC showed appreciable accumulation of c-di-GMP, which was stimulated by added citrate. (D) The effect of R139E CACHE mutation on interaction with LapD in the presence and absence of citrate. For panels B, C, D, and E, all experiments were performed in triplicate ($\pm$ SD), with an indicated P value of either <0.05 (*) or <0.01 (**) with a student's t-test comparing each strain without versus with citrate.

453   **Fig. 4. LapD interacts with multiple DGCs**. (A) Interaction map of DGCs that interact with LapD
454   by 20 hours by bacterial-two-hybrid assay. Interaction map constructed using XXXXX. (B) A
455   model of a GcbC, LapD, LapG signaling "basket" complex (23).