1
Title:
REMAP: An online remote sensing application for land cover classification and monitoring

Authors:
Nicholas J. Murray[1]   murr.nick@gmail.com
David A. Keith[1,2]   david.keith@unsw.edu.au
Daniel Simpson[1]   danielksimpson@gmail.com
John H. Wilshire[1]   j.wilshire@unsw.edu.au
Richard M. Lucas[1]   richard.lucas@unsw.edu.au

Author details:
[1] Centre for Ecosystem Science, School of Biological, Earth and Environmental Science,
University of New South Wales, Sydney, Australia
[2] New South Wales Office of Environment and Heritage, Hurstville, New South Wales,
Australia

Corresponding Author:
Dr Nicholas Murray, Centre for Ecosystem Science, University of New South Wales,
Sydney, Australia 2052 | Phone: +61 414 815 788 | Email: murr.nick@gmail.com

Running head: REMAP: An online remote sensing application
19

ABSTRACT

1. Recent assessments of progress towards global conservation targets have revealed a paucity of indicators suitable for assessing the changing state of ecosystems. Moreover, land managers and planners are often unable to gain timely access to maps they need to support their routine decision-making. This deficiency is partly due to a lack of suitable data on ecosystem change, driven mostly by the considerable technical expertise needed to make ecosystem maps from remote sensing data.

2. We have developed a free and open-access online remote sensing and environmental modelling application, REMAP (*the remote ecosystem monitoring and assessment pipeline*; https://remap-app.org) that enables volunteers, managers, and scientists with little or no experience in remote sensing to develop high-resolution classified maps of land cover and land use change over time.

3. REMAP utilizes the geospatial data storage and analysis capacity of the Google Earth Engine, and requires only spatially resolved training data that define map classes of interest (e.g., ecosystem types). The training data, which can be uploaded or annotated interactively within REMAP, are used in a random forest classification of up to 13 publicly available predictor datasets to assign all pixels in a focal region to map classes. Predictor datasets available in REMAP represent topographic (e.g. slope, elevation), spectral (Landsat Archive image composites) and climatic variables (precipitation, temperature) that can inform on the distribution of ecosystems and land cover classes.

4. The ability of REMAP to develop and export high-quality classified maps in a very short (<10 minute) time frame represents a considerable advance towards globally accessible and free application of remote sensing technology. By enabling access to

data and simplifying remote sensing classifications, REMAP can catalyse the

monitoring of land use and change to support environmental conservation, including

developing inventories of biodiversity, identifying hotspots of ecosystem diversity,

ecosystem-based spatial conservation planning, mapping ecosystem loss at local

scales, and supporting environmental education initiatives.

KEYWORDS

Ecosystem monitoring, GIS, Google Earth Engine, Image classification, Landsat

Archive, Land cover mapping, Remote sensing, Satellite mapping

INTRODUCTION

Maps of land use and land cover change have been a central component of

environmental management and conservation planning for decades (Margules & Pressey

2000). Land cover maps enable the depiction of the distribution of ecosystems and land cover

types, assessments of biodiversity and identification of areas undergoing loss, fragmentation

and degradation (Haddad *et al.* 2015; Potapov *et al.* 2017). As well as supporting spatial

conservation planning, including mapping threats to nature, they are often used as surrogates

for species distributions. However, existing methods for mapping land cover extent and

changes over time are often based on remote sensing and rely on expert implementation and

comprehensive knowledge of space borne or airborne sensor data, analytical methods and

data uncertainties. This ÷capacity gapø has been a severe constraint in obtaining information

on the status of the worldøs natural environment and has hindered environmental conservation

programs across a range of spatial scales (Pereira, Brevik & Trevisani 2018; Murray *et al.* in

press).

Recent advances in geospatial data access, storage and analysis have vastly improved

our ability to utilize satellite sensor data archives in studies of land cover and land cover

change (e.g. Lewis *et al.* 2016; Gorelick *et al.* 2017). Moderate (< 30 m) resolution remote

sensing analyses are now possible at the global extent and have enabled the development of

complex remote sensing analyses (Gong *et al.* 2013; Hansen *et al.* 2013; Pekel *et al.* 2016).

At the same time, increases in satellite revisit frequencies, reductions in the time between

data acquisition and delivery to users, and increasing access to data archives have led to the

development of near real-time alert systems that can rapidly identify land cover loss and

change in areas where no ground observations can be obtained. These systems mainly focus

on automatic detection and analysis of land cover change for groups of related biomes (e.g.

forests) and have vastly improved the ability of non-specialists, environmental managers and

policy makers to access and use remote sensing data (Asner *et al.* 2009; Hansen *et al.* 2016;

Lucas & Mitchell 2017).

In this manuscript, we present a new online geospatial application that enables

volunteers, managers, students and scientists with little or no experience in remote sensing to

develop classified maps of land cover at Landsat spatial resolutions. The *Remote Sensing*

Monitoring and Assessment Pipeline (REMAP) utilizes the geospatial data storage and analysis

capacity of the Google Earth Engine (GEE; https://earthengine.google.com), a cloud-based

analysis platform, to allow users to interactively develop machine learning classifications of

land cover within an area of interest anywhere in the world for which there is sufficient

archival Landsat data. The REMAP application additionally allows monitoring and analysis of

land cover change by enabling users to map ecosystem distributions at two points in time (i.e.

2003 and 2017), quantify area change of each map class, and report the standard distribution

size metrics used by the International Union for the Conservation of Nature (IUCN) Red List

of Ecosystems (Keith *et al.* 2013).

REMAP was developed to complement a range of other applications that support the

conservation of biodiversity, including GeoCAT (Bachman *et al.* 2011), Global Forest Watch

(www.globalforestwatch.org), the Map of Life (www.mol.org) and R packages such as

'redlistr' (Lee & Murray 2017) and 'rCat' (Moat & Bachman 2017). Potential uses of REMAP

include mapping the distributions of ecosystem types (Murray *et al.* in press), developing

land cover maps for protected areas (Lucas *et al.* 2015), assessing the performance of

protected areas over multi-decadal time frames (Green *et al.* 2013; Murray & Fuller 2015),

and identifying areas where degradation of ecosystems has occurred (Bhagwat *et al.* 2017).

REMAP was also developed to support the global effort to assess the status of all ecosystem

types on earth under the IUCN Red List of Ecosystems criteria (Keith *et al.* 2015; Rodríguez

et al. 2015) and can contribute to monitoring progress towards addressing the 2020

Convention on Biological Diversity Aichi Targets (CBD 2014). We describe here the

rationale for design, methodological considerations and analytical framework of REMAP, and

demonstrate its utility and limitations with four case studies (see *Case Studies*).

REMAP: REMOTE ECOSYSTEM MONITORING & ASSESSMENT PIPELINE

REMAP (https://remap-app.org) is a free and open-source web application that classifies land

cover according to user-supplied training data and a set of globally available remote sensing

datasets as predictor variables (Figure 1). We followed six design principles to develop

REMAP:

1. *Provide the ability to develop high quality maps from remote sensing data in a short time*

frame and without the need for high performance computers. Maps can be developed in

REMAP within a few minutes and, because REMAP completes classifications online by

accessing the GEE, the only prerequisites are an internet connection and web browser.

2. *Reduce the need to download, pre-process and process remote sensing data for use in*
environmental mapping. The system offers access to 13 publicly available geospatial
predictors that represent spectral, topographic and climatic variables that may influence
the distribution of different land cover types. Default predictors were selected to enable
the development of high quality maps of the widest range of land cover types possible,
and users are provided with options to explore different combinations of predictors in the
production of their classified map.

3. *Simplify implementation of machine learning classification approaches.* REMAP conducts
its classifications using the random forest algorithm (Breiman 2001) with a single execute
button. This approach allows users to implement a widely used machine learning method
known to achieve high classification accuracy from large amounts of potentially
correlated predictor variables (Rodriguez-Galiano *et al.* 2012).

4. *Permit the production of maps for at least two time periods to enable the quantification of*
any detectable spatial change. REMAP can be used to measure the impacts of, for
example, deforestation (Hansen & Loveland 2012), coastal reclamation (Murray *et al.*
2014), and many other land cover changes that can be reliably observed with Landsat
sensors.

5. *Enable estimation of standard spatial metrics used for assessing the status of ecosystems.*
Metrics that are useful for environmental conservation, including area, change in area,
extent of occurrence (EOO) and area of occupancy (AOO) can be calculated by users to
assess ecosystem change and contribute to global efforts to assess the status of
ecosystems.

6. *Implement free and open access software design principles.* Source code for REMAP is
available and we will maintain open access to the application (see *Data Accessibility*).

DATA

The 13 publically available gridded datasets that were selected for inclusion in REMAP

(Table 1) met the requirement of (i) full global extent, (ii) free availability with sufficient

open access to be included in the GEE public data archive, and (iii) sufficiently high spatial

resolution to permit identification of ecosystem distributions and common land cover classes.

The final set of predictors includes spectral variables and derived indices from archival

Landsat sensor data for two time periods, climate data (mean annual rainfall and mean annual

temperature; Hijmans *et al.* 2005) and topographic data (derived from Shuttle Radar

Topography Mission data).

To obtain the required global coverage of cloud-free Landsat sensor data for two

periods, referred to here as historical (1999-2003) and ÷currentø (2014-2017), we developed

two global Landsat image composites. We produced image stacks of all Landsat scenes for

each period ($N_{1999\text{-}2003}$ = 340,658 images; $N_{2014\text{-}2017}$ = 375,674 images) and applied the GEE

implementation of the FMASK cloud masking algorithm (Gorelick *et al.* 2017). From these,

the median pixel of Landsat Enhanced Thematic Mapper (ETM+; bands 2-5) bands 2 5

(visible blue to shortwave infrared) and Operational Land Imager (OLI; bands 1-4) was used

to generate the two 4-band global image composites. From these composites, Normalized

Differenced Vegetation Index (Pettorelli 2013), Normalized Difference Water Index

(McFeeters 1996) and several other index layers were generated for use as spectral predictors

(Table 1). The provision of spectral data for two time periods facilitates the estimation of

change in land cover extent, which is important for monitoring of the impact of threatening

processes such as deforestation (Hansen *et al.* 2013), fragmentation (Haddad *et al.* 2015),

coastal reclamation (Murray *et al.* 2014), aquaculture (Thomas *et al.* 2017) and water

extraction (Tao *et al.* 2015).

USER INPUT

Users of REMAP generally follow a 7 step procedure to map, assess and monitor

ecosystem types or land cover classes (Table 2). Initially, users are required to define their

region of interest interactively (focus region) or to upload a vector file (.kml). This enables

REMAP to clip input data to a region of interest and limit the extent of the classification. The

maximum size of the region of interest is presently 100,000 km^2 due to limitations applied to

users of the GEE (Gorelick *et al.* 2017). Future versions of REMAP may increase this size

limit, although for larger regions or more complex map classifications we recommend users

directly utilise the GEE (https://earthengine.google.com).

Spatially resolved training data that define map classes of interest, which can include

ecosystem types, land cover classes, areas of change (e.g. deforestation) or anthropogenic

areas (e.g. urban areas) are used to assign a class membership to all pixels within a focal

region. If developing land cover maps, we recommend that users adopt land cover

classification taxonomies that are internationally recognized and confirm to International

Organisation for Standards (ISO) such as the Food and Agricultural Organisation¢s (FAO)

Land Cover Classification System (LCCS). Training data can be provided interactively by

adding training points via the user interface with reference to the predictor layers or by

uploading data which identify the location of observation points and their class membership

(.csv file). These may be sourced from field observations, external data archives, expert

opinion, literature or existing maps. In general, classifications with larger numbers of training

points will achieve higher class accuracies and we recommend users supply a minimum of 50

points per class to develop an initial map.

CLASSIFICATION APPROACH

REMAP uses a random forest classifier to assign pixels to user-defined map classes

(Breiman 2001). With sufficient training data that are representative of the classes of interest,

REMAP implements the classification on the predictor data and returns a classified image to

the browser window. In many cases, use of the default predictors (Table 2) will yield

classification accuracies that are acceptable to the user. To allow users to assess classification

accuracy, REMAP returns a confusion matrix that compares classification results with a

random subset of points held-out of the training dataset. Users can tune their classifications to

maximize accuracies, either overall or for the class(es) of interest, (ideally to >85%;

Congalton & Green 2008) by providing more training data for the classifier or by selecting a

custom set of predictors (Table 2).

ECOSYSTEM MONITORING AND ASSESSMENT

Once a classified map of acceptable accuracy has been produced, REMAP can conduct

the spatial analyses required to assess Criteria A (change in distribution size) and B (range

size) of the IUCN Red List of Ecosystems (Keith *et al.* 2013; Bland *et al.* 2017). To assess

Criterion A, REMAP computes the area of each class by summing the number of pixels in each

class. Criterion A requires assessors to estimate change in area over time, which can be

achieved by repeating the workflow for the second time period. To account for potential

changes in land cover between the two time periods, users should develop a new training set

or modify the existing set to ensure accurate representation of land cover in the second time

period. Once area estimates are completed for two time periods, assessors can follow the

IUCN Red List of Ecosystems guidelines to estimate area change manually (Bland *et al.*

2017) or with the recently developed ‒redlistr‒ R package (Lee & Murray 2017). To assess

criterion B of the IUCN Red List of Ecosystems, REMAP applies a minimum convex polygon

to a class of interest and reports its area, representing the Extent of Occurrence (EOO) of the

map class. Finally, the Area of Occupancy (AOO) of a map class is calculated by applying a

10×10 km grid and counting the number of grid cells occupied by the map class (Bland *et al.*

2017; Murray *et al.* 2017).

To support further analyses of the classified map data, users can export each classified

map as a georeferenced raster file (.tif). Furthermore, training data can be exported as a .csv

file with fields ‒latitudeø, ‒longitudeø and ‒classø suitable for import into a GPS unit or GIS

software. Training data can also be saved as a JSON file, which is analogous to ‒save

workspaceø functions in other software. This allows users to return to their analysis at a later

time by uploading the JSON file (see Appendix 1 for examples).

CASE STUDIES

Classifications of remote sensing data enable the measurement and monitoring of an

enormous range of environmentally relevant variables. To demonstrate the use of REMAP, we

developed case studies for (i) mapping a single ecosystem type (e.g. Murray *et al.* 2012;

Nascimento *et al.* 2013), (ii) generating a comprehensive land cover map for a region of

interest (e.g.,Malatesta *et al.* 2013; Connette *et al.* 2016), and (iii) quantifying land cover

change between two periods (e.g., Sexton *et al.* 2013; Olofsson *et al.* 2016; Thomas *et al.*

2017). All training data (.csv) and REMAP workspace files (.JSON) used to reproduce these

case studies are available in supplementary material (Appendix 1) and can be used in

association with tutorials available on the REMAP website (https://remap-app.org/tutorial).

1. *Mapping single land cover types or ecosystem types.* Mapping the distribution and change

of mangrove ecosystems has been an important focus of ecosystem monitoring programs

for decades due to their provision of ecosystem services (Mumby *et al.* 2004; Spalding *et*

al. 2014) and susceptibility to a wide range of threats (Cavanaugh *et al.* 2014; Asbridge *et*

al. 2016; Duke *et al.* 2017). In this case study, we developed a simple classification of

mangroves and non-mangrove from a set of 150 training points for a small focal region

(8301 ha) in the Gulf of Carpentaria, Australia (Figure 2). Against random subsets of

training data, the resubstitution accuracy reported by REMAP was 99.2%. Furthermore, a

random allocation of 389 points over the focal region indicated a 93.3% agreement with

the 2000 global mangrove map data produced for the year 2000 (Giri *et al.* 2011).

2. *Comprehensive classification of land cover for a focal region.* Land cover maps, which

represent all land types in a region, is a common aim of remote sensing programs (Lucas

& Mitchell 2017). We used REMAP to develop a land cover map with classes *semi-*

deciduous vine forest, *eucalypt woodland* and *human settlement* for a focal region in the

dry tropics of Northern Australia (Figure 3; Figure S1). A comparison with ecosystem

maps produced by the state-wide regional ecosystem mapping program, which develops

regulatory land cover maps through manual interpretation of aerial photography and

Landsat TM and SPOT satellite imagery, indicated good agreement between the two

mapping methods (Figure 3; Neldner *et al.* 2017; Queensland Department of Natural

Resources and Mines 2017). We provide a second land cover example that covers a larger

area with more land-cover classes in the Supplementary Material (Cheduba Island,

Myanmar, Figure S2).

3. *Quantifying land cover change.* To demonstrate capacity to detect changes in land and

water, REMAP was applied to the two Landsat composite images available (2003) and

OLS (2017) data acquired over Dubai, United Arab Emirates. The resulting maps provide

quantitative information on the extent of marine ecosystem loss as a result of large-scale

coastal reclamation projects (Figure 4). REMAP's use for change mapping is also

demonstrated with a deforestation example at Roraima, Brazil (Figure 1, Figure S3,

Appendix A).

DISCUSSION

REMAP is a fast, user-friendly approach to developing land cover maps from freely

available remote sensing data and its outcomes can be accepted if the accuracies of

classifications meet the expectations of the users. Our case studies indicate that such

accuracies can be achieved in REMAP but these depend upon the accuracy of the training data.

By utilizing the geospatial storage and analysis capacity of the GEE, REMAP allows users

with no prior knowledge in remote sensing and analysis to develop maps directly within a

web-browser. This enables mapping to be undertaken in regions by locally-responsible

individuals and organisations where computing infrastructure is scarce or the quality of

internet connections do not allow the download of remote sensing data for local analyses.

Indeed, REMAP is particularly useful for participatory mapping projects, expert elicitation and

engagement with a wide-range of environmental stakeholders.

We acknowledge that REMAP has several limitations. Most notably, the ability of

REMAP to produce accurate maps is limited by the quality of the training data, the accuracy of

the predictors, and the suitability of the predictor set for distinguishing land cover classes

and. Further development of the REMAP application will therefore include incorporating a

greater number of relevant predictor data layers, such as climate maxima and minima. Future

work will also focus on (i) incorporating new global image composites from the same or

different years to allow monitoring of land use and cover change with higher temporal

resolution or selection of specific time frames by users, (ii) utilizing all relevant and available

satellite imagery (e.g. Sentinel 2), (iv) improving the user experience through the provision of

more analysis tools (e.g. image differencing), and (v) improving the application for use in

collecting field data and producing maps in mobile devices.

In conclusion, we have developed REMAP to make remote sensing accessible to a very

wide audience with the aim of broadening the use of classified maps in ecosystem monitoring

and conservation programs, and to help support the conservation of natural environments. We

expect REMAP to extend the ability of volunteers, students, scientists and managers to assess

the extent of land cover changes and implement conservation actions to reduce the loss of

natural ecosystems.

ACKNOWLEDGEMENTS

The project was supported by a Google Earth Engine Research Award and an Australian Research Council Linkage Grant LP130100435, co-funded by the International Union for the Conservation of Nature, MAVA Foundation, NSW Office of Environment and Heritage, and the South Department of Environment, Water and Natural Resources. We particularly wish to thank David Thau for advice throughout the project and the Google Earth Engine team for developing the Google Earth Engine (https://earthengine.google.com), without which this application would not be possible.

AUTHOR CONTRIBUTIONS

N.J.M, D.A.K and R.M.L conceived the project. N.J.M and J.H.W. developed the remote sensing classification approach. J.H.W., D.S. and N.J.M wrote the application code and website. N.J.M wrote the manuscript with input from all coauthors.

DATA ACCESSIBILITY

REMAP: the remote ecosystem monitoring and assessment pipeline is a freely accessible and open-source web application available at: www.remap-app.org. Data used to develop the figures in this manuscript are available at figshare (https://figshare.com/s/7125654aded6d9235f08). A snapshot of the remap source code at the time of publication is available (www.github.com/REMAPApp/REMAP). Nearmap aerial imagery courtesy of Nearmap Pty. Ltd. (© 2017 Nearmap Australia Pty. Ltd.).

REFERENCES

Asbridge, E., Lucas, R., Ticehurst, C. & Bunting, P. (2016) Mangrove response to

environmental change in Australia's Gulf of Carpentaria. *Ecology and Evolution,* **6,**

3523-3539.

Asner, G.P., Knapp, D.E., Balaji, A. & Páez-Acosta, G. (2009) Automated mapping of

tropical deforestation and forest degradation: CLASlite. *Journal of Applied Remote*

Sensing, **3,** 033543.

Bachman, S., Moat, J., Hill, A.W., de Torre, J. & Scott, B. (2011) Supporting Red List threat

assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys***,** 117-

126.

Bhagwat, T., Hess, A., Horning, N., Khaing, T., Thein, Z.M., Aung, K.M., Aung, K.H., Phyo,

P., Tun, Y.L., Oo, A.H., Neil, A., Thu, W.M., Songer, M., LaJeunesse Connette, K.,

Bernd, A., Huang, Q., Connette, G. & Leimgruber, P. (2017) Losing a jewelô Rapid

declines in Myanmarøs intact forests from 2002-2014. *PLoS ONE,* **12,** e0176364.

Bland, L.M., Keith, D.A., Miller, R.M., Murray, N.J. & Rodríguez, J.P. (2017) Guidelines for

the application of IUCN Red List of Ecosystems Categories and Criteria, Version 1.1.

International Union for the Conservation of Nature, Gland, Switzerland.

Breiman, L. (2001) Random forests. *Machine learning,* **45,** 5-32.

Cavanaugh, K.C., Kellner, J.R., Forde, A.J., Gruner, D.S., Parker, J.D., Rodriguez, W. &

Feller, I.C. (2014) Poleward expansion of mangroves is a threshold response to

decreased frequency of extreme cold events. *Proceedings of the National Academy of*

Sciences, **111,** 723-727.

CBD (2014) Strategic plan for biodiversity 2011-2020. Secretariat of the Convention on

Biological Diversity.

Congalton, R.G. & Green, K. (2008) *Assessing the Accuracy of Remotely Sensed Data:*

Principles and Practices. CRC press.

Connette, G., Oswald, P., Songer, M. & Leimgruber, P. (2016) Mapping Distinct Forest

Types Improves Overall Forest Identification Based on Multi-Spectral Landsat

Imagery for Myanmarø Tanintharyi Region. *Remote Sensing,* **8,** 882.

Duke, N.C., Kovacs, J.M., Griffiths, A.D., Preece, L., Hill, D.J.E., van Oosterzee, P.,

Mackenzie, J., Morning, H.S. & Burrows, D. (2017) Large-scale dieback of

mangroves in Australiaø Gulf of Carpentaria: a severe ecosystem response,

coincidental with an unusually extreme weather event. *Marine and Freshwater*

Research, **68,** 1816-1829.

Giri, C., Ochieng, E., Tieszen, L., Zhu, Z., Singh, A., Loveland, T., Masek, J. & Duke, N.

(2011) Status and distribution of mangrove forests of the world using earth

observation satellite data. *Global Ecology and Biogeography,* **20,** 154-159.

Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H. & Liu,

S. (2013) Finer resolution observation and monitoring of global land cover: first

mapping results with Landsat TM and ETM+ data. *International Journal of Remote*

Sensing, **34,** 2607-2654.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D. & Moore, R. (2017)

Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote*

Sensing of Environment.

Green, J.M., Larrosa, C., Burgess, N.D., Balmford, A., Johnston, A., Mbilinyi, B.P., Platts,

P.J. & Coad, L. (2013) Deforestation in an African biodiversity hotspot: Extent,

variation and the effectiveness of protected areas. *Biological Conservation,* **164,** 62-

72.

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy,

T.E., Sexton, J.O., Austin, M.P., Collins, C.D., Cook, W.M., Damschen, E.I., Ewers,

R.M., Foster, B.L., Jenkins, C.N., King, A.J., Laurance, W.F., Levey, D.J., Margules,

C.R., Melbourne, B.A., Nicholls, A.O., Orrock, J.L., Song, D.-X. & Townshend, J.R.

(2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science*

Advances, **1,** e1500052.

Hansen, M.C., Alexander, K., Alexandra, T., Peter, V.P., Svetlana, T., Bryan, Z., Suspense,

I., Belinda, M., Fred, S. & Rebecca, M. (2016) Humid tropical forest disturbance

alerts using Landsat data. *Environmental Research Letters,* **11,** 034008.

Hansen, M.C. & Loveland, T.R. (2012) A review of large area monitoring of land cover

change using Landsat data. *Remote Sensing of Environment,* **122,** 66-74.

Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A.,

Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A.,

Chini, L., Justice, C.O. & Townshend, J.R.G. (2013) High-resolution global maps of

21st-century forest cover change. *Science,* **342,** 850-853.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very High

Resolution Interpolated Climate Surfaces for Global Land Areas. *International*

journal of Climatology, **25,** 1965-1978.

Keith, D.A., Rodríguez, J.P., Brooks, T.M., Burgman, M.A., Barrow, E.G., Bland, L., Comer,

P.J., Franklin, J., Link, J., McCarthy, M.A., Miller, R.M., Murray, N.J., Nel, J.,

Nicholson, E., Oliveira-Miranda, M.A., Regan, T.J., Rodríguez-Clark, K.M., Rouget,

M. & Spalding, M.D. (2015) The IUCN Red List of Ecosystems: Motivations,

Challenges, and Applications. *Conservation Letters,* **8,** 214-226.

Keith, D.A., Rodríguez, J.P., Rodríguez-Clark, K.M., Nicholson, E., Aapala, K., Alonso, A.,

Asmussen, M., Bachman, S., Basset, A., Barrow, E.G., Benson, J.S., Bishop, M.J.,

Bonifacio, R., Brooks, T.M., Burgman, M.A., Comer, P., Comín, F.A., Essl, F.,

Faber-Langendoen, D., Fairweather, P.G., Holdaway, R.J., Jennings, M., Kingsford,

R.T., Lester, R.E., Nally, R.M., McCarthy, M.A., Moat, J., Oliveira-Miranda, M.A.,

Pisanu, P., Poulin, B., Regan, T.J., Riecken, U., Spalding, M.D. & Zambrano-

Martínez, S. (2013) Scientific foundations for an IUCN Red List of Ecosystems. *PLoS*

ONE, **8,** e62111.

Lee, C. & Murray, N. (2017) redlistr: Tools for the IUCN Red List of Ecosystems and

Species.

Lewis, A., Lymburner, L., Purss, M.B., Brooke, B., Evans, B., Ip, A., Dekker, A.G., Irons,

J.R., Minchin, S. & Mueller, N. (2016) Rapid, high-resolution detection of

environmental change over continental scales from satellite dataóthe Earth

Observation Data Cube. *International Journal of Digital Earth,* **9,** 106-111.

Lucas, R., Blonda, P., Bunting, P., Jones, G., Inglada, J., Arias, M., Kosmidou, V., Petrou,

Z.I., Manakos, I., Adamo, M., Charnock, R., Tarantino, C., Mücher, C.A., Jongman,

R.H.G., Kramer, H., Arvor, D., Honrado, J.P. & Mairota, P. (2015) The Earth

Observation Data for Habitat Monitoring (EODHaM) system. *International Journal*

of Applied Earth Observation and Geoinformation, **37,** 17-28.

Lucas, R. & Mitchell, A. (2017) Integrated Land Cover and Change Classifications. *The*

Roles of Remote Sensing in Nature Conservation: A Practical Guide and Case Studies

(eds R. Díaz-Delgado, R. Lucas & C. Hurford), pp. 295-308. Springer.

Malatesta, L., Attorre, F., Altobelli, A., Adeeb, A., De Sanctis, M., Taleb, N.M., Scholte, P.T.

& Vitale, M. (2013) Vegetation mapping from high-resolution satellite images in the

heterogeneous arid environments of Socotra Island (Yemen). *Journal of Applied*

Remote Sensing, **7,** 073527-073527.

Margules, C.R. & Pressey, R.L. (2000) Systematic conservation planning. *Nature,* **405,** 243-

253.

McFeeters, S.K. (1996) The use of the Normalized Difference Water Index (NDWI) in the

delineation of open water features. *International Journal of Remote Sensing,* **17,**

1425-1432.

Moat, J. & Bachman, S. (2017) rCAT: Conservation Assessment Tools.

Mumby, P.J., Edwards, A.J., Ernesto Arias-Gonzalez, J., Lindeman, K.C., Blackwell, P.G.,

Gall, A., Gorczynska, M.I., Harborne, A.R., Pescod, C.L., Renken, H., C. C. Wabnitz,

C. & Llewellyn, G. (2004) Mangroves enhance the biomass of coral reef fish

communities in the Caribbean. *Nature,* **427,** 533-536.

Murray, N.J., Clemens, R.S., Phinn, S.R., Possingham, H.P. & Fuller, R.A. (2014) Tracking

the rapid loss of tidal wetlands in the Yellow Sea. *Frontiers in Ecology and the*

Environment, **12,** 267-272.

Murray, N.J. & Fuller, R.A. (2015) Protecting stopover habitat for migratory shorebirds in

East Asia. *Journal of Ornithology,* **156,** 217-225.

Murray, N.J., Keith, D.A., Bland, L.M., Ferrari, R., Lyons, M.B., Lucas, R., Pettorell, N. &

Nicholson, E. (in press) The role of satellite remote sensing in structured ecosystem

risk assessments. *Science of the Total Environment.*

Murray, N.J., Keith, D.A., Bland, L.M., Nicholson, E., Regan, T.J., Rodríguez, J. &

Bedward, M. (2017) The use of range size to assess risks to biodiversity from

stochastic threats. *Diversity and Distributions,* **23,** 474-483.

Murray, N.J., Phinn, S.R., Clemens, R.S., Roelfsema, C.M. & Fuller, R.A. (2012)

Continental scale mapping of tidal flats across East Asia using the Landsat Archive.

Remote Sensing, **4,** 3417-3426.

Nascimento, W.R., Souza-Filho, P.W.M., Proisy, C., Lucas, R.M. & Rosenqvist, A. (2013)

Mapping changes in the largest continuous Amazonian mangrove belt using object-

based classification of multisensor satellite imagery. *Estuarine, Coastal and Shelf*

Science, **117,** 83-93.

Neldner, V.J., Wilson, B.A., Dillewaard, H.A., Ryan, T.S. & Butler, D.W. (2017)

Methodology for Survey and Mapping of Regional Ecosystems and Vegetation

Communities in Queensland. Version 4.0. . Queensland Herbarium, Queensland

Department of Science, InformationTechnology and Innovation, Brisbane, Brisbane.

Olofsson, P., Holden, C.E., Bullock, E.L. & Woodcock, C.E. (2016) Time series analysis of

satellite data reveals continuous deforestation of New England since the 1980s.

Environmental Research Letters, **11,** 064002.

Pekel, J.F., Cottam, A., Gorelick, N. & Belward, A.S. (2016) High-resolution mapping of

global surface water and its long-term changes. *Nature,* **540,** 418-422.

Pereira, P., Brevik, E. & Trevisani, S. (2018) Mapping the environment. *Science of the Total*

Environment, **610,** 17-23.

Pettorelli, N. (2013) The normalized difference vegetation index. Oxford University Press.

Potapov, P., Hansen, M.C., Laestadius, L., Turubanova, S., Yaroshenko, A., Thies, C., Smith,

W., Zhuravleva, I., Komarova, A. & Minnemeyer, S. (2017) The last frontiers of

wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Science*

Advances, **3,** e1600821.

Queensland Department of Natural Resources and Mines (2017) Regional ecosystem and

remnant map version 8.0. Queensland, Australia.

Rodriguez-Galiano, V.F., Ghimire, B., Rogan, J., Chica-Olmo, M. & Rigol-Sanchez, J.P.

(2012) An assessment of the effectiveness of a random forest classifier for land-cover

classification. *ISPRS Journal of Photogrammetry and Remote Sensing,* **67,** 93-104.

455 Rodríguez, J.P., Keith, D.A., Rodríguez-Clark, K.M., Murray, N.J., Nicholson, E., Regan,

456   T.J., Miller, R.M., Barrow, E.G., Bland, L.M., Boe, K., Brooks, T.M., Oliveira-

457   Miranda, M.A., Spalding, M. & Wit, P. (2015) A practical guide to the application of

458   the IUCN Red List of Ecosystems criteria. *Philosophical Transactions of the Royal*

459   *Society B: Biological Sciences,* **370,** 20140003.

460 Sexton, J.O., Urban, D.L., Donohue, M.J. & Song, C. (2013) Long-term land cover dynamics

461   by multi-temporal classification across the Landsat-5 record. *Remote Sensing of*

462   *Environment,* **128,** 246-258.

463 Spalding, M.D., Ruffo, S., Lacambra, C., Meliane, I., Hale, L.Z., Shepard, C.C. & Beck,

464   M.W. (2014) The role of ecosystems in coastal protection: Adapting to climate

465   change and coastal hazards. *Ocean & Coastal Management,* **90,** 50-57.

466 Tao, S., Fang, J., Zhao, X., Zhao, S., Shen, H., Hu, H., Tang, Z., Wang, Z. & Guo, Q. (2015)

467   Rapid loss of lakes on the Mongolian Plateau. *Proceedings of the National Academy*

468   *of Sciences*.

469 Thomas, N., Lucas, R., Bunting, P., Hardy, A., Rosenqvist, A. & Simard, M. (2017)

470   Distribution and drivers of global mangrove forest change, 1996́2010. *PLoS ONE,*

471   **12,** e0179302.

Tables

Table 1. List of predictor layers available for use in land cover classifications using REMAP. Short name refers to the name given to each layer in the REMAP user interface. REMAP default indicates whether the predictor is used in a default classification. Raw data for all predictors used in REMAP are available for download from the Google Earth Engine.

Long Name	Short Name	REMAP Default?	Earth Engine ID
Topographic			
Shuttle Radar Topography Mission (SRTM) Elevation	Elevation		USGS/SRTMGL1_003
SRTM Slope	Slope		USGS/SRTMGL1_003
Climatic			
Mean Annual Temperature	Mean Annual Temperature		WORLDCLIM/V1/BIO
Annual Precipitation	Annual Precipitation		WORLDCLIM/V1/BIO
Spectral			
Normalised Difference Vegetation Index (NDVI)	NDVI		LANDSAT/LC8_SR
Normalised Difference Water index (NDWI)	NDWI		LANDSAT/LC8_SR
Water Band Index (WBI)	WBI		LANDSAT/LC8_SR
Blue band minus Red band (BR)	BR		LANDSAT/LC8_SR
Normalised Difference Blue Green (BG)	BG		LANDSAT/LC8_SR
Blue band	Blue		LANDSAT/LC8_SR
Green band	Green		LANDSAT/LC8_SR
Red band	Red		LANDSAT/LC8_SR
Near Infrared band (NIR)	NIR		LANDSAT/LC8_SR

Table 2. Descriptions of major analysis steps required to develop classified maps in REMAP. Analysis step refers to button in the sidebar of the REMAP user interface.

Analysis steps		Purpose	Options
1	Focus Region	Define the boundary of the analysis (region of interest)	Move vertices or supply by .kml file.
2	Build Training Set	Define the map classes to be used in the classification and provide georeferenced locations for each class	Uploading a training set (.csv, .kml or .JSON) or train interactively using Landsat image mosaics and predictor base layers
3	Select Predictors	Select predictor layers to be used in the classification.	Custom selection or use default settings (Table 1)
4	Classify	Run the random forest classification and return the classified map.	Run the classification on either the 2017 (present) or 2003 (historical) Landsat image mosaic.
5	Results	Obtain map accuracy statistics and area of each map class in hectares	
6	Assessment	Obtain area, AOO and EOO estimates for a single map class	
7	Export Data	Export training data or the classified image	Export training data as a .csv (for mapping or using in a GPS), a .JSON file (for saving the current workspace) or a georeferenced .tif file (for map making and further analysis).

Figures

Figure 1. Simplified process chart of REMAP: the remote ecosystem assessment and monitoring pipeline. REMAP requires spatially resolved training data, and estimates class membership of all pixels in a region of interest using global remote sensing predictor layers and the random forests classification algorithm. To facilitate observations of land cover change, classifications in REMAP can be implemented on Landsat data obtained in the year 2003 or data obtained in the year 2017.

Figure 2. Workflow demonstrating the use of REMAP to map of a single ecosystem type, mangroves of the Gulf of Carpentaria, Australia. The panels show (a) the Landsat 8 OLI 3-year composite base layer from which all Landsat indices available in REMAP are calculated, (b) the Normalized Differenced Water Index (NDWI), (c) the Normalized Differenced Vegetation Index (NDVI), and (d) the final classified map of the distribution of mangroves in the region of interest (red box).

Figure 3. Demonstration of the use of Remap to classify ecosystem types, Mount Stuart, Queensland, Australia. (a) High resolution aerial photograph, (b) the 2017 Landsat OLI image composite, (c) training data used to produce the final 3-class map, and (d) the final classified map of the distribution of ecosystems in the focal region. Aerial photography in panel (a) copyright 2017 Nearmap Australia Pty Ltd.

Figure 4. The use of REMAP to identify cover change between 2003 and 2017, Dubai. The classified land-water maps developed from (a) the 2003 global Landsat mosaic and (b) 2003 land-water classification (c) 2017 global Landsat mosaic and (d) 2017 land-water classification. (e) image differencing allows areas of coastal reclamation to be mapped and quantified. Refer to Figure 1 and Appendix A for a deforestation example.

SUPPLEMENTARY MATERIAL

Appendix A: Supplementary Data

Data used to produce Figure 1 and Figure S3

- training points (remap_points_roraimaForest.csv)
- remap workspace (remap_training_roraimaForest.json)

Data used to produce Figure 2

- training points (remap_points_carpentariaMangroves.csv)
- remap workspace (remap_training_carpentariaMangroves.json)

Data used to produce Figure 3

- training points (remap_points_mtStuart.csv)
- remap workspace (remap_training_mtStuart.json)

Data used to produce Figure 4

- training points (remap_points_Dubai_2003.csv)
- remap workspace (remap_training_Dubai_2003.json)
- training points (remap_points_Dubai_2017.csv)
- remap workspace (remap_training_Dubai_2017.json)

Data used to produce Figure S2

- training points (remap_points_chedubaMyanmar.csv)
- remap workspace (remap_training_chedubaMyanmar.json)

Appendix B: Land cover example 2

Figure S1. Comparison of land cover map produced by the Queensland State Government with the REMAP map shown in Figure 3, Mount Stuart, Queensland, Australia. (a) Queensland government regional ecosystem map produced from aerial photography and satellite image interpretation (Neldner *et al.* 2017; Queensland Department of Natural Resources and Mines 2017), (b) the classified map of the distribution of major ecosystems in the focal region produced with REMAP.

Figure S2. Demonstration of the use of REMAP to classify land cover types in Cheduba Island, Myanmar. The focal region for which the classification is implemented is shown by the red polygon.

Figure S3. Demonstration of the use of REMAP to map deforestation in the Roraima area of Brazil.