IDENTIFICATION OF AN EARLY SUBSET OF CEREBELLAR NUCLEI NEURONS IN MICE

Maryam Rahimi-Balaei[1,2], Shayan Amiri[1,3], Thomas Lamonerie[4], Sih-Rong Wu[5,6,7], Huda Y. Zoghbi[6,7,8],

G. Giacomo Consalez[9,10], Daniel Goldowitz[2], and Hassan Marzban[1,*]

[1]Department of Human Anatomy and Cell Science, The Children's Hospital Research Institute of

Manitoba (CHRIM), Max Rady College of Medicine, Rady Faculty of Health Sciences, University of

Manitoba, Winnipeg, Manitoba, Canada

[2]Department of Medical Genetics, Centre for Molecular Medicine and Therapeutics, BC Children's

Hospital Research Institute, University of British Columbia, Vancouver, BC, Canada

[3]Department of Pharmacology and Therapeutics, Division of Neurodegenerative Disorders, St Boniface

Hospital Albrechtsen Research Centre, University of Manitoba, Winnipeg, MB, Canada

[4]Université Côte d'Azur, CNRS, Inserm, iBV, France

[5]University of California San Francisco (UCSF), San Francisco, California, USA

[6]Department of Neuroscience, Baylor College of Medicine, Houston, TX, 77030, USA

[7]Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital, Houston, TX, 77030, USA

[8]Howard Hughes Medical Institute, Baylor College of Medicine, Houston, TX, 77030, USA

[9]Division of Neuroscience, IRCCS Ospedale san Raffaele

[10]Università Vita-Salute San Raffaele, 20132 Milan, Italy

Address correspondence to: Dr. H. Marzban, Department of Human Anatomy & Cell Science, Rm 129

BMSB, 745 Bannatyne Avenue, Max Rady College of Medicine, Rady Faculty of Health Sciences,

University of Manitoba, Winnipeg, Manitoba, R3E 0J9, Canada

Tel: 204-789-3467

Fax: 204-789-3920

Email: hassan.marzban@umanitoba.ca

26	**Manuscript:**	29 Pages; 9 Figures; 0 Tables
27	**Key Words:**	Cerebellar nuclei, SNCA, MEIS2, OTX2, Atoh1
28	**Running Title:**	Rostroventral subset of cerebellar nuclei neurons

ABSTRACT

Cerebellar nuclei (CN) neurons serve as the primary output of the cerebellum and originate from the

cerebellar primordium at early stages of cerebellar development. Employing various methodologies, we

have characterized a specific subset of CN neurons that do not originate from the rhombic lip or ventricular

zone of the cerebellar primordium. Embryos were collected at early stages of development and processed

for immunohistochemistry (IHC), Western blotting, *in situ* hybridization (ISH), embryonic culture, DiI

labeling, and flow cytometry analysis (FCM). Our findings indicate that a subset of CN neurons expressing

α-synuclein (SNCA), OTX2, MEIS2, and p75NTR (NGFR) are located in the rostro-ventral (rv) region

of the nuclear transitory zone (NTZ), while CN neurons derived from the rhombic lip are positioned in

the caudo-dorsal (cd) area of the NTZ in the cerebellar primordium. Utilizing Otx2-GFP and Atoh1$^{-/-}$

mice, we have determined that these cells do not originate from the germinal zone of the cerebellar

primordium. These results suggest the existence of a novel extrinsic germinal zone for the cerebellar

primordium, possibly the mesencephalon, from which early CN neurons originate.

SIGNIFICANCE STATEMENT

The cerebellum contains a variety of distinct neuronal populations, each playing a significant role in its

function within the brain. This research demonstrates that a particular subset of cerebellar nuclei neurons

originates from a previously unrecognized germinal zone specific to the cerebellar primordium,

independently of Atoh1's influence.

INTRODUCTION

In the mouse, the cerebellar primordium emerges at approximately embryonic day (E) 7–E8 as a neuroepithelial swelling on the rostral lip of the fourth ventricle, which is part of the alar plate of the metencephalon (rhombomere-1) (Kebschull et al., 2023, Goldowitz and Hamre, 1998, Sotelo, 2004, Wang and Zoghbi, 2001a, Marzban et al., 2015). The cerebellar primordium is rostrally limited by the isthmus, the mesencephalon-rhombencephalon (rhombomere-1, r1) or midbrain-hindbrain boundary. Orthodenticle homeobox 2 (*Otx2*) and gastrulation brain homeobox 2 (*Gbx2*) are among the earliest genes expressed in the neuroectoderm, dividing it into anterior and posterior domains with a border that marks the isthmus. *Otx2* is required for the development of the forebrain and midbrain, and *Gbx2* for the anterior hindbrain (Millet et al., 1999). The isthmus, at the junction of *Otx* and *Gbx* expressing areas, is a multi-signaling center regulating genes essential for the early development of both the mesencephalon and cerebellar primordium (Joyner et al., 2000, Wurst and Bally-Cuif, 2001).

The cerebellar primordium contains two distinct germinal zones, which are responsible for cerebellar neurogenesis and gliogenesis; the ventrally-located ventricular zone (VZ) and dorsally-located rhombic lip (Fink et al., 2006). Early studies on the development of CN indicated that both glutamatergic and GABAergic CN neurons originate from the cerebellar VZ (Voogd and Glickstein, 1998). Further studies highlighted the involvement of the rhombic lip as an origin for glutamatergic CN neurons (between E9-E12) (Fink et al., 2006, Wang and Zoghbi, 2001a, Green and Wingate, 2014, Machold and Fishell, 2005, Kebschull et al., 2023). LIM homeobox transcription factor 1 alpha (LMX1A) regulates the development of glutamatergic CN neurons originating from the rhombic lip. It is also expressed in the nuclear transitory zone (NTZ) and is considered to be a marker for the majority of rhombic lip-derived CN neurons in addition to a subset of LMX1A positive cells that do not originate from the rhombic lip migratory stream (Chizhikov et al., 2006, Chizhikov et al., 2010).

In this report, we provide evidence that a subset of CN neurons expressing OTX2, SNCA, MEIS2,

and p75 Neurotrophin Receptor (p75NTR)/Nerve Growth Factor Receptor (NGFR) originate from the

rostral end of the cerebellar primordium and are positioned in the rostroventral region of the NTZ during

early cerebellar development. Our data suggest that the mesencephalon could represent a third germinal

zone and, as such, the origin of previously unrecognized neurons that contribute to CN formation and

development.

RESULTS

In mouse embryos, the earliest neuronal populations, which are located at the rostral end of the

cerebellar primordium, are immunopositive for neurofilament-associated antigens (NAA) 3A10, a marker

typically expressed in neuronal somata and axons (Marzban et al., 2008, Marzban et al., 2019) (Fig. 1).

To further explore these cells in the rostral end of the cerebellar primordium, we used an antibody to detect

SNCA, which is expressed in CN neurons during embryogenesis (Zhong et al., 2010). In sagittal sections

of the E9 cerebellar primordium, we identified SNCA$^+$ neurons at the rostral end of the NTZ (Fig. 1E-F

and Fig. Suppl. 4). To determine the location of the SNCA$^+$ cells and compare it to that of the rhombic

lip-derived LMX1A$^+$ CN neurons (E10-12) in the NTZ, double immunofluorescence labeling of SNCA

and LMX1A was performed on sagittal sections of E12 cerebellar primordium (Fig. 2 A-D, medial section

and Fig. E-H, lateral section). This shows that an LMX1A$^+$ population of CN neurons, originating from

the rhombic lip, either flanks SNCA$^+$ neurons in the medial section (Fig. 2A-D) or is located dorsally in

the lateral section (Fig. 2E-H) in the NTZ. To further explore the characteristics of SNCA$^+$ cells, double

immunofluorescence labeling with OTX2 and SNCA was performed during early cerebellar development.

OTX2 is highly expressed in the mesencephalon and its caudal limit coincides with the boundary of the

rhombencephalon (i.e., the isthmus) (Kurokawa et al., 2004). Interestingly, results from double

immunofluorescence experiments in sagittal sections of E12 cerebella showed that OTX2 was highly

expressed in the mesencephalon (Fig. 3A, B), and was co-expressed with SNCA$^+$ cells in the rostral region

of the cerebellar primordium (Fig. 3 A-E). Next, we used quantitative flow cytometry analysis to

determine whether OTX2$^+$ cells co-express SNCA and/or LMX1A (Fig. 3F, G). An increase in the number

of OTX2 and SNCA positive cells was observed at E14 in comparison to E12 (*p<0.05). The majority of

neural cells were not SNCA$^+$/LMX1A$^+$, consistent with our IHC findings, or SNCA$^+$/OTX2$^+$, with the

exception of a few cells that did show co-labeling. Of note, more SNCA$^+$/OTX2$^+$ than SNCA$^+$/LMX1A$^+$

cells were detected at E12 compared to E14.

IHC performed on sagittal sections of E12 to E15 cerebella showed that OTX2$^+$ cells are present

at the rostral end of the cerebellar primordium (Fig. 4 A-F). To evaluate the number of OTX2$^+$ cells in the

developing rostral cerebellum, the total number of OTX2$^+$ cells in the cerebellar primordium sections were

counted from E12 to E15 (Fig. 4G). A comparison between these embryonic stages indicated a slight

increase in the number of OTX2+ cells between E12 and E15. Significant differences were observed

between E12 and E14 (**p<0.01), E12 and E15 (***p<0.001), as well as E13 and E15 (#p<0.05).

To confirm the presence of OTX2$^+$ cells in the rostral end of the cerebellar primordium, we used

Otx2-GFP transgenic mice and performed double immunofluorescence labeling with OTX2 and GFP on

E13 sagittal sections (Fig. 5). Results showed that an extension of OTX2-GFP positive cells, highly

prevalent in the mesencephalon, crossed the isthmus and terminated in the rostral end of the cerebellar

primordium in the NTZ (Fig. 5B, C, and D). To further validate our observations, we employed RNAscope

fluorescence *in situ* hybridization (FISH) at E12 to detect the presence of *Otx2* mRNA in the cerebellar

primordium (Fig. 5 E-J). *Otx2* mRNA was highly expressed in the mesencephalon and caudally extended

to the rostral cerebellar primordium in the NTZ (Fig. 5 E-J and Suppl. Fig. 4).

To evaluate whether OTX2$^+$ cells are present in the absence of *Snca* gene expression, we used *Pap*$^-$

$^{/-}$; *Snca*$^{-/-}$ mice. OTX2 immunoperoxidase staining showed that even in the absence of *Snca* expression,

OTX2$^+$ cells still terminated in the rostral cerebellar primordium (Suppl. Fig. 1). Nerve growth factor

receptor (p75NTR; Ngfr), which regulates neuronal proliferation and differentiation (Jiang et al., 2008,

Bernabeu and Longo, 2010, Dechant and Barde, 2002), has also been shown to be expressed in the rostral

end of the cerebellar primordium in Pap$^{-/-}$; Snca$^{-/-}$ mice (Rahimi-Balaei et al., 2019b). To determine

whether SNCA$^+$ neurons are p75NTR immunopositive, double immunolabeling with SNCA and p75NTR

was performed. Section IHC revealed that SNCA$^+$ cells in the NTZ exhibited cell membrane expression

of p75NTR (Fig. 6A–D). This data was further confirmed by Western blot analysis of SNCA and p75NTR

protein expression in embryonic cerebellar lysates (E11, 13 and 15; Fig. 6d). In addition, to conclusively

determine that p75NTR is expressed in the membranes of SNCA$^+$ cells, we utilized primary dissociated

cultures of cerebellum at E10, DIV 4 (Fig. 6E-H). IHC demonstrated that p75NTR was indeed expressed

in the membranes of SNCA$^+$ cells (with punctate appearance, Fig. 6E–H (Dechant and Barde, 2002)).

To determine whether OTX2 and SNCA immunopositive cells exist in the absence of rhombic lip-

derived CN neurons, we used Atoh1$^{-/-}$ mice (Fig. 7 A-C). Immunostaining revealed the presence of SNCA$^+$

and OTX2$^+$ cells at the rostral end of the cerebellar primordium in these animals. To validate our findings,

we evaluated MEIS2, a transcription factor typically expressed in the NTZ, in Atoh1$^{+/+}$ and Atoh1$^{-/-}$

sagittal sections of the cerebellar primordium at E12. Notably, our results show the presence of two distinct

sets of MEIS2$^+$ cells in the NTZ of Atoh1$^{+/+}$ embryos: rhombic lip-derived CN neurons located in the

caudodorsal region, which do not express SNCA (MEIS2$^+$/SNCA$^-$), and a subset of MEIS2$^+$/SNCA$^+$ CN

neurons situated in the rostroventral region of the NTZ (Fig. 7D-I). In Atoh1$^{-/-}$ sections, MEIS2$^+$/SNCA$^+$

cells were still present in the rostroventral region of the NTZ, whereas the subpopulation of

MEIS2$^+$/SNCA$^-$ rhombic lip-derived CN neurons was absent in the caudodorsal region of the NTZ (Fig.

7 J-O).

To explore if the rostroventral subset of CN neurons derived from an external cerebellar germinal

zone, possibly the mesencephalon, we utilized FAST DiI as a neuronal tracer to mark cells within a

potential source region. FAST DiI was applied to the dorsum of the caudal mesencephalon at E9, where

it was maintained for 4 days (Fig. 8 A, a). At DIV 4, cells stained with DiI in the dorsum of the

mesencephalon were present in both rostral (Fig. 8B) and caudal (Fig. 8C) directions. Sections from the

cerebellar primordium revealed the presence of DiI-labeled cells within the cultured embryo (Fig. Suppl.

2). To investigate the earliest DiI-positive cell population in the mesencephalon and avoid unwanted cell

staining due to long DiI exposure, we focused on early cerebellar development at E9; it has been reported

that one of the premier neuronal populations in the CNS is present at this time in the mesencephalon and

projects caudally (Stainier and Gilbert, 1990, Easter et al., 1993). To determine if an early generation of

mesencephalic DiI-labeled cells is present within the cerebellar primordium, we limited cellular DiI

exposure to only 24 hours (Fig. 8 D, d), after which it was removed (Fig. Suppl. 3). Intriguingly, almost

all DiI$^+$ cells migrated caudally and presented in the cerebellar primordium (Fig. 8E, e and Fig. Suppl. 3).

DiI staining was clear and strong in cells as shown in Fig. 8 E, e and Fig. Suppl. 3 (arrowhead); it should

be noted, however, that some DiI staining was evident in other cells in the rostral end of the cerebellar

primordium, although the signal appeared relatively weak (Suppl. Fig. 3).

DISCUSSION

Previously, it was commonly held that all CN neurons had a singular origin in the ventricular zone

(Pierce, 1975, Miale and Sidman, 1961). However, recent discoveries have challenged this notion,

revealing a dual origin for CN neurons. Some originate from the ventricular zone, while others arise from

the rhombic lip (Wang and Zoghbi, 2001b, Wang et al., 2005, Ben-Arie et al., 1997). Genetic fate mapping

further suggested that most glutamatergic CN projection neurons may arise from the rhombic lip (Ben-

Arie et al., 1997, Machold and Fishell, 2005, Fink et al., 2006). Transcription factor expression patterns

indicate that CN neurons migrate from the rhombic lip to the NTZ through a subpial stream pathway,

while sequentially expressing the genes *Lmx1a*, *Pax6*, *Tbr2,* and *Tbr1* (Fink et al., 2006).

In this study, we characterized a subset of CN neurons that do not originate from the cerebellar

primordium germinal zones during early cerebellar development. We demonstrated a newly identified

subset of CN neurons, which are SNCA$^+$/ OTX2$^+$/ MEIS2$^+$/ p75NTR$^+$/ LMX1A$^-$, in the rostroventral

region of the NTZ in the rostral end of the cerebellar primordium. This suggests the existence of a new

germinal zone during cerebellar neurogenesis.

The exact origin of the SNCA$^+$/ OTX2$^+$/ MEIS2$^+$/ p75NTR$^+$/ LMX1A$^-$ CN neurons is currently

unclear; however, they do not originate from the rhombic lip or ventricular zones. Rhombic lip-derived

Tbr1$^+$/Lmx1a$^+$ CN neurons are born at E9, but do not reach the NTZ until ~E11 (Kebschull et al., 2023,

Machold and Fishell, 2005, Wang et al., 2005, Manto et al., 2012, Fink et al., 2006, Rahimi-Balaei et al.,

2018). Our results revealed that SNCA$^+$ cells are a group of differentiating neurons (NAA 3A10$^+$) present

in the NTZ at E9, before the arrival of any neurons that originate from the rhombic lip. The majority of

SNCA$^+$ CN neurons are not LMX1A$^+$; however, a small proportion of cells seems to exhibit co-

expression. This suggests that in the early stages of CN neurogenesis, the pattern of protein expression in

SNCA$^+$ neurons may be changing, possibly with respective down- and upregulation of *Snca* and *Lmx1a$^+$*.

Conversely, this may not occur at all, and 'co-labeling' could have been observed due to overlapping cells.

A similar display of SNCA expression has recently been demonstrated in oligodendrocyte development

and maturation (Djelloul et al., 2015). Our findings indicated that CN neuron somata are highly positive

for SNCA until E14, after which SNCA expression is refined to PCs at P0 (Zhong et al., 2010) and axonal

projections, indicating that SNCA$^+$ cells at the rostral end of the cerebellar primordium are associated with

CN establishment in the NTZ. Our results from studies using Pap$^{-/-}$; *Snca$^{-/-}$* mice showed that lack of

SNCA during cerebellar development did not affect the rostroventral subset of CN neurons (Rahimi-

Balaei et al., 2019a), which still expressed OTX2 and terminated in the rostral cerebellar primordium. In

adult mice, postnatal expression of SNCA has been reported in medial CN neurons (Fujita et al., 2020).

Furthermore, our observations reveal the expression of MEIS2 in the NTZ (Fig. 7 and (Morales and Hatten, 2006)), along with several other NTZ markers such as POU3F1, BRN2, IRX3, and MEIS1 as documented in (Kebschull et al., 2023, Wu et al., 2022), c-Ret and Tlx3 (Fig. Suppl. 4), indicating the existence of discrete subpopulations of precursors within the CN. MEIS2 (homeobox transcription regulator) is expressed in the mesencephalic alar plate of both mice (Cecconi et al., 1997) and chicks (Agoston and Schulte, 2009, Bobak et al., 2009). MEIS2 plays a crucial role in the development of cranial nerves and craniofacial structures (Machon et al., 2015). It has been suggested that mesencephalon development is regulated by Meis2 cross-talk with Otx2 (Agoston and Schulte, 2009). Notably, the expression of MEIS2 in CN neurons is remarkably distinct, allowing for a clear demarcation of the caudodorsal (SNCA$^-$/MEIS2$^+$) and rostroventral (SNCA$^+$/MEIS2$^+$) regions of the NTZ, whether originating from the rhombic lip or not (Fig. 7D-I). CN neurons derived from the rhombic lip require ATOH1 expression, as they are absent in an Atoh1-null mutant (Wang et al., 2005, Machold and Fishell, 2005, Zordan et al., 2008, Florio et al., 2012, Ben-Arie et al., 1997). In the Atoh1-null cerebellar primordium, the caudodorsal (SNCA$^-$/MEIS2$^+$) subset of CN is absent in the NTZ, while the rostroventral (SNCA$^+$/MEIS2$^+$) population is still present, indicating that the Meis2$^+$ rostroventral subset of the CN in the NTZ develops independently from Atoh1.

What are the potential origins of SNCA$^+$/ OTX2$^+$/ MEIS2$^+$/ p75NTR$^+$/ LMX1A$^-$ cells in the NTZ during the early stages of cerebellar development? Our study suggests that the origin of these cells may be guided by a continuous flow of rostroventral cells toward the rostral end (mesencephalon), and caudodorsal cells toward the caudal end (rhombic lip) of the cerebellar primordium. It is well established that *Otx2* is required for the development of the fore- and midbrain, while *Gbx2* is necessary for anterior hindbrain development (Simeone et al., 1992, Alvarado-Mallart, 2005, Li and Joyner, 2001, Joyner et al., 2000). The expression of OTX2 in the rostral and GBX2 in the caudal neural tube, is considered limited to the mesencephalon/metencephalon boundary (isthmus) (Alvarado-Mallart, 2005). However, a study by

Martinez *et al.* on chick/quail chimeras determined that the rostral portion of the cerebellar primordium is

located more rostrally in the so-called 'mesencephalic' alar plate (Martinez and Alvarado-Mallart, 1989).

Recently, *Isl1* positive cells in the anterior cerebellum were shown to exhibit residual OTX2 protein

activity (Wizeman et al., 2019). Several other studies have shown that rostral cerebellar development is

associated with factors expressed in the caudal mesencephalon, such as engrailed family En1 and -2

(Joyner et al., 1991, Hanks et al., 1995, Millen et al., 1995), and Acp2 (Bailey et al., 2013, Bailey et al.,

2014), which are regulated by the multi-signaling center known as the isthmic organizer (Martinez et al.,

2013). Furthermore, by employing embryonic cultures in conjunction with DiI labeling, we determined

that the source of the rostroventral subset of CN neurons appears to be an external cerebellar germinal

zone, possibly originating from the mesencephalon, which is in line with other studies (Nichols and Bruce,

2006, Wizeman et al., 2019). While our current study hints at the possibility of the caudal mesencephalon

serving as a novel extrinsic germinal zone for the cerebellar primordium, further investigation is required.

Genetic inducible fate mapping and long-term follow-up studies are essential to gain a comprehensive

understanding of the origin, fate, and connectivity of, as well as the role played by the SNCA$^+$/ OTX2$^+$/

MEIS2$^+$/ p75NTR$^+$/ LMX1A$^-$ rostroventral subset of CN neurons in the development of the cerebellum.

In conclusion, our study indicates that the SNCA$^+$/ OTX2$^+$/ MEIS2$^+$/ p75NTR$^+$/ LMX1A$^-$

rostroventral subset of CN neurons do not originate from the well-known distinct germinative zones of

the cerebellar primordium. Instead, our findings suggest the existence of a previously unidentified

extrinsic germinal zone, potentially the mesencephalon.

MATERIAL AND METHODS

Animal maintenance

All animal procedures were performed in accordance with institutional regulations and the *Guide*

to the Care and Use of Experimental Animals from the Canadian Council for Animal Care. For this study,

we used embryos from 47 CD1 timed-pregnant mice at embryonic day (E) 9 to E18 (total number of embryos was 409). Our approach to sample size was to include enough samples to reach a sufficient level, which was determined empirically (in most of the experiments, n was equal to 9).Timed-pregnant, prostatic acid phosphatase (PAP) mutant ($Pap^{-/-}$) (Zylka et al., 2008, Hokin and Hokin, 1959) mice were used at E12, because they do not express $Snca$ ($Pap^{-/-}$;$Snca^{-/-}$) and are a valuable experimental tool to assess whether SNCA is required in the development of the CN neurons (Rahimi-Balaei et al., 2019b). In addition, we studied GFP-tagged Otx2 mouse embryos (from Thomas Lamonerie lab at Université Côte d'Azur), a reporter mouse line in which the Otx2 protein is fused to the fluorescence protein GFP ($Otx2^{Otx2-GFP/+}$; (Fossat et al., 2007)). The animal procedures related to GFP-tagged Otx2 mouse embryos were in accordance with Université Côte d'Azur Institutional Animal Care and Use Committee guidelines. *Atoh1* knockout embryos were provided by Dr. Huda Zoghbi at the Baylor College of Medicine. The mice were bred, phenotyped, and genotyped (by the Zoghbi lab) according to the previously described protocol (Wang et al., 2005). Animal procedures involving Atoh1 knockout mice were in accordance with The Baylor college of Medicine Institutional Animal Care and Use Committee guidelines.

All timed-pregnant CD1 mice were obtained from the Central Animal Care Service, University of Manitoba. Animals were housed at room temperature and relative humidity (18–20°C, 50–60%) on a light:dark cycle (12:12 h) with free access to food and water. The embryonic age was determined by referencing the morning after the detection of a vaginal plug following overnight mating, considered to be E 0.5. CD1 timed-pregnant mice were anesthetized at E9, 10, 11, 12, 13, 14, 15, or 18 (+ 0.5) (n=47) using 40% isoflurane (USP, Baxter Co. Mississauga, Ontario, Canada), after which embryos were removed and prepared for Western blotting, embryonic culture, or flow cytometry analysis (FCM) or fixed in 4% paraformaldehyde (PFA) for immunohistochemistry (IHC) or neutral buffered formalin for *in situ* hybridization (ISH).

Section immunohistochemistry

Cryostat sections (20 μm) of 4% PFA fixed samples were utilized for IHC as described in our

previous studies (Bailey et al., 2014, Bailey et al., 2013). Antibody dilutions were used as follows: α-

synuclein 1:500 (sc-69977, Santa Cruz), p75NTR 1:1000 (8238, Cell Signaling), LMX1A 1:500

(AB10533, EMD Millipore Corporation), OTX2 1:1000 (ab114138, Abcam), NAA 1:500 (3A10,

Developmental Studies Hybridoma Bank), Meis2 1:500 (ab244267, Abcam), and GFP 1:1000 (1020,

Aves Labs). Fluorescent detection was performed using antibodies as follows: Streptavidin Alexa Fluor®

488 conjugate, Alexa Fluor® 568 Goat Anti-Rabbit IgG (H+L), Alexa Fluor® 488 Chicken Anti-Mouse

IgG (H+L), Alexa Fluor® 488 Chicken Anti-Rabbit IgG (H + L), and Alexa Fluor® 568 Goat Anti-Mouse

IgG (H+L) (S-11223, A-11036, A21200, A21441, A11004, respectively, from Life Technologies),

AlexaFluro® 568 Donkey anti-goat (A-11057, invitrogen), and AlexaFluro® 647 Donkey anti-mouse (A-

31571, Invitrogen), all at 1:1000. Detection of peroxidase IHC was performed as described previously

(Rahimi Balaei et al., 2016, Bailey et al., 2014, Bailey et al., 2013) using HRP conjugated goat anti-rabbit

IgG and goat anti-mouse IgG (H+L) antibodies (EMD Millipore Corporation, 12-348 and AP308P,

respectively), both at 1:500, and developed with 3,3'-diaminobenzidine solution (DAB, Sigma, St. Louis

MO, USA).

Primary dissociated cerebellum cell cultures

Primary cerebellar cultures were prepared from E10 CD1 mice, and maintained for varying 'days

in vitro' (DIV; 1, 2, 3, 5, and 8), according to published methods (Shabanipour et al., 2019). Briefly, the

entire cerebellum was removed from each embryo and immediately placed into ice cold Ca^{2+}/Mg^{2+}-free

Hank's balanced salt solution (HBSS) containing gentamicin (10 μg/ml) and glucose (6 mM). Next,

cerebella were incubated at 34°C for 12 min in HBSS containing 0.1% trypsin. After washing, the

cerebella were gently triturated with a Pasteur transfer pipette in HBSS containing DNase I (5 U/ml) and

12 mM $MgSO_4$ until the cell mass was no longer visible. Cells were collected by centrifugation (1,200

rpm, 4°C for 5 min) and re-suspended in seeding medium (Dulbecco's modified Eagle's medium and F12,

1:1) supplemented with putrescine (100 μM), sodium selenite (30 nM), L-glutamine (1.4 mM), gentamicin

(5 μg/ml), and 10% heat-inactivated fetal bovine serum. Cells were seeded on poly-L-ornithine coated

glass coverslips (12 mm) at a density of 5×10^6 cells/ml, and each coverslip was placed into an individual

well of a 24-well plate. After 6–8 h incubation in a CO_2 incubator (95% humidity, 37°C, 5% CO_2) to

accomplish cell adherence, 500 μl of culture medium supplemented with transferrin (200 μg/ml), insulin

(20 μg/ml), progesterone (40 nM), and triiodothyronine (0.5 ng/ml) was added to each culture well. After

7 days, half of the medium in each dish was replaced with fresh medium that was additionally

supplemented with cytosine arabinoside (4 μM) and bovine serum albumin (100 μg/ml) (Bailey et al.,

2013, Marzban and Hawkes, 2007, Marzban et al., 2003).

Embryonic cultures and DiI labeling of cells within the mesencephalon

Embryonic cultures were prepared from E9 and E10 CD1 timed-pregnant mice, and maintained

for various DIV (4 and 6). Each embryo was removed from the amniotic sac and immediately placed into

ice cold Ca^{2+}/Mg^{2+}-free HBSS containing gentamicin (10 μg/ml) and glucose (6 mM). Embryos were

placed into 24-well plates in culture medium plus 10% fetal bovine serum and incubated in a CO_2

incubator (95% humidity, 37°C, 5% CO_2) (Marzban and Hawkes, 2007). During incubation, embryos

were monitored every 6 h to evaluate the heartbeat (as an indicator of survival). On DIV 4 or 6, each well

was fixed with 4% PFA and prepared for whole mount IHC.

For neuronal tracing and labeling, we used FAST DiI crystal (FAST DiI™ solid; DiIΔ9,12-C18(3),

CBS (1,1'-Dilinoleyl-3,3,3',3'-Tetramethylindocarbocyanine, 4-Chlorobenzenesulfonate, D7756, Fisher

Scientific). Briefly, FAST DiI was inserted into the mesencephalon at E9 using a sharp-ended needle

(30g). After insertion of FAST DiI, images were captured by a stereomicroscope to assess the location of

DiI at DIV 0. After placing the embryos into 24-well plates in culture medium, they were monitored every

6 hours and fixed with 4% PFA on the desired day. Next, whole mount IHC with neurofilament-associated

antigen (NAA 3A10) was performed to visualize neural fiber growth, followed by sectioning and imaging

of the DiI positive cells in the mesencephalon and cerebellar primordium.

Fluorescent in situ hybridization (FISH)-RNAscope

All of the ISH experiments were carried out using E12 CD1 mice, in which the cerebellar

primordium is well established, using RNAscope ACD HybEZ™ II Hybridization System and

RNAscope® Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics, Hayward, CA, USA).

Briefly, embryos were fixed in 10% (vol/vol) neutral buffered formalin at room temperature for 24 h,

dehydrated, and embedded in paraffin. Tissue sections cut at 10 μm thickness were processed for RNA *in*

situ detection according to the manufacturer's user manual. The sequence of the probe used in this study

was Mm-*Otx2*-C3 (444381, ACD); cyanine 3 (NEL744E001KT; TSA® Plus, Perkin Elmers, Waltham,

MA, USA) was used as the fluorophore.

Western blot analyses

Equal amounts of total protein were separated by SDS/PAGE in 10–15% precast gels (Bio-Rad,

Hercules, CA, USA) and transferred onto PVDF-membranes. For Western blot analysis, membranes were

blocked in 5% nonfat dry milk (NFDM) in TBS containing 0.02% Tween 20 (TBST) for 1hour at RT and

subsequently incubated overnight at 4°C with primary antibodies [α-synuclein (sc-69977, Santa Cruz,

1:2000) or p75NTR (8238, Cell Signaling, 1:1000), all dilutions in blocking buffer]. After washing with

TBST, membranes were exposed to secondary antibodies [HRP conjugated goat anti-mouse IgG

(AP308P, Millipore, 1:6000) or HRP conjugated goat anti-rabbit IgG (12-348, Millipore, 1:6000)]. Bands

were visualized using the enhanced chemiluminescence (ECL) protocol on scientific imaging film. All

bands were normalized to β-actin expression.

Counting of OTX2 positive cells
338
To assess the number of OTX2-positive cells, we conducted immunohistochemistry (IHC) labeling

on slides containing serial sections from embryonic days 12, 13, 14, and 15 (n=3 at each timepoint). Under

the microscope, we systematically counted OTX2-positive cells within the cerebellar primordium. This

analysis encompassed a minimum of 10 sections, spread across at least 3 slides, ensuring comprehensive

coverage of OTX2 expression along the mediolateral axis of the cerebellar primordium. For each slide,

the counts of OTX2-positive cells from all sections were cumulatively calculated to determine the total

number of positive cells per slide. Subsequently, statistical analysis was employed to compare the results

obtained at different developmental time points.

Flow cytometry analysis

After dissociation (see primary dissociated cerebellar cell culture section), cells were fixed in 4%

PFA. Each group consisted of 350,000 cells and all procedures before sorting were performed at room

temperature. Primary antibodies used: α-synuclein 1:500 (sc-69977, Santa Cruz), LMX1A 1:1000

(AB10533, EMD Millipore Corporation), and OTX2 1:500 (ab114138, Abcam). Fluorescent detection

was performed using antibodies as follows: Alexa Fluor 488 Chicken Anti-Rabbit IgG (H / L) and Alexa

Fluor 568 Goat Anti-Mouse IgG (H+L) (A21441 and A11004, respectively, from Life Technologies),

both at 1:750. Cells were pelleted by centrifugation, washed, and resuspended in 200 μL staining buffer

(1X PBS contains 1% FBS and 1mM EDTA). Data were acquired on a CytoFlex-LX flow cytometer

(Beckman Coulter) equipped with 355, 375, 405, 488, 561, 638 and 808 Laser lines using CytExpert

software, and analyzed with Flow Jo (version 10, Treestar, San Carlos, CA) at the University of Manitoba

flow cytometry core facility. Cellular debris was excluded using forward light scatter/side scatter plot.

Imaging and figure preparation

For bright field microscopy, images were captured using a Zeiss Axio Imager M2 microscope

and subsequently analyzed with Zeiss Microscope Software (Zen Image Analyses software; Zeiss,

Toronto, ON, Canada). For fluorescence microscopy of entire cerebellar sections, a Zeiss Lumar V12

Fluorescence stereomicroscope (Zeiss, Toronto, ON, Canada) equipped with a camera was used; captured

images were analyzed using Zen software. For high magnification fluorescence microscopy, a Zeiss Z1

and Z2 Imager and a Zeiss LSM 700 confocal microscope (Zeiss, Toronto, ON, Canada) equipped with

camera and Zen software were used to capture and analyze images. Images were cropped, corrected for

brightness and contrast, and assembled into montages using Adobe Photoshop CS5 Version 12.

Statistical analysis

Data on OTX2 positive cell counts are presented as the mean ± standard error of mean (SEM) from

n separate experiments. Statistical significance of differences was evaluated by One-Way ANOVA

followed by a Tukey`s multiple comparison test. Differences were considered statistically significant

when $P<0.05$. All statistical analyses were performed using GraphPad Prism 6 software for Windows.

ACKNOWLEDGEMENTS:

The authors would like to acknowledge Dr. Christine Zhang and Mr. Gerald Stelmack for their

technical assistance, and Science Impact (Winnipeg, Canada) for (post-)editing the manuscript.

COMPETING INTEREST:

The authors have no conflicts of interest.

FUNDING:

This study was supported by grants from the Natural Sciences and Engineering Research Council

(HM: NSERC Discovery Grant # RGPIN-2018-06040), The Children's Hospital Research Institute of

Manitoba (HM: CHRIM Grant # 320035), Research Manitoba Tri-Council Bridge Funding Program (HM:

Grant # 47955), and University Collaborative Research Program (UCRP).

419 **REFERENCES**
420
AGOSTON, Z. & SCHULTE, D. 2009. Meis2 competes with the Groucho co-repressor Tle4 for binding

to Otx2 and specifies tectal fate without induction of a secondary midbrain-hindbrain boundary

organizer. *Development,* 136**,** 3311-22.

ALVARADO-MALLART, R. M. 2005. The chick/quail transplantation model: discovery of the isthmic

organizer center. *Brain Res Brain Res Rev,* 49**,** 109-13.

BAILEY, K., RAHIMI BALAEI, M., MANNAN, A., DEL BIGIO, M. R. & MARZBAN, H. 2014.

Purkinje cell compartmentation in the cerebellum of the lysosomal Acid phosphatase 2 mutant

mouse (nax - naked-ataxia mutant mouse). *PLoS One,* 9**,** e94327.

BAILEY, K., RAHIMI BALAEI, M., MEHDIZADEH, M. & MARZBAN, H. 2013. Spatial and

temporal expression of lysosomal acid phosphatase 2 (ACP2) reveals dynamic patterning of the

mouse cerebellar cortex. *Cerebellum,* 12**,** 870-81.

BEN-ARIE, N., BELLEN, H. J., ARMSTRONG, D. L., MCCALL, A. E., GORDADZE, P. R., GUO,

Q., MATZUK, M. M. & ZOGHBI, H. Y. 1997. Math1 is essential for genesis of cerebellar

granule neurons. *Nature,* 390**,** 169-72.

BERNABEU, R. O. & LONGO, F. M. 2010. The p75 neurotrophin receptor is expressed by adult mouse

dentate progenitor cells and regulates neuronal and non-neuronal cell genesis. *BMC*

neuroscience, 11**,** 1.

BOBAK, N., AGOSTON, Z. & SCHULTE, D. 2009. Evidence against involvement of Bmp receptor 1b

signaling in fate specification of the chick mesencephalic alar plate at HH16. *Neurosci Lett,* 461**,**

223-8.

CECCONI, F., PROETZEL, G., ALVAREZ-BOLADO, G., JAY, D. & GRUSS, P. 1997. Expression of

Meis2, a Knotted-related murine homeobox gene, indicates a role in the differentiation of the

forebrain and the somitic mesoderm. *Dev Dyn,* 210**,** 184-90.

CHIZHIKOV, V., STESHINA, E., ROBERTS, R., ILKIN, Y., WASHBURN, L. & MILLEN, K. J.

2006. Molecular definition of an allelic series of mutations disrupting the mouse Lmx1a (dreher)

gene. *Mammalian genome,* 17**,** 1025-1032.

CHIZHIKOV, V. V., LINDGREN, A. G., MISHIMA, Y., ROBERTS, R. W., ALDINGER, K. A.,

MIESEGAES, G. R., CURRLE, D. S., MONUKI, E. S. & MILLEN, K. J. 2010. Lmx1a

regulates fates and location of cells originating from the cerebellar rhombic lip and telencephalic

cortical hem. *Proceedings of the National Academy of Sciences,* 107**,** 10725-10730.

DECHANT, G. & BARDE, Y.-A. 2002. The neurotrophin receptor p75NTR: novel functions and

implications for diseases of the nervous system. *Nature neuroscience,* 5**,** 1131-1136.

DJELLOUL, M., HOLMQVIST, S., BOZA-SERRANO, A., AZEVEDO, C., YEUNG, M. S.,

GOLDWURM, S., FRISÉN, J., DEIERBORG, T. & ROYBON, L. 2015. Alpha-synuclein

expression in the oligodendrocyte lineage: an in vitro and in vivo study using rodent and human

models. *Stem cell reports,* 5**,** 174-184.

EASTER, S. S., JR., ROSS, L. S. & FRANKFURTER, A. 1993. Initial tract formation in the mouse

brain. *J Neurosci,* 13**,** 285-99.

FINK, A. J., ENGLUND, C., DAZA, R. A., PHAM, D., LAU, C., NIVISON, M., KOWALCZYK, T. &

HEVNER, R. F. 2006. Development of the deep cerebellar nuclei: transcription factors and cell

migration from the rhombic lip. *The Journal of neuroscience,* 26**,** 3066-3076.

FLORIO, M., LETO, K., MUZIO, L., TINTERRI, A., BADALONI, A., CROCI, L., ZORDAN, P.,

BARILI, V., ALBIERI, I., GUILLEMOT, F., ROSSI, F. & CONSALEZ, G. G. 2012.

Neurogenin 2 regulates progenitor cell-cycle progression and Purkinje cell dendritogenesis in

cerebellar development. *Development,* 139**,** 2308-20.

FOSSAT, N., LE GRENEUR, C., BEBY, F., VINCENT, S., GODEMENT, P., CHATELAIN, G. &

LAMONERIE, T. 2007. A new GFP-tagged line reveals unexpected Otx2 protein localization in

retinal photoreceptors. *BMC Dev Biol,* 7**,** 122.

FUJITA, H., KODAMA, T. & DU LAC, S. 2020. Modular output circuits of the fastigial nucleus for

diverse motor and nonmotor functions of the cerebellar vermis. *Elife,* 9.

GOLDOWITZ, D. & HAMRE, K. 1998. The cells and molecules that make a cerebellum. *Trends in*

neurosciences, 21**,** 375-382.

GREEN, M. J. & WINGATE, R. J. 2014. Developmental origins of diversity in cerebellar output nuclei.

Neural Dev, 9**,** 1.

HANKS, M., WURST, W., ANSON-CARTWRIGHT, L., AUERBACH, A. B. & JOYNER, A. L. 1995.

Rescue of the En-1 mutant phenotype by replacement of En-1 with En-2. *Science,* 269**,** 679-82.

HOKIN, M. R. & HOKIN, L. E. 1959. The synthesis of phosphatidic acid from diglyceride and

adenosine triphosphate in extracts of brain microsomes. *Journal of Biological Chemistry,* 234**,**

1381-1386.

JIANG, X., GWYE, Y., MCKEOWN, S. J., BRONNER-FRASER, M., LUTZKO, C. & LAWLOR, E.

R. 2008. Isolation and characterization of neural crest stem cells derived from in vitro–

differentiated human embryonic stem cells. *Stem cells and development,* 18**,** 1059-1071.

JOYNER, A. L., HERRUP, K., AUERBACH, B. A., DAVIS, C. A. & ROSSANT, J. 1991. Subtle

cerebellar phenotype in mice homozygous for a targeted deletion of the En-2 homeobox.

Science, 251**,** 1239-43.

JOYNER, A. L., LIU, A. & MILLET, S. 2000. Otx2, Gbx2 and Fgf8 interact to position and maintain a

mid–hindbrain organizer. *Current opinion in cell biology,* 12**,** 736-741.

KEBSCHULL, J. M., CASONI, F., CONSALEZ, G. G., GOLDOWITZ, D., HAWKES, R., RUIGROK, T. J. H., SCHILLING, K., WINGATE, R., WU, J., YEUNG, J. & UUSISAARI, M. Y. 2023. Cerebellum Lecture: the Cerebellar Nuclei-Core of the Cerebellum. *Cerebellum*.

KUROKAWA, D., KIYONARI, H., NAKAYAMA, R., KIMURA-YOSHIDA, C., MATSUO, I. & AIZAWA, S. 2004. Regulation of Otx2 expression and its functions in mouse forebrain and midbrain. *Development*, 131**,** 3319-3331.

LI, J. Y. & JOYNER, A. L. 2001. Otx2 and Gbx2 are required for refinement and not induction of mid-hindbrain gene expression. *Development*, 128**,** 4979-4991.

MACHOLD, R. & FISHELL, G. 2005. Math1 is expressed in temporally discrete pools of cerebellar rhombic-lip neural progenitors. *Neuron*, 48**,** 17-24.

MACHON, O., MASEK, J., MACHONOVA, O., KRAUSS, S. & KOZMIK, Z. 2015. Meis2 is essential for cranial and cardiac neural crest development. *BMC Dev Biol*, 15**,** 40.

MANTO, M., GRUOL, D. L., SCHMAHMANN, J. D., KOIBUCHI, N. & ROSSI, F. 2012. *Handbook of the cerebellum and cerebellar disorders*, New York, Springer.

MARTINEZ, S. & ALVARADO-MALLART, R. M. 1989. Rostral cerebellum originates from the caudal portion of the so-called 'mesencephalic' vesicle: a study using chick/quail chimeras. *European Journal of Neuroscience*, 1**,** 549-560.

MARTINEZ, S., ANDREU, A., MECKLENBURG, N. & ECHEVARRIA, D. 2013. Cellular and molecular basis of cerebellar development. *Front Neuroanat*, 7**,** 18.

MARZBAN, H., DEL BIGIO, M. R., ALIZADEH, J., GHAVAMI, S., ZACHARIAH, R. M. & RASTEGAR, M. 2015. Cellular commitment in the developing cerebellum. *Frontiers in cellular neuroscience*, 8**,** 450.

MARZBAN, H. & HAWKES, R. 2007. Fibroblast growth factor promotes the development of deep cerebellar nuclear neurons in dissociated mouse cerebellar cultures. *Brain research*, 1141**,** 25-36.

MARZBAN, H., KHANZADA, U., SHABIR, S., HAWKES, R., LANGNAESE, K., SMALLA, K. H.,

BOCKERS, T. M., GUNDELFINGER, E. D., GORDON-WEEKS, P. R. & BEESLEY, P. W.

2003. Expression of the immunoglobulin superfamily neuroplastin adhesion molecules in adult

and developing mouse cerebellum and their localisation to parasagittal stripes. *Journal of*

Comparative Neurology, 462**,** 286-301.

MARZBAN, H., KIM, C.-T., DOORN, D., CHUNG, S.-H. & HAWKES, R. 2008. A novel transverse

expression domain in the mouse cerebellum revealed by a neurofilament-associated antigen.

Neuroscience, 153**,** 1190-1201.

MARZBAN, H., RAHIMI-BALAEI, M. & HAWKES, R. 2019. Early trigeminal ganglion afferents

enter the cerebellum before the Purkinje cells are born and target the nuclear transitory zone.

Brain Struct Funct, 224**,** 2421-2436.

MIALE, I. L. & SIDMAN, R. L. 1961. An autoradiographic analysis of histogenesis in the mouse

cerebellum. *Exp Neurol,* 4**,** 277-96.

MILLEN, K. J., HUI, C. C. & JOYNER, A. L. 1995. A role for En-2 and other murine homologues of

Drosophila segment polarity genes in regulating positional information in the developing

cerebellum. *Development,* 121**,** 3935-45.

MILLET, S., CAMPBELL, K., EPSTEIN, D. J., LOSOS, K., HARRIS, E. & JOYNER, A. L. 1999. A

role for Gbx2 in repression of Otx2 and positioning the mid/hindbrain organizer. *Nature,* 401**,**

161-4.

MORALES, D. & HATTEN, M. E. 2006. Molecular markers of neuronal progenitors in the embryonic

cerebellar anlage. *J Neurosci,* 26**,** 12226-36.

NICHOLS, D. H. & BRUCE, L. L. 2006. Migratory routes and fates of cells transcribing the Wnt-1

gene in the murine hindbrain. *Developmental dynamics,* 235**,** 285-300.

PIERCE, E. T. 1975. Histogenesis of the deep cerebellar nuclei in the mouse: an autoradiographic study.

Brain Res, 95**,** 503-18.

RAHIMI-BALAEI, M., BERGEN, H., KONG, J. & MARZBAN, H. 2018. Neuronal Migration During

Development of the Cerebellum. *Front Cell Neurosci,* 12**,** 484.

RAHIMI-BALAEI, M., BUCHOK, M., VIHKO, P., PARKINSON, F. E. & MARZBAN, H. 2019a.

Loss of prostatic acid phosphatase and alpha-synuclein cause motor circuit degeneration without

altering cerebellar patterning. *PLoS One,* 14**,** e0222234.

RAHIMI-BALAEI, M., BUCHOK, M., VIHKO, P., PARKINSON, F. E. & MARZBAN, H. 2019b.

Loss of prostatic acid phosphatase and α-synuclein cause motor circuit degeneration without

altering cerebellar patterning. *PloS one,* 14.

RAHIMI BALAEI, M., JIAO, X., ASHTARI, N., AFSHARINEZHAD, P., GHAVAMI, S. &

MARZBAN, H. 2016. Cerebellar expression of the neurotrophin receptor p75 in naked-ataxia

mutant mouse. *International journal of molecular sciences,* 17**,** 115.

SHABANIPOUR, S., DALVAND, A., JIAO, X., RAHIMI BALAEI, M., CHUNG, S. H., KONG, J.,

DEL BIGIO, M. R. & MARZBAN, H. 2019. Primary Culture of Neurons Isolated from

Embryonic Mouse Cerebellum. *J Vis Exp*.

SIMEONE, A., ACAMPORA, D., GULISANO, M., STORNAIUOLO, A. & BONCINELLI, E. 1992.

Nested expression domains of four homeobox genes in developing rostral brain. *Nature,* 358**,**

687-90.

SOTELO, C. 2004. Cellular and genetic regulation of the development of the cerebellar system.

Progress in neurobiology, 72**,** 295-339.

STAINIER, D. & GILBERT, W. 1990. Pioneer neurons in the mouse trigeminal sensory system.

Proceedings of the National Academy of Sciences, 87**,** 923-927.

VOOGD, J. & GLICKSTEIN, M. 1998. The anatomy of the cerebellum. *Trends in cognitive sciences,* 2**,**

307-313.

WANG, V. Y., ROSE, M. F. & ZOGHBI, H. Y. 2005. Math1 expression redefines the rhombic lip

derivatives and reveals novel lineages within the brainstem and cerebellum. *Neuron,* 48**,** 31-43.

WANG, V. Y. & ZOGHBI, H. Y. 2001a. Genetic regulation of cerebellar development. *Nature Reviews*

Neuroscience, 2**,** 484.

WANG, V. Y. & ZOGHBI, H. Y. 2001b. Genetic regulation of cerebellar development. *Nat Rev*

Neurosci, 2**,** 484-91.

WIZEMAN, J. W., GUO, Q., WILION, E. M. & LI, J. Y. 2019. Specification of diverse cell types

during early neurogenesis of the mouse cerebellum. *eLife,* 8**,** e42388.

WU, J. P. H., YEUNG, J., RAHIMI-BALAEI, M., WU, S. R., ZOGHBI, H. & GOLDOWITZ, D. 2022.

The Transcription Factor Pou3f1 Sheds Light on the Development and Molecular Diversity of

Glutamatergic Cerebellar Nuclear Neurons in the Mouse. *Front Mol Neurosci,* 15**,** 921901.

WURST, W. & BALLY-CUIF, L. 2001. Neural plate patterning: upstream and downstream of the

isthmic organizer. *Nat Rev Neurosci,* 2**,** 99-108.

ZHONG, S.-C., LUO, X., CHEN, X.-S., CAI, Q.-Y., LIU, J., CHEN, X.-H. & YAO, Z.-X. 2010.

Expression and subcellular location of alpha-synuclein during mouse-embryonic development.

Cellular and molecular neurobiology, 30**,** 469-482.

ZORDAN, P., CROCI, L., HAWKES, R. & CONSALEZ, G. G. 2008. Comparative analysis of

proneural gene expression in the embryonic cerebellum. *Dev Dyn,* 237**,** 1726-35.

ZYLKA, M. J., SOWA, N. A., TAYLOR-BLAKE, B., TWOMEY, M. A., HERRALA, A., VOIKAR,

V. & VIHKO, P. 2008. Prostatic acid phosphatase is an ectonucleotidase and suppresses pain by

generating adenosine. *Neuron,* 60**,** 111-122.

581
582

Fig. 1. The presence of neurons in the cerebellar primordium at E9 and spatial expression of SNCA. The
cerebellar primordium immunostained with NAA and SNCA, shows that a distinct subset of neurons
located at the rostral end of the developing cerebellum.
A: Dorsal view of the schematic illustration of the cerebellar primordium, mesencephalon, isthmus, and
4th ventricle. The red line indicates the sagittal plane about which the sections shown in **B-D** were taken.
B-D: Sagittal section through the cerebellar primordium at early E9 immunoperoxidase-labeled with NAA
3A10 shows the presence of neurons in the cerebellar primordium that cross the isthmus (i) and continue
to the mesencephalon. **C**. A higher magnification of **B**. **D**. Differentiated neurons at E9.5 are visible; a
higher magnification is shown in the inset, **d**.
E: Sagittal section through the cerebellar primordium at late E9. Immunofluorescence labeling of SNCA
shows SNCA$^+$ (green) expressing CN neurons in the mesencephalon, at the isthmus (i) and in the rostral
part of the cerebellar primordium (cb). **F**. A higher magnification of **E**.
Abbreviations: 4thv, 4th ventricle; cb, cerebellum; c, caudal; d, dorsal; i, isthmus; m, mesencephalon; r,
rostral; v, ventral
Scale bar = 100 µm in B; 50 µm in C and E; 20 µm in D

Fig. 2. SNCA expression relative to Lmx1a in the early developing mouse cerebellum. The sagittal section
through the cerebellar primordium at E12.5 reveals double immunofluorescence labeling with SNCA and
LMX1A antibodies in both medial (panels A-D) and lateral (panels E-H) sections.
A–G: SNCA (green, **A** and **E**) and LMX1A (red, **B** and **F**) immunopositive cells are located at the NTZ.
SNCA$^+$ cells continue to the mesencephalon, and LMX1A$^+$ cells toward the rhombic lip. Merged images
show that the SNCA$^+$ cells form a population of CN neurons distinct from the rhombic lip-derived cells
(LMX1A$^+$) in the NTZ (**C** and **G**). **D** and **H** show a higher magnification of **C** and **G**, respectively.
Abbreviations: 4thV, 4th ventricle; cb, cerebellum; c, caudal; d, dorsal; i, isthmus; m, mesencephalon; rl,
rhombic lip; NTZ, nuclear transitory zone; r, rostral; v, ventral.
Scale bar = 100 µm.

Fig. 3. Expression of SNCA and OTX2 in the developing cerebellar primordium. Double
immunofluorescence labeling with SNCA and OTX2 antibodies in sagittal sections of the cerebellar
primordium at E12.5.
A–E: Double immunolabeling of SNCA (green) and OTX2 (red) of a sagittal section of the cerebellar
primordium at E12.5 **A**. High OTX2 immunoreactivity was detected in the mesencephalon. SNCA$^+$ cells
in the mesencephalon accompany OTX2$^+$ cells across the isthmus (i) and in the NTZ. **B**. higher
magnification of **A**. **C-E**. Immunolabeling with SNCA (green, **C**) and OTX2 (red, **D**) and merged (**E**)
shows co-expression in the NTZ.
F-G: Cerebellar cells at E12 and E14 were dissociated and immunolabeled for SNCA + OTX2, SNCA +
LMX1A, or each antibody individually (F). Quantitative analysis was performed by flow cytometry to

detect SNCA$^+$, OTX2$^+$, LMX1A$^+$, SNCA$^+$/ LMX1A$^+$ and SNCA$^+$/ OTX2$^+$ cells (**G**). No significant
differences were observed for the number of LMX1A$^+$ cells between E12 and E14. An increase in SNCA
and OTX2 positive cells was shown at E14 in comparison to E12 (*$P<0.05$). The number of cells positive
for SNCA/LMX1A or SNCA/OTX2 was very low at E12 and E14.
Abbreviations: 4thv, 4th ventricle; cb, cerebellum; c, caudal; d, dorsal; i, isthmus; m, mesencephalon; rl,
rhombic lip; NTZ, nuclear transitory zone; r, rostral; v, ventral.
Scale bar = 200 µm in A; 50 µm in B; 20 µm in C, D, and E.
629

Fig. 4. The spatial and temporal expression of OTX2 in the developing cerebellum of mice. Sagittal
sections through the cerebellar primordia at E12, 13, 14, and 15 were analyzed for peroxidase
immunoreactivity of OTX2.
A–F: Sagittal section through cerebellar primordium at E12 (**A**; higher magnification shown in **B**), E13
(**C**; higher magnification shown in **D**), E14 (**E**), and E15 (**F**). High OTX2 immunoreactivity at the
mesencephalon is evident, and a few OTX2+ cells cross the isthmus and position at the rostral part of the
cerebellar primordium at the NTZ. Examples of Otx2 expressing cells are indicated by arrowheads in
panels B, D, E, and F. In panels E and F, the isthmus and adjacent cerebellar area are showed in the same
orientation as panels A-D (I; isthmus). The orientation label (r-c and d-v; rostral caudal and dorsal ventral)
in panel A applies to all panels.
G: Comparison of the number of OTX2 positive cells in the cerebellar primordium from E12 to E15.
Results indicate a slight increase in the number of OTX2 positive cells over time. Significant differences
were observed between E12 and E14 (**$P<0.01$), E12 and E15 (***$P<0.001$), as well as E13 and E15
(#$P<0.05$). Data were analyzed by One-Way ANOVA followed by a Tukey's multiple comparison test.
Abbreviations: 4thv, 4th ventricle; cb, cerebellum; c, caudal; d, dorsal; i, isthmus; m, mesencephalon; rl,
rhombic lip; r, rostral; v, ventral.
Scale bar = 200 µm in A and C; 50 µm in F (applies to B, D, E, and F).
647

Fig. 5. Otx2 expression in developing cerebella of Otx2-GFP transgenic and control mice. Sagittal sections
through the cerebellar primordium of Otx2-GFP transgenic mice at E13 are shown in panels A-D, and
RNAscope fluorescence in situ hybridization (FISH) of Otx2 at E12 in wild-type mice in panels E-J.
A-D: DAPI labels the outline of the cerebellar primordium and mesencephalon (**A**, blue). GFP expression,
which is enhanced by immunofluorescence labeling with anti-GFP (**B**, green), and immunofluorescence
labeling for OTX2 (**C**, red), reveal co-labeled cells (**D**, merged) in the mesencephalon and NTZ at the
rostral end of the cerebellar primordium. Arrows indicate the isthmus.
E-G: Merged channels of the *in situ* hybridization of *Otx2* mRNA probe counterstained with DAPI at low
(**E**) and high (**F** and **G**) magnification captured by confocal microscopy.
H-J: The Otx2 mRNA signal was strong in the mesencephalon and extended as a tail through the isthmus
to the rostral cerebellar primordium in the NTZ. The isthmus is indicated by arrows in (**E, F, H, I**) and a
line in (**J**).
Abbreviations: cb, cerebellum; c, caudal; d, dorsal; m, mesencephalon; rl, rhombic lip; ntz, nuclear
transitory zone; r, rostral; v, ventral.

Scale bar = 200 µm in D (applies to A-D); 100 µm in H (applies to E and H); 20 µm in J (applies to F -
J).
664

Fig. 6. SNCA immunopositive cells express p75NTR (NGFR) in developing cerebellum in vivo and in
vitro.
A–D: Double immunofluorescence labeling with SNCA (**A**, green) and p75NTR (**B**, red) antibodies
reveals co-labeled cells (**C**, merged) in the NTZ; a higher magnification of the NTZ is shown in **D**.
d. Western blot analysis of SNCA and p75NTR expression during cerebellar development. Immunoblots
of total cerebellar lysates from embryos at E11, E13 and E15 indicate an increase in expression of SNCA
and p75NTR from E11 to E15. Equal protein loading was confirmed by β-actin expression.
E–H: Primary dissociated cultures of cerebellum obtained from an E10 mouse embryo (DIV 4), double
immunofluorescence stained for SNCA (**E**: green) and p75NTR (**F**: red) (merged image shown in **G**). **H**
is a higher magnification of **G**; punctuate cellular p75NTR immunoreactivity is marked by arrow heads.
Abbreviations: cb, cerebellum; c, caudal; d, dorsal; NTZ, nuclear transitory zone; r, rostral; v, ventral.
Scale bar = 50 µm in A–C and D–F; 20 µm in H; 10 µm in G.
677

Fig. 7. SNCA, OTX2, and MEIS2 expression in the developing cerebella of Atoh1 null and control mice.
Sagittal sections through the cerebellar primordia of Atoh1$^{+/+}$ and Atoh1$^{-/-}$ embryos at E12.5; double
immunofluorescence labeling with SNCA + OTX2 and SNCA + MEIS2 antibodies is demonstrated.
A–C: SNCA (green, **A**) and OTX2 (red, **B**) immunopositive cells are present in the mesencephalon and
NTZ of Atoh1 KO mouse (merged channels, **C**).
D-F: In E12.5 Atoh1$^{+/+}$ sagittal sections, SNCA$^+$ (green, **D**) cells are observed in the mesencephalon and
NTZ. MEIS2$^+$ (red, **E**) cells are present in the mesencephalon and NTZ in addition to another population
of MEIS2$^+$ cells in the dorsal region of the NTZ which extends to the rhombic lip. Merged image (**F**)
confirms the presence of two distinct sets of MEIS2$^+$ cells in the NTZ: rhombic lip-derived CN neurons
located in the caudodorsal (cd) region, which do not express SNCA (MEIS2$^+$/SNCA$^-$), and a subset of
MEIS2$^+$/SNCA$^+$ CN neurons situated in the rostroventral (rv) region of the NTZ.
G-I: Higher magnification of **D-F.**
J-L: In E12.5 Atoh1$^{-/-}$ sagittal sections, the expression of SNCA (green, **J**) shows no change as compared
to Atoh1$^{+/+}$, whereas Meis2 abundance (red, **K**) exhibited a different pattern. Thus, MEIS2$^+$/SNCA$^+$ cells
were still present in the rostroventral region of the NTZ. However, the subpopulation of MEIS2$^+$/SNCA$^-$
rhombic lip-derived CN neurons were absent in the caudodorsal region of the NTZ.
M-O: Higher magnification of **J-I**.
Abbreviations: 4thV, 4th ventricle; cb, cerebellum; cd, caudodorsal; i, isthmus; m, mesencephalon; rl,
rhombic lip; NTZ, nuclear transitory zone; rv, rostroventral.
Scale bar = 100 µm
698

699 **Fig. 8.** Labeling of mesencephalic cells during early development in mice using embryonic culture. Fast
700 DiI was applied to an embryo at E9 and kept for 4 days (DIV 4, panels A-C) or removed after 24 hours
701 (DIV 1, panels D-E).

A, a: Fast DiI was inserted in the mesencephalon at E9 (DIV 0), arrowhead shows insertion location of DiI crystal in the mesencephalon. The arrow indicates the isthmus. Inset "a" shows the whole embryo and indicates the site of DiI insertion.

B-C: 4 days post DiI insertion, DiI positive cells directed both rostrally to the mesencephalon (**B**) and caudally to the rostral cerebellar primordium (**C**).

D, d: Fast DiI was inserted in the mesencephalon at E9 (DIV 0) (indicated by arrowhead) and removed at DIV1, arrow shows the isthmus. Inset "d" shows the whole embryo and indicates the site of DiI insertion.

E, e: DiI positive cells present in cerebellar primordium at DIV 6 after removal of Fast DiI at DIV1.

Abbreviations: cb, cerebellum; c, caudal; d, dorsal; m, mesencephalon; r, rostral; v, ventral.

Scale bar = 500 µm in a, d; 200 µm in A, D; 100 µm in B-C, E.

Fig. 9. A schematic illustration of the developing cerebellum at ~E12, created by BioRender.com. The expression patterns of SNCA, MEIS2, and OTX2 within the cerebellar primordium indicate a subset of CN neurons with origins distinct from the rhombic lip (rl).

Abbreviations: c, caudal; d, dorsal; m, mesencephalon; ntz, nuclear transitory zone; r, rostral; rl, rhombic lip; v, ventral.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. OTX2 expression persists in Snca null mice. OTX2 immunoperoxidase staining in the cerebellar primordium of Pap$^{-/-}$; Snca$^{-/-}$ mice at E12. Results show OTX2$^+$ cells are present in the ntz and terminate in the rostral cerebellar primordium.

Abbreviations: cb, cerebellum; I, isthmus; ntz, nuclear transitory zone

Scale bar = 50 µm

Supplementary Figure 2. Fast DiI was applied to an embryo at E9 and kept for 4 days in vitro (DIV 4), revealing labeled cells within the cerebellar primordium.

A-C: Low and high magnifications show only a few positive cells in the rostral cerebellar primordium in the NTZ at the level of the medial cerebellar section.

D-F: Low and high magnifications indicate the presence of just a few positive cells in the rostral cerebellar primordium in the NTZ at the level of the lateral cerebellar section.

Abbreviations: cb, cerebellum; c, caudal; d, dorsal; m, mesencephalon; NTZ, nuclear transitory zone; r, rostral; v, ventral.

Scale bar = 100 µm in A applies to D; 50 µm in B applies to E; 20 µm in C applies to I as well.

Supplementary Figure 3. Fast DiI was applied to the embryo at E9 and removed after 24 hours, resulting in the visualization of labeled cells within the cerebellar primordium (higher magnification in Fig. 8 E,e).

A: DiI positive cells present in cerebellar primordium at DIV 6 after removal of Fast DiI at DIV1 (indicated by arrowhead), arrow points to the isthmus.

741   **B:** A higher magnification from the caudal to mesencephalon and rostral rhombencephalon shows DiI
742   positive cells in the cerebellar primordium, arrow points to the isthmus.
743   Abbreviations: cb, cerebellum; c, caudal; d, dorsal; m, mesencephalon; r, rostral; v, ventral.
744   Scale bar = 500 µm in B; 100 µm in C.
745

746   **Supplementary Figure 4.** *In situ* hybridization demonstrates expression pattern of Otx2, Snca, C-Ret,
747   and Tlx3. These images show distribution of Otx2, Snca, C-Ret and Tlx3 in the cerebellar primordium at
748   E11.5, E13.5 and E15.5; all markers are present in the rostroventral region of the NTZ. All images show
749   cerebellar primordia. Image credit: Allen Institute. © 2008 Allen Institute for Brain Science. Allen
750   Developing Mouse Brain Atlas. Available online at: https://developingmouse.brain-map.org/.
751
752

A DiI at E9 (DIV0)→(DIV4)

D DiI at E9 (DIV0)→removed (DIV1)→(DIV6)