Nearly Neutral Evolution Across the *Drosophila melanogaster* Genome

David Castellano[1]

Jennifer James[2]

Adam Eyre-Walker[2,3]

1. Bioinformatics Research Centre

Aarhus University

C.F. Møllers Allé 8

DK-8000 Aarhus C

Denmark

2. School of Life Sciences

University of Sussex

Brighton

BN1 9QG

United Kingdom

3. Correspondence: a.c.eyre-walker@sussex.ac.uk

Abstract

Under the nearly neutral theory of molecular evolution the proportion of effectively neutral mutations is expected to depend upon the effective population size (N_e). Here we investigate whether this is the case across the genome of *Drosophila melanogaster* using polymorphism data from 128 North American lines. We show that the ratio of the number of non-synonymous and synonymous polymorphisms is negatively correlated to the number of synonymous polymorphisms, even when the non-independence is accounted for. The relationship is such that the proportion of effectively neutral non-synonymous mutations increases by ~45% as N_e is halved. However, we also show that this relationship is steeper than expected from estimates of the distribution of fitness effects from the site frequency spectrum. We investigate

a number of potential explanations for this and show, using simulation, that this is consistent with a model of genetic hitch-hiking: genetic hitch-hiking depresses diversity at neutral and weakly selected sites, but has little effect on the diversity of strongly selected sites. Thus, the rate at which the proportion of effectively neutral mutations changes with increasing N_e also depends on the factors that cause the variation in N_e.

Introduction

In 2015, Tomoko Ohta was awarded the Crafoord prize, along with Richard Lewontin, for her contributions to evolutionary biology and population genetics. Among her many contributions, she is best known for her nearly neutral theory of molecular evolution in which she proposed that there is a class of mutations that are substantially influenced by both random genetic drift and natural selection (Ohta and Kimura 1971; Ohta 1972b, 1973, 1977, 1992). In species with large effective population sizes selection is effective and in species with small effective population sizes random drift dominates the fate of this class of mutations. As a consequence, the proportion of mutations that are effectively neutral varies with effective population size. The theory has had a profound influence on how we think about molecular evolution and processes such as the rate of evolution (Lanfear, et al. 2014), the evolution of the mutation rate (Lynch 2010; Lynch, et al. 2016) and genome size (Lynch and Conery 2003).

There is now considerable evidence that there is a class of mutations that are nearly neutral, the vast majority of which are thought to be slightly deleterious. First, putatively functional mutations segregate at lower frequencies in the population than putatively neutral mutations; for example, in many species non-synonymous mutations segregate at lower frequencies than synonymous mutations (Akashi 1999; Cargill, et al. 1999; Hughes 2005), and it has also been shown that mutations in conserved non-coding sequences segregate at lower frequencies than those in neighbouring sequences (Andolfatto 2005; Drake, et al. 2006; Asthana, et al. 2007). Second, the rate of non-synonymous to synonymous substitution is higher along lineages leading to species with

small effective population size (note that in most cases the effective population is not directly measured but a surrogate measure, such as body size, is used instead) (Ohta 1993, 1995; Moran 1996; Woolfit and Bromham 2003, 2005; Popadin, et al. 2007). In fact, Ohta produced the very first evidence for her theory by noting that the rate of divergence in protein coding sequences relative to overall rate of DNA divergence, as measured by DNA-DNA hybridisation, was positively correlated to generation time, a proxy for population size (Ohta 1972a). Third, the ratio of non-synonymous or synonymous polymorphism appears to be greater in those species with smaller effective population size, as measured by synonymous diversity (Piganeau and Eyre-Walker 2009; Elyashiv, et al. 2010; Galtier 2016; James, et al. 2017).

Ohta used her theory to explain how the rate of protein evolution could be relatively constant even if the mutation rate varied between taxa; she reasoned that species with large population sizes might have high mutation rates per year, but this would be offset by having a small proportion of effectively neutral mutations (Ohta and Kimura 1971; Ohta 1972b, 1992). However, despite all the qualitative evidence that nearly neutral mutations exist, there are few robust estimates how rapidly the proportion of effectively neutral mutations changes with effective population size.

We can potentially investigate how the proportion of effectively neutral mutations changes as a function of N_e by considering how the number of non-synonymous (p_N) relative to synonymous polymorphisms (p_S) changes as a function of the number of synonymous polymorphisms (p_S). The logic is as follows. Let us assume that synonymous mutations are neutral so that $p_S = kN_eu$ where k is Watterson's coefficient (=Sum $1/i$), N_e is the effective population size and u is the mutation rate. Let us further assume that non-synonymous mutations are either effectively neutral or deleterious so that $p_N = kN_euf$, where f is the proportion of effectively neutral mutations and k is Watterson's constant (Watterson 1975). We can therefore investigate how rapidly the proportion of effectively neutral mutations changes as N_e increases by considering the correlation between p_N/p_S and p_S. However, there are two

issues. First, the two variables are not independent. We circumvent this problem by dividing p_S into two halves using a hypergeometric distribution (Piganeau and Eyre-Walker 2009; James, et al. 2017); this means that p_{S1} and p_{S2} have uncorrelated sampling errors. We then consider the correlation between p_N/p_{S1} and p_{S2}. Second, the theory outlined above assumes that the population is at equilibrium. Non-equilibrium processes can potentially mimic the effects of variation in N_e. Brandvain and Wright (Brandvain and Wright 2016) give an instructive example. Imagine a population, initially with no diversity. The population will accumulate genetic diversity through mutation, but the sites subject to deleterious mutation will approach their equilibrium frequency faster than neutral mutations because the equilibrium frequency is lower (Gordo and Dionisio 2005; Do, et al. 2015; Brandvain and Wright 2016). Hence, if we were to sample a locus through time then we would observe a negative relationship between p_N/p_S and p_S. However, N_e has not changed since the population was founded.

The precise relationship between p_N/p_S and p_S is potentially predictable. Ohta showed that the rate of evolution is proportional to $1/N_e$ over a substantial range of N_e values if the DFE is an exponential distribution (Ohta 1977). Kimura subsequently showed that under a gamma distribution with a shape parameter of 1/2 the rate was proportional to $1/N_e^{0.5}$, which suggests, given that the exponential distribution is a gamma distribution with shape parameter of one, that the rate of evolution under a gamma distribution is proportional to $1/N_e^{\beta}$, where β is the shape parameter. This general result was confirmed by Welch et al.(Welch, et al. 2008). Welch et al. also demonstrated that the number of polymorphisms and the nucleotide diversity are proportional to $N_e^{1-\beta}$; hence the nucleotide diversity at selected sites divided by the nucleotide diversity at neutral sites should be proportional to $N_e^{-\beta}$ and hence there should be a linear relationship, with a slope of $-\beta$ between the log of p_N/p_S and the log of p_S if the DFE is a gamma distribution.

Several groups have shown that $\log(p_N/p_S)$ is negatively correlated to $\log(p_S)$ across animal and plant species for nuclear DNA (Galtier 2016; Chen, et al.

2017) and across animal species for mtDNA (James, et al. 2017). However, no investigation has been made as to whether this relationship also holds within a species across a genome. This might be expected since the effective population size also varies across the genome in many species due to the effects of recombination, genetic hitch-hiking and background selection (Charlesworth 2009; Ellegren and Galtier 2016); regions of the genome with low rates of recombination, high mutations rates or gene density, both, have lower effective population sizes than regions with high rates of recombination and a low density of selected mutations.

Results

To investigate the relationship between the proportion of effectively neutral mutations and the effective population size, we compiled polymorphism data from 7918 autosomal genes from *D. melanogaster*. We also use a smaller data set of 4676 autosomal genes with short introns (< 66 bp, refs) as an alternative neutral reference. Because we are interested in the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ and p_N and p_S can be zero for individual genes, we grouped genes into 10 groups of 791 genes (10 groups of 467 genes when using introns) and considered the correlation between $\log(\text{sum}(p_N)/\text{sum}(p_{S1}))$ and $\log(\text{sum}(p_{S2}))$ (similar results were obtained with other groupings – see below).

As expected, given previous work, there is a positive correlation between p_S and the rate of recombination (RR) (Spearman's correlation $r_s = 0.99$, p<0.001) (Figure 1)(Begun and Aquadro 1992; Presgraves 2005; Langley, et al. 2012; Mackay, et al. 2012; Campos, et al. 2014). This positive correlation might be due to variation in the mutation rate or variation in N_e across the genome. Since, there is no correlation between the synonymous site divergence (d_S) between *D. melanogaster* and *D. yakuba* and RR ($r_s = -0.10$, p = 0.78), we conclude as others have done, that recombination is not mutagenic (Begun and Aquadro 1992; Betancourt and Presgraves 2002; Mackay, et al. 2012; Campos, et al. 2014), and that there is variation in N_e due to either genetic hitch-hiking or background selection. The relationship

between p_S and RR is non-linear with a tendency for the number of synonymous polymorphisms to be substantially depressed in regions of low recombination (Figure 1).

Figure 1. The correlation between the number of synonymous polymorphisms per site and the rate of recombination. Each point represents the average of 791 genes grouped according to their mutation rate.

We find that $\log(p_N/p_S)$ is negatively correlated to $\log(p_S)$ (slope = -0.53 (95% CIs = -0.48, -0.57), $p < 0.001$) (figure 2A). The relationship is almost perfectly linear, consistent with the underlying distribution of fitness effects being a gamma distribution. The slope is such that halving the effective population size increases the proportion of effectively neutral mutations by ~45%. The results are unaffected by using different grouping schemes (20 groups slope = -0.54 (-0.60, -0.48), 50 groups slope = -0.53 (-0.59, -0.48)).

We have assumed that synonymous mutations are neutral, whereas we know that there is selection on synonymous codon use in Drosophila species (Shields, et al. 1988; Akashi 1995). To investigate whether this affects our

results we repeated the analysis using data from short introns as our neutral sites. The relationship between $\log(p_N/p_{I1})$ and $\log(p_{I2})$ is highly significant ($p < 0.001$) (figure 2B), almost perfectly linear and has a slope which is very similar to that found using synonymous sites (slope = -0.55 (-0.47, -0.65)).

In many species, we do not have a recombination rate map, so we also investigated the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ grouping genes by their p_S or p_I values rather than the RR. To do this we followed the method of James et al. (James, et al. 2017) in which p_S (p_I) is split three-ways using a hypergeomtric distribution and p_{S1} (p_{I1}) is used to rank and group genes, p_{S2} (p_{I2}) is used to measure p_S (p_I) and p_{S3} (p_{I3}) is used to calculate p_N/p_S (p_N/p_I). The slopes using this method are -0.52 (-0.48, -0.56) and -0.64 (-0.53, -0.77) for synonymous and intron sites respectively (Figure 2C, 2D), confirming that this is a satisfactory method for investigating the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ when recombination rate data are not available and there is sufficient polymorphism data although the results are subject to higher levels of sampling error.

Figure 2. The relationship between p_N/p_S and p_S or p_i grouping genes by either recombination rate or p_S or p_i. A) p_s grouping genes by recombination rate, B) p_i grouping genes by recombination rate, C) p_s grouping genes by p_s and D) p_i grouping genes by p_i.

If the DFE is gamma distribution then we expect the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ to be equal to the negative of the shape parameter of the DFE (Welch, et al. 2008). To investigate whether this is the case we estimated the shape parameter of the DFE using the site frequency spectrum (SFS) at non-synonymous and synonymous sites using the method of Keightley and Eyre-Walker (Keightley and Eyre-Walker 2007). We find that the shape parameter is significantly smaller (shape = 0.33 (0.30, 0.38)) than the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ ($p < 0.001$ bootstrapping the data by gene). This discrepancy has been noted before in animal mtDNA (James, et al. 2017) and although not explicitly mentioned, is evident in animal and plant nuclear DNA (Chen, et al. 2017).

There are a number of potential explanations for why the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ is greater than the shape parameter of the DFE estimated using the SFS. First, there might be a negative relationship between N_e and the mutation rate. This seems unlikely within the Drosophila genome, given that there is no correlation between d_S, a measure of the mutation rate, and the rate of recombination (see above). However, to investigate further, we split the synonymous (and intron divergence) into two halves using the hypergeometric distribution then used d_{S1} (d_{I1}) to calculate the average N_e for a group of genes (grouped by RR) as p_S/d_{S1} (p_I/d_{I1}) and used the other d_S (d_I) value as our estimate of the mutation rate. For synonymous sites, we find no correlation between our estimate of N_e and d_S (slope = -0.017, $p = 0.54$) but for introns we observe a positive and significant relation (slope = 0.10, $p < 0.001$) (Supplementary Figures 1A and 1B); this latter correlation is consistent with the fact that d_I is positively correlated to RR at the gene level ($r_s = 0.09$, $p < 0.001$). These patterns are consistent with there being a positive correlation between RR and the mutation rate, a positive relationship which is offset at synonymous sites by greater efficiency of selection against deleterious mutations in highly recombining areas, leading to a lack of a correlation between synonymous divergence and RR. It therefore seems likely that there is a positive correlation between the mutation rate and N_e in *Drosophila* and hence no evidence that they are negatively correlated. Similar results are obtained for

the relationship between $\log(p_N/p_S)$ versus $\log(p_S/d_S)$ and $\log(p_N/p_i)$ versus $\log(p_S/d_i)$ grouping by RR or neutral polymorphism levels (results not shown).

Second, the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ might be steeper than the estimate of the shape parameter of the DFE because the DFE is affected by N_e; the shape parameter or the mean selection coefficient of new deleterious mutations might increase with increasing N_e. To investigate this, we estimated the DFE using the SFS for each group of genes separately, grouping genes by the RR. The method estimates the shape parameter and the mean value of $N_e s$ for the DFE. Since we expect mean $N_e s$ to increase as N_e increases we divided our estimate of mean $N_e s$ by p_S to yield an estimate of mean strength of selection (equivalent results are obtained if $N_e s$ is divided by p_S/d_S). There is a marginally significant positive correlation between p_S and the shape parameter (r = 0.72, p = 0.020) (Figure 3a), however the slope of this relationship is very shallow, and the shape parameter never approaches the (negative) of the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$. The mean strength of selection is uncorrelated to recombination rate (r = 0.51, p = 0.13)(Figure 3b), but this is not a powerful analysis since the estimate of the mean value of $N_e s$ is subject to great uncertainty, even in a large sample, such as the one we have used.

Third, the difference between the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ and the shape parameter of the DFE estimated from the SFS might simply be due to the fact that the DFE is not well described by the gamma distribution and hence there is no expectation that the two quantities should be similar. However, the linearity between $\log(p_N/p_S)$ and $\log(p_S)$ suggests that the gamma distribution is a good description of the DFE, at least for mutations that are relatively weakly selected.

Figure 3. The relationship between the A) shape parameter and the B) estimated mean strength of selection and the rate of recombination.

Fourth, linked selection might affect the effective population size of neutral and deleterious mutations to different extents. We have previously shown, using simulation, that background selection does not affect the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ (James, et al. 2017), however hitch-hiking may affect the slope. It has been demonstrated, using simulation, that deleterious genetic diversity recovers more rapidly after a bottleneck than neutral genetic diversity (Gordo and Dionisio 2005; Brandvain and Wright 2016) and this is also evident in HIV data after a hard-selective sweep (Pennings, et al. 2014).

To investigate whether hitch-hiking is responsible for the increase in the slope, we ran two sets of simulations. In the first, we simulated a non-recombining locus composed of neutral sites and blocks of sites subject to varying levels of natural selection (i.e. the locus is composed equal number of sites subject Ns = -1, -2, -4…etc). This locus was also subject to adaptive evolution, the rate of which was varied between simulations. We find, as expected that nucleotide diversity decreases at the locus as we increase the rate of adaptive for neutral and weakly selected sites (Figure 4a). However, the effect is attenuated for mildly deleterious mutations and there is no effect on diversity for strongly selected deleterious mutations, under the conditions of our simulation.

This effect could potentially explain why the slope of $\log(p_N/p_S)$ and $\log(p_S)$ is steeper than expected from the estimate of the shape parameter of the DFE; regions with high levels of hitch-hiking will have low diversity, but the diversity of non-synonymous sites will be depressed less than that of synonymous sites; hence the slope between $\log(p_N/p_S)$ and $\log(p_S)$ will be steeper. To investigate this, we ran a second set of simulations in which a locus composed of neutral sites and deleterious mutations drawn from a gamma distribution, were linked to a locus undergoing adaptive evolution. For each rate of adaptive evolution, we averaged the number of neutral and selected polymorphisms per site across multiple sampling points and runs. We find the slope of $\log(p_N/p_S)$ and $\log(p_S)$ is substantially and significantly steeper (slope

= -0.67 (0.029)) than 0.4, the shape parameter of the gamma distribution used to generate the deleterious mutations (Figure 4B).

A)

B)

Figure 4. The effects of hitch-hiking. A) The diversity at sites subject to different levels of negative selection, but all part of the same non-recombining locus, plotted against the frequency of advantageous mutation in the locus. B) p_N/p_S plotted against p_S both on log-scales, for simulations in which

deleterious mutations are drawn from a gamma distribution. Each point represents the mean from a simulation run at a certain rate of beneficial mutation. The slope of the line, -0.67, is significantly steeper than the shape parameter of the gamma distribution used to simulate the data, which was 0.4.

Discussion

We have shown that the proportion of mutations that are effectively neutral, as measured by the ratio of the number of non-synonymous and synonymous polymorphisms, decreases as the effective population size, as measured by the number of synonymous polymorphisms, increases across the *D. melanogaster* genome. On a log-log scale the relationship is linear and the slope of the relationship is such that halving the effective population size increases the proportion of effectively neutral mutations by ~45%. The slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ is similar to that found across plant and animal species for nuclear DNA (Galtier 2016; Chen, et al. 2017) and for animal mtDNA (James, et al. 2017).

We find, as others have (Chen, et al. 2017; James, et al. 2017), that the slope of the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ is steeper than we expect given an estimate of the DFE from the site frequency spectrum. We find no evidence that this is due to a negative correlation between N_e and the mutation rate across the genome or a change in the shape parameter or mean of the DFE. Instead we find evidence that genetic hitch-hiking affects the rate at which the proportion of effectively neutral mutations declines with N_e. We demonstrate, by simulation that hitch-hiking depresses the level of genetic diversity at neutral and slightly deleterious sites, but has little or no effect on the diversity of strongly deleterious mutations. As a consequence, regions of the genome with high rates of hitch-hiking have low neutral diversity, but deleterious genetic diversity is not as affected. This effect is likely due to the fact that genetic diversity recovers more rapidly at negatively selected sites than neutral sites after a bottleneck or hitch-hiking event (Gordo and Dionisio 2005; Pennings, et al. 2014; Brandvain and Wright 2016).

The correlation between $\log(p_N/p_S)$ and $\log(p_S)$ is usually interpreted in terms of the nearly neutral theory, as being due to variation in the effective population size and the effect this variation has on the efficiency of selection. However, as we have shown, the slope of this relationship is steeper than expected and this is consistent with a hitch-hiking model in which the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ could emerge through non-equilibrium dynamics. Testing whether the relationship between $\log(p_N/p_S)$ and $\log(p_S)$ can be explained entirely in terms of non-equilibrium dynamics will require knowledge of the rate and strength of selection acting upon advantageous mutations and extensive simulation.

Our results suggest that the proportion of effectively neutral mutations varies across the *Drosophila* genome, and declines as a function of synonymous diversity; this is likely to be due to two factors; variation in the effective population size and non-equilibrium dynamics. Interestingly, we show that genetic hitch-hiking has little or no effect on strongly deleterious genetic variation.

Materials and methods

Datasets

This study was carried out on the four large autosomes (2L, 2R, 3L and 3R) of *D. melanogaster,* using release 5 of the Berkeley Drosophila Genome Project (BDGP 5, http://www.fruitfly.org/sequence/release5genomic.shtml) as the reference genome.

The population genomic data came from Raleigh, North Carolina. The details of their provenance and breeding are in Mackay et al. (Mackay, et al. 2012), Freeze 1.0 *Drosophila melanogaster* Genetic Reference Panel (DGRP) project. Sites with residual heterozygosity and low-quality values were excluded from the analyses. The method to estimate the distribution of fitness effects of new mutations (DFE) requires all sites to have been sampled in the same number of chromosomes and since some sites were not successfully

sampled in all samples we reduced the original data set to 128 chromosomes by randomly sampling the polymorphisms at each site without replacement. To estimate divergence out to *D. yakuba* we randomly sampled one *D. melanogaster* chromosome per site.

Coding exon and short intron (≤65 bp) annotations from *D. melanogaster* were retrieved from FlyBase (release 5.50, http://flybase.org/, last accessed March 2013). Genes 1:1 orthologs across *D. yakuba – D. melanogaster* were obtained from FlyBase (http://flybase.org/). We obtained a multiple genome alignment between the DGRP isogenic lines (Mackay, et al. 2012) and the *D. yakuba* genome (Drosophila 12 Genomes, et al. 2007) using the BDGP 5 coordinates. This alignment is publicly available at http://popdrowser.uab.cat/ (Ramia, et al. 2012). For each gene, we took all non-overlapping coding exons, independently of their inclusion levels. When two exons overlapped, the largest was chosen for subsequent analyses. Only exons without frameshifts, gaps or early stop codons were retained. In this way, we tried to avoid potential alignment errors which would inflate our mutation rate estimates. Our final data set fulfilling all these criteria contains 7,918 coding genes.

Exonic sequences were trimmed in order to contain only full codons. We define our sites "physically", so we estimated the rates of substitution at sites of different degeneracy separately. Only zero-fold and 4-fold degenerate sites in exon core codons (as described by Warnecke and Hurst (Warnecke and Hurst 2007)) were used. To estimate the rate of synonymous substitutions, we restricted our analysis to those triplets coding the same amino acid in the two species (*D. melanogaster – D. yakuba*). In restricting our analysis to codons not exhibiting nonsynonymous differences we assume that the codon has undergone no amino acid substitution — this avoids having to compute the different pathways between two codons, which differ by more than one change and it is a reasonable assumption given the level of amino acid divergence. For 4-fold degenerate sites we used the method of Tamura (Tamura 1992) to correct for multiple hits; this method allows for unequal GC content and ts/tv bias. We calculated the number of substitutions and the

folded site frequency spectrum (SFS) for 4-fold degenerate sites and zero-fold degenerate sites, using an *ad hoc* Perl Script.

We used positions 8–30 of introns ≤65 bp in length as an alternative neutral reference for some analyses (Halligan and Keightley 2006). For intron sequences, the invariant GT and AG dinucleotides at the 5' and 3' splice junctions, respectively, are excluded before calculating divergence. Only genes with at least one short intron and with less than 10% of gaps in the aligned sequences were kept. 4,676 orthologous genes pass the intron quality criteria in our final data set. We use an *ad hoc* Perl Script to estimate the number of short intron substitutions and to compute the folded SFS. Multiple hits were corrected using Jukes and Cantor method (Jukes and Cantor 1969).

Recombination rates were taken from Comeron et al. (Comeron, et al. 2012) (www.recobinome.com). They estimated the rate of crossovers in 100 kb non-overlapping windows in cM/Mb units. The rate of crossing-over for a gene is the rate in the 100kb that overlapped the mid-point of the gene.

Hypergeometric sampling
To investigate the correlation between p_N/p_S and p_S ranking genes by its recombination rate we split p_S into 2 independent variables (similar to the splitting done in Smith and Eyre-Walker (Smith and Eyre-Walker 2003); Piganeau and Eyre-Walker (Piganeau and Eyre-Walker 2009); Castellano et al. (Castellano, et al. 2016)). This was done by generating a random multivariate hypergeometric variable as follows:

$$P_{s1} = \text{Hypergeometric}(P_s, \tfrac{1}{2} \times L_s), \qquad (1)$$
$$P_{s2} = P_s - P_{s1}, \qquad\qquad\qquad (2)$$

where L_s is the total number of synonymous 4-fold degenerated sites and P_s is the total number of 4-fold segregating sites. We divide ΣP_{s1} and ΣP_{s2} by ½ of ΣL_s to get p_{S1} and p_{S2} at the bin level, respectively. We used p_{S1} to estimate p_N/p_{S1} and p_{S2} as a statistically independent estimate of the level of neutral diversity at the bin level.

To investigate the correlation between p_N/p_S and p_S ranking genes by its level of neutral diversity we split p_S into 3 independent variables following the scheme described by James et al. (James, et al. 2017). This was done by generating a random multivariate hypergeometric variable as follows:

P_{s1} = Hypergeometric(P_s, ⅓ x L_s),   (3)

$P_{s2-3} = P_s - P_{s1}$,   (4)

P_{s2} = Hypergeometric(P_{s2-3}, ⅓ x L_s),   (5)

$P_{s3} = P_s - P_{s1} - P_{s2}$ (6)

We divided ΣP_{s2} and ΣP_{s3} by ⅓ of ΣL_s to get p_{S2} and p_{S3} at the bin level, respectively. We used p_{S1} to rank genes and assign genes to bins, we then used p_{S2} to estimate p_N/p_{S1} and p_{S3} as a statistically independent estimate of the level of neutral diversity at the bin level.

Gene binning strategy

To estimate the ratio between non-synonymous (p_N) to synonymous polymorphisms (p_{S1}) (or short intron, p_{i1}) it is necessary to combine data from several genes because estimates from a single gene are noisy and often undefined because of the lack of segregating synonymous (or short intron) sites. We therefore group genes into bins according to their rate of recombination or neutral polymorphism. Three different grouping schemes are used: 10 bins, 20 bins and 50 bins.

To investigate the relationship between the log of p_N/p_{S1} (or p_N/p_{I1}) and the log of our proxy for the $N_e \sim p_{S3}/d_S$ (or p_{I3}/d_I) we used the substitution rate at exon core 4-fold degenerate sites (or at short intron sites) as a proxy for the neutral mutation rate.

Distribution of fitness effects Estimates using the SFS

DFE-alpha (Eyre-Walker and Keightley 2009) models the DFE at functional sites by assuming a gamma distribution, specified by the mean strength of selection, $\gamma = -N_e s$, and a shape parameter β, allowing the distribution to take

on a variety of shapes ranging from leptokurtic to platykurtic. We ran DFE-alpha assuming a single instantaneous change in population size from an ancestral size N_1 to a present-day size N_2 having occurred t_2 generations ago and the folded SFS mode since the results are more robust to both demographic changes and linked selection (Messer and Petrov 2013). Provided the SFS at both neutral and functional sites, DFE-alpha infers γ, β, N_2/N_1 and t_2. We ran DFE-alpha for each bootstrap replicate independently (see below) using the local version provided at: http://www.homepages.ed.ac.uk/pkeightl//software.

Confidence intervals and p-value estimates

To calculate the 95% confidence intervals (CIs) for the slope (b) of the relationship between $\log(p_N/p_S)$ vs $\log(p_S)$, we bootstrapped the data by gene 1,000 times. We split each bootstrap dataset into N bins (see Gene Binning Strategy above) and reestimated the slope for each replicate independently. To estimate the statistical significance of the difference between the slope (b) and the shape parameter (β) of the DFE using the SFS (Eyre-Walker and Keightley 2009) we estimated β and b for each bootstrap dataset. The p-value was the proportion of replicates in which $-b > \beta$.

Statistical analyses

All statistical analyses are performed using the R statistical software (Team 2013). Linear regressions are carried out calling the R function "lm" (from the R package "base"). We calculate Spearman rank correlations (ρ) using the R function "cor.test" (from the R package "base"). The random hypergeometric variable is obtained through the R function "rhyper" (from the R package "stats"). All the code used to perform the analyses are available upon request from DC.

Forward simulations

We used the forward simulation software SLiM, version 2.4.1 (Haller and Messer 2017) to simulate the behaviour of a single, 20 kb non-recombining locus with a mutation rate of 1×10^{-6}, evolving over time in a small population

of 100 individuals. Parameter values were chosen to generate substantial variation in the locus.

We ran two types of simulations. In the first, we investigated the impact of selective sweeps on different types of site, from strongly deleterious to neutral. Our locus was composed of 9 equally common categories of site at which mutations were subject to negative selection with $4Ns$ values of 0, -1, -2, -4, -8, -16, -32, -64, -128. We also allowed the locus to undergo adaptive evolution varying the total proportion of adaptive mutations from 0 to 0.5% We then investigated how the genetic diversity at each category of site changed as we increased the frequency of selective sweeps, such that adaptive mutations constituted between 0.05% and 0.5% of the total number of mutations that occurred in a simulation run. The adaptive mutations were subject to selection such that $4Ns = 400$. For each simulation run, after an initial burn-in of 1000 generations we took samples of all mutations segregating in the population every 500 generations, corresponding to $5N$ generations so that samples should be largely independent. The simulations, were run for 25500 generations in total. For each set of parameters, we ran our simulations 30 times. We then averaged values of diversity over simulation runs. To compare the impact of hitch-hiking on sites under different levels of selection we divide the diversity for a particular set of sites (e.g. $Ns = -4$) by the mean diversity from the simulation with no adaptive evolution.

In the second set of simulations, the locus was divided into two, with half of the mutations being neutral and the other half deleterious mutations which were sampled from a gamma distribution with a shape parameter 0.4 and a mean s of 10, which is roughly in line with what we observe in estimates of the DFE from real data. As before, we altered the frequency at which a locus experienced selective sweeps by changing the frequency with which adaptive mutations occurred. We used 12 different frequencies of adaptive mutations, such that adaptive mutations constituted between 0 and 1% of the total mutations that occurred in a simulation run. We used the same burn-in and sampling procedure as in our previous set of simulations, and again for each

set of parameters we ran our simulations 30 times. We then calculated average values of p_S and p_N/p_S across our simulation runs.

Acknowledgements

We are grateful to Meg Woolfit for early work on this project many years ago and for Matt Webster for reinvigorating our interest in this problem. Funding for this study was provided by the University of Sussex and NERC, grant number NE/L502042/1.

Literature cited

Akashi H. 1999. Inferring the fitness effects of DNA mutations from polymorphism and divergence data: statistical power to detect directional selection under stationarity and free recombination. Genetics 151:221-238.

Akashi H. 1995. Inferring weak selection from patterns of polymorphism and divergence at "silent" sites in Drosophila DNA. Genetics 139:1067-1076.

Andolfatto P. 2005. Adaptive evolution of non-coding DNA in Drosophila. Nature 437:1149-1152.

Asthana S, Noble WS, Kryukov G, Grant CE, Sunyaev S, Stamatoyannopoulos JA. 2007. Widely distributed noncoding purifying selection in the human genome. Proc Natl Acad Sci U S A 104:12410-12415.

Begun DJ, Aquadro CF. 1992. Levels of naturally occuring DNA polymorphism correlate with recombination rates in *D. melanogaster*. Nature 356:519-520.

Betancourt AJ, Presgraves DC. 2002. Linkage limits the power of natural selection in Drosophila. Proc. Natl. Acad. Sci. USA 99:13616-13620.

Brandvain Y, Wright SI. 2016. The Limits of Natural Selection in a Nonequilibrium World. Trends Genet 32:201-210.

Campos JL, Halligan DL, Haddrill PR, Charlesworth B. 2014. The relation between recombination rate and patterns of molecular evolution and variation in Drosophila melanogaster. Mol Biol Evol 31:1010-1028.

Cargill M, Altshuler D, Ireland J, Sklar P, Ardlie K, Patil N, Lane CR, Lim EP, Kalyanaraman N, Nemesh J, et al. 1999. Characterization of single-nucleotide polymorphisms in coding regions of human genes. Nature Genetics 22:231-238.

Castellano D, Coronado-Zamora M, Campos JL, Barbadilla A, Eyre-Walker A. 2016. Adaptive Evolution Is Substantially Impeded by Hill-Robertson Interference in Drosophila. Mol Biol Evol 33:442-455.

Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. Nat Rev Genet 10:195-205.

Chen J, Glemin S, Lascoux M. 2017. Genetic Diversity and the Efficacy of Purifying Selection across Plant and Animal Species. Mol Biol Evol 34:1417-1428.

Comeron JM, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in Drosophila melanogaster. PLoS Genet 8:e1002905.

Do R, Balick D, Li H, Adzhubei I, Sunyaev S, Reich D. 2015. No evidence that selection has been less effective at removing deleterious mutations in Europeans than in Africans. Nat Genet 47:126-131.

Drake JA, Bird C, Nemesh J, Thomas DJ, Newton-Cheh C, Reymond A, Excoffier L, Attar H, Antonarakis SE, Dermitzakis ET, et al. 2006. Conserved noncoding sequences are selectively constrained and not mutation cold spots. Nat Genet 38:223-227.

Drosophila 12 Genomes C, Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M, Gelbart W, et al. 2007. Evolution of genes and genomes on the Drosophila phylogeny. Nature 450:203-218.

Ellegren H, Galtier N. 2016. Determinants of genetic diversity. Nat Rev Genet 17:422-433.

Elyashiv E, Bullaughey K, Sattath S, Rinott Y, Przeworski M, Sella G. 2010. Shifts in the intensity of purifying selection: an analysis of genome-wide polymorphism data from two closely related yeast species. Genome Res 20:1558-1573.

Eyre-Walker A, Keightley PD. 2009. Estimating the rate of adaptive molecular evolution in the presence of slightly deleterious mutations and population size change. Mol Biol Evol 26:2097-2108.

Galtier N. 2016. Adaptive Protein Evolution in Animals and the Effective Population Size Hypothesis. PLoS Genet 12:e1005774.

Gordo I, Dionisio F. 2005. Nonequilibrium model for estimating parameters of deleterious mutations. Phys Rev E Stat Nonlin Soft Matter Phys 71:031907.

Haller BC, Messer PW. 2017. SLiM 2: Flexible, Interactive Forward Genetic Simulations. Mol Biol Evol 34:230-240.

Halligan DL, Keightley PD. 2006. Ubiquitous selective constraints in the Drosophila genome revealed by a genome-wide interspecies comparison. Genome Res 16:875-884.

Hughes AL. 2005. Evidence for abundant slightly deleterious polymorphisms in bacterial populations. Genetics 169:533-538.

James J, Castellano D, Eyre-Walker A. 2017. DNA sequence diversity and the efficiency of natural selection in animal mitochondrial DNA. Heredity (Edinb) 118:88-95.

Jukes TH, Cantor CR. 1969. Evolution of protein molecules. In: Munro NH, editor. Mammalian protein metabolism. New York: Academic Press. p. 121-123.

Keightley PD, Eyre-Walker A. 2007. Joint inference of the distribution of fitness effects of deleterious mutations and population demography based on nucleotide polymrophism frequencies. Genetics 177:1-11.

Lanfear R, Kokko H, Eyre-Walker A. 2014. Population size and the rate of evolution. Trends Ecol Evol 29:33-41.

Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Pool JE, Langley SA, Suarez C, Corbett-Detig RB, Kolaczkowski B, et al. 2012. Genomic Variation in Natural Populations of Drosophila melanogaster. Genetics 192:533-+.

Lynch M. 2010. Evolution of the mutation rate. Trends Genet 26:345-352.

Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection and the evolution of the mutation rate. Nat Rev Genet 17:704-714.

Lynch M, Conery JS. 2003. The origins of genome complexity. Science 302:1401-1404.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu DH, Casillas S, Han Y, Magwire MM, Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference Panel. Nature 482:173-178.

Messer PW, Petrov DA. 2013. Frequent adaptation and the McDonald-Kreitman test. Proc Natl Acad Sci U S A 110:8615-8620.

Moran NA. 1996. Accelerated evolution and Muller's rachet in endosymbiotic bacteria. Proc Natl Acad Sci U S A 93:2873-2878.

Ohta T. 1993. Amino acid substitution at the *Adh* locus of *Drosophila* is facilitated by small population size. Proc. Natl. Acad. Sci. USA 90:4548-4551.

Ohta T. 1972a. Evolutionary rate of cistrons and DNA divergence. J. Mol. Evol. 1:150-157.

Ohta T. 1977. Extension of the neutral mutation drift hypothesis. In: Kimura M, editor. Molecular Evolution and Polymorphism. Mishima: National Institute of Genetics. p. 148-167.

Ohta T. 1992. The nearly neutral theory of molecular evolution. Ann. Rev. Ecol. Syst. 23:263-286.

Ohta T. 1972b. Population size and rate of evolution. J. Mol. Evol. 1:305-314.

Ohta T. 1973. Slightly deleterious mutant substitutions in evolution. Nature 246:96-98.

Ohta T. 1995. Synonymous and nonsynonymous substitutions in mammalian genes and the nearly neutral theory. J. Mol. Evol. 40:56-63.

Ohta T, Kimura M. 1971. On the constancy of the evolutionary rate of cistrons. J. Mol. Evol. 1:18-25.

Pennings PS, Kryazhimskiy S, Wakeley J. 2014. Loss and recovery of genetic diversity in adapting populations of HIV. PLoS Genet 10:e1004000.

Piganeau G, Eyre-Walker A. 2009. Evidence for variation in the effective population size of animal mitochondrial DNA. PLoS ONE 4:e4396.

Popadin K, Polishchuk LV, Mamirova L, Knorre D, Gunbin K. 2007. Accumulation of slightly deleterious mutations in mitochondrial protein-coding genes of large versus small mammals. Proc Natl Acad Sci U S A 104:13390-13395.

Presgraves DC. 2005. Recombination enhances protein adaptation in Drosophila melanogaster. Current biology : CB 15:1651-1656.

Ramia M, Librado P, Casillas S, Rozas J, Barbadilla A. 2012. PopDrowser: the Population Drosophila Browser. Bioinformatics 28:595-596.

Shields DC, Sharp PM, Higgins DG, Wright F. 1988. "Silent" sites in *Drosophila* are not neutral: evidence of selection among synonymous codons. Mol. Biol. Evol. 5:704-716.

Smith NG, Eyre-Walker A. 2003. Partitioning the variation in mammalian substitution rates. Mol Biol Evol 20:10-17.

Tamura K. 1992. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases. Mol. Biol. Evol. 9:678-687.

Team RC. 2013. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

Warnecke T, Hurst LD. 2007. Evidence for a trade-off between translational efficiency and splicing regulation in determining synonymous codon usage in Drosophila melanogaster. Mol Biol Evol 24:2755-2762.

Watterson GA. 1975. On the number of segregating sites. Theor. Popul. Biol. 7:256-276.

Welch JJ, Eyre-Walker A, Waxman D. 2008. Divergence and Polymorphism Under the Nearly Neutral Theory of Molecular Evolution. J Mol Evol 67:418-426.

Woolfit M, Bromham L. 2003. Increased rates of sequence evolution in endosymbiotic bacteria and fungi with small effective population sizes. Mol Biol Evol 20:1545-1555.

Woolfit M, Bromham L. 2005. Population size and molecular evolution on islands. Proc Biol Sci 272:2277-2282.