The Hummingbird and the Hawk-moth: Species Distribution, Geographical Partitioning,

and Macrocompetition across the United States

Abdel Halloway[1], Christopher J. Whelan[1], and Joel S. Brown[2]

[1]Department of Biological Sciences, University of Illinois at Chicago

845 W. Taylor St. (M/C 066) Chicago, IL 60607

[2]Integrated Mathematical Oncology, Moffitt Cancer Center

SRB-4, 12902 USF Magnolia Drive Tampa, FL 33612

<u>**Corresponding Author**</u>

Abdel Halloway

Department of Biological Sciences, University of Illinois at Chicago

845 W. Taylor St. (M/C 066) Chicago, IL 60607

ahallo2@uic.edu

ABSTRACT

We introduce a new concept called macrocompetition – defined as the mutual

suppression of diversity/species richness of competing clades – and investigate evidence for its

existence. To this end, we analyzed the distribution of two convergent nectarivorous families,

hawk-moths and hummingbirds, over the continental United States to determine whether there is

geographic partitioning between the families and its potential causes. Using stepwise regression,

we tested for latitudinal and longitudinal biases in the species richness of both taxa and the

potential role of 10 environmental variables in their distribution pattern. Hawk-moth species

richness increases with longitude (eastward-bias) while that of hummingbirds declines

(westward-bias). Similar geographic patterns can be seen across Canada, Mexico and South

America. Hawk-moth species richness is positively correlated with higher overall temperatures

(especially summer minimums), atmospheric pressure, and summer precipitation; hummingbird

species richness is negatively correlated with atmospheric pressure and positively correlated with

winter daily maxima. The species richness patterns reflect each family's respective anatomical

differences and support the concept of macrocompetition between the two taxa. Hawk-moth

species richness was highest in states with low elevation, summer-time flowering, and warm

summer nights; hummingbird species richness is highest in the southwest with higher elevation,

greater cool season flowering and high daytime winter temperatures. Hawk-moths and

hummingbirds as distinct evolutionary technologies exhibit niche overlap and geographical

partitioning. These are two of three indicators suggested by Brown and Davidson for inter-

taxonomic competition. We intend the patterns revealed here to inspire further exploration into

competition and community structuring between hawk-moths and hummingbirds.

Keywords: Biogeography, Sphingidae, Trochilidae, Competition, Scale

INTRODUCTION

Competitive interactions help shape distribution (Hutchinson 1978), origination (Rosenzweig 1978; Hutchinson 1978; Schluter 2000; Ripa et al. 2009) and extinction of species (Gause 1934). Competition affects small-scale interactions among species yet also drives larger scale phenomena and lies at the core of processes like competitive speciation (Rosenzweig 1978) and incumbent replacement (Rosenzweig and McCord, 1991; Silvestro et al., 2015). It is most often studied at the local scale, either between individuals within a population mutually suppressing fitness, or between populations mutually suppressing each other's population size.

Competition may also operate at higher taxonomic levels. By occupying potential niches space of another, one taxonomic group may limit the species diversification or adaptive radiation of another. In this case, competition suppresses species richness rather than fitness or population size. We propose that competition thus acts on three levels:

- Microcompetition operates between individuals and suppresses access to resources
- Mesocompetition operates between populations and suppresses population sizes
- Macrocompetition operates between higher order taxa and suppresses species richness

These three forms of competition should occur on different temporal, spatial and taxonomic scales. Macrocompetition, which suppresses species diversity and the radiation of species within taxonomic groups, must occur over large temporal and spatial scales and at taxonomic levels higher than the species. Because of this link between spatial, temporal, and organizational scales, macrocompetition must be studied at its own appropriate scale (Jablonski, 2008). Just as population level mesocompetition is not studied by aggregating individual microcompetitive interactions, macrocompetition cannot be studied through the aggregation of mesocompetitive and microcompetitive interactions.

When studying macrocompetition, the adaptations specific to each clade are most

important. These clade-specific adaptations, which we refer to as evolutionary technologies, are

the tools that allow each clade to exploit environments and resources and form the basis of

macrocompetition. For macrocompetition, each taxa must exhibit one or more derived traits that

are shared among the members of the taxa but distinct from members of the competing taxa. The

evolutionary feasibility of these traits to the members of the taxa; and their unavailability to

members of other taxa defines the evolutionary technology (Vincent and Brown 2005). While

not originally intended as such, Families may represent a rough, but good first cutoff for

describing different evolutionary technologies (Pintor et al. 2011); and certainly members of

different Orders, Classes and Phyla represent different taxa for the purposes of

macrocompetition.

Mesocompetition between populations of different taxa has been well-documented.

Examples include tadpoles and aquatic insects (Morin et al., 1988) and insect larvae (Mokany

and Shine, 2003), granivorous rodents and ants (Brown and Davidson, 1977; Brown and

Davidson, 1979), granivorous birds and rodents (Brown et al., 1997), frugivorous birds and bats

(Palmeirim et al., 1989), insectivorous lizards and birds (Wright, 1979), and insectivorous birds

and ants (Haeming, 1994; Jedlicka et al. 2006). Mesocompetition may even exist between

species of separate phyla, such as the competition between scavenging vertebrates and microbes

for detritus (Janzen, 1977; Shivik 2006) or vertebrates and fungi for rotting fruit (Cipollini and

Stiles 1993; Cipollini and Levey 1997). Brown and Davidson (1977) identified three key

indicators to determine potential intertaxonomic mesocompetition: 1) shared extensive use of the

same particular resource, 2) reciprocal increases in population size when one competing species

is excluded, and 3) partitioning along a geographic or climatic gradient. We propose analogous

indicators as signals of macrocompetition: 1) shared extensive use of the same class of resources,

2) reciprocal increases in species richness via adaptive radiation when a competing taxon is

excluded, and 3) partitioning along geographical and climatic gradients across the shared taxa's

range.

Pollination systems provide ample opportunities for intertaxonomic competition. Both

Primack and Howe (1975) and Thomas et al. (1986) reported competition between

hummingbirds and butterflies, and Laverty and Plowright (1985) reported competition between

hummingbirds and bumblebees. Due to many convergent characteristics, competition between

hawk-moths (Sphingidae) and hummingbirds (Trochilidae) seem just as likely. Both taxa are

highly-specialized nectar feeders and pollinators as adults. They have similar sizes, hover when

feeding, and some species in each taxon possess tongues and other features that are often adapted

to a single species of plant (Johnsgard, 1997; Tuttle, 2007). Despite their remarkable similarity

and strong niche overlap, competition between these two Families has seldom been investigated.

Only Carpenter (1979) explored the possibility of direct competition between hawk-moths and

hummingbirds. Her study documented spatial and temporal partitioning between hawk-moths

and hummingbirds. Hawk-moths dominated *Ipomopsis* feeding sites through depletion of nectar

resources. Hummingbirds exhibited aggressive behaviour towards hawk-moths, suggesting

hummingbirds perceive hawk-moths as competitors.

Differences in morphology and physiology can shape broad scale biogeographical

patterns (Buckley et al., 2012). With this in mind, we compared species richness of

hummingbirds and hawk-moths at the continental scale of North America with the goal of

evaluating hallmarks of macrocompetition as modified from Brown and Davidson (1977). We

seek broad scale geographic and climatic correlations of diversity that might provide insights

into the patterns of diversity of hawk-moths and hummingbirds that may result from inter-taxon

competition. Do diversity patterns of these two families covary positively or negatively? As

nocturnal ectotherms, does hawk-moth diversity increase with summer rain and temperatures?

As diurnal endotherms, do hummingbirds gain a competitive edge with colder temperature and

cool season flowering? Do hawk-moths suffer more from low oxygen and elevation than do

hummingbirds? Ultimately, to what extent can large-scale biogeography provide insights and

clues into competition and geographical partitioning?

MATERIALS AND METHODS

Study Families

Hawk-moths (Order Lepidoptera, Family Sphingidae) and Hummingbirds (Order

Apodiformes, Family Trochilidae) are nectarivores exhibiting morphological convergence.

Worldwide, the approximately 953 species of hawk-moths (Kitching, and Cadiou, 2000) are

moderate to large sized insects with wingspans that range from 25 to 200 mm (Kitching and

Cadiou, 2000) and body weights ranging from 0.1 to 7 g (Janzen 1984). Hawk-moths outside of

the tribe Smerinthini typically possess enhanced proboscides for nectar feeding and water

drinking, allowing a longer lifespan than species which survive on fat reserves during their adult

phase of the life cycle (Janzen, 1984). Locally, they seem to distinguish between visited and

unvisited flowers. Over the course of days, they efficiently revisit flowers and patches on a

regular basis and, over seasons, exhibit well directed long distance movements and migrations

(Janzen, 1984). Hawk-moths evolved unique flight skills, including the ability to hover and a

capacity for quick, long distance flight (Scoble, 1992). For instance, about half of the hawk-moth

species at Santa Rosa National Park in Costa Rica migrate out of the park (Janzen, 1986). Many

North America species disperse across continents, though the consistency and regularity of such

dispersals are unknown (Tuttle, 2007). Some North American species likely migrate between

North and South America as such cross-continental migration is known for many hawk-moths of

the Western Palearctic (Pittaway, 1993).

All 328 hummingbird species reside in the New World (Schumann, 1999). The family

includes the smallest known bird species with wing lengths from 29 to ≥ 90 mm (Johnsgard,

1997) and body masses ranging from 2 to 21 g (Schumann, 1999). Hummingbirds, like hawk-

moths, possess specialized features for nectar-feeding, including elongated bills and extensible

bitubular tongues for reaching and extracting nectar. Large breast muscles (30% of body weight)

and specialized wings giving them the ability to hover and fly backwards. Hummingbirds are

capable of long distance flight, with 13 of the 15 species of the United States exhibiting some

degree of long distance migration (Johnsgard, 1997).

Many New World species of flowers exhibit distinct pollination syndromes that favor the

morphology and behavior of hawk-moths or hummingbirds, respectively. Phalaenophilic (moth-

pollinated) flowers typically open at night and use odor instead of visual cues as attractants,

resulting in strongly scented but pale flowers. Comparatively narrow nectar tubes match the thin

probosci typical of moths. Sex organs of phalaenophilic flowers are typically recessed, with the

anther and stigma inserted within the corolla tube.

Ornithophilic (hummingbird-pollinated) flowers typically open during the day, are

vividly colored (usually red), and offer little to no scent. Relatively wide nectar tubes match the

bill width typical of hummingbirds (Faegri and van der Pijl, 1979). Exserted sex organs with the

anther and stigma extending beyond the corolla characterize ornithophilous flowers (Kulbaba

and Morley, 2008).

Despite these differences in flower morphology, both phalaenophilic and ornithophilic

flowers share traits through convergent evolution. Both offer abundant nectar sources contained

deep within long nectar tubes. Visual guides for pollinators are relatively absent in both flower

types, with moths using the contours of the blossom as a guide (Faegri and van der Pijl, 1979).

Individuals of both hawk-moths and hummingbirds prefer high sugar and abundant nectar, and

each family readily feeds on the other's flowers (Cruden et al., 1983; Cruden et al., 1976; Haber

and Frankie, 1989). This extensive niche overlap offers ample opportunities for competition.

Distribution Analysis

We determined the species richness of hummingbirds and hawk-moths across the

continental USA. Using range maps and text descriptions provided by Johnsgard (1997) and

Tuttle (2007), we determined the species richness for the 49 states of the continental USA. We

used states as our scale of resolution as finer scale distribution data for hawk-moths does not

exist. For our purpose of looking at large-scale biogeographic patterns, coarse but complete data

is more important than fine but incomplete data. Our analysis included rare native species but

excluded species non-native to the United States. First, we tested for geographic gradients of

each Family's diversity by creating a general linear model in which each state's longitude and

latitude – based upon the centroid of the state – were independent variables and species richness

as the dependent variable. Once confirmed, we investigated environmental variables as potential

determinants of the pattern. These climatic, seasonal and elevational variables were selected *a*

priori based on our hypotheses.

For each of the 48 contiguous states, we determined mean daily, maximum, and

minimum summer and winter temperatures; mean summer and winter precipitation; and average

atmospheric pressure. To eliminate potential bias due to total rainfall, we also calculated the -

difference between mean winter and summer precipitation (winter-summer). We obtained

weather data using the Monthly Station Normals 1971-2000 CLIM 81 from NOAA and

averaging across all weather stations within each state. Winter variables were calculated with

data from December, January, and February, and summer variables used June, July, and August.

Precipitation was used as a proxy for time of flowering: winter/early spring (cool season) vs.

summer/early fall (warm season). Since changes in elevation also lead to changes in both

temperature and atmospheric pressure, we used the barometric formula (eq. 1) with the annual

mean temperature and the mean elevation of the state to determine mean atmospheric pressure

(SI Table 1). Here, P_h is mean atmospheric pressure (in atmospheres), P_0 is atmospheric pressure

at sea level, g is the gravitational constant, M is molar mass, T is absolute temperature, R is the

universal gas constant, and h is elevation in meters.

$$P_h = P_0 e^{\frac{-g*M*h}{R*T}}$$ (1)

Mean elevation per state was taken from the 2004-2005 Statistical Abstract of the United States,

Section 6.

General linear modelling was used to determine which variables correlated significantly

with species richness. For each Family separately, we used a step-wise regression, eliminating at

each step the least significant variables based upon their p-values. This left a linear model with

the remaining significant variables at a level of $p < 0.05$. Because of redundancy among some

variables, three different permutations of tests were performed. The first permutation used

summer daily mean temperature, winter daily mean temperature, winter precipitation, summer

precipitation, and atmospheric pressure. Since the west is drier with lower overall precipitation,

we ran a second test using precipitation difference in lieu of winter and summer precipitation. A

third test used daily maximum temperatures for hummingbirds and daily minimum temperatures

for hawk-moths as the two Families are diurnal and nocturnal respectively.

Additional Species Richness Data: We can also provide figures for the species richness of

hummingbirds by Canadian Provinces, Mexican States, and South American countries. We can

provide species richness data for Canadian Provinces, six regions of Mexico (data by states is not

available), Provinces of South Africa, and Australian Provinces.

Additional bird and insect Families of the continental USA: We specifically chose the

hummingbird and hawk-moth Families as likely candidates for macrocompetition. Yet, what we

find for each may simply be a more generally property of birds and insects in the 48 contiguous

states, USA. So, we haphazardly selected 12 bird Families and 7 insect Families that exhibited

sufficient species richness and data quality to test for within Family latitudinal and longitudinal

trends (Table 3).

RESULTS

Fifteen hummingbird species and 101 hawk-moth species inhabit the continental United

States. Figures 2a and 2b show the species richness of hawk-moths and hummingbirds,

respectively, in the United States, Canada, and Mexico by state, province, territory, and region.

Inspection of these graphs reveals that hummingbird species increase from north to south and

from east to west (Figure 2b). Hawk-moth species likewise increase from north to south, but in

contrast to hummingbirds, hawk-moth diversity increases from west to east (Figure 2a). The

result of the latitudinal and longitudinal GLM confirm the directional bias seen on the map

(Table 1). In summary, hummingbird diversity peaks in the Southwestern United States, while

hawk-moth richness peaks in the Southeastern United States.

We examined the association between environmental variables and the diversities of

hawk-moths and hummingbirds with the results shown in Table 2. Model permutation 1 shows

hawk-moth species richness increasing with summer precipitation and summer daily average

temperatures, and hummingbird species richness decreasing with winter and summer

precipitation and increases with winter daily average temperature. With permutation 2, the

relationship for hawkmoths now showed an increase with winter daily average temperature and

atmospheric pressure and a decrease with precipitation difference, indicating an association with

summer rains; hummingbird species richness showed a positive correlation with winter daily

average temperature and a negative correlation with atmospheric pressure. The third permutation

of tests, which used the minimum and maximum temperatures for hawkmoths and hummingbirds

respectively, showed similar correlations for hummingbirds as the second test (but with new

coefficients) but that hawk-moths were only significantly correlated with summer daily

minimum temperature. In summary, hawk-moth diversity is higher in states with higher

summertime precipitation and temperatures, particularly the minimum temperature (consistent

with nocturnal activity), and states with overall low elevations. Hummingbird diversity is higher

in states with higher wintertime temperatures, particularly the winter highs (consistent with

diurnal activity), and states with higher elevation.

Additional Species Richness Data: The latitudinal trends in species richness of hawk-

moths and hummingbirds is pervasive throughout North America, from Canada to Mexico (Fig.

2). Only one species of hummingbird is found east of the province of Alberta, Canada, while

hawk-moth species richness is greater in eastern than in western Canada. The small eastern

province of Prince Edward Island, for instance, has the same number of species as the large

western province of British Columbia. The geographical trend is more difficult to see in Mexico,

because hawk-moth species richness could only be resolved to 6 regions. Nonetheless,

hummingbird species richness in the Yucatan is low compared to close-by and neighboring states

to the west. Each state (Yucatan, Campeche, and Quintana Roo) has only 9 species each – 10

combined – compared with 14 in neighboring Tabasco, 20 in Michoacán, and 26 in Guerrero. In

contrast, there are 133 hawk-moth species present in the state of Veracruz alone compared to 120

species in the states of Nayarit, Jalisco, Colima, Michoacán, Oaxaca, and Guerrero combined.

Species richness of hawk-moths and hummingbirds appears to exhibit a similar reversal

in directional dominance in South America as we found in North America. Hummingbird species

richness is greater in western than in eastern South America (Johnsgard, 1997; NatureServe,

2010). Ecuador has approximately twice as many species as Brazil, and hummingbird species

richness per area is greater in Chile than in Argentina despite having fewer species overall.

Though the data for hawk-moths in South America are less comprehensive, species

richness appears greater in French Guiana, Argentina, Bolivia, and Venezuela (CATE, 2010)

than in countries to the west. Together with the results from North America, climate and

topography appear to exert opposing effects on the continental distributions of hawk-moths and

hummingbirds. Finally, both South Africa and Australia (Fig. 3) show support for hawk-moth

species richness increasing longitudinally from west to east, and latitudinally towards the

equator. What we do not known is whether these north-south and east-west patterns hold up

within the very large Cape and Gauteng Provinces of South Africa and the Western Province of

Australia.

Additional bird and insect Families of the continental USA: Of the seven additional insect

Families two showed no significant latitudinal trends, four showed significantly increasing

richness from east to west (opposite hawk-moths), and only one shared the same pattern as the

hawk-moths (Table 3). The species richness of Libellulidae (largest family of dragonfly)

increases from west to east. Of the twelve bird Families, two showed no significant latitudinal

patterns, one showed a significant increase in richness from west to east (the Parulidae), and nine

exhibit the same pattern as the hummingbirds with more species as one moves westwards.

Despite shared trends the hummingbirds do stand out in having just a single species in 36 of 48

states of the contiguous USA. All of the other bird families of Table 2 have at least three or four

species reaching east coast states, even those with fewer total numbers of species than

hummingbirds.

DISCUSSION

Our analyses for evaluating evidence for macrocompetition between hummingbirds and

hawk-moths has its limitations. Firstly, correlation does not necessarily illuminate causation.

Yet, given the degree to which hawk-moths and hummingbirds have been ignored as potential

shapers of each other's biodiversity and distributions, correlations will shed light on some of our

hypotheses and suggest new ones. Secondly, using states of the USA as our unit of replication is

geographically crude; they vary in size by more than two orders of magnitude, have diverse and

irregular shapes, and adjacent states will have some degree of spatial autocorrelation. A more

fine-grained and detailed level of division, such as the county, or the use of GIS data would be

preferable, and in many cases possible for hummingbirds. Unfortunately, hawk-moth diversity

and distribution data are coarse; and this is the first systematic analysis of hawk-moth species

richness. Fine grain data on hawk-moth species' ranges and presence/absence are deficient

throughout the world and digital range maps are scarce to non-existent with the range maps from

Tuttle (2007) not digitizable. We feel our scale best balances the need for accurate data and

diverse sampling units. We have high confidence in state-wide inventories of hawk-moths.

Perhaps, intriguing results will inspire more detailed interest and work. Thirdly, due to the

irregularities of species boundaries, states, especially those along the border with Mexico, could

contain species with well-established populations that occupy just a fraction of the state. While

this inflates the numbers of species within the state, any boundary drawing would necessarily

have this problem. We felt it best to accept the current haphazard sizes and irregularities of states

rather than create more regular spatial sampling schemes that would amplify uncertainties in the

presence/absence of hawk-moth species.

Despite these limitations, our results show clearly divergent trends in broad scale

geographical patterns of hawk-moths and hummingbirds and are striking and consistent with the

hypothesis of intertaxon competition and a phenomenon we call macrocompetition.

Environmental correlates of the geographic trends for each of the two taxa further support the

potential for geographical partitioning based on elevation, precipitation, and temperature.

Though only suggestive, our investigation highlights an important yet understudied interaction at

higher levels of taxonomy. The pattern for hummingbirds is well known, but the pattern that

emerged for hawk-moths is novel and quite unexpected.

Basic Geography

The 15 hummingbird species and 101 hawk-moth species in the US exhibit a clear and

opposite latitudinal bias in species richness. Most hummingbird species are found in the western

United States, with only 1 species breeding east of the Rocky Mountains. This result is generally

consistent with the expectation that species diversity should be greater in mountainous areas due

to habitat heterogeneity and reproductive isolation. The distribution of hawk-moths on the other

hand offers some striking and initially counter-intuitive patterns. Despite their relatively large

size and habitat heterogeneity, western states are conspicuously depauperate in hawk-moth

species. States on the eastern seaboard just a fraction of the size of California have much higher

hawk-moth diversities. Similar diversity asymmetries appear when noting Maine's (extreme

northeast) high diversity in contrast to low diversity in the state of Washington (far northwest).

Even neighboring states show this pattern with North Dakota having over 25% more hawk-moth

species than Montana despite being smaller in area, sharing a biome, and offering much less

environmental heterogeneity. This clear portioning invites explanation by way of climatic

analysis.

Climate Analysis

Being a large, nocturnal insect poses challenges and opportunities for such an

evolutionary technology. Accordingly, the analyses showed how hawk-moth species richness

increases with higher temperatures – especially high summer minimums – higher summer

precipitation (associated with warm season flowering), and high atmospheric pressure. In studies

along mountain gradient, hawk-moth pollination activity fell off once summer minimums

reached below 15ºC in mountainous areas (Harlington, 1968; Cruden et al., 1976). With respect

to elevation, insects likely suffer more than birds from a drop in the partial pressure of oxygen;

the hawk-moth tracheal system requires diffusion of O_2 into and CO_2 out of spiracles located on

the exoskeleton. Adult insects, as shown by a study with the tobacco hornworm, *Manduca sexta*,

are smaller when reared under hypoxic conditions (Harrison et al., 2010). And, smaller size is

not favorable for hawk-moths that must maintain thoracic heat for flight (Dorsett, 1962). As seen

in North America, South Africa, Australia, and likely South America, the eastern sides of

continents are more likely to offer summer rains and higher nighttime temperatures during the

summer relative to west sides.

Relative to hawk moths, the evolutionary technology of hummingbirds should allow for a

higher fitness in colder temperatures, cool season flowering, and lower barometric pressures both

in regard to temperatures and oxygen partial pressures. Species richness of hummingbirds

correlated with all of these in the predicted direction. In the Andes of South America,

hummingbird richness is highest in the 1800-2500m range (Schuchmann, 1999) as well as

highest between 500-1800m in southwestern United States (Wethington et al., 2005). The

respiratory system of hummingbirds maximizes the intake of O_2. Correcting for body size,

hummingbirds have the largest heart, fastest heart and breathing rates, and densest erythrocyte

concentrations among all bird families (Johnsgard, 1997). For example, *Colibri coruscas*

increases oxygen consumption by only 6 to 8% when hovering under hypoxic conditions

equivalent to an altitude of 6000m (Berger, 1974). Hummingbirds are better pollinators than

insects in the cloudy, windy, and rainy conditions often found at high elevations (Cruden, 1972).

The species richness of hummingbirds at these altitudes may result from their physiological

adaptations (McGuire, 2014). For somewhat similar reasons of inter-taxonomic, ant diversity

may steadily decline with elevation while in the tropics small mammal diversity often peaks at

intermediate elevations (Nor 2001, Heaney 2001, McCain 2005, Sanders et al. 2007).

Different evolutionary technologies should influence methods of resource exploitation as

well as the costs. We can expand the notion of fundamental and realized niches to whole

Families. It is clear that hawk-moth characteristics favor areas of warm growing season

flowering that are low in elevation and oxygen rich. Presumably these are features that would

favor hummingbirds as well. Though endotherms, hummingbirds are extremely small and easily

lose heat. Many hummingbird species undergo nightly torpor to conserve energy. Moreover,

though suffering only a small uptick in energetic costs at higher elevations, hummingbirds are

still more efficient at lower elevations. Furthermore, migratory hummingbird species, like hawk

moths, time their activity in the United States for the summer. It seems that the fundamental

niche of hummingbirds contains that of hawk-moths, and extends further into colder

temperatures and higher elevations. So, why does the species richness and hummingbird

Family's realized niche not match its fundamental niche? *Why only one hummingbird species*

east of the Rocky Mountains?

Whither Macrocompetition?

Hummingbirds and hawk-moths are two phylogenetically distant, yet morphologically

convergent families of nectarivores (Cruden et al., 1983; Carpenter, 1979; Kessler et al., 2010;

Aigner and Scott, 2002; Fulton, 1999; Willmott and Burquez, 1996). Prior work supports high

niche overlap and definitive micro-competition and the opportunity for meso-competition. Our

analyses support geographical partitioning and inter-taxonomic macrocompetition. That hawk-

moths contribute to an unusually low diversity of hummingbirds in the east and vice-versa in the

west provides an open but intriguingly viable hypothesis.

The evolution and distribution of nectar feeding bats in the family Phyllostomidae may

also shed light on the possibility of macrocompetition between hummingbirds and hawk-moths.

These bats use the same class of resources, taking from flowers with similar – if not higher –

amounts of sugar and nectar to those pollinated by hawk-moths and hummingbirds (Cruden et

al., 1983). The biogeography of these bats resembles that of hummingbirds, existing primarily in

tropical regions (none inhabit temperate North America) with species richness highest in the

Andes. Furthermore, Phyllostomid bats underwent a radiation in the mid-Miocene around the

same time as the hummingbird radiation. Though diversifying in a similar manner to

hummingbirds, there are many fewer species of true nectarivores -- sixteen genera with 38

species adapted for nectar feeding (Fleming et al., 2009). This relatively limited species diversity

of bat nectarivores may reflect competition from both hummingbirds and hawk-moths for

available niche space. These nectarivorous Phyllostomid bats show a similar pollination

syndrome to hawk-moths – nocturnal foraging with dependence on scent to guide them to

flowers – with some plant species relying on both bats and hawk-moths for pollination

(Hernandez-Montero and Sosa, 2015). We conjecture that bat nectarivore diversity is also

constrained by competition from hawk-moths at lower elevations and by hummingbirds at higher

elevations, severely restricting their potential niche space to high-elevation areas at night.

The alterative to macrocompetition would view the geographic partitioning of

hummingbirds and hawk-moths as apparent and not causally related. Indeed, a number of North

American bird Families show east to west increases in diversity, and perhaps this is the norm. In

particular, we do not suggest that showing geographic partitioning by haphazard pairs of

Families provides evidence for macrocompetition. Here the two Families were selected *a priori*,

because of their convergence in morphology and ecological niche similarities. They do exhibit

microcompetition (Carpenter 1979). In terms of mesocompetition, experiments are lacking as to

whether within communities hawk-moths depress the population sizes of hummingbirds and

vice-versa. But, given the capacity of hawk-moths and hummingbirds to deplete nectar, it seems

likely. The geographic partitioning of diversity was striking. No bird Families with comparable

total continental diversities plunge to just one species as one hits the longitudinal midsection of

the United States; and the results for the hawk moths are novel and striking in the opposite

direction. Of seven other insect Families only one exhibited a significant increase in species

richness from west to east.

In terms of Brown and Davidson's (1979) requirements for inter-taxonomic competition

the first and third indicators are met. While not conclusive, the evidence for macrocompetition

between hawk-moths and hummingbirds is tantalizing. The all but ignored eco-evolutionary

relationships between hawk-moths and hummingbirds seems ripe for study. Profitable avenues of

studies could include 1) better local data on hummingbird and hawk-moth diversities and

population sizes, and 2) mesocompetition studies on smaller scales to verify that each Family can

influence local abundances and diversities. Looking more generally, we hope our analyses

inspire further study into local and even continent-wide distributions of competition and niche

partitioning.

ACKNOWLEDGEMENTS

Abdel Halloway wishes to thank the NSF for funding his graduate studies. This material is based

upon work supported by the National Science Foundation Graduate Research Fellowship under

Grant Nos. DGE-0907994 and DGE-1444315. Any opinion, findings, and conclusions or

recommendations expressed in this material are those of the authors(s) and do not necessarily

reflect the views of the National Science Foundation.

REFERENCES

Aigner PA, Scott PE (2002) Use and pollination of a hawkmoth plant, *Nicotiana attenuata*, by
   migrant hummingbirds. Southwest Nat 47:1-11

Berger M (1974) Energiewechsel von Kolibris beim Schwirrflug unter Hoehenbedingungen.
   Journal fuer Ornithologie 115:273-288

Brown JH, Davidson DW (1977) Competition between seed-eating rodents and ants in desert
   ecosystems. Science 196:880-882

Brown JH, Davidson DW (1979) An experimental study of competition between seed-eating
   desert rodents and ants. Am Zool 19:1129-1143

Brown JS, Kotler BP, and Mitchell WA (1997) Competition between birds and mammals: a
   comparison of giving-up densities between crested larks and gerbils. Evol Ecol 11:757-
   771

Buckley L, Hurlbert AH, and Jetz W (2012) Broad-scale ecological implications of ectothermy
   and endothermy in changing environments. Glob Ecol and Biogeogr 21:873-885

Carpenter F (1979) Competition between hummingbirds and insects for nectar. Am Zool
   19:1105-1114

Cipollini ML, Stiles EW (1993) Fruit rot, antifungal defense, and palatability of fleshy fruits for
   frugivorous birds. Ecology 74:751-762

Cipollini ML, Levey DJ (1997) Secondary metabolites of fleshy vertebrate-dispersed fruits:
   adaptive hypotheses and implications for seed dispersal. Am Nat 150:346-372.

Cruden RW (1972) Pollinators in high-elevation ecosystems: relative effectiveness of birds and
   bees. Science 176:1439–1440

Cruden RW, Kinsman S, Stockhouse II RE, Linhart YB (1976) Pollination, fecundity, and the

distribution of moth-flowered plants. Biotropica 8:204-210

Cruden RW, Hermann SM, and Peterson S (1983) Patterns of nectar production and plant-

pollinator coevolution. In: Bentley B, Elias T (eds) The biology of nectaries. Columbia

University Press, New York, pp 80-125

Dorsett DA (1962) Preparation for flight by hawk-moths. J Exp Biol 39:579-588

Faegri K, van der Pijl L (1979) The principles of pollination ecology. Pergamon Press Inc.,

Elmsford, New York, USA.

Fleming TH, Geiselman C, Kress WJ (2009) The evolution of bat pollination: a phylogenetic

perspective. Ann Bot 104:1017-1043

Fulton M, Hodges SA (1999) Floral isolation between *Aquilegia Formosa* and *Aquilegia*

pubescens. Proc R Soc Lond B Biol Sci 266:2247

Gause GF (1934) The struggle for existence. Williams and Wilkins, Baltimore

Haeming PD (1994) Effects of ants on the foraging of birds in spruce trees. Oecologia 97:35-40.

Harrison JF, Kaiser A, VandenBrooks JM (2010) Atmospheric oxygen level and the evolution of

insect size. Proc R Soc Lond B Biol Sci 277:1937–46.

Heaney LR (2001) Small mammal diversity along elevational gradients in the Philippines: an

assessment of patterns and hypotheses. Glob Ecol Biogeogr 10:15-39

Hernández-Montero J, Sosa VJ (2015) Reproductive biology of *Pachira aquatic* Aubl.

(Malvaceae: Bombacoideae): a ropical tree pollinated by bats, sphingid moths, and honey

bees. Plant Species Biol 31:125-134

Haber WA, Frankie GW (1989) A tropical hawkmoth community: Costa Rican dry forest

Sphingidae. Biotropica 21:155-172

Hutchinson GE (1978) An introduction to population ecology. Yale University Press, New

Haven

Janzen DH (1977) Why fruits rot, seeds mold, and meat spoils. Am Nat 980:691-713

Janzen DH (1984) Two ways to be a tropical big moth: Santa Rosa saturniids and sphingids. Oxf

Surv Evol Biol 1:85-140

Janzen DH (1986) Biogeography of an unexceptional place: what determines the saturniid and

sphingid moth fauna of Santa Rosa National Park, Costa Rica, and what does it mean to

conservation biology? Brenesia 25/26:51-87

Jedlicka JA, Greenberg R, Perfecto I, Philpott SM, Dietsch TV (2006) Seasonal shift in the

foraging niche of a tropical avian resident: resource competition at work? J Trop Ecol

22:385-395.

Jablonski D (2008) Scale and hierarchy in macroevolution. Paleontology 50:87-109

Johnsgard PA (1997) The hummingbirds of North America. Smithsonian Institution Press,

Washington, D.C.

Kessler D, Diezel C, Baldwin IT (2010) Changing Pollinators as a Means of Escaping

Herbivores. Curr Biol 20:237-242

Kitching IJ (2017) Sphingidae Taxonomic Inventory (formerly CATE-Sphingidae),

http://sphingidae.myspecies.info/, accessed in 2014

Kitching IJ, Cadiou J (2000) Hawkmoths of the world: an annotated and illustrated revisionary

checklist (Lepidoptera: Sphingidae). Cornell University Press, Ithaca

Kulbaba MW, Worley AC (2008) Floral design in *Polemonium brandegei* (Polemoniaceae):

genetic and phenotypic variation under hawk-moth and hummingbird pollination. Int J

Plant Sci 169:509-522

Laverty TM, Plowright RC (1985) Competition between hummingbirds and bumble bees for

nectar in flowers of *Impatiens biflora*. Oecologia 66:25-32

McCain CM (2005) Elevational gradients in diversity of small mammals. Ecology 86:366-372.

McGuire JA, Witt CC, Remsen Jr. JV, Corl A, Rabosky DL, Altshuler DL, Dudley R (2014)

Molecular phylogenetics and the diversification of hummingbirds. Curr Biol 24:910-916

Mokany A, Shine R (2003) Competition between tadpoles and mosquito larvae. Oecologia

135:615-620

Morin PJ, Lawler SP, Johnson EA (1988) Competition between aquatic insects and vertebrates:

interaction strength and higher order interactions. Ecology 69:1401-1409

Natureserve (2018) NatureServe Web Service. Arlington, VA. U.S.A. Available

http://services.natureserve.org. (Accessed: 2010)

Nor S (2001) Elevational diversity patterns of small mammals on Mount Kinabalu, Sabah,

Malaysia. Glob Ecol Biogeogr 10:pp.41-62.

Palmeirim JM, Gorchov DL, Stoleson S (1989) Trophic structure of a neotropical frugivore

community: is there competition between birds and bats? Oecologia 79:403-411

Pintor LM, Brown JS, Vincent TL (2011) Evolutionary game theory as framework for studying

biological invasions. Am Nat 177:410-423

Pittaway AR (1993) The hawk-moths of the western palaearctic. Harley Books, Colchester

Primack RB, Howe HF (1975) Interference competition between a hummingbird (*Amazilia*

tzacatl) and skipper butterflies (Hesperiidae). Biotropica 7:55-58

Ripa J, Storlind L, Lundberg P, Brown JS (2009) Niche-coevolution in consumer resource

communities. Evol Ecol Res 11:305-323

Rosenzweig ML (1978) Competitive speciation. Biol J Linn Soc Lond 10:275-289.

Rosenzweig ML, McCord RD (1991) Incumbent replacement: evidence for long-term

evolutionary progress. Paleobiology 17:202-213

Sanders NJ, Lessard JP, Fitzpatrick MC, Dunn RR (2007) Temperature, but not productivity or

geometry, predicts elevational diversity gradients in ants across spatial grains. Glob Ecol

Biogeogr 16:640-649.

Schluter D (2000) The ecology of adaptive radiation. Oxford University Press, Oxford

Schuchmann KL (1999) Family Trochilidae (hummingbirds). In: del Hoyo J, Elliot A, Sargatal J.

(eds) Handbook of the birds of the world, vol. 5. Lynx Edicions, Barcelona, pp 468-535

Scoble MJ (1992) The lepidoptera – form, function, and diversity. Oxford University Press,

Oxford

Shivik JA (2006) Are vultures birds, and do snakes have venom, because of macro- and

microscavenger conflict? BioSience 56:819-823.

Silvestro D, Antonelli A, Salamin N, Quental TB (2015) The role of clade competition in the

diversification of North American canids. Proc Natl Acad Sci U S A 112:8684-8689

Thomas CD, Lackie PM, Brisco MJ, Hepper DN (1986) Interactions between hummingbirds and

butterflies at a *Hamelia patens* bush. Biotropica 18:161-165

Tuttle JP (2007) The hawk moths of North America: a natural history study of the Sphingidae of

the United States and Canada. The Wedge Entomological Research Foundation,

Washington, D.C.

Wethington SM, West GC, Carlson BA (2005) Hummingbird conservation: discovering diversity

patterns in southwest USA. Proc RMRS 36:162-168

Willmott AP, Burquez A (1996) The pollination of *Merremia palmeri* (Convonulaceae): can

hawk-moths be trusted? Am J Bot 83:1050-1056

Wright SJ (1979) Competition between insectivorous lizards and birds in central Panama. Am

Zool 19:1145-1156

Vincent TL, Brown JS (2005) Evolutionary game theory, natural selection, and darwinian

dynamics. Cambridge University Press, Cambridge

Table 1: GLM of Latitude and Longitude per state with species richness per family

Family	S	Latitude	Longitude
Sphingidae	101	-1.301[a]	0.264[c]
Trochilidae	15	-0.143[a]	-0.116[a]

a: $p<0.001$, b: $p<0.01$, c: $p<0.05$

Table 2: Coefficients, total R-squared, and AIC for each final linear model. Permutation signifies

the permutation of variables used for each regression as seen in the methods. PRECIP indicates

precipitation, DIFF difference, SUM summer, WINT winter, AVG average daily mean

temperature, MIN average daily minimum temperature, MAX average daily maximum

temperature, and ATM atmospheric pressure

Permutation	Family	Model	R^2	AIC
1	Sphingidae	$S = -7.701 + 0.116 \times SUM.PRECIP\,[a] + 1.470 \times SUM.AVG\,[a]$	0.6448	169.37
	Trochilidae	$S = 8.215 - 0.021 \times WINT.PRECIP\,[b] - 0.052 \times SUM.PRECIP\,[a] + 0.249 \times WINT.AVG\,[a]$	0.651	47.64
2	Sphingidae	$S = -20.73 - 0.094 \times PRECIP.DIFF\,[a] + 0.87 \times WINT.AVG\,[a] + 57.25 \times ATM\,[a]$	0.6187	173.76
	Trochilidae	$S = 31.04 + 0.177 \times WINT.AVG\,[a] - 30.63 \times ATM\,[a]$	0.5149	62.96
3	Sphingidae	$S = -5.90 + 1.89 \times SUM.MIN\,[a]$	0.6035	173.8
	Trochilidae	$S = 28.54 + 0.164 \times WINT.MAX\,[a] - 28.98 \times ATM\,[a]$	0.5148	62.97

545   a: p<0.001, b: p<0.01

546   Table 3: Linear model of Latitude and Longitude with species richness per insect and bird family

Family	S	Latitude	Longitude
Acrididae	215	-0.722	-1.165[a]
Hesperiidae	216	-1.963[b]	-0.114
Libellulidae	109	-1.130[a]	0.206[b]
Nymphalidae	166	0.379	-0.367[b]
Papilionidae	24	-0.126[b]	-0.104[a]
Riodinidae	23	-0.270[b]	-0.078[b]
Saturniidae	69	-0.798[a]	0.035

Family	S	Latitude	Longitude
Accipitridae	23	-0.142[b]	-0.084[a]
Anatidae	47	-0.093	0.012
Caprimuglidae	8	-0.139[a]	-0.018[b]
Corvidae	18	0.009	-0.111[a]
Emberizidae	42	-0.035[a]	-0.177[a]
Hirundinidae	8	0.077[b]	-0.028[b]
Icteridae	21	-0.167[b]	-0.069[b]
Parulidae	51	-0.128	0.313[a]
Picidae	23	-0.130[b]	-0.087[a]
Turdidae	15	0.128[b]	-0.044[a]
Tyrannidae	34	-0.059	-0.210[a]
Vireonidae	12	0.009	-0.015

a: $p<0.001$, b: $p<0.01$

Fig. 1 (a) *Macroglossum stellatarum*, the Hummingbird Hawk-moth, hovering by lavender

flowers (b) *Amazilia tzacatl*, the Rufous-tailed Hummingbird, feeding in Costa Rica. As seen in

the image below, hawk-moths have extremely long extensile probosces for collecting nectar and

the ability to hover in front of flowers. Convergent features are seen in the image of the

hummingbird below with long bills and extensile tongues and the ability to hover. Image of *M.*

stellatarum by Thorsten Denhard, CC-BY-SA-3.0. Image of *A. tzacatl* by T. R. Shankar Rama,

CC-BY-SA-4.0

Fig. 2 Species richness per state, province, territory, and region of in the United States, Canada,

and Mexico (a) hawk-moths (order Lepidoptera, family Sphingidae) and (b) hummingbirds

(order Apodiformes, family Trochilidae). Greater color intensity reflects greater proportional

species richness per family. As one can see, hawk-moths are more species rich in the eastern half

of the northern North American continent while hummingbirds are more species rich in the

western half of the northern North American continent. Both species show increasing species

richness moving from north to south.

Fig. 3 Species richness of a) hawk-moths per pre-1994 South Africa province and b) hawk-moths

per state and territory of Australia, and c) hummingbirds per country in South America. One can

clearly see the same eastern bias for hawk-moths and western bias for hummingbirds.

Fig. 1a

Fig. 1b

Fig. 2

Fig. 3