The MYC antagonist MNT autoregulates its expression and supports proliferation in MAX deficient cells

M Carmen Lafita-Navarro[1,4], Judit Liaño-Pons[1], Andrea Quintanilla[1,5], Ignacio Varela[1], Fabiana Ourique[1,6], Gabriel Bretones[1,7], Rosa Blanco[1], Julia Aresti[1], Patrick Carroll[2], Peter Hurlin[3], Robert N. Eisenman[2], M. Dolores Delgado[1] and Javier León[1*]

[1]Instituto de Biomedicina y Biotecnología de Cantabria (IBBTEC), Universidad de Cantabria-CSIC, and Department of Molecular Biology, Universidad de Cantabria, Santander, Spain

[2]Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, USA.

[3]Shriners Hospitals for Children Research Center, and Department of Cell, Developmental and Cancer Biology, Knight Cancer Institute, Oregon Health and Science University, Portland, USA.

[4]Present address: Sorrell lab, Department of Cell Biology UT Southwestern Medical Center, Dallas, USA

[5]Present address: Edinburgh Cancer Research UK Centre, Institute of Genetics and Molecular Medicine, University of Edinburgh, United Kingdom

[6]Present address: Department of Biochemistry, Universidade Federal de Santa Catarina (UFSC), Florianópolis, Brazil

[7]Present address: Departamento de Bioquímica y Biología Molecular, Facultad de Medicina, Instituto Universitario de Oncología (IUOPA), Universidad de Oviedo, Spain

*Corresponding author

Running title: MNT functions in the absence of MAX

Key words: MNT, MAX, MYC, MLX, transcription, promoter

ABSTRACT

MNT is a transcription factor of the MXD family. MNT-MAX dimers down-regulate genes
by binding to E-box sequences, which can also be bound by MYC-MAX to activate
transcription. MNT has been described as a modulator of MYC activity but little is known
about MNT regulation and whether MNT has MAX-independent functions. Using a MAX
deficient cell line and siRNA-mediated silencing of MAX, we show that in the absence of
MAX, the total MNT levels are elevated and that MNT localizes both in the cytoplasm
and the nucleus. In contrast, MNT is predominantly nuclear when MAX is expressed.
MNT is required for optimal cell proliferation even in the absence of MAX, being the first
report of a MAX-independent function of MNT. Interestingly, MNT forms homodimers
and autoregulates its expression by repressing its own promoter. The tight MNT
regulation and its activity in absence of MAX suggest its importance on cell homeostasis.

INTRODUCTION

MNT (also called MXD6) is a basic helix-loop-helix leucine zipper (bHLHLZ) protein and
it is the most divergent member of the MXD family, that also includes MXD1, MXI1, MXD3
and, MXD4. MNT forms heterodimers with MAX through the bHLHLZ domain, and binds
to E-box DNA sequences. MYC is one of the most prevalent human oncoproteins. MYC
can also interact with MAX and bind E-boxes *(Conacci-Sorrell et al 2014, Dang 2012)*.
Whereas MYC-MAX, upon binding to E-boxes, acts primarily as transcriptional activator,
the typical effect of MNT-MAX is the transcriptional repression *(Hurlin et al 1997, Meroni*
et al 1997). MNT can bind not only to MAX but also to the MAX-like HLH protein MLX
(Billin et al 1999, Meroni et al 2000) which alternatively can interact with MLXIP
(MONDOA) and MLXIPL (MONDOB) proteins. Therefore, MNT participates both in the
MAX- and MLX-centred networks [reviewed in *(Diolaiti et al 2015, Yang & Hurlin 2017)*]
serving as the link between the MAX-MYC and MLX-MONDO networks.

MNT is expressed constitutively in proliferating and quiescent cells and protein
levels do not show major fluctuations when quiescent cells are mitotically stimulated
(Hurlin et al 1997 , Hurlin et al 2003, Walker et al 2005) . MNT[-/-] mice are not viable
(Hurlin et al 2003, Toyo-oka et al 2004) while MXD1[-/-], MXI1[-/-] and MXD3[-/-] mice survive,
suggesting that MNT function is not redundant with that of the other MXD proteins *(Foley*
et al 1998, Queva et al 1999, Schreiber-Agus & DePinho 1998). Moreover, MNT is the
only MXD protein in invertebrates *(Diolaiti et al 2015)*.

Consistent with MNT functioning as a MYC transcriptional antagonist, enforced
MNT expression inhibits cell proliferation and impairs MYC-dependent transformation

(Hurlin et al 1997, Walker et al 2005). The deficiency or down-regulation of MNT in

fibroblasts leads to increased proliferation (i.e., similarly to MYC-overexpression) and

partially rescues the proliferative arrest caused by MYC deficiency *(Dezfouli et al 2006,*

Hurlin et al 2003, Nilsson et al 2004, Walker et al 2005). MNT ablation *in vivo* leads to

breast and T-cell tumors *(Dezfouli et al 2006, Hurlin et al 2003, Toyo-oka et al 2004)*,

and partial or total MNT deficiency in mice models impairs MYC-dependent

tumorigenesis *(Link et al 2012; Campbell et al 2017),* suggesting that *MNT* may function

as a tumor suppressor gene. On the other hand, *MNT* knock-out in some cell models

inhibits proliferation and promotes apoptosis *(Dezfouli et al 2006, Hurlin et al 2003,*

Nilsson et al 2004). Thus, as MYC and MNT coexist in proliferating cells, it has been

suggested a role of MNT as a "buffer" of MYC activity.

In the present work we studied possible MAX-independent functions of MNT

using as the main model UR61 cells. These rat pheochromocytoma cells that do not

express a functional MAX protein but a truncated form (termed MAX^{PC12}) that lacks the

second helix and leucine zipper region of the bHLH-LZ domain which are the regions

responsible for dimerization with MYC and MNT *(Hopewell & Ziff 1995)*. Here we

describe a change of MNT subcellular localization depending on MAX expression and

that MNT is required for optimal cell proliferation even in the absence of MAX, being the

first example of a MAX-independent function of MNT. Finally, we show that MNT

transrepresses its own promoter and MNT ability to regulate gene expression in the

absence of MAX.

RESULTS

MNT levels depend on MAX

We first studied MNT levels in proliferating cells from eleven cell lines derived from

different tissues and species, including two cell lines lacking MAX: UR61 and human

small cell lung carcinoma H1417 cells *(Romero et al 2014)*. The results showed that,

although MNT expression varies among cell lines, the two MAX-deficient cell lines

expressed high MNT levels (*Figure 1A*). MNT was expressed in all cell lines as a protein

doublet, due to a slower migrating phosphorylated MNT form *(Popov et al 2005)*.

The high MNT expression in the two MAX-deficient cell lines led us to explore

whether MAX influenced MNT levels. For this purpose, we transfected UR61 cells with

a construct carrying human MAX cDNA driven by the metallothionein promoter, which is

activated by Zn^{2+} cations *(Canelles et al 1997)*. Several clones were isolated and two of

them with robust MAX induction were mixed and the resulting cell line was termed
URMax34. We also generated a cell line transfected with the empty vector, termed
URMT. The induction of MAX in response to Zn^{2+} in URMax34 cells was confirmed by

Figure 1. High MNT expression in MAX-deficient cells. (A) Cell lysates of the indicated cell lines were analyzed by immunoblot to determine the levels of MNT and MAX. β-actin levels were determined for protein loading control. The MAX-deficient cell lines analyzed were: UR61 (rat pheochromocytoma) and H1417 (human small cell lung carcinoma). The rest are MAX-expressing cells: HEK293T (human embryonic kidney, 293T), K562 (human chronic myeloid leukaemia), HeLa (human cervical cancer), Ramos (human B cell lymphoma), SH-SY-Y5 (SHSY, human neuroblastoma), T98G (human glioblastoma), MEFs (mouse embryonic fibroblasts), Neuro-2a (N2a, mouse neuroblastoma) and C6 (rat glioma). (B) Control URMT and URMax34 cells were treated for 24 h with 50 and 100 μM Zn^{+2} and the MNT and MAX protein expression were determined by immunoblot. α-tubulin levels were determined as a control for protein loading. The experiment was repeated three times. (C) Levels of MNT and MAX determined by immunoblot in UR61 cells 24 h after transfection with a MAX expression vector or the empty vector pCEFL (EV). β-actin levels are shown as a protein loading control. The result was reproduced in three experiments. (D) mRNA expression determined by RT-qPCR in URMT and URMax34 cells treated for 24 h with 100 μM Zn^{2+}. Data are represented as mean ±s.d. from three independent experiments, *P =0.012.

immunoblot (*Figure 1B*). We examined the effect of MAX induction on MNT levels in
URMax34 cells. The results showed that MNT was down-regulated upon MAX induction
by Zn^{2+} (*Figure 1B*). To confirm this result and rule out effects potentially related to the
generation of stably transfected clones (as URMax34 system), UR61 cells were
transiently transfected with a MAX expression vector and the results showed a decrease
in MNT protein levels in MAX-transfected cells (*Figure 1C*). It is noteworthy that the r-
expression of MAX in these cells provoked the down-regulation of MNT at the mRNA
levels, as determined by RT-qPCR (*Figure 1D*).

We sought to confirm this in a different cell type and species. For this purpose, a

MAX expression vector was transfected into human myeloid leukemia K562 cells. The

immunoblot results showed lower MNT levels in the cells with MAX overexpression

(*Figure 2A*). Next, we used the Kmax12 cell line *(Canelles et al 1997)*, a K562 derivative

carrying a MAX transgene which expression is induced by Zn^{2+}. Induction of MAX

Figure 2. MAX regulates MNT expression in K562 cells. (A) MNT and MAX levels determined by immunoblot in K562 cells 24 h after transfection with a MAX expression vector. The results were reproduced in three biological replicates (B) MNT and MAX levels determined by immunoblot in Kmax12 cells treated for 48 h with 100 µM Zn^{2+} to induce MAX expression. The results were reproduced in three biological replicatess. (C) MNT and MAX protein expression analyzed by immunoblot in K562 48 h after transfection with siRNA against *MAX* gene. β-actin levels are shown as a protein loading control. The results were reproduced in two experiments. (D) MNT and MAX mRNA expression analyzed by RT-qPCR 48 h after transfection with siRNA against *MAX* gene. Data are shown as mean ± s.d. of three biological replicates. *P <0.005

expression in Kmax12 cells resulted in a concomitant MNT downregulation (*Figure 2B*).

We also used the opposite approach, i.e. depleting cells of MAX and analyzing the

expression of MNT. As shown in *Figure 2C*, MNT protein expression was upregulated

when MAX was silenced with siRNA in K562. Interestingly, MNT mRNA was also up-

regulated in MAX-silenced cells (*Figure 2D*). Altogether, the results showed that low MAX

levels result in MNT up-regulation, likely at the transcriptional level.

MNT binds and repress its own promoter in the presence of MAX

The above results showing that MNT down-regulation took place at the mRNA

level prompted us to investigate if MNT impairs the activity of its own promoter.

Figure 3. MNT regulates its own promoter. (A) Schematic representation of human, rat and mouse MNT promoters showing two conserved E-boxes. The coordinates correspond to the 5' nucleotide of each E-box, using the UCSC genome browser (http://genome.ucsc.edu/, release GRCh37/hg19). (B) Schematic representation of luciferase reporters driven by human MNT promoter generated in this work. The longest reporter (850 bp upstream the transcription start site) contains two conserved E-boxes. The coordinates are as in USCS browser (release GRCh37/hg19). (C) Luciferase assays in the indicated cell lines transfected with the reporters described above plus MNT and/or MAX expressing vectors or the corresponding empty vectors (EV). Luciferase activity was assayed 36 h or 24 h after transfection of UR61 and HEK293T cells, respectively. The data are shown as mean ± s.e.m. of 9 (for 850MNT-luc) or 4 independent transfections (for the rest of experimental points). **$P < 0.005$; *$P < 0.01$.

Bioinformatic analysis of human, mouse and rat MNT promoter regions revealed that

there are two E-box sequences within 1 kb upstream the transcriptional start site of *MNT*

(one canonical E-box, CACGTG (E-box 1), and one non-canonical, CATGTG (E-box 2)

that are conserved among these three different species (*Figure 3A*).

We then constructed a luciferase reporter carrying the 850 bp upstream region

of the transcription start site from the human MNT gene. The construct was termed

850MNT-Luc (*Figure 3B*). HEK293T cells (which express MAX) were transfected with

the 850MNT-Luc and MNT expression vectors. The results showed that MNT

overexpression led to a reduction in the luciferase activity (*Figure 3C*), suggesting that

MNT-MAX negatively regulates the MNT promoter. To determine the contribution of the
two E-boxes in the MNT-mediated negative autoregulation we constructed two reporters
containing each of the E-boxes, termed 220MNT-Luc (containing the last 220 bp of the
850MNT-Luc reporter which includes the E-box 2) and 570MNT-Luc (containing the first
570 bp of the 850MNT-Luc reporter which includes the E-box 1) (*Figure 3B*). The
luciferase assays in HEK293T cells showed that the E-box 2, mapping at −788 was
critical for MNT-mediated downregulation of its own promoter (*Figure 3C*, left panel).

We also investigated the activity of the MNT promoter in UR61 cells, which do
not express MAX. UR61 cells were transfected with the 850MNT-Luc vector together
with MNT and MAX expression vectors. The results also showed a decrease in the
luciferase activity although less than in HEK293T cells (*Figure 3C*, right panel). The
expression of MNT alone also led to a decrease in the luciferase activity, suggesting that
MNT can downregulate MNT promoter in UR61 cells in the absence of MAX. The
repressive effect of MNT was stronger in HEK293T cells than in UR61 cells, which may

be explained by the limited
overexpression of MNT protein achieved
in transfected UR61 cells (shown
below). Co-transfection of MNT and
MAX resulted in a decrease in the
activity of both promoter constructs
(850MNT-Luc and 220MNT-Luc,
carrying fragments of the human MNT
promoter of 842 and 223 bp,
respectively). Since MNT-MAX dimers
bind E-boxes in the promoters to repress
transcription, we analysed the ChIP-seq
data published by the ENCODE project
(http://genome.ucsc.edu/ENCODE).
The data revealed two regions bound by
MAX in the human *MNT* promoter that
encompass the two E-boxes (*Figure
4A*). We also analysed the ChIP peaks
for MAX, MYC and MXI1 proteins on the
MNT promoter in other human cell lines
as H1ES, HeLa, NB4, A549, GM78,
HEPG, SKSM and IMR90. The analysis
revealed that the three proteins

Figure 4. MNT binds to *MNT* promoter. (A) Schematic representation of human and rat MNT promoters showing the peaks for MAX of the ChIP experiments with anti-MAX antibody on human K562 cell line as published by the ENCODE project. (B) ChIP with anti-MNT (left) and anti-MAX (right) antibodies on the MNT promoter of URMT and URMax34 cells, both treated with 100 μM Zn^{2+} for 24 h. MNT and MAX binding to *MNT* gene promoter in the region containing the E-box 2 (-593 in rat gene) was studied by qPCR. Amplicons from Atrogin-1 (*FBX032*) were used as positive control for MNT binding and a region 1389 bp upstream from the TSS without E-boxes was used as negative control. The data are means ±s.e.m. from four independent experiments. * $P<0.05$.

presented peaks at the same positions near the transcription start site of the human MNT

gene (*Figure 4-figure supplement 1*), suggesting that members of the MYC and MXD

family bind and possibly regulate the promoter of *MNT*.

As MNT was downregulated when MAX was re-expressed, we hypothesized that

MNT-MAX heterodimers might bind to the *MNT* promoter and down-regulate its own

expression. To explore this hypothesis,

a ChIP assay was performed with MNT

and MAX antibodies in URMT and

URMax34 treated with Zn^{2+} to induce

MAX. We studied the -593 E-box of the

MNT promoter, as well as a region of the

Atrogin-1 (FBXO32) promoter as a

positive control for MNT binding

(Terragni et al 2011). A region mapping

at 1369 bp upstream on MNT

transcription start site with no E-boxes

was used as negative control. As shown

in *Figure 4B*, MNT and MAX bound to

the *MNT* promoter in URMax34 cells.

However, in the MAX-deficient control

URMT cells, neither MNT nor MAX were

bound to the *MNT* promoter. In the same

way, MNT and MAX were bound to the

positive control *Atrogin-1* promoter in

URMax34 but not in URMT control cells

(*Figure 4B*). These data show that MNT

most likely binds to its own promoter as

a heterodimer with MAX and suggest a

possible negative regulation of MNT own

expression.

Figure 5. MNT subcellular localization depends on MAX. (A) Cell extracts were subjected to cytoplasmic/nuclear fractionation in H1417 (human lung cells deficient in MAX), HEK293T, URMT and URMax34 treated for 24 h with 100 μM Zn^{2+} to induce MAX. The levels of MNT and MAX were determined in each fraction by immunoblot. SIN3B and RhoGDI (ARHGDIA) were used as nuclear and cytoplasmic markers respectively. "C" refers to cytoplasmic fraction and "N" to nuclear fraction. Similar results were obtained in three biological replicates. (B) K562 were transfected with siRNA against the *MAX* gene and 48 h later cell extracts were prepared and subjected to cytoplasmic/nuclear fractionation. The expression of MAX, MNT and MYC were analyzed by immunoblot. The expression of CTCF and α-tubulin were analyzed as a control for nuclear and cytoplasmic proteins respectively. scrRNA, control scrambled siRNA. The results were reproduced in three biological replicates.

MNT localizes in the cytoplasm of MAX-deficient cells

We next examined the subcellular localization of the excess MNT present in MAX-

deficient cells. URMT and URMax34 cells were treated with Zn^{2+} to induce MAX and we

performed a cytoplasmic and nuclear fractionation to analyse MNT levels. In control

URMT cells, MNT protein was found in the nuclear fraction, as expected, but also at

similar levels in the cytoplasm. In contrast, in URMax34 expressing MAX, MNT protein

was found only in the nucleus (*Figure 5A*), the localization typically found for MNT *(Lafita-*
Navarro et al 2016). We were not able to use untreated URMax34 cells as a control as
they express some MAX even in the absence of Zn^{2+} (*Figure 1B*). Cytoplasmic MNT
was also observed in H1417 cells, human cells deficient in MAX. As a control, the
localization of MNT was analysed in HEK293T cells, that express the MAX protein.
Cytoplasm/nucleus fractionation revealed that MNT and MAX were localised in the
nucleus but not in the cytoplasm of HEK293T cells (*Figure 5A*). To test if the localization
of MNT depended on MAX, we silenced the MAX protein in K562 cells with siRNA and
we carried out cytoplasm/nucleus fractionation. In control K562 cells, MNT and MAX
were mainly localised in the nucleus. In contrast, in MAX-depleted cells, a significant
amount of MNT was found in the cytoplasm of K562 cells (*Figure 5B*). It is noteworthy
that nuclear MNT levels were similar in URMT and URMax34, as well as in control K562
and MAX-depleted K562 cells. As controls for nuclear proteins, we used MYC and CTCF
(*Figure 5B*). Taken together, the data indicates that MAX absence leads to an
upregulation of MNT protein and that the excess of MNT accumulates in the cytoplasm.
222

MNT knock-down impairs cell proliferation in MAX-deficient cells
We next asked for a possible biological effect of MNT on cell proliferation in the UR61
MAX-deficient cells. We decided to knock-down MNT through siRNAs. For this, we used
lentiviral particles containing two short-hairpin constructs against the rat MNT gene. Two
shMNT constructs that efficiently reduced MNT levels were used, termed shMNT-1 and
shMNT-2 (*Figure 6A*). The vectors also carried a puromycin-resistance gene. UR61 cells
were transiently co-transfected with the shMNT constructs (or the empty vector) and a
GFP expression vector in a proportion 1:5 to ensure that the GFP-positive cells also
incorporated the shMNT plasmid. Six days after transfection, the fraction of GFP positive
cells were analysed by flow cytometry. The results showed that the fraction of GFP
positive cells was clearly reduced in cells with depleted MNT (*Figure 6B*), suggesting
that MNT loss resulted in impaired cell proliferation. In a second approach, we
transfected the UR61 cells with the sh-MNT constructs and counted viable cells after 3
and 7 days of transfection. As shown in *Figure 6C,* MNT-silenced UR61 cells grew slower
than controls. We also performed clonogenic assays in UR61 transfected with the
shMNT and/or a MAX expression vector (in 1:3 proportions to ensure that the shMNT
containing cells had also incorporated the MAX vector) as well as the empty vectors. 24
h after transfection puromycin was added and after selection the colonies were stained
with crystal violet, the dye was solubilised and quantified. The results showed that MNT
depletion provoked a dramatic growth inhibition. These results of MNT depletion

Figure 6. MNT knock-down impairs cell proliferation in MAX-deficient cells. (A) Silencing of MNT by short-hairpin constructs. URMT cells were transfected with vectors encoding sh-MNT-1 and shMNT-2. 24 h after transfection the cells were treated with puromycin (0,3 µg/ml) and 72 h after transfection the cells were lysed and the levels of MNT analyzed by immunoblot. β-actin levels were determined as a control for protein loading. (B) Fraction of GFP-expressing UR61 cells assessed by flow cytometry 7 days after co-transfection with GFP and shMNT vectors (in proportion 1:5) and analyzed by flow cytometry. The data are shown as mean ± s.e.m. of three independent experiments, *P =0.006, **P =0.008. (C) Cell proliferation determined by cell counting at 3, 7 and 12 days after transfection of shMNT-1 or the empty vector pLKO1. The data are shown as relative mean values ± s.e.m. of three independent experiments, *P = 0.01, **P = 0.008 (D) Cell growth determined by crystal violet staining in UR61 cells transfected with the indicated vectors. After 15 days of puromycin selection the colonies were stained with crystal violet and the dye was solubilised and quantified by absorbance at 595 nm. EV, empty vector (pLKO for shMNTs and pCEFL for MAX). **P < 0.005. Data show mean values ± s.d. from three independent experiments. (E) Fraction of sub G_0 UR61 cells transfected with the shMNT-1 vector and empty vector (EV). Cells were fixed and stained with propidium iodide at day 6 post-transfection and puromycin selection. The percentage of cells containing less than 2C DNA content was determined by flow cytometry. The data are mean values of three independent experiments ± s.d., *P =0.014. (F) Levels of MNT determined by immunoblot in URMT and URMax34 cells 72 h after transfection with the shMNT vector and treated for 12 h with 100 µM Zn^{2+}. MAX, cyclin A2 and survivin (BIRC5) were also determined. β-actin levels were determined for protein loading control. Two experiments were performed with similar results.

impairing cellular proliferation in a MAX-independent manner are unprecedented. These

results also showed that UR61 cells overexpressing the MAX protein grew more slowly

than the control *(Figure 6D),* confirming the effect previously reported for the MAX-
deficient PC12 pheochromocytoma cells (UR61 parental cells) *(Hopewell & Ziff 1995)*.
In addition, when MNT silencing was accompanied by MAX enforced expression, the
inhibition of UR61 cell proliferation was stronger (*Figure 6D*). Consistent with the anti-
proliferative effects of MNT depletion, we failed to generate stable MNT-silenced UR61
cell lines (not shown). In order to generate UR61 cells with MNT overexpression we
transfected UR61 with MNT expression vectors but we were not able to detect a
significant increase in MNT protein levels upon transfection. This is likely due to the
autoregulation of MNT protein levels, as the overexpression of MNT protein was readily
detected by immunoblot when the transfected cells were treated with the proteasome
inhibitor bortezomib (*Figure 6-figure supplement 1A*). In contrast, some increase in the
levels of MNT was detected when MNT was transfected in MAX-expressing URMax34
cells (*Figure 6-figure supplement 1B)* or when MNT was co-transfected with MAX in
UR61 cells (*Figure 6-figure supplement 1C*).

Since the depletion of MNT impairs cell proliferation of UR61 cells, we wondered
whether this was accompanied by cell death. To study this, the levels of survivin (BIRC5)
in URMT and URMax34 were analysed by immunoblot (*Figure 6F*) and the results
showed a decrease in survivin and cyclin A (a marker of proliferation) in cells with
depleted MNT, both in MAX-deficient cells (compare lanes 1 vs 2 in *Figure 6F*) and in
MAX-expressing cells (compare lanes 3 vs 4 in *Figure 6F*), although the effect was
stronger in these latter cells. To check the contribution of apoptosis, we also measured
DNA content flow cytometry. We found that the fraction of cells with a sub-G0 amount of
DNA was higher in cells transfected with the shMNT vector (*Figure 6E*). The results
suggest that the depletion of MNT leads to cell proliferation arrest and apoptosis in UR61.

MNT modifies gene expression in MAX-deficient cells
Since the data shown above indicated that MNT knock-down in UR61 cells leads to
growth arrest and cell death, and considering that MNT is a transcription factor, we set
out to study if MNT regulates the transcriptional program in the absence of MAX. For this
purpose, we investigated the effects of MNT depletion on the transcriptome of cells with
and without MAX. URMT and URMax34 cells were transfected with the short-hairpin
RNA construct against the rat MNT gene (shMNT-1) or the empty vector as control in
two different biological replicates. We first checked the appropriate conditions to
efficiently silence the expression of MNT upon transfection of the shMNT constructs in
URMT and URMax34 cells. A significant MNT depletion was consistently achieved at 72
h post-transfection (not shown). Thus, for the RNAseq experiment, the URMT and

URMax34 cells were transfected with the shMNT-1 construct and two days after
transfection, cells were treated with Zn²⁺ for 24 h to achieve significant MAX-induced
expression (in URMax34) as well as significant MNT depletion (*Figure 7A*). Two

Figure 7. Gene expression changes in MNT knock-down cells. (A) mRNA expression of MNT and MAX in URMT and URMax34 cells lysed 72 h after transfection with shMNT and 24 h of treatment with 100 μM Zn²⁺. The experiments were performed in duplicated (dup.1 and dup.2) biological replicates. These RNAs were subjected to RNA-seq. (B) Percentage of upregulated and downregulated genes in cells transfected with the shMNT or pLKO (empty) vectors in URMT and URMax34 cells, as indicated at the bottom. The number of genes regulated comparing the different samples are indicated at the top of the figure. The gene inclusion criteria were a change in mRNA level ≥0.7 or ≤-0.7 log2 FPKM fold change. The RNA-seq data are from two independent experiments. The genes regulated in the different comparisons are listed in the *Supplementary Files 1 to 4*. (C) Venn diagram showing the genes that are regulated by MNT in MAX-deficient (URMT) or MAX-expressing cells (URMax34). The graph at the right shows the number of up- or down-regulated genes in the group of overlapping genes. (D) mRNA expression regulation due to MNT silencing in genes selected after the RNA-seq data analysis. URMT and URMax34 were transfected with shMNT-1 (upper panel) and shMNT-2 (lower panel) or empty vector pLKO. After 48 h cells were treated for 24 h with 100 μM Zn²⁺, total RNA was prepared and mRNA levels of the indicated genes were determined by RT-qPCR . Most of the selected genes are involved in DNA repair or cell cycle as indicated at the bottom. The data are represented as mean ±s.e.m. from four (shMNT-1) or three (shMNT-2) independent transfection experiments. In all cases, *P* <0.05..

independent transfections and RNA preparations of each cell line were submitted for
next generation sequencing RNA-seq and the data was processed bioinformatically as
escribed in the Methods section.

Once we obtained the expression values of each experimental replicate we
compared the different conditions. For all of the different comparison we grouped the
genes that were up-regulated and down-regulated in both biological replicates
considering a log2 RPKM fold change higher than 0.7 or smaller than -0.7 of the
corresponding control and a p-value < 0.1. The heatmaps of the gene expression
signatures clearly showed that the depletion of MNT in both URMT and URMax34
induced gene expression changes (*Figure 7-figure supplement 1*) indicating that MNT
can be involved in transcriptional regulation without the concourse of MAX. Specifically,
281 genes were regulated upon MNT depletion in MAX-deficient URMT cells (*Figure 7-*
figure supplement 2) and 537 genes in MAX-expressing cells (i.e., URMax34 treated with
Zn^{2+}) (*Figure 7-figure supplement 3*). Of those, approximately 62 % of the genes were
down-regulated upon MNT depletion in both cell lines (*Figure 7B*). In addition, the
URMax34 vs URMT gene expression signatures heatmaps also showed that the
expression of MAX induced gene transcriptional changes independently of MNT
depletion (*Figure 7-figure supplement 1*). Specifically, 377 genes were found to be
differentially expressed when comparing URMax34 and URMT cells (*Figure 7-figure*
supplement 4) and 361 genes when comparing URMax34 and URMT cells depleted of
MNT (*Figure 7-figure supplement 5*). Among *MAX*-regulated genes, roughly half of the
genes were downregulated (*Figure 7B*). Importantly, the comparison between the lists of
differentially expressed genes upon MNT-depletion in cells without MAX (URMT) and
with MAX (URMax34) revealed 158 shared genes in the two biological replicates (*Figure*
7-figure supplement 6). This indicated that 56% and 30% of MNT-induced transcriptional
changes in each cell line respectively are regulated by MNT in a MAX-independent
manner (*Figure 7C*).

The genes showing expression changes in the MNT-depleted URMT and
URMax34 cells were analyzed with the GSEA and the MsigDB platform
(http://software.broadinstitute.org/gsea/msigdb). The genes regulated upon MNT knock-
down were compared with the gene sets derived from the biological process gene
ontology and the comparison showed that genes belonging to cell cycle processes are
the most enriched set in MNT-depleted cells both in MAX expressing and not expressing
cells (*Figure 7-figure supplement 7)*. This functional analysis of the gene transcriptional
signatures are consistent with the biological effects of MNT-depletion in cell proliferation.

Thus, in order to explain and confirm the effects of MNT ablation on UR61 cell
growth, we selected several genes involved in cell cycle and DNA replication which

expression was changed according to the RNA-seq analysis. URMT and URMax34 cells

were transfected with the two shMNT constructs and the empty vector following the same

conditions as the conditions used for the RNAseq experiment. The expression of the

selected genes was analyzed by RT-qPCR. *Figure 7D* shows that genes involved in cell

Figure 8. MNT and MLX interaction in UR61 cells. (A) URMT cells were transfected with a MLX expression vector or the empty vector (EV). 48 h after transfection, total cell lysates were prepared and precipitated with anti-MNT or anti-MLX antibodies, as well as unspecific IgG. The presence of MNT and MLX were detected in the immunoprecipitates by immunoblot. The results were reproduced in two experiments (B) URMax34 cells were transfected with a MLX expression vector and 48 h after transfection cells were treated with 100 µM Zn^{2+} for 24 h. Then total cell lysates were prepared and immunoprecipitated with anti-MNT. As a control, a mixture of lysates from cells treated and untreated with Zn^{2+} was immunoprecipitated with unspecific IgG. The presence of MNT, MLX and MAX in the immunoprecipitates were analysed by immunoblot. The results were reproduced in three experiments immunoprecipitations. (C) Schematic representation of the full-length (FL) and the deletion mutant ΔHLH MNT-HA used in subsequent experiments. The SID (Sin3 Interacting Domain), bHLH, LZ domains, HA tag and amino acids of the murine protein are indicated. (D) URMT cells were co-transfected with a vector expressing MLX and the constructs shown in (C) as indicated at the top. 48 h after transfection, cell lysates were prepared and the cells were immunoprecipitated with mouse anti-HA antibody to pull down the exogenous MNT proteins constructs. As a control, lysates (a mixture of lysates from cell transfected with FL MNT and ΔHLH MNT) were also immunoprecipitated with unspecific IgGs. The immunoblot analysis using a rat monoclonal anti-HA antibody revealed the presence of FL MNT and the smaller ΔHLH MNT The presence of MLX in the immunoprecipitates was analysed with an anti-MLX antibody. (E) MLX localization in UR61 cells. Nuclear ("N") and cytoplasmic ("C") extracts were prepared from URMT cells as described in Materials and Methods and the levels of MNT and MLX were determined in each fraction by immunoblot. SIN3B and RhoGDI (ARHGDIA) were used as nuclear and cytoplasmic markers respectively. The antibodies used are as described in the *Supplementary File 1*. A picture of the gel stained with Coomasie Blue after transference is also shown as an indicator of the total amount of proteins in the cytoplasmic and nuclear fraction. (F) Interaction between MNT and MLX in the nucleus and cytoplasm. URMT cells were transfected with a MLX expression vector and 48 h later nuclear and cytoplasmic extracts were prepared and immunoprecipitated with anti-MNT or anti-MLX antibodies. The levels of MNT and MLX in the immunoprecipitates were assayed by immunoblot. The results were reproduce in two independent experiments

cycle and DNA replication (*E2F7*, *CDKN1C*, *CDC6*, *ORC1*) and, DNA repair (*BRCA2*, *TRIP13*, *NEIL3*, *EXO1*, *PARPB*), were down-regulated upon silencing of *MNT* with the two shMNT constructs, except *CDKN1C* (p57, a cell cycle inhibitor) which was up-regulated. Altogether, the results are consistent with the negative effect of MNT depletion on cell proliferation in UR61 cells (*Figure 6*).

We then wondered how could MNT act as a transcription factor in cells lacking MAX. Besides MAX, MNT can also bind the HLH protein MLX *(Cairo et al 2001, Meroni et al 2000)*. To show if this interaction also takes place in UR61, we transfected MLX-Flag into URMT cells and 48 h later lysates were immunoprecipitated with anti-MLX antibody and anti-MNT antibodies. The immunoprecipitates were analysed by immunoblot and the results showed the interaction between MNT and MLX (*Figure 8A*). We next studied this interaction in URMax34 cells treated with Zn^{2+}, i.e. cells expressing MAX. The immunoblot results showed that MNT and MLX also interacted but the interaction was weaker when MAX expression was induced by Zn^{2+}. Thus, the data suggest that, at least in our experimental conditions, MNT-MAX dimers were formed more efficiently than MNT-MLX dimers (*Figure 8B*). To confirm the MNT-MLX interaction in URMT cells, we prepared a HA-tagged MNT mutant with a deletion of most of the HLH domain of mouse MNT, termed ΔHLHMNT-HA (*Figure 8C*). This mutant or the wild-type counterpart was transfected into URMT and the corresponding lysates were immunoprecipitated with anti-HA. The results showed that HLH domain is required for the MNT-MLX interaction (*Figure 8D*).

In contrast to MAX, MLX is expressed in both nucleus and cytoplasm *(Billin et al 2000)*. By immunoblot analysis it was found that MLX was present in nucleus and, at a lesser extent, in the cytoplasm of URMT cells (*Figure 8E*). Next, we prepared nuclear and cytoplasmic fractions of URMT cells and studied by immunoprecipitation with anti-MNT and anti-MLX antibodies if MNT and MLX interact. The results showed that this was indeed the case in both nuclear and cytoplasmic fractions (*Figure 8F*). The proximity ligation assay also showed that MNT and MLX interacted in both nucleus and cytoplasm (not shown).

Finally, it is described that MNT can form homodimers in two-hybrid experiments *(Hurlin et al 1997, Meroni et al 1997)*. Thus, a possibility is that MNT acts a transcription factor as a homodimer. This homodimerization has not been demonstrated *in vivo* so we tested it in HEK293T and UR61 cells. We first co-transfected HEK293T cells with GFP-MNT and Flag-MNT constructs and immunoprecipitated with anti-GFP antibody. The immunoblot analysis demonstrated the presence of the smaller Flag-MNT protein in the material immunoprecipitated with anti-GFP (*Figure 9A*). As expected, both big and small MNT forms were detected when the immunoblots were analysed with anti-MNT antibody

362   (*Figure 9A*). The results suggested the presence of homodimers between GFP-MNT and

363   MNT-Flag in HEK293T cells. As a control, the lysates with anti-MAX antibody were

364   imunoprecipitated, and both MNT and GFP-MNT were found to be bound to MAX (not

365   shown). Next, we investigated the MNT homodimerization in the UR61 system. URMT

Figure 9. MNT homodimerization in UR61 cells. (A) HEK293T cells were co-transfected with a GFP-MNT and MNT-Flag expression vectors, and 24 h later lysates were prepared and immunoprecipitated with anti-GFP antibody. The presence of MNT-GFP and MNT-Flag were revealed by immunoblotting the antibodies indicated at the right. As a control, lysates were precipitated with unspecific IgG. The antibodies used in each immunoblot are indicated at the right. The results were reproduced in two immunoprecipitations. (B) URMT cells were infected with lentivirus encoding a MNT-GFP and 72 h later were transfected with MNT-Flag and treated with 15 nM bortezomib for 12 h before harvesting. 48 h after transfection lysates were prepared and immunoprecipitated with anti-GFP antibody. The presence of MNT-GFP, MNT and MLX in the immunoprecipitates were assessed by immunoblot. The antibodies used in each immunoblot are indicated at the right. The results were reproduced in two experiments. (C) URMT cells were infected with lentivirus encoding a GFP-MNT and 72 h later transfected with expression vectors for full length (FL) MNT or ΔHLH MNT (Figure 8C) as indicated at the top of each lane. 48 h after transfection, cell lysates were prepared and immunoprecipitated with anti-HA to pull down the transfected MNT proteins. The levels of MNT-GFP and MNT were determined by immunoblot using anti-GFP and polyclonal anti-MNT antibody. (D) HEK293T cells were co-transfected with expression vectors for MNT-GFP and expression vectors for FL MNT or ΔHLH MNT (Figure 8C) as indicated at the top of each lane. 24 h later lysates were prepared and immunoprecipitated with anti-MNT antibody. The levels of MNT-GFP and MNT were determined by immunoblot using anti-GFP and polyclonal anti-MNT antibody. The results were reproduced in two experiments. (E) Schematic summary of the interactions of MNT found in URMT (MAX-less) and URMax34 (expressing MAX when treated with Zn^{2+}).

cells were infected with lentiviral particles containing the GFP-MNT construct, immunoprecipitated, and the lysates analysed by immunoblot with anti-GFP antibody and anti-MNT antibodies. The results showed that endogenous MNT was present in the immunoprecipitates with anti-GFP (*Figure 9B*). The results suggest that MNT forms homodimers in human HEK293T cells and in MAX-deficient rat URMT cells. As MNT dimerization in yeast two hybrid assays depends on the HLH-LZ *(Hurlin et al 1997, Meroni et al 1997)*, we asked whether HLH was involved in the homodimerization *in vivo*. We transfected the MNT mutant lacking the bHLH region (ΔbHLH MNT) (*Figure 8C*) as well as the wild-type form and we carried out immunoprecipitations with anti-HA antibodies which should only recognize the exogenous proteins. The immunoblot analysis of the immunoprecipitates revealed that ΔbHLH MNT was unable to bind GFP-MNT (*Figure 9C*). The same experiment was performed in URMT cells and the same result was observed, i.e, GFP-MNT bound to wild-type MNT-HA but not to the ΔbHLH form (*Figure 9D*). We conclude MNT homodimerize through the HLH domain, as expected. Altogether, the results indicate that in URMT cells, MNT can form homodimers or heterodimers with MLX, whereas in URMax34 (in the presence of Zn^{2+}), MNT can also form heterodimers with MAX, as schematized in *Figure 9E*.

DISCUSSION

In this study, we report several novel findings: (i) MNT is required for optimum proliferation even in the absence of MAX; (ii) in the absence of MAX, MNT expression is elevated and a significant fraction localizes in the cytoplasm; (iii) MNT represses its own transcription in a MAX-dependent manner; (iv) MNT is able to regulate the expression of genes in the absence of MAX and (v) MNT forms homodimers in vivo. Previous reports suggest that MNT functions as a "MYC buffer", curbing excessive MYC activity. For instance, *in vivo* MNT knockdown impedes MYC-driven lymphomagenesis *(Campbell et al 2017)* whereas MNT silencing leads to MYC-like phenotypes *(Hurlin et al 2003, Nilsson et al 2004, Walker et al 2005)*. Our data are in line with this critical MNT function, as they show a tight control of MNT expression by which MNT limits its own mRNA expression. MNT protein levels are also under strict control in UR61 cells, as MNT cannot be overexpressed at high levels upon transfection unless the proteasome is inhibited. Given the relevance of MNT to modulate MYC activity and its central position between the MYC-MAX and MLX-MONDO networks, the activities and regulation of MNT are a key issue on MYC-dependent oncogenesis.

MNT expression was high in some MAX-deficient cells like the rat UR61 cells and human lung carcinoma H1417 cells. The absence of MAX was in part responsible for this

effect because (i) MAX re-expression in UR61cells results in decreased MNT mRNA and

protein, and (ii) MAX overexpression in K562 cells results in MNT down-regulation

whereas MAX knock-down results in MNT up-regulation. The mechanism for this MAX

effect on MNT levels depended, at least partially, on the auto-repression of MNT

expression by MNT-MAX dimers. Luciferase reporter experiments showed that MNT

represses its own promoter (which contains two conserved E-boxes) and ChIP assays

showed that MNT binds to its own promoter when MAX is ectopically expressed in UR61

cells. In agreement with these results, significant levels of MNT localize to the cytoplasm

of MAX-deficient cells whereas MNT remains nuclear in MAX-expressing cells. This

would explain why MAX re-expression leads to a decrease in MNT levels: MNT-MAX

dimers would be formed to repress MNT expression in the nucleus. On the contrary, in

the absence of MAX, MNT is expressed at higher levels since there would be no negative

MNT autoregulation. The model is schematized in *Figure 10*.

We explored possible MAX-independent effects of MNT using the UR61 model.

Although originally MAX was defined as an obligate dimerization partner of MYC, work

carried out in the PC12 model and in *Drosophila* indicated that MYC can function in a

MAX-independent manner, for example in inhibition of differentiation *(Gallant 2013,*

Maruyama et al 1987, Vaque et al 2008). MYC overexpression blocks RAS-induced

differentiation of UR61 cells *(Vaque et al 2008)* however, we did not detect a significant

effect of MNT depletion on differentiation (not shown). In contrast, depletion of MNT in

MAX-deficient UR61 cells impairs cell proliferation. This is the first report on a MAX-

independent function of a MXD protein. In contrast, in rat and mouse fibroblasts, MNT

knock-out leads to increased proliferation and transformation capacities *(Hurlin et al*

2003, Nilsson et al 2004). The fact that MNT depletion impairs UR61 proliferation (in a

MAX-independent manner) adds complexity to the MNT-MYC functional interactions.

Actually, in gastric cancer high MNT expression correlates with shorter survival whereas

MYC overexpression has the opposite effect (http://kmplot.com/analysis/).

Since MNT is a transcription factor, we compared the transcriptomes of parental

UR61 cells versus cells with depleted MNT. Upon MNT depletion, a number of genes

involved in cell cycle progression and DNA damage response were downregulated in

cells with and without MAX. This is concordant with the decrease in proliferation exerted

by MNT silencing in UR61. It is open to discussion how MNT regulates genes in the

absence of MAX. MNT can form heterodimers with the HLH protein MLX *(Cairo et al*

2001, Meroni et al 2000) and we have confirmed the ability of MNT to bind MLX in UR61

cells, but when MAX is expressed, the dimers MNT-MAX are favoured versus MNT-MLX.

As a relevant fraction of MLX is cytoplasmic and we have shown that MNT and MLX also

interact in the cytoplasm, this interaction could help to explain the increase in cytoplasmic MNT observed in MAX-depleted cells. The model is depicted in the *Figure 10*. Moreover, we have shown that MNT homodimerizes in UR61 cells, and therefore it is likely that MNT regulates genes as a homodimer. Further work is required to clarify this point. The results reported here show a strict autoregulation of MNT, supporting a pivotal role of MNT in the control of cell proliferation even in the absence of MAX. Moreover, we showed that MNT has activities in the absence of MAX. Given that MNT participates in both the MAX-MYC and MLX-MONDO networks, the regulation of MNT is a relevant issue in cell biology and in tumor development promoted by MYC.

Figure 10. Model of MNT fates depending on MAX. In MAX-expressing cells, most of the MNT is retained in the cell nucleus where it limits its own expression. In MAX-deficient cells as UR61, MNT is distributed in nucleus and cytoplasm and is unable to bind the promoter and regulate its transcriptional activity. The model includes the presence of MNT-MNT homodimers and the interaction MNT-MLX in the cytoplasm, which might be responsible for the MNT partial localization in the cytoplasm in MAX-deficient cells.

MATERIALS AND METHODS

Cell lines and transfections

Cell lines were obtained from ATCC and grown in either RPMI-1640 or DMEM (Lonza) supplemented with 10% fetal bovine serum (Lonza), 150 µg/ml of gentamicin and 2 µg/ml of ciprofloxacin. To generate the URMT and URMax34 cell line, UR61 cells *(Guerrero et al 1988)* were electroporated (260 V, 1 mFa, BioRad apparatus) with a pHeBo-MT or pHeBo-MT-Max vector which carries a human MAX cDNA under the control of metallothionein promoter *(Canelles et al 1997)*. Cells were selected with 0.2 mg/ml of hygromycin (Life Technologies) and cell clones were isolated by limiting dilution Transient transfections were carried out using the Ingenio Electroporation solution (Mirus) in an Amaxa nucleofector. The cells were transfected or transduced with expression vectors for MAX (pCEFL-MAX*(Mauleon et al 2004)*, MLX (β isoform, (pMS18-

MLX,*(Billin et al 1999)*, human MNT (pCMVSport6-MNT, Origene Technologies), MNT-Flag (human MNT with FLAG at the C-ter lentiviral Lv158 vector Genecopoeia), MNT-GFP (human MNT with GFP at the N-ter, lentiviral Lv103 vector Genecopiea,), FL MNT-HA (full-length murine MNT tagged at the C-ter with hemagglutinin epitope, HA), ΔHLH MNT-HA (murine MNT carrying a deletion of amino acids 221-272 amino acids and tagged with HA) (both in pcDNA3 with the zeocin resistance gene inserted), short-hairpin human MNT-1 (shMNT-1, pLKO-shMNT from Sigma Mission, TRCN0000085733), or MNT-2 (shMNT-2, pLKO-shMNT from Sigma Mission, TRCN0000235815), siRNA for human MAX (Sigma, SASI_Hs01_00011941).

Immunoblot and immunoprecipitation

Cell lysis, immunoblots and immunoprecipitations were performed as described.*(Garcia-Sanz et al 2014)* .The antibodies used are described in *Supplementary File 1.*

Luciferase reporters and assays

To generate the 850MNT-luc, 220MNT-luc, 570MNT-luc reporter vectors, two pair of primers were designed for each construct (*Supplementary File 2*), targeting sequences of the human genome corresponding to 850 bp upstream the transcription start site (TSS) of *MNT* gene. The amplified DNA was inserted into the pBV-luc reporter vector at the EcoRV and HindIII sites *(He et al 1998)*. The sequence of MNT promoter was from the UCSC genome browser (http://genome.ucsc.edu/; GRCh37/hg19). Cells were transfected with jetPEI reagent (Polyplus) and luciferase assays were performed as described *(Garcia-Sanz et al 2014)*.

Cell proliferation and cell cycle analysis

Cell proliferation was monitored with a cell counter (NucleoCounter NC-100, Chemometec) or a cytometer (Guava PCA, Merck Millipore). For the clonogenic assays, 1-2 x 10^6 cells/ml were seeded in a 6-well or 60 mm plate after transfection by electroporation. 48 h post-transfection, cells were selected with puromycin at 0.2-0.5 µg/ml final concentration. After 8-17 days the cells were stained with crystal violet and the dye measured by absorbance at 595 nm as described *(Ferrandiz et al 2009)*. To determine the sugG0-G1 population, cells were transfected and 7 days post-transfection the DNA concentration was analyzed by propidium iodide staining and flow cytometry as described *(Albajar et al 2011)*.

RNA analysis

For qPCR, total RNA was isolated using the TRI Reagent® Solution (Invitrogen). cDNA was generated by reverse transcription (RT) using the iScript (Bio-Rad). Quantitative polymerase chain reaction (qPCR) was performed with specific primers (*Supplementary File 2*) using the iTaq™ Universal SYBR® Green Supermix (Bio-Rad) and CFX ConnectTM Real-Time PCR Detection System (Bio-Rad). RNA was converted into cDNA and analysed as described *(Bretones et al 2011)*. Levels of mRNA were normalized against actin and RPS14 mRNA levels. For RNA-seq, total RNA was isolated with the RNeasy kit (Qiagen) from two independent experiments of MNT silencing in URMT and URMax34 cells, both treated with 100 μM $ZnSO_4$ for 24 h. mRNA libraries were prepared using the Illumina TruSeq RNA Sample Prep Kit v2 (kit RS-122-2002, Illumina). A minimum of 40 million 50 bases single-end reads per sample were obtained. Tophat algorithm *(Trapnell et al 2009)* was used to align the data using a set of gene model annotations and/or known transcripts of rat genome (rn4) obtained from RefSeq database. Cufflinks software *(Trapnell et al 2010)* was run to estimate transcripts abundances represented in RPKM units (reads per kilobase per million reads) as described (Mortazavi et al 2008). The gene expression of the genes are represented as RPKM values (*Figure 8 –Supplement 2 to 5)*.

Chromatin immunoprecipitation

Fractionation of total cell extracts into cytoplasmic and nuclear fractions and chromatin immunoprecipitation (ChIP) were performed essentially as described *(Garcia-Sanz et al 2014)*. Immunoprecipitated DNA was purified with the QIAquick PCR Purification Kit (Qiagen) and analysed by qPCR. The antibodies and primers used are described in *Supplementary Files 1 and 2,* respectively.

Nuclear/cytoplasmic fractionation

Cytoplasmic and nuclear extracts were prepared essentially as described *(Conacci-Sorrell et al 2010)*. The hypotonic buffer to lyse cells was 10 mM HEPES pH 7, 10 mM KCl, 0.25 mM EDTA pH 8, 0.125 mM EGTA pH 8, 0.5 mM spermidin, 0.1% NP40, 1 mM DTT and proteases inhibitors The hypertonic buffer for nuclear extracts was 20 mM HEPES, 400 mM NaCl, 0.25 mM EDTA, 1.5 mM $MgCl_2$, 0.5 mM DTT and proteases inhibitors.

Statistical analysis

Student's t-test was used to evaluate the significance of differences between control and experimental groups. A p-value of less than 0.05 was considered as significant. The threshold for expression changes in the RNA-seq analysis was set as log2(RPKM fold change) ≥ 0.7 or ≤ -0.7 and a p-value <0.1.

Bioinformatic analysis

Genomic sequences of 1000 bp upstream TSS of human, mouse and rat MNT genes were obtained from *H. sapiens* (GRCh37/hg19), *M. musculus* (GRCm38/mm10) and *R. norvegicus* (RGSC 5.0/rn5) assemblies. For the ChIP-seq ENCODE analysis, the data for MAX, MYC and MXI1 was obtained for the indicated cell lines. The SYDH ENCODE project was chosen for representation of the binding peaks in the promoter of the human MNT gene. The sequences corresponding to those peaks were analysed confirming the presence of the E-boxes identified in the human MNT promoter

Source data: the RNA-seq data is loaded in the European Nucleotide Archive with the accession number PRJEB23604

ACKNOWLEDGEMENTS

We are grateful to Montserrat Sánchez-Céspedes and Don Ayer for cell lines and constructs, Maralice Conacci-Sorrel for comments on the manuscript, and Sandra Zunzunegui and Maria Aramburu for technical help.

FUNDING

The work was supported by grant SAF2014-53526-R from MINECO, Spanish Government, to JL and MDD (partially funded by FEDER program from European Union) RNE is supported by grant CA57138/CA/NCI from National Institutes of Health. PJH is supported by grants from Shriners Hospitals for Children. MCL-N and JL-P were recipients of F.P.U. fellowships from Spanish Government.

CONFLICT OF INTEREST

The authors declare no conflict of interests

AUTHOR ORCID

Javier León, https://orcid.org/0000-0001-5803-0112

REFERENCES

Albajar M, Gomez-Casares MT, Llorca J, Mauleon I, Vaque JP, Acosta JC, Bermudez
A, Donato N, Delgado MD, Leon J. 2011. MYC in chronic myeloid leukemia:
induction of aberrant DNA synthesis and association with poor response to
imatinib. *Mol Cancer Res* **9**: 564-76
Billin AN, Eilers AL, Coulter KL, Logan JS, Ayer DE. 2000. MondoA, a novel basic
helix-loop-helix-leucine zipper transcriptional activator that constitutes a positive
branch of a max-like network. *Mol Cell Biol* **20**: 8845-54
Billin AN, Eilers AL, Queva C, Ayer DE. 1999. Mlx, a novel Max-like BHLHZip protein
that interacts with the Max network of transcription factors. *J Biol Chem* **274**:
36344-50
Bretones G, Acosta JC, Caraballo JM, Ferrandiz N, Gomez-Casares MT, Albajar M,
Blanco R, Ruiz P, Hung WC, Albero MP, Perez-Roger I, Leon J. 2011. SKP2
oncogene is a direct MYC target gene and MYC down-regulates p27(KIP1)
through SKP2 in human leukemia cells. *J Biol Chem* **286**: 9815-25
Cairo S, Merla G, Urbinati F, Ballabio A, Reymond A. 2001. WBSCR14, a gene
mapping to the Williams--Beuren syndrome deleted region, is a new member of
the Mlx transcription factor network. *Hum Mol Genet* **10**: 617-27
Campbell KJ, Vandenberg CJ, Anstee NS, Hurlin PJ, Cory S. 2017. Mnt modulates
Myc-driven lymphomagenesis. *Cell Death Differ* **24**: 2117-26
Canelles M, Delgado MD, Hyland KM, Lerga A, Richard C, Dang CV, Leon J. 1997.
Max and inhibitory c-Myc mutants induce erythroid differentiation and resistance
to apoptosis in human myeloid leukemia cells. *Oncogene* **14**: 1315-27
Conacci-Sorrell M, McFerrin L, Eisenman RN. 2014. An Overview of MYC and Its
Interactome. *Cold Spring Harb Perspect Med* **4**: 1-24
Conacci-Sorrell M, Ngouenet C, Eisenman RN. 2010. Myc-nick: a cytoplasmic
cleavage product of Myc that promotes alpha-tubulin acetylation and cell
differentiation. *Cell* **142**: 480-93
Dang CV. 2012. MYC on the path to cancer. *Cell* **149**: 22-35
Dezfouli S, Bakke A, Huang J, Wynshaw-Boris A, Hurlin PJ. 2006. Inflammatory
disease and lymphomagenesis caused by deletion of the Myc antagonist Mnt in
T cells. *Mol Cell Biol* **26**: 2080-92
Diolaiti D, McFerrin L, Carroll PA, Eisenman RN. 2015. Functional interactions among
members of the MAX and MLX transcriptional network during oncogenesis.
Biochim Biophys Acta **1849**: 484-500
Ferrandiz N, Martin-Perez J, Blanco R, Donertas D, Weber A, Eilers M, Dotto P,
Delgado MD, Leon J. 2009. HCT116 cells deficient in p21(Waf1) are
hypersensitive to tyrosine kinase inhibitors and adriamycin through a
mechanism unrelated to p21 and dependent on p53. *DNA Repair* **8**: 390-9
Foley KP, McArthur GA, Queva C, Hurlin PJ, Soriano P, Eisenman RN. 1998. Targeted
disruption of the MYC antagonist MAD1 inhibits cell cycle exit during
granulocyte differentiation. *EMBO J* **17**: 774-85
Gallant P. 2013. Myc function in Drosophila. *Cold Spring Harb Perspect Med* **3**:
a014324
Garcia-Sanz P, Quintanilla A, Lafita MC, Moreno-Bueno G, Garcia-Gutierrez L, Tabor
V, Varela I, Shiio Y, Larsson LG, Portillo F, Leon J. 2014. Sin3b interacts with
myc and decreases myc levels. *J Biol Chem* **289**: 22221-36
Guerrero I, Pellicer A, Burstein DE. 1988. Dissociation of c-fos from ODC expression
and neuronal differentiation in a PC12 subline stably transfected with an
inducible N-ras oncogene. *Biochem Biophys Res Commun* **150**: 1185-92
He TC, Sparks AB, Rago C, Hermeking H, Zawel L, da Costa LT, Morin PJ, Vogelstein
B, Kinzler KW. 1998. Identification of c-MYC as a target of the APC pathway.
Science **281**: 1509-12

Hopewell R, Ziff EB. 1995. The nerve growth factor-responsive PC12 cell line does not
express the Myc dimerization partner Max. *Mol Cell Biol* **15**: 3470-8
Hurlin PJ, Queva C, Eisenman RN. 1997. Mnt, a novel Max-interacting protein is
coexpressed with Myc in proliferating cells and mediates repression at Myc
binding sites. *Genes Dev* **11**: 44-58.
Hurlin PJ, Zhou ZQ, Toyo-oka K, Ota S, Walker WL, Hirotsune S, Wynshaw-Boris A.
2003. Deletion of Mnt leads to disrupted cell cycle control and tumorigenesis.
EMBO J **22**: 4584-96
Lafita-Navarro MC, Blanco R, Mata-Garrido J, Liano-Pons J, Tapia O, Garcia-Gutierrez
L, Garcia-Alegria E, Berciano MT, Lafarga M, Leon J. 2016. MXD1 localizes in
the nucleolus, binds UBF and impairs rRNA synthesis. *Oncotarget* **7**: 69536-48
Maruyama K, Schiavi SC, Huse W, Johnson GL, Ruley HE. 1987. myc and E1A
oncogenes alter the responses of PC12 cells to nerve growth factor and block
differentiation. *Oncogene* **1**: 361-7
Mauleon I, Lombard MN, Munoz-Alonso MJ, Canelles M, Leon J. 2004. Kinetics of
myc-max-mad gene expression during hepatocyte proliferation in vivo:
Differential regulation of mad family and stress-mediated induction of c-myc.
Mol Carcinog **39**: 85-90
Meroni G, Cairo S, Merla G, Messali S, Brent R, Ballabio A, Reymond A. 2000. Mlx, a
new Max-like bHLHZip family member: the center stage of a novel transcription
factors regulatory pathway? *Oncogene* **19**: 3266-77
Meroni G, Reymond A, Alcalay M, Borsani G, Tanigami A, Tonlorenzi R, Nigro CL,
Messali S, Zollo M, Ledbetter DH, Brent R, Ballabio A, Carrozzo R. 1997. Rox,
a novel bHLHZip protein expressed in quiescent cells that heterodimerizes with
Max, binds a non-canonical E box and acts as a transcriptional repressor.
EMBO J **16**: 2892-906.
Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. 2008. Mapping and
quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* **5**: 621-8
Nilsson JA, Maclean KH, Keller UB, Pendeville H, Baudino TA, Cleveland JL. 2004.
Mnt loss triggers Myc transcription targets, proliferation, apoptosis, and
transformation. *Mol Cell Biol* **24**: 1560-9
Popov N, Wahlstrom T, Hurlin PJ, Henriksson M. 2005. Mnt transcriptional repressor is
functionally regulated during cell cycle progression. *Oncogene* **24**: 8326-37
Queva C, McArthur GA, Ramos LS, Eisenman RN. 1999. Dwarfism and dysregulated
proliferation in mice overexpressing the MYC antagonist MAD1. *Cell Growth*
Differ **10**: 785-96
Romero OA, Torres-Diz M, Pros E, Savola S, Gomez A, Moran S, Saez C, Iwakawa R,
Villanueva A, Montuenga LM, Kohno T, Yokota J, Sanchez-Cespedes M. 2014.
MAX inactivation in small cell lung cancer disrupts MYC-SWI/SNF programs
and is synthetic lethal with BRG1. *Cancer Discov* **4**: 292-303
Schreiber-Agus N, DePinho RA. 1998. Repression by the Mad(Mxi1)-Sin3 complex.
Bioessays **20**: 808-18
Terragni J, Nayak G, Banerjee S, Medrano JL, Graham JR, Brennan JF, Sepulveda S,
Cooper GM. 2011. The E-box binding factors Max/Mnt, MITF, and USF1 act
coordinately with FoxO to regulate expression of proapoptotic and cell cycle
control genes by phosphatidylinositol 3-kinase/Akt/glycogen synthase kinase 3
signaling. *J Biol Chem* **286**: 36215-27
Toyo-oka K, Hirotsune S, Gambello MJ, Zhou ZQ, Olson L, Rosenfeld MG, Eisenman
R, Hurlin P, Wynshaw-Boris A. 2004. Loss of the Max-interacting protein Mnt in
mice results in decreased viability, defective embryonic growth and craniofacial
defects: relevance to Miller-Dieker syndrome. *Hum Mol Genet* **13**: 1057-67
Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with
RNA-Seq. *Bioinformatics* **25**: 1105-11
Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL,
Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq

reveals unannotated transcripts and isoform switching during cell differentiation.
Nat Biotechnol **28**: 511-5
Vaque JP, Fernandez-Garcia B, Garcia-Sanz P, Ferrandiz N, Bretones G, Calvo F,
Crespo P, Marin MC, Leon J. 2008. c-Myc inhibits Ras-mediated differentiation
of pheochromocytoma cells by blocking c-Jun up-regulation. *Mol Cancer Res* **6**:
325-39
Walker W, Zhou ZQ, Ota S, Wynshaw-Boris A, Hurlin PJ. 2005. Mnt-Max to Myc-Max
complex switching regulates cell cycle entry. *J Cell Biol* **169**: 405-13
Yang G, Hurlin PJ. 2017. MNT and Emerging Concepts of MNT-MYC Antagonism.
Genes 8: pii: E83