The phylogeography of westslope cutthroat trout

Michael K. Young[1*], Kevin S. McKelvey[1], Tara Jennings[2], Katie Carter[2,4], Richard Cronn[2],

Ernest R. Keeley[3], Janet L. Loxterman[3], Kristine L. Pilgrim[1], and Michael K. Schwartz[1]

[1]U.S. Forest Service, National Genomics Center for Wildlife and Fish Conservation, Rocky

Mountain Research Station, 800 E. Beckwith Avenue, Missoula, MT 59801 USA

[2]U.S. Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis,

OR 97331 USA

[3]Department of Biological Sciences, Idaho State University, 921 S. 8th Avenue, Mail Stop 8007,

Pocatello, ID 83209 USA

[4]Current address: Department of Horticulture, Oregon State University, Corvallis, OR 97331

USA

*Corresponding author: mkyoung@fs.fed.us; 1.406.542.3254

Keywords: distinct population segments, phylogeny, evolutionarily significant units,

microsatellites, mitochondrial, mtDNA, *Oncorhynchus clarkii lewisi*, SNP

Abstract.-Identifying units of conservation of aquatic species is fundamental to informed natural

resources science and management. We used a combination of mitochondrial and nuclear

molecular methods to identify potential units of conservation of westslope cutthroat trout, a

taxon native to montane river basins of the northwestern U.S. and southwestern Canada.

Mitogenomic sequencing identified two major lineages composed of nine monophyletic clades,

and a well-supported subclade within one of these, largely delineated by river basins. Analyses

of microsatellites and single nucleotide polymorphisms corroborated most of these groupings,

sometimes with less resolution but demonstrating more complex connections among clades. The

mitochondrial and nuclear analyses revealed that Pleistocene glacial cycles profoundly

influenced the distribution and divergence of westslope cutthroat trout, that this taxon crossed the

Continental Divide in two separate events, and that genetically pure but nonindigenous fish were

widely distributed. Herein, we recognize nine geographically discrete, cytonuclear lineages

largely circumscribed by major river basins as potential units of conservation: 1) John Day; 2)

Coeur d'Alene; 3) St. Joe; 4) North Fork Clearwater; 5) Salmon; 6) Clearwater headwaters; 7)

Clearwater-eastern Cascades; 8) neoboreal, consisting of most of the Columbia upstream from

central Washington, the Fraser in British Columbia, and the South Saskatchewan in Alberta; and

9) Missouri.

───────────────────────────────────────────────

If the biota, in the course of aeons, has built something we like but do not understand, then who

but a fool would discard seemingly useless parts? To keep every cog and wheel is the first

precaution of intelligent tinkering.

45                                                              —Aldo Leopold (1953)

───────────────────────────────────────────────

Leopold's (1953) admonition has long served as a core paradigm of conservation biology. In

his time, the focus was on preserving species, but species often comprise distinct lineages that go

by many names—subspecies, distinct population segments, evolutionarily significant units,

operational taxonomic units, or stocks—and constitute potential units of conservation (Avise

2000) that can be accorded taxonomic or legal standing, e.g., under the U.S. Endangered Species

Act (Waples 1991) or Canada's Species at Risk Act (Mee et al. 2015). Historically, evidence for

the presence of these lineages was largely based on variation in morphology, behavior, life

history, or ecology, usually associated with a geographically discrete distribution, but

increasingly sophisticated molecular tools have revolutionized their identification and spawned a

branch of science dedicated to their study, phylogeography (Avise 2000). There is particular

urgency to identify units of conservation for aquatic ectotherms such as fishes, mussels, crayfish,

and amphibians. These aquatic groups are disproportionately represented among at-risk taxa

(Williams et al. 2011), and shifts in climate are beginning to alter their distributions (Comte and

Grenouillet 2013; Eby et al. 2014). Correctly inferring the existence and extent of lineages within

a taxon also relies on an understanding of the geological and climatic histories of current and

former habitats. For aquatic species, especially fishes in western North America, this means

appreciating the variation in hydrological connections over evolutionary timescales (McPhail and

Lindsey 1986; Minckley et al. 1986), particularly with respect to the influence of Pleistocene

glaciations on hydrologic networks (Bernatchez and Wilson 1998). A robust intraspecific

phylogeny also demands that a taxon be sampled across its entire range with sufficient intensity

to represent all potential units of conservation, and be examined with tools of sufficient

sensitivity to recognize them (Heath et al. 2008). It also requires awareness of the potential for

human-assisted relocation of nonlocal fish populations (Metcalf et al. 2012; Pritchard et al.

2015).

The systematics and distribution of cutthroat trout (*Oncorhynchus clarkii*) have been a source

of contention since the species was first formally described in the 19[th] century (Richardson

1836). Various forms have been described as full species or as subspecies with little regard to

previous work (Thorgaard et al., this volume), and the origin and affiliation of some type

specimens has been obscure (Metcalf et al. 2012). Moreover, our understanding of the historical

distributions of putative taxa is far from perfect, in part because of the broad-scale husbandry and

translocation of some forms of cutthroat trout (Wiltzius 1985) and the rapid extirpation of others

(Young and Harig 2001). A valiant attempt to resolve this ambiguity was made by the

preeminent taxonomist of the 20[th] century on western North American trout, Robert Behnke,

who divided cutthroat trout into 14 extant and 2 extinct subspecies (Behnke 1992). Among these

subspecies was the westslope cutthroat trout *O. c. lewisi*, first observed by Lewis and Clark in

1805 at the eventual type location, the Great Falls of the Missouri River (Girard 1856). Primarily

relying on morphological evidence, Behnke (1992) asserted that the cutthroat trout of the interior

Columbia River basin also represented this subspecies, an observation broadly corroborated by

phylogenetic work (Leary et al. 1987; Wilson and Turner 2009; Loxterman and Keeley 2012;

Houston et al. 2012). Whether there is substantial phylogenetic diversity within this taxon across

its range, however, has not been resolved. Because of the absence of substantial morphological

differences among westslope cutthroat trout in different river basins, the U.S. Fish and Wildlife

Service concluded that the taxon constituted a single unit of conservation in response to a

petition for listing under the U.S. Endangered Species Act (U.S. Fish and Wildlife Service 2003).

Yet subsequent molecular studies indicated the presence of marked geographic divergence

within westslope cutthroat trout (Drinan et al. 2011; Loxterman and Keeley 2012) that was likely

driven by the complex geological and climatic history of the region (McPhail and Lindsey 1986).

Although Behnke (1992) was uncertain about the ultimate source of the lineage that led to

westslope cutthroat trout, he posited that most of its present distribution could be attributed to

events during the Wisconsinan glaciation 100–15 ka ago. He proposed that major waterfalls

developed during this time in the Pend Oreille, Spokane, and Kootenai River basins that may

have isolated westslope cutthroat trout upstream. Thereafter, the Clark Fork River basin served

as the source for fish dispersing—across divides via stream capture—to the east and north to the

upper Missouri River and South Saskatchewan River basins in Montana, Wyoming, and Alberta

and to the south to the Clearwater and Salmon River basins in Idaho. He postulated that outwash

floods from Glacial Lake Missoula carried westslope cutthroat trout to disjunct river basins along

the eastern face of the Cascade Range in Washington and to the John Day River basin in Oregon,

and may have been an alternate route for populations that colonized the Clearwater and Salmon

River basins. Recession of continental glaciers, along with the temporary proglacial lakes at their

trailing edges, presumably permitted westslope cutthroat trout to colonize waters farther north in

western Montana, northern Idaho, southern British Columbia, and southern Alberta.

Our goal was to test this phylogeographic hypothesis (cf. Crisp et al. 2011). We did so by

assessing the phylogenetic structure of westslope cutthroat trout across the bulk of its historical

distribution, with an additional aim to identify and geographically delineate potential units of

conservation, using a combination of mitochondrial and nuclear molecular information on

different samples of westslope cutthroat trout, including those used to establish hatchery

broodstocks.

**Methods**

*The historical range and its geological history*

The historical range of westslope cutthroat trout is thought to constitute portions of the Fraser,

South Saskatchewan, Missouri, and Columbia River basins (Shepard et al. 2005; McPhail 2007;

Figure 1). Its presence in the Fraser River basin in British Columbia is limited to a few

headwater streams (McPhail 2007), whereas it once occupied most of the tributaries in the Bow

River and Oldman River basins, part of the South Saskatchewan River basin in Alberta (Yau and

Taylor 2013; Fisheries and Oceans Canada 2014). In the Missouri River basin, its range was

thought to include the main stem and most tributaries from the Judith River to the headwaters,

excluding those blocked by waterfalls (e.g., the Sun River basin above present-day Gibson

Dam). In the upper Columbia River basin, the historical range includes 1) the entire Pend Oreille

and Kootenai River basins in British Columbia, Idaho, Montana, and Washington; 2) scattered

populations in the Kettle River basin in British Columbia (McPhail 2007) and presumably

Washington; 3) the Spokane River basin above Spokane Falls, primarily in Idaho, and perhaps

the Sanpoil River basin in Washington (Trotter et al. 2001b). Along the eastern Cascades,

westslope cutthroat trout are found from the Methow River basin in the north to the Yakima

River basin in the south, although there is uncertainty with respect to whether it is indigenous to

these and all intervening basins (Trotter et al. 1999, 2001a). In Oregon, westslope cutthroat trout

are present in tributaries of the main-stem and North Fork John Day River, but believed to be

introduced to the latter (Gunckel 2002). Westslope cutthroat trout are also native to the bulk of

two large river basins in the Snake River in Idaho, the Clearwater and Salmon, but are curiously

absent from all other basins tributary to the Snake River in Oregon, Washington, and Idaho

(except where introduced; Neville and Dunham 2011). Although the four major basins are

presently hydrologically isolated, they were presumably connected in the distant or recent past.

For example, the upper Columbia and Missouri Rivers were connected by the dual outlets of a

lake at Marias Pass in Montana until 1913, when the Union Pacific Railroad diverted all flow to

a tributary of the Two Medicine River of the Missouri River basin (Schultz 1941).

The modern distribution has been substantially reduced because of habitat alteration and

introductions of nonnative species (Shepard et al. 2005). Moreover, hatchery strains of westslope

cutthroat trout have been widely introduced throughout the 20th century and could affect both the

phylogenetic interpretation and evolutionary integrity of wild populations. Three broodstocks are

commonly used at present. In Montana, the primary broodstock was developed from fish in the

upper Flathead and middle Clark Fork River basins (Drinan 2010). In Idaho, westslope cutthroat

trout were domesticated from populations in the Priest River basin, as was the King's Lake

broodstock in Washington (Crawford 1979; M. Campbell, Idaho Department of Fish and Game,

personal communication). A second broodstock in Washington, Twin Lakes, was presumably

developed from fish collected in the Lake Chelan basin or Methow or Wenatchee River basins

(Crawford 1979).

The interior Columbia River basin constitutes the western portion of the distribution of

westslope cutthroat trout and has been present in some form for at least 15 Ma, roughly the age

of the lineage representing cutthroat trout and rainbow trout *O. mykiss* (Stearley and Smith

2016). The eruption and deposition of flood basalts from the mid-Miocene to the Pliocene led to

repeated re-routing and re-entrenchment of the major tributaries (Fecht et al. 1985). This

coincided with the orogeny of the Cascade Range, whereas the topography of the headwaters

was set by the orogeny of the Rocky Mountains 60 Ma earlier (Alt and Hyndman 1995). The

Pleistocene was characterized by the cyclic advance and retreat of the Cordilleran and Laurentide

ice sheets, which buried northerly portions of the current range of westslope cutthroat trout under

hundreds of meters of ice. The repeated advance of the Purcell lobe of the Cordilleran ice sheet

near Sandpoint, Idaho also led to the damming of the Clark Fork River and filling of Glacial

Lake Missoula (Waitt 1985). The repeated failure of this ice dam led to catastrophic jökulhlaups

lasting 1–2 weeks in which the discharge from the draining glacial lake temporarily exceeded

that of all rivers on Earth combined. These glacial outburst floods, perhaps up to 100 at multi-

decadal intervals during the Wisconsinan glaciation (Booth et al. 2003) and an unknown number

from earlier Pleistocene glacial intervals (Bader et al. 2016), produced the Columbia Basin

scablands. The scablands are characterized by floodwater-scoured surfaces, deep sedimentary

deposits from temporary ponding as water backed up at topographic chokepoints, e.g., Wallula

Gap in Washington, and huge nickpoints in many river channels that resulted in waterfalls likely

to constitute migration barriers to aquatic taxa, e.g., Spokane Falls on the Spokane River,

Palouse Falls on the Palouse River, Metalline Falls on the Pend Oreille River, and falls on many

lesser basins (Waitt 1985). The Okanagan and Columbia River lobes of this ice sheet also led to

the formation of Glacial Lake Columbia, which at its highstand extended upstream to the Purcell

lobe and portions of the Coeur d'Alene and St. Joe River valleys (Atwater 1987; Hanson and

Clague 2016). Contemporaneous mountain glaciers (Pierce 2003) would have forced headwater

taxa to lower portions of many watersheds to escape habitats that were cold, turbid, and

unproductive. Similar conditions—rerouted and glacially derived streams and temporary glacial

lakes—prevailed in the upper Missouri River basin. Until the Pleistocene (and perhaps during

portions of it), the Missouri River drained northward to Hudson Bay (Howard 1958). By the Last

Glacial Maximum 23–19 ka ago (Hughes et al. 2013), the Laurentide ice sheet had forced the

Missouri River south to near its present course within the Mississippi River basin. As that ice

sheet retreated, it created a series of proglacial lakes (Alden 1932; Colton et al. 1961) that

persisted as recently as 11.5 ka ago (Davis et al. 2006) and may have served as springboards for

colonization by aquatic taxa farther north. The retreating Cordilleran ice sheet also left a trail of

temporary lakes that could have served as stepping stones for aquatic species to colonize basins

farther north, such as the Fraser River (McPhail and Lindsey 1986).

Two other major events in the history of the Columbia River basin merit attention. About 3

Ma ago, the Columbia River captured the Snake River, which previously drained farther south to

the Klamath or Sacramento River basins (Wood and Clemens 2002; Stearley and Smith 2016).

Although the connection between the Snake River and lower Columbia River did not promote

extensive faunal transfer (at least of salmonids), it did provide a route for another Pleistocene

flood: the draining of Lake Bonneville. Unlike the ice dam failures of Glacial Lake Missoula,

Lake Bonneville drained by overtopping and rapidly downcutting at Red Rock Pass 17.5 ka ago

(Amidon and Clark 2015; Oviatt 2015), delivering 4,750 km$^3$ of water in what represents the

largest (by volume) known freshwater flood (O'Connor 1993). Like those of Glacial Lake

Missoula, this flood backed up many of the tributary rivers and may have conveyed aquatic

species across large distances. Because it was less violent and longer lasting—a couple months

to one year (Malde 1968; Jarrett and Malde 1987)—the draining of Lake Bonneville may been

more conducive to fish survival and translocation. Interpreting its possible phylogeographic

effect is difficult because it was both preceded and followed by outwash floods from Glacial

Lake Missoula, the last of which was 14.7 ka ago (Waitt 1985; McDonald et al. 2012; Balbas et

al. 2017), with additional floods still later from smaller proglacial lakes that formed as the

Purcell lobe of the Cordilleran ice sheet retreated (Waitt et al. 2009; Peters 2012) and perhaps

from the longer-persisting Glacial Lake Columbia (Balbas et al. 2017).

*Conservation units*

We define units of conservation (sensu distinct population segments; Waples 1991) as

populations or groups thereof that are substantially reproductively isolated and that constitute an

important component of the evolutionary legacy of a species. Ideally, units of conservation

would be identified based on an array of phylogenetically informative ecological, life history,

morphological, and genetic characteristics (Fraser and Bernatchez 2001). Variation in

morphology has been related to ecotypic differences in westslope cutthroat trout (Seiler and

Keeley 2009), and such differences have been shown to have a genetic basis and contribute to

identifying units of conservation in some salmonids (Keeley et al. 2007). To our knowledge,

however, there has been no comparison of morphological or ecological traits across the entire

range of westslope cutthroat trout (cf. Bestgen et al. 2013). In addition, relative to other Pacific

salmon and trout, cutthroat trout are ecological generalists capable of expressing a wide array of

life history strategies (Young 1995; Waples et al. 2001), rendering these characteristics less

definitive for delineating units of conservation. Given this background of uncertainty, we used

well-supported, reciprocally monophyletic mitochondrial clades (Moritz et al. 1995) as working

hypotheses about the units of conservation within westslope cutthroat trout, particularly when

geographic structuring reinforced those distinctions. We also sought support from analyses of

two suites of nuclear markers to corroborate the mitochondrial results.

*Laboratory analyses*

We obtained three sets of genetic information from two groups of samples of westslope

cutthroat trout: 1) mitogenomic sequences of specimens from most of the historical range in the

U.S. and Canada ($n$ = 96 fish); 2) allele frequencies of 10 microsatellite loci (Vu and Kalinowski

2009) in the aforementioned specimens (save one from Buck Creek, WA, sample 44); and 3)

allele frequencies of 52 variable nuclear single nucleotide polymorphisms (SNPs) from

specimens collected across a large portion of the U.S. range ($n = 524$ fish from 55 sites; Table 1;

Figure 1). Specimens from wild populations that were representative of those used to establish

each of the hatchery broodstocks were included in one or both sample groups. Some of the sites

in the first two analyses also had additional individuals analyzed using SNPs ($n = 29$), but spatial

overlap in these datasets was not complete.

Whole genomic DNA was extracted from tissue samples using the QIAGEN DNeasy Tissue

Kit (Qiagen, Valencia, CA, USA). Extracts were converted into indexed genomic libraries using

the Illumina TruSeq DNA HT kit (version 2). To enrich samples for mitochondrial DNA targets,

we developed hybridization-enrichment baits based on the complete mitochondrial genome

sequence for Lahontan cutthroat trout *O. c. henshawi* (NCBI accession AY886762). Baits and

blocking probes were designed and synthesized by MYcroarray, LLC (version 2; Ann Arbor,

MI), and hybridization-enrichment reactions followed their protocol with a 12-cycle final

amplification. Enriched libraries were pooled to 48-plex and sequenced using two lanes of 101-

base-pair, single-end reads on the Illumina HiSeq2000 at the Center for Genome Research and

Biocomputing at Oregon State University (http://cgrb.oregonstate.edu/). Indices were

demultiplexed using CASAVA v. 1.8.2, and individual sample mitochondrial genomes were

assembled using the *O. c. henshawi* reference and the CLC-Bio v. 7.0 assembler, with the

following quality and alignment parameters: quality score limit of 0.05, maximum number of

ambiguities per read = 2; mismatch cost = 3, indel cost = 3, indel open cost = 6, indel extend cost

= 1, length and similarity fraction = 0.95, and non-specific matches mapped randomly.

Individual samples were represented by an average of 3.674 million microreads (minimum =

281,755; maximum = 13,176,436); for assembly, we used a maximum of 4 million reads per

sample. Short read sequences from this study are available under BioProject ID PRJNA389467

from National Center for Biotechnology Information.

In the microsatellite analysis, we conducted analyses of 10 loci used in previous studies

of westslope cutthroat trout: OclMSU14, OclMSU15, OclMSU17, OclMSU21, OclMSU23,

OclMSU24, OclMSU25, OclMSU27, OclMSU30, OclMSU33 (Vu and Kalinowski 2009; Drinan

et al. 2011). The reaction volume (10 µl) contained 1.0 µl DNA, 1× reaction buffer (Applied

Biosystems), 2.0 mM $MgCl_2$, 200 µM of each dNTP, 1 µM reverse primer, 1 µM dye-labeled

forward primer, 1.5 mg/ml BSA, and 1U Taq polymerase (Applied Biosystems). The PCR

profile was 94 °C/5 min, [94 °C/1 min, 55 °C/1 min, 72 °C/30 s] × 40 cycles. The resultant

products were visualized on a LI-COR DNA analyzer (LI-COR Biotechnology).

For the SNP analyses, we used Competitive Allele Specific PCR (KASPar) assays

(KBiosciences, Hoddesdon, Herts, England) to amplify 52 variable nuclear loci (Harwood and

Phillips 2011; Kalinowski et al. 2011; Amish et al. 2012; Campbell et al. 2012; Pritchard et al.

2012). We used a suite of 44 diagnostic SNP alleles to ensure no fish introgressed with rainbow

trout or Yellowstone cutthroat trout *O. c. bouvieri* alleles were included in the analysis

(McKelvey et al. 2016). The PCR touchdown profile contained an initial annealing temperature

of 65 °C and decreased by 0.80 °C per cycle until most cycles ran at 57 °C. We visualized PCR

products on an EP1 Reader (Fluidigm) and determined individual genotypes using Fluidigm SNP

Genotyping Software.

*Phylogenetic analyses*

We conducted phylogenetic analyses on the three aforementioned sources of data, plus a

fourth, to assess intraspecific diversity in westslope cutthroat trout.

Mitogenomic sequences were aligned in MAFFT version 7 (Katoh and Standley 2013),

followed by minor manual adjustments. We used MitoAnnotator (Iwasaki et al. 2013) to order

and identify gene regions. We followed Satoh et al. (2016) and the mitotRNAdb database

(http://mttrna.bioinf.uni-leipzig.de/mtDataOutput/) to identify paired and unpaired regions in

rRNAs and tRNAs and evaluate whether nucleotide insertions or deletions were likely to

represent genotyping errors; we subsequently made 12 adjustments to the 1.6 million nucleotides

in the dataset. Unique haplotypes were identified using DAMBE version 6 (Xia 2017) and used

to construct neighbor-joining and maximum-likelihood phylogenetic trees. To portray

evolutionary distances among and between clades, we used MEGA 7.0 (Kumar et al. 2016) to

construct a neighbor-joining tree based on the MCL model with gamma-distributed evolutionary

rates and some invariant positions, bootstrapped 1,000 times. We included sequences from

Lahontan cutthroat trout, greenback cutthroat trout *O. c. stomias*, and rainbow trout as outgroups

(GenBank accessions NC006897, KP013107, KP013117, KP013084, DQ288268–71,

KP085590). Pairwise genetic distances (as a percentage) between samples and groups were

based on the number of nucleotide differences. For the maximum-likelihood analysis, we used

PartitionFinder 2.0 (Lanfear et al. 2016) to select the best-fitting partitioning scheme as

measured by AICc, constrained to the suite of evolutionary models considered by RAxML

(Stamatakis 2014) and excluding all outgroups. Data subsets included 1) codon position by gene

for the protein-coding genes (39 subsets); 2) paired and unpaired regions among tRNAs and the

origin of replication for the light strand (2 subsets) and between rRNAs (4 subsets); 3) intergenic

spacers and the non-coding portions of the control region (1 subset); and 4) conserved sequence

blocks of the control region, nucleotides coding for more than one gene, and tRNA anticodons (1

subset). Because RAxML will only consider a single evolutionary model for the entire suite of

partitions, we then compared AICc scores among maximum likelihood models using the GTR

and GTRGAMMA evolutionary models, and chose the model with the best score. We then ran

RAxML version 8.1.21 implemented through the RAxMLGUI (Silvestro and Michalak 2012)

and set for rapid bootstrapping (1,000 bootstraps) and a thorough ML search. We recognized

potential units of conservation as reciprocally monophyletic clades with > 70% bootstrap support

in the mitogenomic maximum-likelihood phylogeny. We did not single out subclades therein

with high levels of support unless: 1) they differed from all other subclades by more than the

average intra-clade divergence (0.04%) or 2) they had been previously recognized as a potential

unit of conservation, i.e., fish in the Missouri River basin (Drinan et al. 2011).

Further evaluation of these clades as potential units of conservations was undertaken by

analyzing a dataset combining all ND2 sequences of westslope cutthroat trout from Loxterman

and Keeley (2012) and one additional specimen in GenBank ($n = 93$, accessions EU186799,

JQ747580–596; Table 1) with those examined in the mitogenomic analyses (including the

outgroups). Some of these specimens ($n = 26$) were used in both analyses, reducing the number

of unique specimens ($n = 163$). We built a neighbor-joining tree using unpartitioned data

bootstrapped 1,000 times in MEGA 7.0, for which the Tamura-Nei model with gamma-

distributed rates was the best-fitting evolutionary model based on AICc scores.

Because the microsatellites and variable SNPs for westslope cutthroat trout were developed

(Campbell et al. 2012) almost entirely from specimens representing the two most northeasterly

and least genetically divergent clades identified in the mitogenomic analysis in the Missouri,

Kootenai, and Clark Fork River basins, ascertainment bias was more likely to prevent the

phylogenetic tree topology from representing the species tree (Lachance and Tishkoff 2013) but

less likely to incorrectly assign individuals to groups (Bradbury et al. 2011). Consequently, we

evaluated whether putative groups identified in STRUCTURE 2.3.4 (Hubisz et al. 2009), using

either the microsatellite or SNP datasets, matched those of the best-supported clades in the

mitogenomic analyses. We applied settings recommended by Falush et al. (2003) for detecting

subtle population subdivision by using the admixture model, correlated allele frequencies among

populations, and an allele frequency distribution parameter ($\lambda$) set to 1. We allowed STRUCTURE

to infer the value of the model's Dirichlet parameter ($\alpha$) for the degree of admixture from the

data. In the SNP analysis, we used $K = 15$ (the number of major river basins) as the maximum

number of potential groups and completed 20 replicate runs for each value of $K$. In the

microsatellite analysis, we used $K = 10$ (the number of mitogenomic clades) as the maximum

number. In both analyses, we set the burn-in period to 100,000 MCMC repetitions and the post-

burn-in period to 100,000, ignoring the user-defined population-of-origin or sampling location

for each individual (i.e., adopting non-informative priors). We attempted to identify an optimal

value of $K$ by calculating the maximum likelihood value and $\Delta K$ (Evanno et al. 2005) using the

program STRUCTURE HARVESTER (Earl and vonHoldt 2012).

Stocking of genetically pure westslope cutthroat trout was inferred from the appearance of

mitochondrial haplotypes outside their river basin of origin, or of individuals in nuclear analyses

that assigned to groups related to those that founded the primary broodstocks in Idaho and

Montana. For Montana, we regarded samples from the South Fork Flathead River (sample codes

35, 94, and 179; Table 1) as representative of broodstock genotypes. Samples from the Priest

River basin (codes 24, 137, and 138) were deemed to represent fish from the Idaho broodstock,

as well as the King's Lake broodstock in Washington. The Falls Creek sample (code 60)

represented Washington's Twin Lakes broodstock, but these genotypes were not found outside

the eastern Cascades river basins.

**Results**

*Mitochondrial analyses*

Hybridization-capture probes allowed us to enrich mitochondrial genome sequences to a very

high level (mean mitochondrial representation = 57.5% of all sequences), resulting in

mitochondrial genome assemblies represented by extremely high mean depth (range: 997—

20,208 reads), and complete coverage for all positions in the genome. Complete mitogenomes

(16,676 nucleotides) included 13 protein-coding genes, 2 rRNAs, 22 tRNAs, and several

noncoding regions (Table 2), and revealed 89 haplotypes among the 96 specimens. The

mitogenomic maximum-likelihood phylogeny recovered two main lineages (mean difference

between lineages based on number of nucleotide differences, 0.42%; mean difference within

lineages, 0.06–0.11%) that contained nine highly supported, monophyletic clades and one

subclade (Figure 2; Table 3). The topology and levels of support in the mitogenomic neighbor-

joining analysis were effectively identical (results not shown). One lineage (hereafter, the

southern lineage) contained six clades (1–5b) largely associated with individual river basins: the

John Day River (1) in Oregon; the upper Coeur d'Alene (2) and St. Joe Rivers (3) in the Spokane

River basin in Idaho; the North Fork Clearwater River (4) in Idaho; and the lower and middle

Salmon River (5a) and upper Salmon River (5b) in Idaho. The latter two clades, however,

overlapped geographically within the basin. The four clades (6–9) in the other main lineage (the

northern lineage) were less geographically circumscribed: the Clearwater headwaters (6), which

included specimens from the upper Selway and South Fork Clearwater Rivers in Idaho; the

Clearwater-eastern Cascades (7), with representatives from tributaries to the Middle Fork

Clearwater River in Idaho and from the Wenatchee River, Lake Chelan, and Methow River

basins on the eastern side of the Cascade Range in Washington; and a neoboreal (8) clade, which

included all river basins in Alberta, British Columbia, Montana, northern Idaho, and northeastern

Washington. Most of these constitute basins that were covered by continental ice sheets or the

proglacial lakes at their margins. Disjunct representatives of this clade, however, were found in

the Salmon, Clearwater, and Spokane River basins in Idaho. Also recovered in these analyses

was a well-supported subclade within the neoboreal group that represented all specimens (save

one) from the Missouri River (9) basin in Montana.

The analysis of ND2 sequences from fish genotyped for this study and from public databases

had 27 haplotypes, including 8 not observed in the mitogenomic analysis (Table 1). Consistent

with the previous analysis, the ND2 neighbor-joining tree resolved the two main lineages and

recovered similar geographic structure (Figures 3), but with less resolution and support of

additional clades despite greater geographic coverage. This analysis diagnosed the John Day,

Coeur d'Alene, and St. Joe River clades, and grouped but did not distinguish among the North

Fork Clearwater and Salmon River specimens. It also pooled all Clearwater River and eastern

Cascade specimens, and did not resolve Missouri River specimens as a distinct group within the

neoboreal clade.

*Nuclear analyses*

In the STRUCTURE analysis of SNPs, measures of the optimal number of groups were not

concordant. The log-likelihood estimate peaked at the maximum number of groups ($K = 15$)

whereas $\Delta K$ identified $K = 2$ as the optimum number of clusters. Given this outcome, we

examined how assignment to groups changed across the entire range of values (Figure 4 depicts

$K = 4$, 8, and 15). The intermediate value ($K = 8$) was equivalent to the number of mitogenomic

clades geographically represented in the SNP analysis (samples equivalent to the North Fork

Clearwater River and Clearwater River headwater samples were not available). Although these

results broadly corroborated the mitogenomic classification, there were important differences.

Specimens from the John Day River were the first group to be segregated as distinct (at $K = 3$).

Specimens from the Salmon River formed a distinct group at intermediate and higher levels of $K$,

as did those from the Missouri River. The Clearwater River and eastern Cascades clade grouped

at the lower levels of $K$, but split into distinct clusters at the highest levels of resolution. More

complex were patterns among the Coeur d'Alene River, St. Joe River, and neoboreal clades.

Specimens from the Coeur d'Alene and St. Joe Rivers largely assigned to a single group at all

levels of $K$, but this group also included specimens from the neoboreal clade from the lower

Clark Fork, Pend Oreille, and lower Kootenai River basins for lower levels of $K$. The rest of the

neoboreal clade tended to split along major watershed boundaries at higher levels of $K$,

particularly the upper Kootenai River.

In the microsatellite analyses, $K = 5$ (Figure 5) had the largest log-likelihood estimate and the

third-highest $\Delta K$ (which peaked at $K = 2$). The smaller sample sizes in this analysis afforded less

resolution but nonetheless highlighted groups consistent with those identified previously: 1) the

John Day and Clearwater Rivers and eastern Cascades basins; 2) the Salmon River; 3) the Coeur

d'Alene and St. Joe Rivers, some samples from the Pend Oreille and lower Kootenai Rivers, one

from downstream on the Columbia River, and one from the Fraser River; 4) the Clark Fork and

upper Kootenai Rivers in the Columbia basin and the Oldman and Bow Rivers in the South

Saskatchewan basin; and 5) the Missouri River.

*Stocked fish*

The genotypes of one or more fish at as many as 18 sites were suspected of being influenced

by introductions of nonintrogressed but nonindigenous westslope cutthroat trout (Table 4). The

bulk of these observations were from Idaho, and involved the appearance of mitochondrial

haplotypes typical of the neoboreal group in the Salmon and Clearwater River basins. Although

we regarded specimens with neoboreal haplotypes in the St. Joe and Coeur d'Alene River basins,

primarily from the lowermost portions of both basins, as being nonindigenous, this interpretation

in uncertain. Fish with either the neoboreal or southern haplotypes tended to share a nuclear

genotype that was also typical of fish found farther north in the Pend Oreille, lower Clark Fork,

lower Kootenai, middle Columbia, and Fraser River basins. Thus, the St. Joe or Coeur d'Alene

River specimens may have been indigenous representatives of fish that colonized basins to the

north, including that from which Idaho's broodstock was derived, rather than reflect recent

introductions of that hatchery stock. In Montana, the two likely examples of nonindigenous

specimens were from disparate parts of the Missouri River basin that harbored fish having

mitochondrial or nuclear genotypes akin to those from west of the Continental Divide.

**Discussion**

The phylogeographic structure of westslope cutthroat trout is consistent with unglaciated

basins serving as refugia during glacial cycles, followed by extensive colonization from some of

those refugia during interglacial intervals (Bernatchez and Wilson 1998; Hampe and Petit 2005).

The deepest divergence among populations involved the John Day, Salmon, Clearwater, and

Spokane River basins, which were south of the maximum advance of the Cordilleran ice sheet.

Much lower levels of divergence characterized areas formerly occupied by the ice sheets or the

glacial lakes at their margins, collectively represented by the neoboreal mitogenomic clade (and

including the Missouri River basin). Although we have foregone estimating dates of divergence

among clades, much of the diversity seems to be associated with recent climatic events. The

radiation following the Last Glacial Maximum likely reflects the divergence within most

mitogenomic clades (mean, 0.04%), whereas previous interglacials over 100 ka ago may have

led to divergence between the Coeur d'Alene and St. Joe River clades, the neoboreal and the

Clearwater-eastern Cascade clades, and the North Fork Clearwater and Salmon River clades, and

still older events to the split between the southern and northern major lineages. Continental

glaciation during previous glacial maxima is likely to have extirpated groups that advanced

northward during earlier interglacial intervals. That all samples from the previously glaciated

area form a single mitogenomic clade suggests that no individuals representing earlier colonizers

are present in this area. Below, we consider the phylogeographic relevance of these river basins
to westslope cutthroat trout in more detail.

The clade in the John Day River basin, though clearly a member of the southern major
lineage, exhibited the greatest divergence from all other groups. This watershed constitutes the
downstreammost extent of westslope cutthroat trout in the Columbia River basin and is separated
from the next nearest populations in the eastern Cascade Range in Washington by hundreds of
kilometers, and by hundreds more from the nearest members of the southern lineage in the
Clearwater and Salmon Rivers in Idaho. There has been uncertainty with regard to whether these
fish were indigenous or introduced from Montana, Idaho, or Washington (Gunckel 2002), but the
evidence indicates that these fish constitute a distinct, relictual population of westslope cutthroat
trout that did not arrive via glacial outwash floods from upstream. Whether their distribution
extends to the North Fork John Day River basin, which also supports westslope cutthroat trout
but for which there are extensive records of stocking of this taxon (Gunckel 2002), has not been
evaluated. As alluded to earlier, the absence of westslope cutthroat trout from nearby basins in
southeastern Washington and northeastern Oregon with abundant suitable habitat is somewhat
surprising. That westslope cutthroat trout may have been present, however, is hinted at by the
presence of their mitochondrial haplotype among anadromous rainbow trout in at least one area,
the Tucannon River in Washington (Brown et al. 2004), although unrecorded stocking of
westslope cutthroat trout is a plausible alternative explanation.

The Salmon River basin constitutes a modern stronghold for westslope cutthroat trout
(Shepard et al. 2005). It also hosts two relatively divergent mitogenomic clades of the southern
lineage which are found exclusively in this basin. Nevertheless, the clades are geographically
interspersed and showed little differentiation in the nuclear analyses. Their origin may have
reflected isolation in different parts of the Salmon River basin during a previous glacial interval
followed by subsequent mixing. Their nearest relatives appear to be fish found in the North Fork
Clearwater River to the north, but whether these groups represent descendants of one another or
of simultaneous divergence in these different locations is uncertain.

The Clearwater River basin in Idaho constitutes a biodiversity hotspot for westslope cutthroat
trout. It hosts a geographically complex mixture of members of both primary mitogenomic
lineages and four of the 10 mitogenomic clades of westslope cutthroat trout, two of which are
endemic (and a third may be introduced). Its position beyond the southerly extent of the

Cordilleran ice sheet, combined with its relatively low elevation and high precipitation, seems to

have contributed to the basin being a refugium for disjunct or locally endemic populations of an

array of plant and animal taxa during the Last Glacial Maximum (summarized in Shafer et al.

2010). That does not suggest that all portions of the basin were amenable to westslope cutthroat

trout throughout this or earlier glacial cycles. Pollen core records from near the headwaters

indicate a cold, arid climate with vegetation dominated by sagebrush, and that western red cedar

*Thuja plicata*, currently a dominant forest overstory species typical of mesic, montane

environments, did not colonize the basin until the last 5 ka (Herring and Gavin 2015). Moreover,

mountain glaciers would have been prevalent in the higher ranges, constraining westslope

cutthroat trout to the lower portions of many watersheds until conditions permitted them to

colonize these areas, perhaps from multiple locations within the basin.

A remarkably close relationship between the populations in the eastern Cascade Range in

Washington and those in Clearwater River basin—especially the Lochsa River basin—in Idaho

is supported in all analyses. This enigmatic pattern raises questions about how westslope

cutthroat trout crossed the majority of the lower Snake and middle Columbia River basins in

recent geological time and in which direction. The simplest explanation would be stocking of the

Twin Lakes broodstock from Washington in the Clearwater River in Idaho, but there are no

records to support this (Evan Brown, Idaho Department of Fish and Game, personal

communication). There is, however, evidence of the disjunct distribution of undescribed species

of sculpins in central Idaho basins and along the eastern Cascade Range (M. Young, unpublished

data). Given that the modern distribution of these sculpins does not include lacustrine or large,

warm riverine environments, it is more likely that they, and thus westslope cutthroat trout, were

transported downstream from central Idaho to their present distribution in central Washington.

The limited divergence in westslope cutthroat trout haplotypes between these locations also

indicates that this transfer was recent, and may have involved the emptying of Lake Bonneville

or the last of the Glacial Lake Missoula outburst floods.

The major forks of the Spokane River basin are the Coeur d'Alene and St. Joe Rivers. In their

headwaters, these basins harbor distinct clades that are nonetheless sister to one another and

represent the southern lineage of westslope cutthroat trout. A comparable pattern of relatedness

and divergence was evident in the cedar sculpin *Cottus schitsuumsh*, a recently described species

that is largely confined to the Spokane River basin (LeMoine et al. 2014). The lower portions of

both the Coeur d'Alene and St. Joe Rivers, however, also hosted members of the mitogenomic

neoboreal clade. Fish from throughout both basins shared a nuclear genotype that also appeared

in fish farther north in the middle Columbia, Pend Oreille, lower Kootenai, lower Clark Fork

River, and Fraser River basins, all presently unconnected to the Spokane River basin. The pattern

could have arisen from 1) stocking of fish from the Priest River basin, the source of Idaho's

broodstock, throughout the upper Spokane River basin, for which there is some evidence (Table

4); or 2) ephemeral connections associated with fluctuating levels of proglacial lakes that

promoted northerly expansion of genotypes from the Spokane River basin. As noted earlier,

Glacial Lake Columbia reached tens of kilometers up the Coeur d'Alene and St. Joe River

valleys and would have facilitated the movement of fish between these basins. This would also

have provided access to the Pend Oreille basin across the low divide in Rathdrum Prairie in

northern Idaho. Much of this watershed was subjected to the extreme conditions associated with

the repeated, catastrophic drainage of Glacial Lake Missoula, but Glacial Lake Columbia

persisted for some time after the last of these floods (Balbas et al. 2017). As the Purcell trench

lobe of the Cordilleran ice sheet receded, it would have exposed the lower Kootenai River valley

but blocked drainage to the north, diverting streams from that area through the trench cut by the

Purcell lobe and connecting the lower Kootenai and Pend Oreille basins (Langer et al. 2011).

Another connection between these basins appears to have existed at the low divide between the

Bull River (lower Clark Fork) and Bull Lake (lower Kootenai) in Montana (Langer et al. 2011).

Each of these connections would have provided access to fish originating from the Spokane

River basin, which may have culminated in their reaching the Fraser River basin via the

succession of glacial lakes trailing the receding ice sheet. That a fish from that basin was

assigned to a microsatellite group representing fish from the upper Spokane River basin argues

for such a connection.

Yet it is also clear that these fish—or this particular wave of colonizers—did not found the

modern populations of westslope cutthroat trout inhabiting the Kootenai River above Kootenai

Falls or the Clark Fork River above present-day Noxon Dam or Thompson Falls. All these fish

constitute a single mitogenomic clade with a broadly shared nuclear genotype that suggests a

limited number of founding individuals or a limited number of sources. Whatever their origin,

this is now the most widely distributed clade of westslope cutthroat trout, and one that also

displays substantial internal structure. For example, the nuclear data (e.g., SNP STRUCTURE plots

for $K = 2$–6) indicate historical connections between populations in upper Kootenai and Flathead

River basins, for which there are a number of plausible pathways that might have existed

depending on the timing of recession of particular lobes of the Cordilleran ice sheet, the presence

or absence of terminal or lateral moraines that affected drainage patterns, or the existence of

multiple outlets of lakes near drainage divides, e.g., between the Little Bitterroot (Flathead), Big

Thompson (Clark Fork), and Fisher (upper Kootenai) Rivers, or the Stillwater (Flathead) and

Tobacco (upper Kootenai) Rivers.

There is also evidence that these fish not only colonized two additional major river basins

across the Continental Divide—the Missouri and South Saskatchewan Rivers—but that these

appear to be independent events. Based on our results and on earlier studies (Drinan et al. 2011),

the Missouri River clade is a recent derivative of the widely distributed neoboreal clade, likely

founded from a single source; all individuals share two diagnostic mutations in the mitogenome

(one each in tRNA for tyrosine and the control region). The nuclear data imply that their closest

relatives are those in the upper Flathead River basin. A likely route is via the Middle Fork

Flathead River and Nyack Creek to Summit Creek, thence to the Two Medicine River basin and

the rest of the Missouri River basin. This invasion was likely to have coincided with the

lingering presence of glacial lakes formed by the Laurentide ice sheet because the Two Medicine

River is tributary to the Marias River, which presently flows into the Missouri River well

downstream of the Great Falls of the Missouri. Because the falls constitute a barrier to upstream

fish migration, this location was likely inundated by a glacial lake (e.g., Glacial Lake Great Falls

had a highstand of 1200 m [Calhoun 1906], about 250 m higher than the river elevation at the

falls) which would have enabled westslope cutthroat trout to access the Missouri River

headwaters. Furthermore, the lack of genetic divergence (mean distance 0.01%, with several

geographically distant pairs of samples with identical mitogenomic haplotypes) among westslope

cutthroat trout in this basin also indicates that their recent arrival may be their first. The upper

Missouri River basin was not covered by the Laurentide Ice Sheet during any portion of the

Pleistocene, thus suitable habitat for cutthroat trout should have persisted throughout much of

this period and permitted greater divergence. That westslope cutthroat trout did not extend their

distribution to the Yellowstone River basin, where Yellowstone cutthroat trout are found, points

to the absence of a migration corridor via the lower Missouri and Yellowstone Rivers for either

subspecies when they colonized. That a single clade of sculpins *Cottus* sp. (Young et al. 2013;

M. Young, unpublished data) and of mountain whitefish *Prosopium williamsoni* (Whiteley et al.

2006) co-occur with both subspecies in these areas, however, demonstrates that interbasin

exchanges were possible at an earlier time.

The westslope cutthroat trout of Alberta in the South Saskatchewan River basin appear to

have arrived via a different route. They share mitogenomic affinities with fish of the Clark Fork

River basin upstream from Thompson Falls and lack the diagnostic mutations seen in the fish

from the Missouri River basin. The microsatellite analyses also group these fish with those in the

upper Clark Fork River, not with those from the Missouri River. That the invasions of the South

Saskatchewan and upper Missouri River basins were independent events is also supported by the

distribution of bull trout *Salvelinus confluentus*, a cold-water salmonid with similar habitat

preferences, which are present in the former and absent from the latter (Ardren et al. 2011).

Moreover, westslope cutthroat trout may have been constrained to a limited portion of the South

Saskatchewan River basin. Their current distribution includes only the Bow and Oldman River

basins (Fisheries and Oceans Canada 2014), and Marnell (1988) argued that they were

historically absent from Waterton Lakes and the Belly River and crossed the Continental Divide

farther north. Based on topography, a plausible route would have been via the Kootenay River to

Summit Lake in British Columbia, thence to Crowsnest Lake in Alberta, which is part of the

Oldman River basin.

Based on our combined analyses, we find evidence for nine units of conservation within this

taxon (Table 5). We acknowledge that different interpretations of the importance of modern-day

isolation, administrative boundaries, ecoregions, or geographic mixing of genotypes (Mee et al.

2015) could alter this total. For example, a reasoned case can be made that fish representing the

neoboreal clade in the lower Kootenai, lower Clark Fork, Pend Oreille, middle Columbia, and

Fraser River basins should be classified separately from those found elsewhere because their

nuclear genotypes are distinct. Similarly, one could contend that fish in the eastern Cascades in

Washington are distinct from those in the Clearwater River basin in Idaho based on geographic

isolation and different administrative authorities. Such arguments are valid, thus we consider our

estimate an initial hypothesis to be revised following additional sampling and more

comprehensive analyses, but would be cautious about divisions at much smaller scales. Leary et

al. (1988) reported that the bulk of genetic variation in westslope cutthroat trout in Montana was

partitioned among individual populations, and consequently advocated for saving as many

populations as possible to preserve the genetic legacy of this taxon. The substantial structuring of

headwater populations has been observed elsewhere among westslope cutthroat trout (Taylor et

al. 2003; Young et al. 2004) and other salmonid species (Ryman 1983; Wenburg and Bentzen

2001; Pritchard et al. 2009), though it may be an artifact of the widespread anthropogenic

isolation of headwater trout populations (Campbell et al., this volume). In contrast, our data (also

see Drinan et al. 2011) representing nearly the entire range of westslope cutthroat trout

demonstrates that the primary divisions within this taxon arise at broader spatial scales, roughly

conforming to major watershed boundaries and cycles of Pleistocene glaciation, which is

consistent with hypotheses about rangewide genetic structure in temperate zone species (Hampe

and Petit 2005). Although we concur with the intent to save as many populations as possible, the

larger groups we identified are more likely to satisfy criteria for recognizing units of

conservation (Waples 1991; Mee et al. 2015).

Our data and that of others confirms that the westslope cutthroat trout constitutes a coherent

taxon that may warrant recognition as a full species. The level of divergence within westslope

cutthroat trout (0.39% in a 640-bp fragment of cytochrome c oxidase subunit I (COI), the

barcode region) is comparable to that of other species of fishes (0.36%) worldwide (Ward 2009),

whereas its divergence (1.25% in the COI fragment, 1.11% across the mitogenome) from its

nearest relative (Lahontan cutthroat trout) attains or exceeds that of many species pairs (April et

al. 2011). Nonetheless, substantial uncertainties remain with regard to the placement of

westslope cutthroat trout within the overall cutthroat trout lineage. Our mitogenomic analyses

affirmed that westslope cutthroat trout haplotypes are highly divergent from those of rainbow

trout (4.80% across the mitogenome), and more similar to Lahontan cutthroat trout than to

greenback cutthroat trout (1.58%), but there is little consensus among the different genomic

regions. Others have suggested that westslope cutthroat trout may be most closely related to

rainbow trout based on variation in isozyme alleles (Leary et al. 1987), to coastal cutthroat trout

based on karyotypes or morphology (Loudenslager and Gall 1980; Behnke 1992), to Lahontan

cutthroat trout based on mitochondrial gene sequences (Wilson and Turner 2009; Loxterman and

Keeley 2012), either of the latter two based on analyses of Y chromosomes (Brunelli et al. 2013),

and to all three based on a panel of variable nuclear SNP markers (Houston et al. 2012). That

each analysis supports a different relationship may be more indicative of the relative strengths

and weaknesses of different markers than of an ambiguous phylogeny. For example, allozymes

can grossly and unpredictably underestimate haplotype variation (Buth 1984). The karyotype

analysis was conducted on a sample of five westslope cutthroat trout from one hatchery

broodstock (Loudenslager and Gall 1980) and its results contradict those of an earlier study with

specimens from elsewhere (Simon and Dollar 1963). Both studies could be correct because

geographical variation in chromosomal characteristics or multiple karyotypes within populations

are common (Thorgaard 1983). Mitochondrial sequences form the basis of a multitude of

projects to identify animal species (Ratnasingham and Herbert 2013), but introgression and

reticulate evolution among lineages weaken relationships based solely on mitochondrial data

(Taylor and Harris 2012). And all genetic analyses are vulnerable to incomplete geographic

sampling, and the geographic origin and ancestry of specimens used to develop nuclear markers

and diagnostic phenotypes can undermine phylogenetic inferences because of ascertainment bias

(Lachance and Tishkoff 2013) or environmentally driven selection. Given the relative youth of

cutthroat trout lineages, more intensive and representative sampling of the nuclear genome and

of populations likely to be indigenous will be required to resolve this issue.

Although our work represents the most comprehensive effort to describe the phylogeography

of westslope cutthroat trout, many details have yet to be resolved. Locations represented by one

or two individuals would benefit from more thorough sampling to reveal the distribution and

mixing of presumptive haplotypes and the discovery of new ones, such as in the John Day River

(and its presumably stocked North Fork) in Oregon, the eastern Cascade River basins in

Washington, and many portions of the Clearwater River in Idaho. Westslope cutthroat trout in

the Yakima River basin are of unknown provenance, whether related to lineages in the John Day,

the eastern Cascades (as implied in Figure 1), or to one as yet undescribed. Similarly, additional

work to delineate hatchery stocks using nuclear markers would clarify the extent to which

introgression of introduced forms affects indigenous populations, and would help avoid the use

of introduced forms when attempting to establish basin-specific stocks. Despite these

uncertainties, the phylogeographic patterns we have observed are concordant with the varying

connections among river basins in the intermountain West throughout the late Pleistocene. These

patterns can be used as a foundation to understand phylogeographic structure in other freshwater

taxa, albeit with an awareness of the nuances associated with the long evolutionary history of

cutthroat trout in western North America.

**Acknowledgements**

We thank M. Small of the Washington Department of Fish and Wildlife and J. Neal of the Oregon Department of Fish and Wildlife for contributing samples. T. Cross of the Rocky Mountain Research Station assisted with the STRUCTURE analysis of SNPs, and D. Horan helped prepare Figure 1. Comments by Kurt Fausch and two anonymous reviewers markedly improved the manuscript. This work was funded by the RIM Board of the U.S. Forest Service's Region 1, the Rocky Mountain Research Station, and the Pacific Northwest Research Station.

**References**

Alden WC (1932) Physiography and glacial geology of eastern Montana. U.S. Geological Survey Professional Paper 174.

Alt DD, Hyndman DW (1995) Northwest exposures: a geologic story of the Northwest. Mountain Press, Missoula, MT.

Amidon WH, Clark AC (2015) Interaction of outburst floods with basaltic aquifers on the Snake River Plain: Implications for Martian canyons. *Geological Society of America Bulletin*, **127**, 688–701.

Amish SJ, Hohenlohe PA, Painter S, Leary RF, Muhlfeld C, Allendorf FW, Luikart G (2012) RAD sequencing yields a high success rate for westslope cutthroat and rainbow trout species-diagnostic SNP assays. *Molecular Ecology Resources*, **12**, 653–660.

April J, Mayden RL, Hanner RH, Bernatchez L (2011) Genetic calibration of species diversity among North America's freshwater fishes. *Proceedings of the National Academy of Sciences*, **108**, 10602–10607.

Ardren WR, DeHaan PW, Smith CT, Taylor EB, Leary R, Kozfkay CC, Godfrey L, Diggs M, Fredenberg W, Chan J, Kilpatrick CW (2011) Genetic structure, evolutionary history, and conservation units of bull trout in the coterminous United States. *Transactions of the American Fisheries Society*, **140**, 506–525.

Atwater BF (1987) Status of glacial Lake Columbia during the last floods from glacial Lake Missoula. *Quaternary Research*, **27**, 182–201.

Avise JC (2000) Phylogeography: the history and formation of species. Harvard University Press, Cambridge.

Bader NE, Spencer PK, Bailey AS, Gastineau KM, Tinkler ER, Pluhar CJ, Bjornstad BN (2016) A loess record of pre-Late Wisconsin glacial outburst flooding, Pleistocene paleoenvironment, and Irvingtonian fauna from the Rulo site, southeastern Washington, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **462**, 57–69.

Balbas AM, Barth AM, Clark PU, Clark J, Caffee M, O'Connor J, Baker VR, Konrad K, Bjornstad B (2017) 10Be dating of late Pleistocene megafloods and Cordilleran Ice Sheet retreat in the northwestern United States. *Geology*. doi: 10.1130/G38956.1.

Behnke RJ (1992) Native trout of western North America. American Fisheries Society, Bethesda, Maryland.

Bernatchez L, Wilson CC (1998) Comparative phylogeography of Nearctic and Palearctic fishes.
*Molecular Ecology*, **7**, 431–452.

Bestgen KR, Rogers KB, Granger R (2013) Phenotype predicts genotype for lineages of native
cutthroat trout in the Southern Rocky Mountains. Final Report to U. S. Fish and Wildlife
Service, Colorado Field Office, Denver Federal Center (MS 65412), Denver, CO. Larval Fish
Laboratory Contribution 177.

Booth DB, Troost KG, Clague JJ, Waitt RB (2003) The cordilleran ice sheet. *Developments in*
*Quaternary Sciences*, **1**, 17–43.

Bradbury IR, Hubert S, Higgins B, Bowman S, Paterson IG, Snelgrove PV, Morris CJ, Gregory
RS, Hardie DC, Borza T, Bentzen P (2011) Evaluating SNP ascertainment bias and its impact
on population assignment in Atlantic cod, *Gadus morhua*. *Molecular Ecology Resources*,
**11(s1)**, 218–225.

Brown KH, Patton SJ, Martin KE, Nichols KM, Armstrong R, Thorgaard GH (2004) Genetic
analysis of interior Pacific Northwest *Oncorhynchus mykiss* reveals apparent ancient
hybridization with westslope cutthroat trout. *Transactions of the American Fisheries Society*,
**133**, 1078–1088.

Brunelli JP, Mallatt JM, Leary RF, Alfaqih M, Phillips RB, Thorgaard GH (2013) Y
chromosome phylogeny for cutthroat trout (*Oncorhynchus clarkii*) subspecies is generally
concordant with those of other markers. *Molecular Phylogenetics and Evolution*, **66**, 592–
602.

Buth DG (1984) The application of electrophoretic data in systematic studies. *Annual Review of*
*Ecology and Systematics*, **15**, 501–522.

Calhoun FHH (1906) The Montana lobe of the Keewatin ice sheet. U.S. Geological Survey
Professional Paper 50.

Campbell MR, Keeley ER, Kozfkay CC, Loxterman JL, Evans PR, Shiozawa DK (this volume)
Describing and preserving the diversity of cutthroat trout in the Yellowstone River, Snake
River and Bonneville Basins. Chapter 12. Evolutionary biology and taxonomy of cutthroat
trout. American Fisheries Society, Bethesda, Maryland.

Campbell NR, Amish SJ, Pritchard VL, McKelvey KS, Young MK, Schwartz MK, Garza JC,
Luikart G, Narum SR (2012) Development and evaluation of 200 novel SNP assays for

population genetic studies of westslope cutthroat trout and genetic identification of related

taxa. *Molecular Ecology Resources*, **12**, 942–949.

Colton RB, Lemke RW, Lindvall RM (1961) Glacial map of Montana east of the Rocky

Mountains. U.S. Geological Survey Miscellaneous Investigations Series Map I-327, scale 1 :

500,000, 1 sheet.

Comte L, Grenouillet G (2013) Do stream fish track climate change? Assessing distribution

shifts in recent decades. *Ecography*, **36**, 1236–1246.

Crawford B (1979) Origin and history of the trout brood stocks of the Washington Department of

Game. Available online: https://www.watrailblazers.org/science/crawford/. Accessed 8 May

2017.

Crisp MD, Trewick SA, Cook LG (2011) Hypothesis testing in biogeography. *Trends in Ecology*

*& Evolution*, **26**, 66–72.

Davis NK, Locke WW, Pierce KL, Finkel RC (2006) Glacial Lake Musselshell: Late Wisconsin

slackwater on the Laurentide ice margin in central Montana, USA. *Geomorphology*, **75**, 330–

345.

Drinan DP (2010) Thermal adaptation of westslope cutthroat trout *Oncorhynchus clarkii lewisi*.

Master's thesis, Montana State University, Bozeman.

Drinan DP, Kalinowski ST, Vu NV, Shepard BB, Muhlfeld CC, Campbell MR (2011) Genetic

variation in westslope cutthroat trout *Oncorhynchus clarkii lewisi*: implications for

conservation. *Conservation Genetics*, **12**, 1513–1523.

Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for

visualizing STRUCTURE output and implementing the Evanno method. *Conservation*

*Genetics Resources*, **4**, 359–361.

Eby LA, Helmy O, Holsinger LM, Young MK (2014) Evidence of climate-induced range

contractions for bull trout in a Rocky Mountain watershed, U.S.A. *PLoS One*, **9**, e98812.

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the

software STRUCTURE: a simulation study. *Molecular Ecology*, **14**, 2611–2620.

Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus

genotype data: linked loci and correlated allele frequencies. *Genetics*, **164**, 1567–1587.

Fecht KR, Reidel SP, Tallman AM (1985) Paleodrainage of the Columbia River system on the

Columbia Plateau of Washington state: A summary. Rockwell International Corp., Richland,

Washington. RHO-BW-SA-318-P.

Fisheries and Oceans Canada (2014) Recovery Strategy for the Alberta populations of Westslope

Cutthroat Trout (*Oncorhynchus clarkii lewisi*) in Canada [Final]. Species at Risk Act

Recovery Strategy Series. Fisheries and Oceans Canada, Ottawa.

Fraser DJ, Bernatchez L (2001) Adaptive evolutionary conservation: towards a unified concept

for defining conservation units. *Molecular Ecology*, **10**, 2741–2752.

Girard CF (1856) Notice upon the species of the genus *Salmo*, of authors, observed chiefly in

Oregon and California. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **8**,

217–220.

Gunckel S (2002) Stocking history of westslope cutthroat trout and rainbow trout in the upper

John Day Basin, Oregon. Oregon Department of Fish and Wildlife, Corvallis.

Hampe A, Petit RJ (2005) Conserving biodiversity under climate change: the rear edge matters.

*Ecology Letters*, **8**, 461–467.

Hanson MA, Clague JJ (2016) Record of glacial Lake Missoula floods in glacial Lake Columbia,

Washington. *Quaternary Science Reviews*, **133**, 62–76.

Harwood AS, Phillips RB (2011) A suite of twelve single nucleotide polymorphism markers for

detecting introgression between cutthroat and rainbow trout. *Molecular Ecology Resources*,

**11**, 382–385.

Heath TA, Hedtke SM, Hillis DM (2008) Taxon sampling and the accuracy of phylogenetic

analyses. *Journal of Systematics and Evolution*, **46**, 239–257.

Herring EM, Gavin DG (2015) Climate and vegetation since the Last Interglacial (MIS 5e) in a

putative glacial refugium, northern Idaho, USA. *Quaternary Science Reviews*, **117**, 82–95.

Houston DD, Elzinga DB, Maughan PJ, Smith SM, Kauwe JS, Evans RP, Stinger RB, Shiozawa

DK (2012) Single nucleotide polymorphism discovery in cutthroat trout subspecies using

genome reduction, barcoding, and 454 pyro-sequencing. *BMC Genomics*, **13**, 724.

Howard AD (1958) Drainage evolution in northeastern Montana and northwestern North Dakota.

*Geological Society of America Bulletin*, **69**, 575–588.

Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with

the assistance of sample group information. *Molecular Ecology Resources*, **9**, 1322–1332.

Hughes PD, Gibbard PL, Ehlers J (2013) Timing of glaciation during the last glacial cycle:

evaluating the concept of a global 'Last Glacial Maximum' (LGM). *Earth-Science Reviews*,

**125**, 171–198.

Iwasaki W, Fukunaga T, Isagozawa R, Yamada K, Maeda Y, Satoh TP, Sado T, Mabuchi K,

Takeshima H, Miya M, Nishida M (2013) MitoFish and MitoAnnotator: a mitochondrial

genome database of fish with an accurate and automatic annotation pipeline. *Molecular*

*Biology and Evolution*, **30**, 2531–2540.

Jarrett RD, Malde HE (1987) Paleodischarge of the late Pleistocene Bonneville Flood, Snake

River, Idaho, computed from new evidence. *Geological Society of America Bulletin*, **99**, 127–

134.

Kalinowski ST, Novak BJ, Drinan DP, Jennings R, Vu NV (2011) Diagnostic single nucleotide

polymorphisms for identifying westslope cutthroat trout (*Oncorhynchus clarki* [sic] *lewisi*),

Yellowstone cutthroat trout (*Oncorhynchus clarkii bouvieri*) and rainbow trout

(*Oncorhynchus mykiss*). *Molecular Ecology Resources*, **11**, 389–393.

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution*, **30**, 772–780.

Keeley ER, Parkinson EA, Taylor EB (2007) The origins of ecotypic variation of rainbow trout:

a test of environmental vs. genetically based differences in morphology. *Journal of*

*Evolutionary*, **20**, 725–736.

Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis

version 7.0 for bigger datasets. *Molecular Biology and Evolution*, **33**, 1870–1874.

Lachance J, Tishkoff SA (2013) SNP ascertainment bias in population genetic analyses: why it is

important, and how to correct it. *Bioessays*, **35**, 780–786.

Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2016) PartitionFinder 2: new

methods for selecting partitioned models of evolution for molecular and morphological

phylogenetic analyses. *Molecular Biology and Evolution*, **34**, 772–773.

Langer WH, Van Gosen BS, Meeker GP, Adams DT, Hoefen TM (2011) The dispersion of

fibrous amphiboles by glacial processes in the area surrounding Libby, Montana, USA.

*Environmental Earth Sciences*, **64**, 157–168.

Leary RF, Allendorf FW, Phelps SR, Knudsen KL (1987) Genetic divergence and identification
of seven cutthroat trout subspecies and rainbow trout. *Transactions of the American Fisheries*
*Society*, **116**, 580–587.

Leary RF, Allendorf FW, Phelps SR, Knudsen KL (1988) Population genetic structure of
westslope cutthroat trout: genetic variation within and among populations. *Proceedings of the*
*Montana Academy of Sciences*, **48**, 57–70.

LeMoine M, Young MK, McKelvey KS, Eby L, Pilgrim KL, Schwartz MK (2014) *Cottus*
*schitsuumsh*, a new species of sculpin (Scorpaeniformes: Cottidae) in the Columbia River
basin, Idaho-Montana, USA. *Zootaxa*, **3755**, 241–258.

Leopold A (1953) Round river. Oxford University Press, New York.

Loudenslager EJ, Gall GA (1980) Geographic patterns of protein variation and subspeciation in
cutthroat trout, *Salmo clarki*. *Systematic Biology*, **29**, 27–42.

Loxterman JL, Keeley ER (2012) Watershed boundaries and geographic isolation: patterns of
diversification in cutthroat trout from western North America. *BMC Evolutionary Biology*, **12**,
38.

Malde HE (1968) The catastrophic late Pleistocene Bonneville Flood in the Snake River Plain,
Idaho. U.S. Geological Survey Professional Paper 596.

Marnell LF (1988) Status of westslope cutthroat trout in Glacier National Park, Montana.
*American Fisheries Society Symposium*, **4**, 61–70.

McDonald EV, Sweeney MR, Busacca AJ (2012) Glacial outburst floods and loess
sedimentation documented during Oxygen Isotope Stage 4 on the Columbia Plateau,
Washington State. *Quaternary Science Reviews*, **29**, 18–30.

McKelvey KS, Young MK, Wilcox TM, Bingham DM, Pilgrim KL, Schwartz MK (2016)
Patterns of hybridization among cutthroat trout and rainbow trout in northern Rocky
Mountain streams. *Ecology and Evolution*, **6**, 688–706.

McPhail JD (2007) The freshwater fishes of British Columbia. The University of Alberta Press,
Edmonton.

McPhail JD, Lindsey CC (1986) Zoogeography of the freshwater fishes of Cascadia (the
Columbia system and rivers north to the Stikine). Pages 615–637 in Hocutt CH, Wiley EO
(editors) The zoogeography of North American freshwater fishes. John Wiley & Sons, New
York.

Mee JA, Bernatchez L, Reist JD, Rogers SM, Taylor EB (2015) Identifying designatable units

for intraspecific conservation prioritization: a hierarchical approach applied to the lake

whitefish species complex (*Coregonus* spp.). *Evolutionary Applications*, **8**, 423–441.

Metcalf JL, Love Stowell S, Kennedy CM, Rogers KB, McDonald D, Epp J, Keepers K, Cooper

A, Austin JJ, Martin AP (2012) Historical stocking data and 19th century DNA reveal human-

induced changes to native diversity and distribution of cutthroat trout. *Molecular Ecology*, **21**,

5194–5207.

Minckley WL, Hendrickson DA, Bond CE (1986) Geography of western North American

freshwater fishes: description and relationships to intracontinental tectonism. Pages 519–613

in Hocutt CH, Wiley EO (editors) The zoogeography of North American freshwater fishes.

John Wiley & Sons, New York.

Moritz C, Lavery S, Slade R (1995) Using allele frequency and phylogeny to define units for

conservation and management. Pages 249–262 in (Nielsen JL, Powers GA, editors) Evolution

and the aquatic ecosystem: defining unique units in population conservation. Symposium 17.

American Fisheries Society, Bethesda, Maryland.

Neville HM, Dunham JB (2011) Patterns of hybridization of nonnative cutthroat trout and

hatchery rainbow trout with native redband trout in the Boise River, Idaho. *North American*

*Journal of Fisheries Management*, **31**, 1163–1176.

O'Connor JE (1993) Hydrology, hydraulics, and geomorphology of the Bonneville flood.

*Geological Society of America Special Papers*, **274**, 1–84.

Oviatt CG (2015) Chronology of Lake Bonneville, 30,000 to 10,000 yr BP. *Quaternary Science*

*Reviews*, **110**, 166–171.

Peters JL (2012) Late Pleistocene evolution of glacial Lake Purcell: a potential floodwater source

to the Channeled Scabland. Master's thesis, Simon Fraser University, Burnaby, BC.

Pierce K (2003) Pleistocene glaciations of the Rocky Mountains. Pages 63–76 in Gillespie AR,

Porter SC, Atwater BF (editors). The Quaternary Period in the United States: Developments

in Quaternary Science, 1. Elsevier, Amsterdam.

Pritchard VL, Abadía-Cardoso A, Garza JC (2012) Discovery and characterization of a large

number of diagnostic markers to discriminate *Oncorhynchus mykiss* and *O. clarkii*. *Molecular*

*Ecology Resources*, **12**, 918–931.

Pritchard VL, Garza JC, Peacock MM (2015) SNPs reveal previously undocumented non-native
introgression within threatened trout populations. *Conservation Genetics*, **16**, 1001–1006.
Pritchard VL, Metcalf JL, Jones K, Martin AP, Cowley DE (2009) Population structure and
genetic management of Rio Grande cutthroat trout (*Oncorhynchus clarkii virginalis*).
*Conservation Genetics*, **10**, 1209.
Ratnasingham S, Hebert PD (2013) A DNA-based registry for all animal species: the Barcode
Index Number (BIN) system. *PloS One*, **8**, e66213.
Richardson J (1836) The Fish. In: Fauna Boreali-Americana; or the zoology of the northern parts
of British America: containing descriptions of the objects of natural history collected on the
late northern land expeditions, under the command of Sir John Franklin, R.N. J. Bentley,
London. Part 3: i-xv + 1-327, Pls. 74-97.
Ryman N (1983) Patterns of distribution of biochemical genetic variation in salmonids:
differences between species. *Aquaculture*, **33**, 1–21.
Satoh TP, Miya M, Mabuchi K, Nishida M (2016) Structure and variation of the mitochondrial
genome of fishes. *BMC Genomics*, **17**, 719.
Schultz LP (1941) Fishes of Glacier National Park Montana. U.S. National Park Service.
Conservation Bulletin 22.
Seiler MB, Keeley ER (2009) Intraspecific taxonomy and ecology characterize morphological
divergence among cutthroat trout (*Oncorhynchus clarkii* ssp. Richardson) populations.
*Biological Journal of the Linnean Society*, **96**, 266–281.
Shafer A, Cullingham CI, Cote SD, Coltman DW (2010) Of glaciers and refugia: a decade of
study sheds new light on the phylogeography of northwestern North America. *Molecular*
*Ecology*, **19**, 4589–4621.
Shepard BB, May BE, Urie W (2005) Status and conservation of westslope cutthroat trout within
the western United States. *North American Journal of Fisheries Management*, **25**, 1426–1440.
Silvestro D, Michalak I (2012) raxmlGUI: a graphical front-end for RAxML. *Organisms*
*Diversity & Evolution*, **12**, 335–337.
Simon RC, Dollar AM (1963) Cytological aspects of speciation in two North American teleosts,
*Salmo gairdneri* and *Salmo clarki lewisi*. *Canadian Journal of Genetics and Cytology*, **5**, 43–
49.

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, **30**, 1312–1313.

Stearley RF, Smith GR (2016) Fishes of the Mio-Pliocene western Snake River Plain and
vicinity. I. Salmonid fishes from Mio-Pliocene lake sediments in the western Snake River
Plain and the Great Basin. Miscellaneous Publications of the Museum of Zoology, University
of Michigan, 204.

Taylor EB, Stamford MD, Baxter JS (2003) Population subdivision in westslope cutthroat trout
(*Oncorhynchus clarki lewisi*) at the northern periphery of its range: evolutionary inferences
and conservation implications. *Molecular Ecology*, **12**, 2609–2622.

Taylor HR, Harris WE (2012) An emergent science on the brink of irrelevance: a review of the
past 8 years of DNA barcoding. *Molecular Ecology Resources*, **12**, 377–388.

Thorgaard GH (1983) Chromosomal differences among rainbow trout populations. *Copeia*,
**1983**, 650–662.

Thorgaard GH, Bestgen KR, Trotter PC, Loudenslager EJ, Wheeler PA (this volume). Historical
studies on cutthroat trout taxonomy based on morphological-meristic characters, allozymes
and chromosomes. Chapter 5. Evolutionary biology and taxonomy of cutthroat trout.
American Fisheries Society, Bethesda, Maryland.

Trotter PC, McMillan B, Gayeski N, Spruell P, Berkley R (1999) Genetic and phenotypic catalog
of native resident trout of the interior Columbia River Basin. Populations of the upper Yakima
basin. Report DOE/BP-07901-1.

Trotter PC, McMillan B, Gayeski N, Spruell P, Cook MK (2001a) Genetic and phenotypic
catalog of native resident trout of the interior Columbia River Basin. Populations in the
Wenatchee, Entiat, Lake Chelan, and Methow River drainages. Report DOE/BP-00004575-2.

Trotter PC, McMillan B, Gayeski N, Spruell P, Whiteley (2001b) Genetic and phenotypic
catalog of native resident trout of the interior Columbia River Basin. Populations of the Pend
Oreille, Kettle, and Sanpoil River basins of Colville National Forest. Report DOE/BP-
00004575-1.

U.S. Fish and Wildlife Service (2003) Reconsidered finding for an amended petition to list the
westslope cutthroat trout as threatened throughout its range. *Federal Register*, **68**, 46989–
47009.

Vu NV, Kalinowski ST (2009) Twelve tetranucleotide microsatellite loci for westslope cutthroat
trout *Oncorhynchus clarki lewisi* (Salmonidae). *Conservation Genetics Resources*, **1**, 249.

Waitt RB (1985) Case for periodic, colossal jökulhlaups from Pleistocene glacial Lake Missoula.
*Geological Society of America Bulletin*, **96**, 1271–1286.

Waitt RB, Denlinger RP, O'Connor JE (2009). Many monstrous Missoula floods down
channeled scabland and Columbia valley. *Geological Society of America Field Guide*, **15**,
775–844.

Waples RS (1991) Pacific salmon, *Oncorhynchus* spp., and the definition of "species" under the
Endangered Species Act. *Marine Fisheries Review*, **53**, 11–22.

Waples RS, Gustafson RG, Weitkamp LA, Myers JM, Jjohnson OW, Busby PJ, Hard JJ, Bryant
GJ, Waknitz FW, Nelly K, Teel D (2001) Characterizing diversity in salmon from the Pacific
Northwest. *Journal of Fish Biology*, **59(sA)**, 1–41.

Ward RD (2009) DNA barcode divergence among species and genera of birds and fishes.
*Molecular Ecology Resources*, **9**, 1077–1085.

Wenburg JK, Bentzen P (2001) Genetic and behavioral evidence for restricted gene flow among
coastal cutthroat trout populations. *Transactions of the American Fisheries Society*, **130**,
1049–1069.

Whiteley AR, Spruell P, Allendorf FW (2006) Can common species provide valuable
information for conservation? *Molecular Ecology*, **15**, 2767–2786.

Williams JE, Williams RN, Thurow RF, Elwell L, Philipp DP, Harris FA, Kershner JL, Martinez
PJ, Miller D, Reeves GH, Frissell CA (2011) Native fish conservation areas: a vision for
large-scale conservation of native fish communities. *Fisheries*, **36**, 267–277.

Wilson WD, Turner TF (2009) Phylogenetic analysis of the Pacific Cutthroat Trout
(*Oncorhynchus clarki* ssp.: Salmonidae) based on partial mtDNA ND4 sequences: A closer
look at the highly fragmented inland species. *Molecular Phylogenetics and Evolution*, **52**,
406–415.

Wiltzius WJ (1985) Fish culture and stocking in Colorado, 1872-1978. Division report 12.
Colorado Division of Wildlife, Denver.

Wood SH, Clemens DM (2002) Geologic and tectonic history of the western Snake River Plain,
Idaho and Oregon. *Idaho Geological Survey Bulletin*, **30**, 69–103.

Xia X (2017) DAMBE6: new tools for microbial genomics, phylogenetics, and molecular

evolution. *The Journal of Heredity*, esx033. doi: 10.1093/jhered/esx033.

Yau MM, Taylor EB (2013) Environmental and anthropogenic correlates of hybridization

between westslope cutthroat trout (*Oncorhynchus clarkii lewisi*) and introduced rainbow trout

(*O. mykiss*). *Conservation Genetics*, **14**, 885–900.

Young MK (1995) Conservation assessment for inland cutthroat trout. U.S. Forest Service

General Technical Report RM-256.

Young MK, Harig AL (2001) A critique of the recovery of greenback cutthroat trout.

*Conservation Biology*, **15**, 1575–1584.

Young MK, McKelvey KS, Pilgrim KL, Schwartz MK (2013) DNA barcoding at riverscape

scales: assessing biodiversity among fishes of the genus Cottus (Teleostei) in northern Rocky

Mountain streams. *Molecular Ecology Resources*, **13**, 583–595.

Young SF, McLellan JG, Shaklee JB (2004) Genetic integrity and microgeographic population

structure of westslope cutthroat trout, *Oncorhynchus clarki lewisi*, in the Pend Oreille Basin in

Washington. *Environmental Biology of Fishes*, **69**, 127–142.

Table 1. Samples used in the phylogeographic analysis of westslope cutthroat trout. Site labels are in Figure 1.

| Site | 4th-code HU[1] | State/ Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|------|------|------|------|------|------|------|------|------|------|
| Albert | Middle Clark Fork | MT | 46.9213 | -114.3678 | 1 | 1 | A | | 1 |
| Beaver | Yaak | MT | 48.8174 | -115.6787 | 1 | 2 | A | | 2 |
| Bostwick | Gallatin | MT | 45.7995 | -110.9981 | 10 | 3 | A | x | 3 |
| Moose | Blackfoot | MT | 46.9227 | -112.8534 | 6 | **4** | A | x | 4 |
| Deer | Bitterroot | MT | 45.5746 | -114.4784 | 13 | 5 | A | x | 5 |
| EF Bull | Lower Clark Fork | MT | 48.1151 | -115.6868 | 11 | 6 | B | x | 6 |
| Ketchikan | North Fork Flathead | MT | 48.9771 | -114.5071 | 1 | 7 | A | | 7 |
| McGuire | Upper Kootenai | MT | 48.6896 | -115.2728 | 10 | 8 | A | x | 8 |
| Martin | Bitterroot | MT | 46.0057 | -113.8161 | 1 | 9 | A | | 9 |
| McCabe | Blackfoot | MT | 47.1324 | -113.0840 | 6 | 10 | A | x | 10 |
| Miller | Fisher | MT | 48.0827 | -115.4642 | 5 | 11 | A | x | 11 |
| Meadow | Lower Selway | ID | 45.7154 | -115.1953 | 12 | 12 | C | x | 12 |
| Norton | Upper Clark Fork | MT | 45.9811 | -112.8041 | 8 | 13 | A | x | 13 |
| Ontario | Upper Clark Fork | MT | 46.4454 | -112.3940 | 8 | 14 | A | x | 14 |
| North | Bitterroot | MT | 46.7343 | -114.6085 | 1 | 15 | A | | 15 |
| Tyler | Flint-Rock | MT | 46.6377 | -113.5047 | 1 | 16 | A | | 16 |
| Wilkes | Lower Clark Fork | MT | 47.4962 | -115.4883 | 1 | 17 | A | | 17 |
| Canuck | Moyie | MT | 48.9988 | -116.0618 | 8 | 18 | A | x | 18 |
| Ball | Lower Kootenai | ID | 48.7712 | -116.6069 | 10 | 19 | A | x | 19 |
| Rampike | Upper Coeur d'Alene | ID | 47.8197 | -115.9619 | 10 | 20 | D | x | 20 |
| Trail | Upper Coeur d'Alene | ID | 47.8291 | -116.3941 | 10 | 21 | D | x | 21 |
| West Gold | Pend Oreille Lake | ID | 47.9259 | -116.5078 | 10 | 22 | A | x | 22 |
| Avalanche | Upper Missouri | MT | 46.7439 | -111.5180 | 7 | **23** | A | x | 23 |
| Beaver | Priest | ID | 48.7489 | -116.9270 | 10 | 24 | A | x | 24 |
| Mokins | Upper Spokane | ID | 47.7826 | -116.6663 | 1 | 25 | A | | 25 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|---|---|---|---|---|---|---|---|---|---|
| NF St. Joe | St. Joe | ID | 47.4055 | -115.8382 | 1 | 26 | B | | 26 |
| Skin | Moyie | MT | 48.7728 | -116.1446 | 1 | 27 | A | | 27 |
| Slowey | Middle Clark Fork | MT | 47.2808 | -114.9731 | 1 | 28 | A | | 28 |
| Dry Wolf | Judith | MT | 46.9492 | -110.5684 | 1 | 29 | A | | 29 |
| Fish | Jefferson | MT | 45.8032 | -112.4725 | 1 | **30** | A | | 30 |
| Heller | St. Joe | ID | 47.0662 | -115.2167 | 1 | 31 | E | | 31 |
| Tributary of Armstrong | Coeur d'Alene Lake | ID | 47.5543 | -116.5993 | 1 | 32 | A | | 32 |
| NF Dupuyer | Two Medicine | MT | 48.0636 | -112.8182 | 1 | 33 | A | | 33 |
| Sawmill | Belt | MT | 47.1013 | -110.7097 | 1 | **30** | A | | 34 |
| Youngs | South Fork Flathead | MT | 47.2838 | -113.3568 | 1 | 35 | A | | 35 |
| French | Upper North Fork Clearwater | ID | 46.5143 | -115.6768 | 1 | **36** | C | | 36 |
| Osier | Upper North Fork Clearwater | ID | 46.8055 | -115.0836 | 1 | 37 | F | | 37 |
| Crooked Fork | Lochsa | ID | 46.6843 | -114.6958 | 1 | 38 | C | | 38 |
| Pete King | Lochsa | ID | 46.1929 | -115.7064 | 1 | 39 | C | | 39 |
| Baldy | South Fork Clearwater | ID | 45.9415 | -115.6750 | 1 | 40 | C | | 40 |
| Bad Luck | Upper Selway | ID | 45.8956 | -114.7553 | 1 | 41 | B | | 41 |
| Flat | Upper Selway | ID | 45.7213 | -114.8575 | 1 | 42 | C | | 42 |
| Twisp | Methow | WA | 48.3839 | -120.4342 | 5 | 43 | G | x | 43 |
| Buck | Wenatchee | WA | 47.9163 | -120.7132 | 1 | **36** | C | | 44 |
| Cedar | Pend Oreille | ID | 48.8333 | -117.4952 | 9 | **45** | A | x | 45 |
| Ninemile | Sanpoil | WA | 48.3596 | -118.7420 | 1 | 46 | A | | 46 |
| Mill | Upper Salmon | ID | 44.4940 | -114.4200 | 10 | **47** | F | x | 47 |
| Morse | Pashimeroi | ID | 44.6222 | -113.7999 | 9 | **47** | F | x | 48 |
| Jacobs Ladder | Lower Middle Fork Salmon | ID | 45.0766 | -115.3373 | 10 | 49 | F | x | 49 |
| MF Little Timber | Lemhi | ID | 44.5867 | -113.4618 | 1 | 50 | F | | 50 |
| American | South Fork Clearwater | ID | 45.7452 | -115.9635 | 1 | 51 | C | | 51 |
| Scotchman | Flint-Rock | MT | 46.3818 | -113.5519 | 1 | 52 | H | | 52 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|---|---|---|---|---|---|---|---|---|---|
| Fourmile | Smith | MT | 46.5447 | -110.7372 | 8 | 53 | A | x | 53 |
| Kraft | Swan | MT | 47.4142 | -113.7778 | 10 | **4** | A | x | 54 |
| Leiberg | Upper Coeur d'Alene | ID | 47.7209 | -116.3834 | 10 | 55 | I | x | 55 |
| NF Grouse | Pend Oreille Lake | ID | 48.4962 | -116.3481 | 1 | 56 | J | | 56 |
| NF Teton | Teton | MT | 47.9905 | -112.8143 | 1 | **23** | A | | 57 |
| SF Red | South Fork Clearwater | ID | 45.6282 | -115.4310 | 10 | 58 | A | x | 58 |
| Yoosa | Clearwater | ID | 46.4100 | -115.5864 | 7 | **45** | A | x | 59 |
| Falls | Lake Chelan | WA | 48.0810 | -120.3577 | 6 | 60 | C | x | 60 |
| Thayer | Big Hole | MT | 45.2054 | -113.2944 | 1 | **61** | A | | 61 |
| Hungery | Lochsa | ID | 46.3758 | -115.5400 | 1 | 62 | C | | 62 |
| Ross | Upper Kootenai | MT | 48.1853 | -115.9790 | 1 | 63 | K | | 63 |
| Ditch | South Fork Clearwater | ID | 45.7712 | -115.3051 | 10 | 64 | L | x | 64 |
| EF Emerald | St. Joe | ID | 46.9871 | -116.4163 | 1 | 65 | B | | 65 |
| Big | Pashimeroi | ID | 44.4416 | -113.5987 | 1 | 66 | F | | 66 |
| Bitter | Kettle HW | BC | 49.0153 | -118.1243 | 1 | 67 | B | | 67 |
| Blackbird | Middle Salmon-Panther | ID | 45.0675 | -114.3487 | 1 | 68 | F | | 68 |
| Blairmore | Oldman | AB | 49.7143 | -114.4630 | 1 | 69 | M | | 69 |
| Bluff | St. Joe | ID | 47.1668 | -115.5088 | 1 | 70 | E | | 70 |
| Boundary | Upper Middle Fork Salmon | ID | 44.5276 | -115.3004 | 1 | 71 | F | | 71 |
| Buffalo | Beaverhead | MT | 45.1646 | -113.2000 | 1 | **61** | A | | 72 |
| Cayuse | Upper North Fork Clearwater | ID | 46.5998 | -114.8931 | 1 | 73 | C | | 73 |
| Colson | Middle Salmon-Panther | ID | 45.3247 | -114.5365 | 1 | 74 | N | | 74 |
| Crazy | Fraser HW | BC | 51.0228 | -118.6053 | 1 | 75 | A | | 75 |
| Duck | Lower Middle Fork Salmon | ID | 44.8522 | -114.5220 | 1 | 76 | F | | 76 |
| Floodwood | Lower North Fork Clearwater | ID | 47.0241 | -115.9553 | 1 | 77 | F | | 77 |
| Hartley | Kootenai HW | BC | 49.5761 | -115.0255 | 1 | 78 | A | | 78 |
| Indian Grave | Lochsa | ID | 46.4787 | -115.0899 | 1 | 79 | C | | 79 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|------|----------------|----------------|----------|-----------|--------|-------|--------|--------|-------|
| John Day | Upper John Day | OR | 44.2978 | -118.5501 | 1 | 80 | O | | 80 |
| Meadow | South Fork Salmon | ID | 44.8950 | -115.3404 | 1 | 81 | P | | 81 |
| Monk | Priest HW | BC | 49.0329 | -116.9780 | 1 | 82 | A | | 82 |
| NF Elkhorn | Upper Middle Fork Salmon | ID | 44.6405 | -115.3324 | 1 | 83 | Q | | 83 |
| Pack | Pend Oreille Lake | ID | 48.6592 | -116.6244 | 1 | 84 | B | | 84 |
| Sheep | Middle Fork Flathead | MT | 48.2321 | -113.6183 | 1 | 85 | A | | 85 |
| Split | South Fork Salmon | ID | 45.0809 | -115.7844 | 1 | 86 | F | | 86 |
| Toby | Columbia HW | BC | 50.3598 | -116.5000 | 1 | 87 | A | | 87 |
| Truman | Flathead Lake | MT | 47.9938 | -114.4016 | 1 | 88 | R | | 88 |
| Twin | Upper Salmon | ID | 44.5792 | -114.4275 | 1 | 89 | B | | 89 |
| Waiparous | Bow | AB | 51.3864 | -115.1686 | 1 | 90 | A | | 90 |
| Warm Springs | Upper Salmon | ID | 44.1029 | -114.7472 | 1 | 91 | F | | 91 |
| Werner | North Fork Flathead | MT | 48.5931 | -114.3679 | 1 | 92 | S | | 92 |
| Withington | Lemhi | ID | 45.0422 | -113.7966 | 1 | 93 | F | | 93 |
| Wounded Buck | South Fork Flathead | MT | 48.2280 | -113.9705 | 1 | 94 | A | | 94 |
| Gravey | Upper North Fork Clearwater | ID | 46.5284 | -115.1336 | 1 | 95 | C | | 95 |
| Twentyfivemile | Middle Fork Flathead | MT | 48.2055 | -113.3799 | 1 | 96 | A | | 96 |
| WF Bitterroot | Bitterroot | MT | 45.5040 | -114.3334 | 1 | | Z | | 97 |
| Beaver | Blackfoot | MT | 47.0190 | -112.7674 | 1 | | Z | | 98 |
| Chimney | Blackfoot | MT | 46.7626 | -113.0219 | 1 | | Z | | 99 |
| Canyon | Bow | AB | 50.8953 | -114.7774 | 1 | | Z | | 100 |
| Picklejar 2 | Bow | AB | 50.5198 | -114.7737 | 1 | | Z | | 101 |
| Picklejar 4 | Bow | AB | 50.5188 | -114.7841 | 1 | | Z | | 102 |
| Prairie | Bow | AB | 50.8783 | -114.8885 | 1 | | W | | 103 |
| Quirk | Bow | AB | 50.8068 | -114.7533 | 1 | | W | | 104 |
| Running Rain | Bow | AB | 50.5045 | -114.9452 | 1 | | Z | | 105 |
| Silvester | Bow | AB | 50.8614 | -114.7257 | 1 | | Z | | 106 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|---|---|---|---|---|---|---|---|---|---|
| Akolkolex | Columbia HW | BC | 50.9151 | -117.8787 | 1 | | Z | | 107 |
| Botts | Columbia HW | BC | 50.7986 | -116.3615 | 1 | | Z | | 108 |
| Cedar | Columbia HW | BC | 51.2685 | -116.9912 | 1 | | K | | 109 |
| Driftwood | Columbia HW | BC | 50.8700 | -116.4734 | 1 | | Z | | 110 |
| Ladybird | Columbia HW | BC | 49.4514 | -117.8259 | 1 | | V | | 111 |
| Neale | Columbia HW | BC | 51.4593 | -117.0917 | 1 | | Z | | 112 |
| West Fisher | Fisher | MT | 48.0537 | -115.5182 | 1 | | Z | | 113 |
| Doctor | Kootenai HW | BC | 50.0911 | -116.0867 | 1 | | Z | | 114 |
| Lodgepole | Kootenai HW | BC | 49.2819 | -114.8143 | 1 | | X | | 115 |
| Lussier | Kootenai HW | BC | 50.0686 | -115.5088 | 1 | | Z | | 116 |
| Tributary to Albert | Kootenai HW | BC | 50.5663 | -115.5596 | 1 | | Z | | 117 |
| Little Eightmile | Lemhi | ID | 44.7685 | -113.4424 | 1 | | F | | 118 |
| Reservoir | Lemhi | ID | 44.7202 | -113.1261 | 1 | | F | | 119 |
| Big | Lower Middle Fork Salmon | ID | 45.0732 | -115.3535 | 1 | | Z | | 120 |
| Flume | Lower Middle Fork Salmon | ID | 44.8181 | -114.5326 | 1 | | F | | 121 |
| MF Salmon | Lower Middle Fork Salmon | ID | 44.9995 | -114.7292 | 1 | | F | | 122 |
| NF Smith | Lower Middle Fork Salmon | ID | 45.1895 | -115.3464 | 1 | | F | | 123 |
| WF Cedar | Lower North Fork Clearwater | ID | 46.9419 | -115.8490 | 1 | | T | | 124 |
| Albert | Middle Clark Fork | MT | 46.9743 | -114.2704 | 1 | | Z | | 125 |
| Second | Middle Clark Fork | MT | 47.1279 | -114.7446 | 1 | | Z | | 126 |
| Siegel | Middle Clark Fork | MT | 47.2967 | -114.7817 | 1 | | Z | | 127 |
| Beaver | Middle Salmon-Panther | ID | 45.2825 | -114.2774 | 1 | | F | | 128 |
| Beaver | Middle Salmon-Panther | ID | 45.2837 | -114.2854 | 1 | | F | | 129 |
| Salmon | Middle Salmon-Panther | ID | 45.3463 | -113.9195 | 1 | | Z | | 130 |
| Weaver | Moyie HW | BC | 49.4034 | -116.0229 | 1 | | Z | | 131 |
| Harvey | NF Flathead HW | BC | 49.2491 | -114.5677 | 1 | | Z | | 132 |
| Lynx | Oldman | AB | 49.4961 | -114.4960 | 1 | | Z | | 133 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|---|---|---|---|---|---|---|---|---|---|
| Oldman | Oldman | AB | 50.0570 | -114.5917 | 1 | | Z | | 134 |
| Twin | Oldman | AB | 50.1885 | -114.4420 | 1 | | Z | | 135 |
| Vicary | Oldman | AB | 49.8014 | -114.4648 | 1 | | Z | | 136 |
| Hunt | Priest | ID | 48.5809 | -116.7722 | 1 | | AA | | 137 |
| Lime | Priest | ID | 48.8966 | -116.9642 | 1 | | Z | | 138 |
| Bear | South Fork Salmon | ID | 44.8332 | -115.5125 | 1 | | F | | 139 |
| Halfway | South Fork Salmon | ID | 44.7791 | -115.5419 | 1 | | F | | 140 |
| NF Coeur d'Alene | Upper Coeur d'Alene | ID | 47.9586 | -116.0987 | 1 | | V | | 141 |
| Ross | Upper Kootenai | MT | 48.1831 | -115.9788 | 1 | | K | | 142 |
| WF Quartz | Upper Kootenai | MT | 48.5255 | -115.7228 | 1 | | Z | | 143 |
| Dagger | Upper Middle Fork Salmon | ID | 44.4676 | -115.3303 | 1 | | F | | 144 |
| EF Sulphur | Upper Middle Fork Salmon | ID | 44.5492 | -115.4636 | 1 | | F | | 145 |
| Garden | Upper Middle Fork Salmon | ID | 44.7427 | -115.1376 | 1 | | F | | 146 |
| MF Salmon | Upper Middle Fork Salmon | ID | 44.7304 | -115.0185 | 1 | | F | | 147 |
| MF Salmon | Upper Middle Fork Salmon | ID | 44.5670 | -115.3007 | 1 | | N | | 148 |
| Alpine | Upper Salmon | ID | 43.9046 | -114.9227 | 1 | | F | | 149 |
| Alpine | Upper Salmon | ID | 43.9068 | -114.9322 | 1 | | Z | | 150 |
| Cinnibar | Upper Salmon | ID | 44.3564 | -114.4992 | 1 | | F | | 151 |
| Island | Upper Salmon | ID | 44.0960 | -114.5929 | 1 | | Y | | 152 |
| Jordan | Upper Salmon | ID | 44.4303 | -114.7272 | 1 | | F | | 153 |
| Morgan | Upper Salmon | ID | 44.6517 | -113.9324 | 1 | | F | | 154 |
| Road | Upper Salmon | ID | 44.1569 | -114.1735 | 1 | | F | | 155 |
| Rough | Upper Salmon | ID | 44.2413 | -114.8006 | 1 | | F | | 156 |
| Salmon | Upper Salmon | ID | 44.2538 | -114.5629 | 1 | | F | | 157 |
| Salmon | Upper Salmon | ID | 44.5447 | -114.1767 | 1 | | F | | 158 |
| Salmon | Upper Salmon | ID | 44.6874 | -114.0524 | 1 | | F | | 159 |
| Salmon | Upper Salmon | ID | 44.4258 | -114.2225 | 1 | | F | | 160 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|---|---|---|---|---|---|---|---|---|---|
| Salmon | Upper Salmon | ID | 44.2482 | -114.5303 | 1 | | U | | 161 |
| Tributary to Challis | Upper Salmon | ID | 44.5037 | -114.4571 | 1 | | F | | 162 |
| Magruder | Upper Selway | ID | 45.6854 | -114.7750 | 1 | | V | | 163 |
| Trapper | Big Hole | MT | 45.6425 | -112.8711 | 9 | | | x | 164 |
| Little Boulder | Bitterroot | MT | 45.7288 | -114.2451 | 10 | | | x | 165 |
| Overwhich | Bitterroot | MT | 45.7176 | -114.0799 | 8 | | | x | 166 |
| WF Bitterroot | Bitterroot | MT | 45.5040 | -114.3334 | 9 | | | x | 167 |
| Morrell | Blackfoot | MT | 47.2992 | -113.4639 | 8 | | | x | 168 |
| Nevada | Blackfoot | MT | 46.8177 | -112.5681 | 9 | | | x | 169 |
| Snowbank | Blackfoot | MT | 47.0561 | -112.6494 | 6 | | | x | 170 |
| Musselshell | Clearwater | ID | 46.3524 | -115.7558 | 7 | | | x | 171 |
| 4th of July | Coeur d'Alene Lake | ID | 47.6106 | -116.5178 | 10 | | | x | 172 |
| Big Cherry | Fisher | MT | 48.1990 | -115.5969 | 9 | | | x | 173 |
| Cow | Fisher | MT | 48.2257 | -115.2143 | 9 | | | x | 174 |
| West Fisher | Fisher | MT | 48.0427 | -115.4746 | 9 | | | x | 175 |
| Schwartz | Flint-Rock | MT | 46.6941 | -113.8228 | 10 | | | x | 176 |
| MF Little Timber | Lemhi | ID | 44.5867 | -113.4618 | 6 | | | x | 177 |
| Big | North Fork Flathead | MT | 48.5269 | -114.3391 | 10 | | | x | 178 |
| Black Bear | South Fork Flathead | MT | 47.7446 | -113.3834 | 10 | | | x | 179 |
| Beaver | St. Joe | ID | 47.0458 | -115.4580 | 9 | | | x | 180 |
| Bird | St. Joe | ID | 47.2889 | -115.5919 | 10 | | | x | 181 |
| Snowshoe | Upper Clark Fork | MT | 46.6860 | -112.4654 | 10 | | | x | 182 |
| Spotted Dog | Upper Clark Fork | MT | 46.4483 | -112.5164 | 6 | | | x | 183 |
| Twin Lakes | Upper Clark Fork | MT | 46.0631 | -113.2235 | 8 | | | x | 184 |
| Independence | Upper Coeur d'Alene | ID | 47.8806 | -116.4098 | 10 | | | x | 185 |
| Laverne | Upper Coeur d'Alene | ID | 47.7075 | -116.3756 | 9 | | | x | 186 |
| WF Eagle | Upper Coeur d'Alene | ID | 47.7367 | -115.8078 | 10 | | | x | 187 |

| Site | 4th-code HU[1] | State/Province | Latitude | Longitude | $N$[2] | MG[3] | ND2[4] | SNP[5] | Label |
|------|----------------|----------------|----------|-----------|--------|-------|--------|--------|-------|
| Rail | Upper John Day | OR | 44.3462 | -118.5664 | 30 | | | x | 188 |
| Roberts | Upper John Day | OR | 44.3332 | -118.5781 | 22 | | | x | 189 |

[1]HU, hydrologic unit. For Canadian sites, this refers to the basin.

[2]$N$, sample size. Values = 1 except sites for which SNP analyses were also performed.

[3]MG, mitogenomic sequence. Bolded labels denote a haplotype is shared. All individuals were used in microsatellite analyses (except sample 44).

[4]ND2, haplotypes of NADH dehydrogenase 2.

[5]SNP, single nucleotide polymorphism analysis.

1001 Table 2. Haplotype diversity and variation by gene for 96 westslope cutthroat trout mitogenomic

1002 sequences.

1003

| Gene | Position | | Length | H[1] | Genetic difference (%) | V | Pi | Non-syn |
| | Start | End | | | | | | |
|---|---|---|---|---|---|---|---|---|
| ATPase 6 | 8113 | 8795 | 683 | 9 | 0.34 (0.15–0.59) | 9 | 3 | 3 |
| ATPase 8 | 7955 | 8122 | 168 | 5 | 1.19 (0.60–2.38) | 4 | 3 | 2 |
| COI | 5475 | 7025 | 1551 | 13 | 0.33 (0.06–0.64) | 17 | 12 | 0 |
| COII | 7189 | 7879 | 691 | 6 | 0.27 (0.14–0.43) | 5 | 1 | 1 |
| COIII | 8796 | 9580 | 785 | 15 | 0.56 (0.13–1.27) | 17 | 11 | 1 |
| Cyt b | 14382 | 15522 | 1141 | 16 | 0.26 (0.09–0.44) | 16 | 10 | 7 |
| ND1 | 2848 | 3822 | 975 | 17 | 0.53 (0.10–1.03) | 20 | 10 | 6 |
| ND2 | 4035 | 5084 | 1050 | 19 | 0.50 (0.10-0.95) | 22 | 10 | 9 |
| ND3 | 9651 | 9999 | 349 | 6 | 0.69 (0.29–1.15) | 6 | 4 | 1 |
| ND4 | 10360 | 11740 | 1381 | 18 | 0.39 (0.07–0.80) | 22 | 11 | 6 |
| ND4L | 10070 | 10366 | 297 | 3 | 0.45 (0.34–0.67) | 2 | 0 | 1 |
| ND5 | 11953 | 13791 | 1839 | 38 | 0.34 (0.05–0.87) | 45 | 17 | 19 |
| ND6 | 13788 | 14309 | 522 | 11 | 0.44 (0.19–0.77) | 9 | 5 | 2 |
| r12S | 69 | 1015 | 947 | 14 | 0.24 (0.11–0.42) | 12 | 4 | |
| r16S | 1088 | 2772 | 1685 | 29 | 0.25 (0.06–0.53) | 30 | 11 | |
| tRNA-Ala | 5159 | 5227 | 69 | 1 | 0 | 0 | 0 | |
| tRNA-Arg | 10000 | 10069 | 70 | 2 | 1.43 | 1 | 0 | |
| tRNA-Asn | 5229 | 5301 | 73 | 1 | 0 | 0 | 0 | |
| tRNA-Asp | 7101 | 7174 | 74 | 2 | 1.35 | 0 | 0 | |
| tRNA-Cys | 5336 | 5402 | 67 | 2 | 1.49 | 1 | 0 | |
| tRNA-Gln | 3896 | 3966 | 71 | 1 | 0 | 0 | 0 | |
| tRNA-Glu | 14310 | 14378 | 69 | 1 | 0 | 0 | 0 | |
| tRNA-Gly | 9581 | 9650 | 70 | 2 | 1.43 | 1 | 1 | |
| tRNA-His | 11741 | 11809 | 69 | 1 | 0 | 0 | 0 | |
| tRNA-Ile | 3827 | 3898 | 72 | 1 | 0 | 0 | 0 | |
| tRNA-Leu | 2773 | 2847 | 75 | 2 | 1.33 | 1 | 0 | |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| tRNA-Leu2 | 11880 | 11952 | 73 | 1 | 0 | 0 | 0 |
| tRNA-Lys | 7880 | 7953 | 74 | 2 | 1.35 | 1 | 0 |
| tRNA-Met | 3966 | 4034 | 69 | 2 | 1.45 | 1 | 0 |
| tRNA-Phe | 1 | 68 | 68 | 5 | 2.35 (1.47–2.94) | 4 | 2 |
| tRNA-Pro | 15594 | 15663 | 70 | 1 | 0 | 0 | 0 |
| tRNA-Ser | 7026 | 7096 | 71 | 1 | 0 | 0 | 0 |
| tRNA-Ser2 | 11810 | 11878 | 69 | 5 | 2.32 (1.45–2.90) | 4 | 1 |
| tRNA-Thr | 15523 | 15594 | 72 | 1 | 0 | 0 | 0 |
| tRNA-Trp | 5086 | 5157 | 72 | 2 | 1.39 | 1 | 1 |
| tRNA-Tyr | 5403 | 5473 | 71 | 3 | 1.41 | 2 | 1 |
| tRNA-Val | 1016 | 1087 | 72 | 2 | 1.39 | 1 | 1 |
| OL | 5302 | 5345 | 44 | 4 | 2.27 | 2 | 1 |
| CR | 15664 | 16676 | 1013 | 25 | 0.36 (0.10–0.89) | 16 | 9 |
| Mitogenome | 1 | 16676 | 16676 | 84 | 0.19 (0.01–0.49) | 273 | 130 |

[1]Gene abbreviations: ATP, ATPase; CO, cytochrome *c* oxidase; Cyt b, cytochrome b; ND, NADH dehydrogenase; r, ribosomal RNA; tRNA, transfer RNA; OL, origin of light-strand replication; CR, control region.

[2]Abbreviations: H, number of haplotypes; V, number of variables sites; Pi, number of parsimony-informative sites; Non-syn, number of non-synonymous mutations.

1012 Table 3. Mean pairwise genetic divergence (%) among clades (below the diagonal) and within clades (on the diagonal).

1013

| | John Day | Coeur d'Alene | St. Joe | North Fork Clearwater | Salmon A | Salmon B | Clearwater headwaters | Clearwater-E. Cascades | Neoboreal | Missouri |
|---|---|---|---|---|---|---|---|---|---|---|
| John Day | NA | | | | | | | | | |
| Coeur d'Alene | 0.20 | 0.05 | | | | | | | | |
| St. Joe | 0.19 | 0.10 | 0.01 | | | | | | | |
| North Fork Clearwater | 0.24 | 0.16 | 0.16 | 0.04 | | | | | | |
| Salmon A | 0.21 | 0.13 | 0.14 | 0.11 | 0.02 | | | | | |
| Salmon B | 0.21 | 0.13 | 0.14 | 0.11 | 0.05 | 0.03 | | | | |
| Clearwater headwaters | 0.36 | 0.41 | 0.40 | 0.44 | 0.42 | 0.42 | 0.11 | | | |
| Clearwater-E. Cascades | 0.34 | 0.39 | 0.39 | 0.43 | 0.40 | 0.40 | 0.14 | 0.04 | | |
| Neoboreal | 0.36 | 0.42 | 0.41 | 0.46 | 0.43 | 0.43 | 0.16 | 0.10 | 0.04 | |
| Missouri | 0.37 | 0.43 | 0.42 | 0.46 | 0.43 | 0.44 | 0.17 | 0.10 | 0.04 | 0.01 |

1014

1015
1016
1017

1018 Table 4. Stocking records associated with sample sites exhibiting geographically incongruent genotypes. Data are ND2 sequences,
1019 mitogenomic (MG) sequences, or SNPs. Stocking information is from the online databases for Idaho
1020 (https://idfg.idaho.gov/fish/stocking) and Montana (http://fwp.mt.gov/fishing/mFish/).
1021

| Site code | Site | State | Data | Expected genotype | Nearby stocking | Date | Number |
|---|---|---|---|---|---|---|---|
| 25 | Mokins | ID | MG | Coeur d'Alene | Hayden Lake | 1969–2007 | 4,853,000 |
| 26 | NF St. Joe | ID | MG | St. Joe | St. Joe River | 1968–1973 | 235,000 |
| 29 | Dry Wolf | MT | MG | Missouri | No data | | |
| 32 | Tributary of Armstrong | ID | MG | Coeur d'Alene | Coeur d'Alene Lake | 1967–1969 | 73,000 |
| 41 | Bad Luck | ID | MG | Clearwater headwaters | Canyon Creek Lakes, White Cap Creek Lakes | 1968–2008 | 83,000 |
| 48 | Morse[2] | ID | SNP | Salmon | No data | | |
| 55 | Leiberg | ID | MG | Coeur d'Alene | Coeur d'Alene River | 1968–1972 | 153,000 |
| 59 | Yoosa[1] | ID | MG | Clearwater-E. Cascades | No data | | |
| 65 | EF Emerald | ID | MG | St. Joe | No data | | |
| 89 | Twin | ID | MG | Salmon B | Twin Creek, WF Bear Creek | 1990–2008 | 1,500 |
| 120 | Big | ID | ND2 | NF Clearwater-Salmon | Belvidere Lakes | 1971–1989 | 26,000 |
| 130 | Salmon | ID | ND2 | NF Clearwater-Salmon | No data | | |
| 141 | NF Coeur d'Alene | ID | ND2 | Coeur d'Alene | LNF Coeur d'Alene River, Spruce Creek | 1968–1972 | 264,000 |
| 150 | Alpine | ID | ND2 | NF Clearwater-Salmon | Alpine Creek Lakes | 1970–2009 | 44,000 |
| 152 | Island | ID | ND2 | NF Clearwater-Salmon | Island Lake | 1969–1999 | 14,000 |
| 163 | Magruder | ID | ND2 | Clearwater-E. Cascades | Gold Pan Lake | 1968–1994 | 18,000 |
| 164 | Trapper | MT | SNP | Missouri | No data | | |
| 177 | MF Little Timber | ID | SNP | Salmon | NF Little Timber Lakes | 1983–1986 | 500 |

1022 [1]All fish in the SNP analysis clustered as expected.
1023 [2]The fish in the mitogenomic analysis clustered as expected.
1024
1025

1026     Table 5. Proposed units of conservation for westslope cutthroat trout.

1027

1028

| Name | Location | Evidence[1] |
|---|---|---|
| John Day | Upper portion of the John Day River basin | G, MG, MS, SNP |
| Coeur d'Alene | Upper portion of the Coeur d'Alene River basin | G, MG |
| St. Joe | Upper portion of the St. Joe River basin | G, MG |
| North Fork Clearwater | Upper portion of the North Fork Clearwater River | G, MG, ND2 (in part) |
| Salmon | Salmon River basin | G, MG, ND2 (in part), MS, SNP |
| Clearwater headwaters | Headwaters of the Selway and South Fork Clearwater Rivers | G, MG |
| Clearwater-eastern Cascades | Lochsa River, lower Selway River, and lower portions of tributaries to the Middle Fork Clearwater River, and the Wenatchee River, Lake Chelan, and Methow River basins | G (in part), MG, MS, SNP |
| Neoboreal | Columbia River basin from the Sanpoil River upstream (excluding the Spokane River), Fraser River basin, and the South Saskatchewan River basin | G, MG, ND2 (in part), MS, SNP |
| Missouri | Missouri River basin | G, MG, MS, SNP |

1029   [1]Evidence is from geographic discreteness (G), mitogenomic sequences (MG), microsatellites (MS), or single nucleotide
1030   polymorphisms (SNP).

1031

1032

1033    Figure captions.

1034

Figure 1. Sample locations, river basins, and units of conservation within the historical range of

westslope cutthroat trout. (A) Sources of westslope cutthroat trout specimens that were

genetically analyzed using mitogenomic sequences and microsatellites (squares), ND2 sequences

(diamonds), or SNPs (circles). Fish from some locations were analyzed using more than one

method. Major river basins are: 1, John Day; 2, Clearwater; 3, North Fork Clearwater; 4,

Salmon; 5, St. Joe; 6, Coeur d'Alene; 7, Wenatchee River, Lake Chelan, and Methow River in

the eastern Cascades; 8, Pend Oreille; 9, Kootenai; 10, Clark Fork; 11, Bitterroot; 12, Blackfoot;

13, Flathead; 14, Bow (north) and Oldman (south) Rivers in the South Saskatchewan; and 15,

Missouri. (B) Extent of proposed units of conservation. Colored polygons correspond to those

identifying mitogenomic clades (Salmon River clades are combined). The dotted red line denotes

the Continental Divide.

1046

1047    Figure 2. Maximum-likelihood phylogeny of individual mitogenomic haplotypes ($n = 89$) of

1048    westslope cutthroat trout from across its historical range. Support (as a percentage, based on

1049    1,000 bootstraps) is given above each node. Only support $> 70\%$ is shown. Mitogenomic clades

1050    discussed in the text are identified by a unique color. Sample labels are in Table 1.

1051

1052    Figure 3. Neighbor-joining phylogeny of individual ND2 haplotypes ($n = 27$) of westslope

1053    cutthroat trout from across its historical range. Support (as a percentage, based on 1,000

1054    bootstraps) is given above each node. Only support $> 70\%$ is shown. Mitogenomic clades

1055    discussed in the text are identified by a unique color. ND2 haplotype labels are in Table 1.

1056

Figure 4. Assignment of westslope cutthroat trout ($n = 524$) to $K = 4$, 8, or 15 genetic clusters

analyzed at 52 variable SNP loci using STRUCTURE. Each individual (denoted by a single bar) is

represented by the proportion of their genotype that was assigned to each of $K$ clusters. Samples

are sorted geographically by basin, but STRUCTURE was run with no geographic priors. The left

vertical axis identifies major drainage basins of each sample and the right vertical axis the

mitogenomic clades. Clades for the North Fork Clearwater River and Clearwater River

headwaters were not included because no fish from these areas were examined in the SNP

analysis.

1065

1066  Figure 5. Assignment of westslope cutthroat trout ($n = 95$) to $K = 5$ genetic clusters analyzed at

1067  10 microsatellite loci using STRUCTURE. Each individual (denoted by a single bar) is represented

1068  by the proportion of their genotype that was assigned to each of $K$ clusters. Samples are sorted

1069  geographically by basin, but STRUCTURE was run with no geographic priors. The left vertical axis

1070  identifies major drainage basins of each sample.

1071

1072

1073

1074

1075

B

Neoboreal

Missouri

Coeur d'Alene

St. Joe

NF Clearwater

Clearwater-
E. Cascades

John
Day

Clearwater
headwaters

Salmon

0    50    100
km

1078

1079

1080

1081

1082

1083

John Day

Salmon

SF Clearwater

MF Clearwater

E. Cascades

Coeur d'Alene

St. Joe

Lower Clark Fork

Pend Oreille

Lower Kootenai

Upper Clark Fork

Bitterroot

Blackfoot

Flathead

Upper Kootenai

Missouri

John Day

Salmon

Clearwater-
E. Cascades

Coeur d'Alene

St. Joe

Neoboreal

Missouri

K = 15          K = 8          K = 4

1087

John Day
Salmon
NF Clearwater
MF/SF Clearwater
E. Cascades
Coeur d'Alene
St. Joe
Lower Columbia
Fraser
Pend Oreille-
Lower Kootenai-
Lower Clark Fork
Upper Clark Fork
Bitterroot
Blackfoot
Flathead
Upper Kootenai-
Upper Columbia
S. Saskatchewan
Missouri

1088