1   **Running head:** Repetition suppression and prediction error

3   Prediction Error and Repetition Suppression Have Distinct Effects

4   on Neural Representations of Visual Information

Matthew F. Tang[1,2], Cooper A. Smout[1,2], Ehsan Arabzadeh[2,3] & Jason B.

Mattingley[1,2,4]

1. Queensland Brain Institute, The University of Queensland, St Lucia, QLD,

Australia

2. Australian Research Council Centre of Excellence for Integrative Brain

Function, Victoria, Australia

3. Eccles Institute of Neuroscience, John Curtin School of Medical Research,

The Australian National University, Canberra, ACT, Australia

4. School of Psychology, The University of Queensland, St Lucia, QLD,

Australia

Corresponding author

Matthew F. Tang

Queensland Brain Institute

The University of Queensland

St Lucia, QLD, Australia

Email: m.tang1@uq.edu.au

24   **Abstract**

Predictive coding theories argue that recent experience establishes expectations in

the brain that generate *prediction errors* when violated. Prediction errors provide a

possible explanation for *repetition suppression*, where evoked neural activity is

attenuated across repeated presentations of the same stimulus. The predictive

coding account argues repetition suppression arises because repeated stimuli are

expected, whereas non-repeated stimuli are unexpected and thus elicit larger neural

responses. Here we employed electroencephalography in humans to test the

predictive coding account of repetition suppression by presenting sequences of

visual gratings with orientations that were expected either to repeat or change in

separate blocks of trials. We applied multivariate forward modelling to determine

how orientation selectivity was affected by repetition and prediction. Unexpected

stimuli were associated with significantly enhanced orientation selectivity, whereas

selectivity was unaffected for repeated stimuli. Our results suggest that repetition

suppression and expectation have separable effects on neural representations of

visual feature information.

Introduction

At any moment in time, the brain receives more sensory information than can

be responded to, creating the need for selection and efficient processing of

incoming signals. One mechanism by which the brain might reduce its information

processing load is to encode successive presentations of the same stimulus in a

more efficient form, a process known as *neural adaptation* (Fairhall, Lewen, Bialek,

& de Ruyter van Steveninck, 2001; Kvale & Schreiner, 2004; Smirnakis, Berry,

Warland, Bialek, & Meister, 1997). Such adaptation has been observed across

different sensory modalities and species, and has been suggested as a potential

mechanism for enhancing the coding efficiency of individual neurons and neuronal

populations (Adibi, McDonald, Clifford, & Arabzadeh, 2013; Benucci, Saleem, &

Carandini, 2013; Maravall, Petersen, Fairhall, Arabzadeh, & Diamond, 2007). A

particular form of neuronal adaptation, known as *repetition suppression*, is

characterised by attenuated neural responses to repeated presentations of the

same stimulus (Diederen, Spencer, Vestergaard, Fletcher, & Schultz, 2016; Gross,

Schiller, Wells, & Gerstein, 1967; Keller et al., 2017; Movshon & Lennie, 1979;

Rasmussen, Schwartz, & Chase, 2017). Here we asked whether predictive coding

theory, which assumes that sensory processing is influenced by prior exposure, can

account for changes in neural representations observed with repetition suppression.

The phenomenon of repetition suppression has been widely exploited to

investigate neural representations of sensory information. Repeated exposures allow

for more efficient representation of subsequent stimuli, as manifested in improved

behavioural performance despite a significant reduction in neural activity (Henson &

Repetition suppression and prediction error

Rugg, 2003; Schacter & Buckner, 1998). Repetition suppression paradigms have

been used extensively in human neuroimaging because they are commonly

considered to be analogous to the single-cell adaptation effects observed in animal

models (see Barron, Garvert, & Behrens, 2016 for review). The exact relationship

between the effects seen in human neuroimaging studies and animal

neurophysiology has, however, yet to be fully established.

The view that repetition suppression observed in human neuroimaging

studies reflects neuronal adaptation has recently been challenged by hierarchical

predictive coding theories (Auksztulewicz & Friston, 2016; Summerfield, Trittschuh,

Monti, Mesulam, & Egner, 2008). These theories argue that the brain interprets

incoming sensory events based on what would be expected from the recent history

of exposure to such stimuli (Friston, 2005; Rao & Ballard, 1999). According to these

theories, predictions are generated within each cortical area, and are bi-directionally

propagated from higher to lower areas, including to primary sensory regions,

allowing for more efficient representation of expected stimuli. When there is a

precise expectation, incoming information can be efficiently represented by

recruiting a small pool of relevant neurons (Friston, 2005). When there is a mismatch

between an expectation and the stimulus presented, i.e., when there is a *prediction*

error, the stimulus is less efficiently represented and thus elicits a larger neural

response.

The majority of evidence for predictive coding comes from human

neuroimaging experiments in which the presentation of an unexpected stimulus

generates a larger response than the presentation of an expected stimulus. In

studies employing electroencephalography (EEG) and magnetoencephalography

(MEG), this effect is known as the *mismatch negativity* (Garrido, Kilner, Stephan, &

Friston, 2009; Näätänen, Paavilainen, Rinne, & Alho, 2007; Wacongne et al., 2011),

where an unexpected stimulus evokes significantly greater negativity than an

expected stimulus. To date, however, no study has tested a key premise of

predictive coding, namely, that expected stimuli are more efficiently encoded in the

brain relative to unexpected stimuli, in terms of their elementary feature

representations. Nor has any previous investigation examined whether the

mismatch negativity response is associated with a change in neural tuning to

stimulus features such as orientation.

To test the hypothesis that prediction error can account for repetition

suppression effects, Summerfield and colleagues (2008) introduced an experimental

paradigm in which the identity of a face stimulus was either repeated in 80% of

trials (making the repetition *expected*) or was changed in 80% of trials (making the

repetition *unexpected*). There was greater attenuation of the BOLD response in the

fusiform face area when a face repetition was expected, relative to when it was

unexpected, suggesting that repetition suppression is reduced by unexpected

stimuli. This attenuation of repetition suppression by failures of expectation has also

been replicated using fMRI (Larsson & Smith, 2012) and M/EEG, using high-level

stimuli such as faces (Summerfield, Wyart, Mareike Johnen, & de Gardelle, 2011),

and simple stimuli such as tones (Todorovic & de Lange, 2012; Todorovic, van Ede,

Maris, & de Lange, 2011).

Repetition suppression and prediction error

A potential reconciliation of the relationship between expectation and

repetition suppression comes from work showing that while expectations decrease

the overall amount of neural activity, they can also yield sharper representations of

sensory stimuli (Kok, Jehee, & de Lange, 2012). This work goes beyond

conventional neuroimaging approaches, which typically only measure overall levels

of neural activity (Buckner et al., 1998; Kourtzi & Kanwisher, 2001; Tootell, Reppas,

Dale, & Look, 1995). Such amplitude changes could in principle be produced by one

or more different types of change in the underlying neural representations. For

instance, both sharpening, where the response to only unpredicted features is

suppressed, and gain reduction, where a multiplicative suppression occurs for all

features, could be associated with decreased population activity, even though the

amount of information carried by the representations will be markedly different.

Recently introduced multivariate pattern analytic approaches to human neuroimaging

– specifically forward encoding modelling – allow for the quantification of stimulus-

selective information contained within patterns of neural activity in human observers

(Brouwer & Heeger, 2009; Garcia, Srinivasan, & Serences, 2013; King, Pescetelli, &

Dehaene, 2016; Kok, Mostert, & de Lange, 2017; Myers et al., 2015; Salti et al.,

2015; Wolff, Jochim, Akyürek, & Stokes, 2017a). This approach goes beyond typical

multivariate pattern analyses (which normally produce only accuracy scores) by

quantifying neural representations evoked by sensory stimuli to reveal both the

accuracy and the *tuning fidelity* for the specific feature-dimension of interest.

Here we used multivariate forward encoding methods to test whether

repetition suppression and expectation have different effects on the way the brain

represents visual information, in this case the orientation of grating stimuli. To

anticipate the results, we found that soon after stimulus onset, repetition

suppression had no effect on visual orientation selectivity, but violated expectations

were associated with a significantly increased orientation-selective response

through gain modulation, with no corresponding change in response fidelity. This

representation was transiently re-activated at around 200 ms post-stimulus onset,

suggesting that feedback influences initial sensory encoding of an unexpected

stimulus, which in turn allows for updating of the sensory prior.

140   **Results**

We used a modified version of the paradigm introduced by Summerfield and

colleagues (2008), replacing the face stimuli used in that study with oriented

Gabors. These low-level stimuli allowed us to quantify the degree of orientation

selectivity in EEG activity to determine how the representation of orientation is

affected by prediction error and repetition suppression. Each of fifteen observers

participated in two EEG sessions. On each trial, two Gabors were presented

sequentially (100 ms presentation, 600 ms stimulus onset asynchrony), and these

stimulus pairs either repeated or alternated in their orientation (Figure 1A, Movie 1).

The predictability of the repeated and alternating pairs was varied in a block-wise

manner to manipulate expectation. In a *repeating* block, the orientations of the two

Gabors in a pair repeated in 80% of trials, and alternated in the remaining 20%.

These contingencies were reversed in the *alternating* block (Figure 1B). The

orientations of successive stimuli across a block were randomized to limit any

accumulated effects of adaptation and prediction. As repetition suppression and

expectation form orthogonal dimensions of the task, the design allowed us to

isolate their respective contributions to neural responses. Participants completed an

unrelated task of discriminating (red vs blue) rare coloured Gabors (which occurred

on 10% of trials).

Figure 1. Example stimulus displays and task design. **(A)** Schematic of the stimuli
and timing used in the experiment. Participants viewed a rapid stream of pairs of
Gabors and monitored for an infrequent coloured target (10% of trials). The stimulus
orientations varied pseudorandomly across trials between 0° and 160° (in 20°
steps), allowing estimation of orientation-selective information contained within
patterns of EEG activity. **(B)** The orientation of the pairs of Gabors could either
repeat or alternate. In one type of block, 80% of trials were orientation repeats and
the remaining 20% alternated (Repeating blocks); in the other type of block these
contingencies were reversed (Alternating blocks).
169

Movie 1. Example of a stimulus sequence of Gabors in a typical repetition block.

Repetition suppression and prediction error affect the overall level of neural

activity

The Gabors elicited a large response over occipital-parietal areas (Figure 2A).

Consistent with previous work (Cui, Wang, Park, Demb, & Butts, 2016; Keller et al.,

2017; Rentzeperis, Nikolaev, Kiper, & Van Leeuwen, 2012; Summerfield et al., 2011;

Todorovic et al., 2011; Todorovic & de Lange, 2012; Tootell et al., 1998), there was

a significant repetition suppression effect (Repeat < Alternating), such that the

response to repeated stimuli was significantly reduced compared with the response

to alternating stimuli (Figure 2A). The repetition suppression effect was evident over

a large cluster of occipital-parietal electrodes at two time intervals: an early effect

from 79 to 230 ms, and a later effect at 250 to 540 ms after the onset of the second

stimulus (cluster $p < .025$; Figure 2B and caption). A large cluster of frontal

electrodes mirrored the repetition suppression effect with a similar time course: the

ERP over these frontal sites had the same pattern, but was reversed in sign,

Repetition suppression and prediction error

187   suggesting it originated from the same dipole as the occipital response.

Repetition suppression and prediction error

Figure 2. Univariate EEG results for the effect of repetition suppression and
expectation on the second stimulus in a pair. Panels A and B show the main effects
of repetition suppression and expectation, respectively, over three post-stimulus
epochs (100-200ms, 200-300 ms, 300-400 ms) and across all electrodes. The main
effect of repetition suppression is displayed as Repeating minus Alternating trials.
The main effect of expectation is displayed as Expected minus Unexpected trials.
Circles indicate clusters of electrodes with significantly reduced activity, and
crosses indicate clusters of electrodes with significantly increased activity (alpha p
< .05, cluster p < .025, N permutations = 1500). **(C)** Bandpass filtered (2-40 Hz)
event-related potentials (ERPs) for the two conditions, averaged over occipital-
parietal electrodes (O1, O2, Oz, POz, PO7, PO3, PO8, PO4, P3, Pz, P2). A peak
analysis was conducted to aid comparison with previous studies. Orange shading
indicates the P1 component; green shading indicates the N1 component. **(D)** Peak
analysis results for P1 and N1 components. Note that the plotted values represent
differences between conditions, as indicated, rather than condition-specific evoked
responses. Asterisks indicate p <.05. Error bars indicate ±1 standard error.

Repetition suppression and prediction error

Also consistent with previous results (Garrido et al., 2009; Summerfield et al.,

2011; Todorovic et al., 2011; Todorovic & de Lange, 2012), there was a significant

expectation effect (Expected < Unexpected). Specifically, there was a significantly

greater negativity for unexpected versus expected stimuli, and this effect was most

prominent over a cluster of occipital-parietal electrodes around 75-150 ms after

stimulus presentation (Figure 2C). As with the repetition suppression result

described above, there was an expectation effect of opposite polarity over

occipital-parietal electrodes. This effect was significant at an early time point post-

stimulus (79-130 ms), but not at later time points (320-390 ms; Figure 2D). Finally,

there was no interaction between repetition suppression and expectation (i.e., no

significant positive or negative clusters, all $p > .05$). Taken together, these results

reveal both repetition suppression and expectation effects in the neural data, which

were indexed separately as shown in Figure 2.

We conducted a further traditional peak analysis, to aid comparison with

previously published studies on the mismatch negativity (Garrido, Sahani, & Dolan,

2013; Näätänen et al., 2007; Saarinen, Paavilainen, Schöger, Tervaniemi, &

Näätänen, 1992). We bandpass filtered the ERPs (2-40 Hz) to recover the

stereotypic waveform (Figure 2C) and examined two classic early components – the

N1 and P1 – averaged across a broad grouping of occipital-parietal electrodes (O1,

O2, Oz, POz, PO7, PO3, PO8, PO4, P3, Pz, P2). As in previous studies (Caharel,

d'Arripe, Ramon, Jacques, & Rossion, 2009; Dehaene et al., 2001), we defined the

P1 as the largest positivity between 80 and 110 ms after stimulus presentation, and

the N1 as the largest negativity between 90 and 130 ms after stimulus presentation.

Repetition suppression and prediction error

A relatively wide temporal window was used to capture inter-individual response

variation. As expected, for the P1 component the repeated stimulus evoked a

significantly smaller positivity ($t(14) = 3.03$, $p = .009$) than the alternating stimulus

(Figure 2D), reflecting a repetition suppression effect. There was no such effect of

expectation on the P1 ($t(14) = 0.26$, $p = .80$). By contrast, as predicted from previous

work (Garrido et al., 2013; Näätänen et al., 2007; Saarinen et al., 1992), analysis of

the N1 component showed that the unexpected stimulus evoked a significantly

greater negativity than the expected stimulus, ($t(14) = 5.75$, $p < .0001$). The

repetition suppression effect was also present in the N1 ($t(14) = 2.39$, $p = .03$), but

critically in the opposite direction as the expectation effect.

Expectations increase orientation-selective information contained within

patterns of EEG activity

We next examined the key question of whether repetition suppression and

expectation differentially affect neural representations of orientation information. To

do this, we used a forward encoding approach to reconstruct orientation-selective

information contained within the multivariate pattern of EEG activity distributed

across the scalp (Figure 3; see Methods for details). Briefly, this technique

transforms sensor-level responses into tuned 'feature' channels (Brouwer & Heeger,

2009; Garcia et al., 2013; Kay, Naselaris, Prenger, & Gallant, 2008; Myers et al.,

2015), in this case, orientation-selective features. For each trial, the presented

orientation was convolved with a canonical, orientation-selective tuning function

and regressed against the pattern of EEG activity across all sensors at each time

point. This created a spatial filter of the multivariate EEG activity that differentiated

orientations (Figure 3D). These weights were then inverted to reconstruct the model,

and multiplied against an independent set of test trials to produce responses in the

modelled orientation channels. These sets of responses were then used to evaluate

the degree of orientation selectivity in those trials. The procedure was repeated for

all time points in the trial, and a cross-validated approach was used until all trials

had been used for both training and testing.

As shown in Figure 3, the forward encoding revealed a strong, orientation-

selective response derived from the multivariate pattern of EEG activity. This

orientation-tuned response was evident from ~50 to ~470 ms after stimulus onset,

and peaked between ~120-250 ms (Figure 3C). Examination of the regression

weights revealed that this response was largely driven by activity centred over

occipital-parietal areas (Figure 3D).

Figure 3. Results of the forward encoding modelling for orientation-selectivity. **(A)** Time-resolved orientation tuning curve across all participants and conditions in response to the second Gabor in the pair. The forward encoding approach resulted in a tuning curve for each of the nine presented orientations. These tuning curves were then centred at each presented orientation (here labelled as 0°) to combine across all orientations. The orientation-selective response is contained within the overall pattern of EEG; activity begins soon after stimulus onset and peaks at around 250 ms before declining. **(B)** Population tuning curve of the stimulus reconstruction across participants, averaged between 50-100 ms and 150-250 ms after stimulus presentation. Each line is a fitted Gaussian response with a variable offset used to quantify orientation selectivity. Error bars indicate ±1 standard error of the mean across participants. **(C)** Amplitude of the channel response over time, averaged across all conditions (black line). The thick black line indicates significant encoding of stimulus orientation based on a cluster-permutation test across participants (cluster $p < .05$, N permutations = 20,000). Encoding accuracy was reliable from 52 to 470 ms post-stimulus onset. The error shading (in grey) indicates bootstrapped 95% confidence intervals of the mean. **(D)** Topographic plots of the weights (averaged across the 9 orientation channels across all participants) derived from forward encoding at the corresponding time points shown in panel B. (a.u. = arbitrary units).

Repetition suppression and prediction error

To examine our central question of whether repetition suppression and

expectation have differential effects on neural representations of orientation, we

split and averaged the results of the forward encoding by trial type, and fitted these

with Gaussians (see Methods) to quantify orientation selectivity (Figure 4).

Repetition suppression did not affect the amount of orientation selectivity contained

within the EEG data, with similar selectivity for repeated and alternating trials. This

was the case even though the repeated trials had a markedly smaller EEG response

over occipital and parietal electrodes (see Figure 2A), where the forward encoding

model was maximally sensitive. This result is consistent with the 'efficient

representation' hypothesis of repetition suppression (Gotts, Chow, & Martin, 2012),

which argues that the overall neural response is smaller with repetition suppression

due to more efficient coding of stimulus information.

Repetition suppression and prediction error

Figure 4. The effect of repetition suppression and expectation on orientation selectivity measured using forward encoding modelling. **(A)** Amount of orientation-selective information (given by the amplitude of the fitted gaussian) from the EEG signal in response to the second Gabor in a pair, shown separately for repetition suppression (upper panel) and expectation (lower panel). The thick black line indicates significant differences between the conditions (two-tailed cluster-permutation, alpha *p* < .05, cluster alpha *p* < .05, N permutations = 20,000). (B) Population tuning curves averaged over the significant time period (79 – 185 ms) depicted in panel A. The curves, shown as fitted Gaussians, illustrate how overall stimulus representations are affected by repetition and expectation. While there was no difference in orientation tuning for repeated versus alternate stimuli (upper panel), the amplitude of the orientation response increased significantly, and the baseline decreased, for unexpected relative to expected stimuli. Error bars indicate ±1 standard error.

Examining the effect of expectation revealed a markedly different pattern of

results. As shown in Figure 4A, at 79 - 185 ms after the onset of the second

stimulus in the pair, orientation-selectivity increased significantly (*p* < .0001) when

the stimulus was unexpected relative to when it was expected, and this effect arose

at the earliest stages of the brain's response to that stimulus. Moreover, the

expectation signal contained enhanced information about the specific features of

the stimulus that violated the expectation, in this case the orientation of the second

grating. We conducted the same statistical tests on the three other parameters

defining the Gaussian function (namely, the width, centre orientation and baseline)

to determine how repetition suppression and expectation might affect other

properties of the neural representation. There was no reliable influence of repetition

suppression on any of these Gaussian parameters (all p > .32). For expectation,

there was a significant decrease in baseline activity over the same time period as

observed for the increase in amplitude (79-185 ms, p = .001), but there were no

significant effects for the other parameters (all ps > .30).

We followed up this initial analysis to ensure we did not miss any small

effects of repetition suppression or expectation on any aspects of stimulus

representation. We increased the signal-to-noise by averaging the stimulus

reconstruction over this early time period (79-185 ms after stimulus presentation),

and fitted Gaussians to each participant's data individually (Figure 4B). This again

showed that the amplitude of the response was significantly ($t(14)$ = 3.34, p = .005)

higher for unexpected (M = 0.67, SE = 0.06) than for expected (M = 0.41, SE = 0.03)

stimuli. By contrast, the width of the representations was similar for unexpected (M

= 29.62°, SE = 4.72°) and expected (M = 26.72°, SE = 2.74°) stimuli, ($t(14)$ = 0.78,

p = .45). There was also a small but non-significant ($t(14)$ = 1.94, p = .06) trend for a

smaller baseline response (i.e., non-orientation tuned activity) in the unexpected

(M = -0.01, SE = 0.07) than in the expected (M = 0.13, SE = 0.02) condition. For

comparison, we also averaged the same time period for the repetition suppression

conditions, and found similar curves for the repeated and alternating trials (*all ps >*

.18). This analysis confirms the previous result, which employed more conservative

nonparametric cluster-based testing.

It might be argued that the particular baseline period we chose for the

encoding analyses - namely from -100 to 0 ms before the onset of the second

Gabor in each pair – biased the results by incorporating a purely top-down

expectation template triggered by the orientation of the first Gabor (Kok et al.,

2017). To rule out this possibility, we performed a further forward encoding analysis

where we baselined the raw EEG data to the mean activity from -100 to 0 ms before

the *first* Gabor in each pair. Critically, this control analysis involved a baseline period

over which it was not possible to form a top-down expectation of the orientation of

the second Gabor based on the orientation of the first. This analysis yielded the

same pattern of results as the original analysis (Supplementary Figure 1), such that

the unexpected stimulus evoked significantly greater orientation selectivity than the

expected stimulus ($p = .02$). Also in line with the original analyses, the width of the

representation was not affected by expectation ($p = .44$), and there was no effect of

repetition suppression on orientation selectivity ($p = .64$). We can thus be confident

that the effect of expectation on orientation selectivity that we report here, based on

our forward encoding analyses, is not an artefact of the baselining procedure.

We also used a number of approaches to determine whether repetition

suppression and expectation interacted to affect orientation selectivity. First, we

took the difference scores between the combination of factors (e.g., expected

repetition minus unexpected repetition, and expected alternation minus unexpected

alternation) and compared these using the same cluster-based permutation testing

outlined above. This analysis revealed no significant interactions between the

factors for any parameter (all *ps* > .10). Second, we found the largest orientation-

selectivity, defined by the amplitude of the fitted Gaussian, across the 600 ms

following stimulus presentation. For each participant, this resulted in a single value

for the four conditions. Each of these values was subjected to a two-way repeated-

measures ANOVA, which again revealed no significant interaction between the

factors (all *ps* > .30)

To further examine whether orientation-selectivity contained within the overall

pattern of EEG activity differed for unexpected and expected stimuli, we used

multivariate discriminant analysis to determine whether more traditional backward

decoding (Grootswagers, Wardle, & Carlson, 2017; Kamitani & Tong, 2005; King,

Gramfort, Schurger, Naccache, & Dehaene, 2014; Marti, King, & Dehaene, 2015)

produces the same pattern of results as that yielded by the forward encoding

approach described above. The same cross-validation procedure was used as in

the forward encoding analysis, but accuracy was now defined as the proportion of

trials labelled with the correct orientation. To facilitate comparison with the results

of Kok et al., (2013), we took the peak classification accuracy within a 600 ms

window after presentation of the second grating within each pair. This analysis

confirmed the results of the forward encoding: orientations shown in unexpected

trials were classified better than orientations shown in expected trials (F(1,14)

76.42, $p <. 00001$). Again, there was no effect of repetition on classification

accuracy (F(1,14) = 0.027, p = .87); nor was there a significant interaction (F(1,14) =

2.52, p = .13). This suggests the finding is not specific to the analysis method but

rather reflects how expectation affects the representation of sensory information in

general.

Figure 5. Peak (naive Bayes) classification accuracy of the presented grating
orientation for expected and unexpected conditions. The dotted line indicates
chance performance (1/9 orientations). The error bars indicate ±1 standard error of
the mean.
397
Expectation affects the temporal stability of stimulus representations

Next, we examined whether repetition suppression and expectation affected

dynamic, ongoing stimulus representations by using cross-temporal generalization

(King & Dehaene, 2014; King et al., 2014; Myers et al., 2015; Spaak, Watanabe,

Funahashi, & Stokes, 2017; Stokes et al., 2013). To do this, we used the same

forward encoding approach as in the previous analysis, but now the weights were

derived from one time-point on one set of trials, and then applied at every time point

in the test trials. Again, a cross-validation approach was used, with all trials serving

as both training and test. This analysis examined whether the same spatial pattern

of EEG activity that allows for orientation selectivity generalizes to other time points,

thus revealing whether the pattern of orientation-selective activity is stable or

changes over time.

410   As shown in Figure 6, optimal orientation selectivity was on-axis (training

411   time equals test time) between 100 ms and 300 ms after stimulus presentation,

412   suggesting that the stimulus representation changed dynamically over time (King &

413   Dehaene, 2014). There was also significant off-axis orientation-selectivity from 100-

414   500 ms after stimulus presentation, suggesting that some aspects of the neural

415   representation of orientation were stable over time.

416
Figure 6. Cross-temporal generalization of the forward encoding model based on
grating orientations for the main effects of repetition suppression (upper panels) and
expectation (lower panels). The maps have been thresholded (indicated by opacity)
to show clusters (black outlines) of significant orientation selectivity (permutation
testing, cluster threshold p < .05, corrected cluster statistic *p* < .05, 5,000
permutations). The difference between the conditions is shown in the right-hand
column (permutation testing, cluster threshold *p* < .05, corrected cluster statistic *p* <
.05). Opacity and outlines indicate significant differences.
425
426

Repetition suppression and prediction error

There was no effect of repetition suppression on temporal generalization of

orientation information (upper panels of Figure 6), suggesting that repetition

suppression did not affect the temporal stability of neural representations of

orientation. Examining the effect of expectation on cross-temporal generalization

confirmed that there was significantly more on-axis orientation selectivity when the

stimulus was unexpected than when it was expected (cluster $p = .02$). This

increased on-axis orientation selectivity generalized off-axis at around 300-400 ms

after stimulus onset (cluster $p = .01$), suggesting that the same representation that is

activated to process the expectation is reactivated later as the stimulus continues to

be processed. Such a signal could constitute the prior of the prediction, as this

should be updated on the basis of incoming sensory evidence, which in turn would

likely require reactivation of the unexpected stimulus.

439   **Discussion**

Our findings demonstrate that repetition suppression and expectation have

distinct effects on neural representations of simple visual stimuli. Repetition

suppression had no effect on orientation selectivity, even though the neural

response to repeated stimuli was significantly reduced over occipito-parietal areas.

Unexpected stimuli, on the other hand, showed significantly increased orientation-

selectivity relative to expected stimuli. This same early representation of the

unexpected stimulus appeared to be reactivated at 200-300 ms after the initial

neural response, supporting the idea that sensory expectations may be updated

through comparison with incoming sensory evidence. These results suggest that

Repetition suppression and prediction error

repetition suppression and expectation are separable and independent neural

computations.

Our work provides a significant advance in understanding how predictions

allow the brain to process incoming sensory information by comparing what is

expected with what actually occurs. How expectations affect neural responses has

been extensively investigated using mismatch negativity paradigms in which an

unexpected stimulus causes a larger neural response than an expected stimulus

(Bekinschtein et al., 2009; Garrido et al., 2009; Näätänen et al., 2007). Such

mismatch responses to an unexpected stimulus have often been attributed to the

generation of a prediction error that updates expectation based on a conflict

between sensory evidence and the prior (Garrido et al., 2009). To date, however,

most studies have focused exclusively on the overall magnitude of neural responses

to unexpected events, rather than assessing the quality of stimulus-specific

information potentially contained within such responses. As noted above, enhanced

neural activity to unexpected visual events could reflect a differential response to

one of a number of possible stimulus features, or simply an increase in baseline

activity associated with a non-selective response. By examining how expectation

affects the representation of an elementary feature dimension – in this case,

orientation – our results imply the operation of at least two distinct neural processes

at separate times following stimulus onset. Incoming sensory information is first

evaluated against the prior (which occurs early after stimulus presentation). When

an unexpected stimulus is detected and generates a prediction error, the

representation is amplified through gain enhancement. Later, around 300 ms after

stimulus presentation, this same representation is reactivated to update the

expectation against the initially predicted representation.

According to predictive coding theory, expected stimuli should be more

efficiently represented than unpredicted stimuli largely because the reduced neural

response still encodes stimuli with the same fidelity (Friston, 2005). A more efficient

response could be due to sharpening of neuronal tuning to stimulus features, or to a

reduction in the gain of evoked neural responses. Our results strongly support the

latter interpretation. Specifically, there was no evidence that a fulfilled expectation

leads to a sharper representation of orientation information. Our findings might

imply that the brain needs to have more information about an unexpected stimulus,

so a correct response can be made. Our findings thus provide a novel insight into

how predictive coding might change neural representations of sensory information.

The lack of evidence for sharpening of neural tuning in the current results is

in contrast to the findings of a previous study (Kok et al., 2012), in which a high-level

prediction error led to 'sharper' multivariate decoding for expected versus

unexpected visual stimuli. In their study, Kok et al. (2012) used an auditory tone to

cue the orientation of a subsequent visual stimulus, and found significantly reduced

off-label classification accuracy for predicted than for unpredicted stimuli. They

concluded that predictions cause sharpening of stimulus representations. More

recently, using the same task combined with a forward encoding approach, Kok et

al. (2017) showed that response gain is increased for a predicted stimulus.

It is natural to ask why the results of the current study differ from those of

Kok and colleagues outlined above. One possible explanation lies in the different

approaches used to generate expectations across the studies. Specifically, whereas

Kok et al. manipulated expectations by pairing an auditory cue with a visual

stimulus, we exploited the properties of the visual stimuli themselves (i.e., their

orientation) to generate expectations within blocks of trials. An intriguing possibility

is that predictions requiring integration of sensory events from two or more

modalities lead to increased gain, whereas predictions made within a single sensory

modality lead to decreased gain. This might in turn relate to the noted differences

between simple 'local' and higher-order 'global' type predictions (Bekinschtein et

al., 2009; King et al., 2014), which lead to distinct patterns of stimulus-selective

decoding. A similar discrepancy relating to the effects of attention on sensory

representations has been widely discussed, with some studies finding sharpening of

stimulus representations with attention, and others showing gain enhancement (Liu,

Larsson, & Carrasco, 2007; Maunsell, 2015; Maunsell & Treue, 2006; Treue &

Trujillo, 1999). The differences between these results may potentially have arisen

because the tasks relied upon different types of attention (e.g., spatial versus

feature-based). Future studies could determine whether this same divergence

occurs for prediction effects.

The current work applied multivariate model-based approaches to EEG data

to determine how prediction and repetition suppression affect neural

representations of perceptual information. We chose to use EEG so we could

recover the temporal dynamics of these effects – something that would not be

possible with the BOLD signal used in fMRI – and because EEG is the most widely-

used tool for measuring expectation effects in human participants (see Garrido, et

al., 2019 and Paavilainen, 2013 for review), thus facilitating comparison of our

findings with those of other studies. We estimated orientation-selectivity using all

EEG electrodes distributed across the scalp for two principal reasons. First, we

wanted to limit experimenter degrees of freedom (Simmons, Nelson, & Simonsohn,

2011) potentially introduced through the post-hoc selection of subsets of

electrodes. Second, given the broad spatial resolution of EEG, we reasoned that

activity recorded from electrodes at any given scalp location could potentially carry

important feature-selective information from a number of neural sources. The results

revealed that orientation-selective information appears largely driven by electrodes

over occipital-parietal regions (Figure 3D), consistent with a number of previous

studies that employed visual decoding of M/EEG data (Cichy, Pantazis, & Oliva,

2014; Cichy, Ramirez, & Pantazis, 2015; Stokes, Wolff, & Spaak, 2015). As noted

above, however, it is entirely possible that the effects we observed here arose from

sources well beyond the occipital and parietal regions, or even potentially outside

the visual cortical hierarchy. Limitations in the temporal and spatial resolution of

current human imaging methods make it impossible to pinpoint the timing and

location of interactions between visual areas that might reflect the cascade of

predictions and prediction errors involved in sensory encoding. By combining the

current paradigm and multivariate modelling with invasive recordings in animal

models – for example using calcium imaging or extracellular electrode recordings –

it should be possible to test some of the key claims of predictive coding theory that

we have examined here, but at the level of individual neurons.

Surprisingly few studies have used invasive recording methods to test how

predictive coding affects stimulus representations at the neuronal level. One study

in macaques (Kaliukhovich & Vogels, 2010) used a design similar to that of

Summerfield and colleagues, but with high-level objects (fractals and real-world

objects) as stimuli. That study found that expectation did not attenuate repetition

suppression in either spiking activity or local field potentials within the inferior

temporal cortex. A later fMRI study in humans (Kovács, Kaiser, Kaliukhovich,

Vidnyánszky, & Vogels, 2013) used a similar stimulus set, and also found no

attenuation of repetition suppression by expectation in the same cortical region. A

follow-up study provided a potential explanation for these findings by showing that

the attenuation of neural responses associated with repetition suppression are

found with familiar stimuli, but not with unfamiliar stimuli (Grotheer & Kovács, 2014).

Viewed in this light, the stimulus sets used by Kaliukhovich and Vogels (2010) might

not have been sufficiently familiar to yield effects of expectation in their non-human

primate observers.

Other work has shown that context plays an important role in determining

the magnitude of neuronal responses to sensory events. Thus, for example,

(Ulanovsky, Las, & Nelken, 2003) found that rare auditory stimuli generate

significantly larger responses in primary auditory cortical neurons than more

commonly occurring stimuli. This result has been interpreted as a single-neuron

analogue of the mismatch negativity, but the design used in the study did not

control for adaptation effects, thus making it difficult to draw an unambiguous

comparison with the current work. In the visual domain, oddball stimuli have also

been found to modulate neuronal activity in rats, characterised by an enhancement

of responses in the higher-order latero-intermediate area (Vinken, Vogels, & Op de

Beeck, 2017). Moreover, Fiser et al. (2016) found that neurons in mouse primary

visual cortex show a greater response when task-irrelevant visual stimuli that had

been presented during training were omitted, suggesting that an established

expectation had been violated. This result is consistent with the literature on the

mismatch negativity, in which the omission of an expected stimulus results in a

large prediction error (Garrido et al., 2009; Wacongne et al., 2011). In non-human

primates, neurons in the inferior temporal cortex show an enhanced response to

unexpected relative to expected stimuli (Kaposvari, Kumar, & Vogels, 2018), and

population decoding accuracy is higher for unexpected compared with expected

stimuli (Kumar, Kaposvari, & Vogels, 2017). Critically, however, no study has

simultaneously recorded neuronal activity in multiple cortical regions to determine

whether predictions generated in one area refine responses in a second area, as

postulated by predictive coding theory (Friston, 2005; Rao & Ballard, 1999). Such a

direct demonstration is necessary to provide a strong test of the central notion that

cortical areas pass signals between themselves in order to generate expectations.

Unlike the effects of expectation, there is a large body of electrophysiological

work showing that sensory adaptation influences neuronal activity (Adibi et al.,

2013b; Adibi, Clifford, & Arabzadeh, 2013a; Felsen et al., 2002; Kohn & Movshon,

2004; Patterson, Wissig, & Kohn, 2013). For instance, there is a sharpening of

stimulus selectivity in MT neurons following 40 s of adaptation to a drifting grating

(Kohn & Movshon, 2004). As we have highlighted, however, prolonged adaptation is

likely also associated with a significant predictio8n that the next stimulus will be the

same as the previous one. Perhaps more relevant to the current results, Patterson

et al. (2013) found that the width of orientation tuning in V1 is only marginally

sharpened following brief (400 ms) periods of adaptation. Again, however, their

study did not control for expectation, so it is impossible to determine the role of

predictive coding in their observations. Our finding that repetition suppression did

not affect the bandwidth of orientation selectivity measured using EEG is also

consistent with models of orientation adaptation based on human psychophysical

data, which suggest that adaptation does not affect the tuning width of the adapted

neural populations (Clifford, 2002; 2014; Dickinson, Almeida, Bell, & Badcock, 2010;

Dickinson, Morgan, Tang, & Badcock, 2017; Tang, Dickinson, Visser, & Badcock,

2015).

598   In summary, we have shown that repetition suppression and expectation

differentially affect the neural representation of simple, but fundamental, sensory

features. Our results further highlight how the context in which a stimulus occurs,

not just its features, affect the way it is represented by the brain. Our findings

suggest encoding priority through increased gain might be given to unexpected

events, which in turn could potentially speed behavioural responses. This prioritized

representation is then re-activated at a later time period, supporting the idea that

feedback from higher cortical areas reactivates an initial sensory representation in

early cortical areas.

Method

Participants

A group of 15 healthy adult volunteers (9 females, median age = 20.5 yr,

range = 18 to 37 yr) participated in exchange for partial course credit or financial

reimbursement (AUD$20/hr). We based our sample size on work that investigated

the interaction between repetition suppression and prediction error (Summerfield et

al., 2008), and that used forward encoding modelling to investigate orientation

selectivity using MEG with a comparable number of trials as the current study

(Myers et al., 2015). Each person provided written informed consent prior to

participation, and had normal or corrected-to-normal vision. The study was

approved by The University of Queensland Human Research Ethics Committee and

was in accordance with the Declaration of Helsinki.

Experimental setup

The experiment was conducted inside a dimly illuminated room with the

participants seated in a comfortable chair. The stimuli were displayed on a 22-inch

LED monitor (resolution 1920 x 1080 pixels, refresh rate 120 Hz) using the

PsychToolbox presentation software (Brainard, 1997; Pelli, 1997) for MATLAB

(v7.3). Viewing distance was maintained at 45 cm using a chinrest, meaning the

screen subtended 61.18° x 36.87° (each pixel 2.4' x 2.4').

Visual task

The stimuli were Gabors (diameter: 5°, spatial frequency: 2 c/°, 100%

contrast) presented centrally in pairs for 100 ms, separated by 500 ms (600 ms

stimulus onset asynchrony) with a variable (650 to 750 ms) inter-stimulus interval

between trials. Across the trials, the orientations of the Gabors were evenly spaced

between 0º and 160º (in 20º steps) so we could reconstruct orientation selectivity

contained within the EEG response using forward encoding modelling. The

relationship of the orientations of the pairs Gabors was also used to construct the

different repetition suppression and prediction conditions. The orientation presented

in the second Gabor in the pair could either repeat or alternate with respect to the

orientation of the first Gabor. In the alternation trials, the orientation of the first

Gabor was drawn randomly, without replacement, from an even distribution of

orientations that was different to the orientation of the second Gabor. To vary the

degree of prediction, in half of the blocks 80% of the trials had repeated

orientations and 20% of the trials had alternating orientations, whereas in the other

half of the blocks these contingencies were reversed. This design allowed us to

separately examine the effects of repetition suppression and prediction because of

the orthogonal nature of the blocked design. The blocks of 135 trials (~3 mins)

switched between the expectation of a repeating or alternating pattern, with the

starting condition counterbalanced across participants.

The participants' task was to monitor the visual streams for rare, faintly

coloured (red or green) Gabors and to discriminate the colour as quickly and

accurately as possible. Any trial with a coloured target was excluded from analysis.

The orientation match between the pairs was made to be consistent with the

dominant contingency (i.e., repeated or alternating) within that block. Pilot testing

was used prior to the main experiment to set the task at approximately threshold, to

ensure that participants focused exclusively on the colour-discrimination task rather

654 than the orientation contingencies associated with prediction and repetition. Only

655 one participant reported being aware of the changing stimulus contingencies across

656 the blocks when asked at the end of the experiment, and excluding this

657 participant's data had no effect on the key results reported here. Self-paced breaks

658 were provided between each of the 20 blocks within a session, at which time

659 feedback was provided on performance in the preceding block. Each participant

660 completed two sessions of 2700 trials each (5400 trials in total), with each session

661 lasting around 70 mins of experimental time and 45 mins of EEG setup.

662 **EEG acquisition and pre-processing**

663  Continuous EEG data were recorded using a BioSemi Active Two system

664 (BioSemi, Amsterdam, Netherlands). The signal was digitised at 1024 Hz sampling

665 rate with a 24-bit A/D conversion. The 64 active scalp Ag/AgCl electrodes were

666 arranged according to the international standard 10–20 system for electrode

667 placement (Oostenveld & Praamstra, 2001) using a nylon head cap. As per BioSemi

668 system design, the common mode sense and driven right leg electrodes served as

669 the ground, and all scalp electrodes were referenced to the common mode sense

670 during recording.

671  Offline EEG pre-processing was performed using EEGLAB in accordance

672 with best practice procedures (Bigdely-Shamlo, Mullen, Kothe, Su, & Robbins,

673 2015; Keil et al., 2014). The data were initially down-sampled to 256 Hz and

674 subjected to a 0.5 Hz high-pass filter to remove slow baseline drifts. Electrical line

675 noise was removed using *clean_line.m,* and *clean_rawdata.m* in EEGLAB (Delorme

676 & Makeig, 2004) was used to remove bad channels (identified using Artifact

Subspace Reconstruction), which were then interpolated from the neighbouring

electrodes. Data were then re-referenced to the common average before being

epoched into segments around each stimulus pair (-0.5 s to 1.25 s from the first

stimulus in the pair). Systematic artefacts from eye blinks, movements and muscle

activity were identified using semi-automated procedures in the SASICA toolbox

(Chaumon, Bishop, & Busch, 2015) and regressed out of the signal. After this stage,

any trial with a peak voltage exceeding ±100 uV was excluded from the analysis.

The data were then baseline corrected to the mean EEG activity from -100 to 0 ms

before the presentation of the second Gabor in the pair. Critically, the orientations

of the first and second gratings were precisely balanced across the conditions to

avoid any systematic bias in orientation information being carried forward by the

first grating within each pair. Specifically, for every unexpected stimulus presented

in the second grating there was an equal number of every other orientation that was

expected to be presented. As the analysis we employed used a regression-based

approach, any carry over of orientation-selective information from the first to the

second grating therefore could not systematically bias the results.

Experimental Design

We used a modified version of a factorial design that has previously been

used to separately examine the effects of repetition suppression and prediction

error (Kaliukhovich & Vogels, 2010; Kovács et al., 2013; Summerfield et al., 2008;

2011; Todorovic et al., 2011; Todorovic & de Lange, 2012). By comparing the two

repeat conditions with the two alternating conditions, we could examine repetition

suppression while controlling for different levels of expectation. Conversely, by

comparing across the expected and unexpected trials, we could examine prediction

error while controlling for repetition suppression.

The relationship between the pairs of orientations for the different

expectation conditions was based on the original study (Summerfield et al., 2008),

and on other studies (Kaliukhovich & Vogels, 2010; Kovács et al., 2013) that

examined the interaction between repetition suppression and expectation. In the

repeating condition, the orientation of the second Gabor is expected to be the same

as the orientation of the first, whereas in the alternating condition the orientation of

the second Gabor is expected to be *different* from that of the first. This relationship

between the expected orientations of the stimuli in the alternating condition is

slightly different to another modification of the paradigm which employed a more

limited range of stimuli (Todorovic et al., 2011; Todorovic & de Lange, 2012).

Specifically, the paradigm introduced by Todorovic and colleagues used two or

three auditory tones of different frequencies. In the alternating condition, the

expectation is that one tone will follow another (i.e. 1000 Hz and then 1032 Hz), then

this is violated when a 1000 Hz tone is repeated. In this paradigm, an exact

frequency is expected in the alternating condition, a design feature that differs from

the paradigm used in the current work where there is no specific expectation of the

orientation of the second Gabor based on the orientation of the first in the

alternating condition. Instead the expectation in the alternating condition is that the

orientation will change, and this can be violated by repeating the orientation. In this

sense, there is no specific expectation about the second orientation in the

alternating condition. Instead, the rule is about alternating or repeating the first

orientation. We did not implement the Todorovic et al. paradigm because the

combinatorial explosion of stimulus conditions needed to measure orientation

selectivity (such that every orientation is predicted by another orientation). Future

work could investigate how this subtle change in paradigm design affects the

encoding of stimulus information.

Forward encoding modelling

We used a forward encoding approach to estimate the amount of orientation-

selective information contained in the EEG data at each time point of the trial. This

approach differs from standard decoding approaches by modelling each presented

orientation as a continuous variable of a set of tuned orientation-selective channels.

The forward-encoding technique has been successfully used to reconstruct colour

(Brouwer & Heeger, 2009), spatial (Sprague & Serences, 2013) and orientation

(Ester, Sutterer, Serences, & Awh, 2016) selectivity in fMRI data. More recently the

same approach has been applied to EEG and MEG data, which have inherently

better temporal resolution than fMRI (Garcia et al., 2013; Kok et al., 2017; Myers et

al., 2015; Wolff, Jochim, Akyürek, & Stokes, 2017b).

We applied forward encoding modelling to determine how repetition

suppression and prediction error affected orientation selectivity. To do this, the

second orientation (Figure 7A) in the Gabor pair in each trial was used to construct a

regression matrix, with separate regressors for the 9 orientations used across the

experiment. This regression matrix was convolved with a set of basis functions (half

cosines raised to the 8^{th} power (Figure 7C), which allowed complete and unbiased

coverage of orientation space) to allow us to pool similar information patterns

across nearby orientations (Brouwer & Heeger, 2009). We used this tuned

regression matrix to estimate time-resolved orientation selectivity contained within

the EEG activity in a 16 ms sliding window, in 4 ms steps (Figure 8B; Myers et al.,

2015). To avoid overfitting, we used a leave-one-out cross-validation procedure

where the regression weights were estimated for a training set and applied to an

independent test set (Figure 8D). All trial types (including target trials) were used in

training and test sets. This was done by solving the linear equation:

$$B_1 = WC_1 \qquad (1)$$

754   Where B_1 (64 sensors x N training trials) is the electrode data for the training set, C_1

755   (9 channels x N training trials) is the tuned channel response across the training

756   trials, and W is the weight matrix for the sensors we want to estimate (64 sensors x

757   9 channels). W can be estimated using least square regression to solve equation (2):

$$W = (C_1 \, C_1^{T})^{-1} \, C_1^{T} \, B_1 \qquad (2)$$

759   The channel response in the test set C_2 (9 channels x N test trials) was estimated

760   using the weights in (2) and applied to activity in B_2 (64 sensors x N test trials).

$$C^2 = (W \, W^{T}) \, W^{T} \, B^2 \qquad (3)$$

We repeated this process by holding one trial out as test, and training on the

remaining trials until all trials had been used in test and training. The procedure was

repeated for each trial within the trial epoch. We then shifted all trials to a common

orientation, meaning that 0° corresponded to the orientation presented on each trial.

The reconstructed channel activations were separated into the four conditions, and

averaged over trials. These responses were then smoothed with a Gaussian kernel

with a 16 ms window, and fitted with a Gaussian function (4) using non-linear least

square regression to quantify the amount of orientation selective activity.

$$G(x) = A\,exp(-\frac{(x-\varphi)^2}{2\sigma^2}) + C \qquad (4)$$

Where *A* is the amplitude representing the amount of orientation selective activity,

is the orientation the function is centred on (in degrees), is the width (degrees) and

C is a constant used to account for non-orientation selective baseline shifts.

Figure 7. A schematic of the forward-encoding approach applied to EEG activity.
(A) Participants viewed individual gratings at fixation, each with a specific
orientation. **(B)** Neural activity evoked by each grating was measured over the entire
scalp. **(C)** Evoked neural responses were convolved with canonical orientation-
selective functions (grey lines in C) to determine coefficients for the different
orientations (coloured dots and lines, which match the colours of the outlined
gratings in A). These coefficients were then used to generate a regression matrix.
(D) General linear modelling was used on a subset of training trials to generate
weights for each channel. These weights were inverted and simultaneously applied
to an independent test set of data to recover orientation selectivity in the EEG
activity. As EEG activity has high temporal resolution, we can apply the procedure
to many epochs following stimulus presentation to determine the temporal
dynamics of orientation processing (see Figure 3).
788

Multivariate pattern analysis

We conducted a multivariate pattern analysis to build upon the initial forward

encoding results which showed that unexpected stimuli elicit greater orientation

selectivity than expected stimuli. This analysis used the same data as the forward

encoding analysis. We used the *classify* function from *Matlab 2017a* with the

'diaglinear' option to implement a Naive Bayes classifier. For each time point, we

used the same cross-validation procedure as the forward encoding modelling with

the same averaging procedure to select train and test sets of data. The classifier

was given the orientations of the training data and predicted the orientation of the

test data. A trial was labelled correct if the presented orientation was produced. To

facilitate comparison of the results with those of (Kok et al., 2012), we found the

peak classification accuracy for each participant in the 600 ms following stimulus

presentation. The same wide time window was used across conditions to

accommodate large inter-individual differences in peak classification without

biasing the results toward one particular condition.

Statistical testing

A non-parametric sign permutation test was used to determine the null

distribution for testing (Wolff, Jochim, Akyürek, & Stokes, 2017b). This method

makes no assumptions about the underlying shape of the null distribution**.** This was

done by randomly flipping the sign of the data for the participants with equal

probability. Fifty thousand (50,000) permutations were used for the time-series data,

whereas only 5000 were used for the temporal generalization plots because of the

significantly greater computational demands.

Cluster-based non-parametric correction (50,000 permutations for timeseries

and 5,000 for temporal generalization) was used to account for multiple

comparisons, and determined whether there were statistical differences between

the contrasting conditions. Paired-samples t-tests were used to follow up the

analysis in Figure 4 within a specified time window, and no correction was applied.

A two-way repeated measures ANOVA (implemented using GraphPad Prism 7.0c,

La Jolla California, USA) was used to analyse the multivariate pattern analysis

results shown in Figure 5.

820   **Acknowledgements**

821   This work was supported by the Australian Research Council (ARC) Centre of

822   Excellence for Integrative Brain Function (ARC Centre Grant CE140100007) to JBM

823   and EA, and by an ARC Discovery Project (DP170100908) to EA. JBM was

824   supported by ARC Australian Laureate Fellowship (FL110100103).

Supplementary Information

Supplementary Figure 1. The effect of a different baseline period (-100 to 0 ms
before onset of the first Gabor) on orientation selectivity for the two main conditions.
Population tuning curves averaged over the significant time period (79 – 150 ms)
shown in Figure 4A. The curves, shown as fitted Gaussians, illustrate how overall
stimulus representations are affected by repetition and expectation. While there was
no difference in orientation tuning for repeated versus alternate stimuli (left panel),
the amplitude of the orientation response increased significantly, and the baseline
decreased, for unexpected relative to expected stimuli (right panel). Error bars
indicate ±1 standard error.
837

838   **Reference**

Adibi, M., McDonald, J. S., Clifford, C. W. G., & Arabzadeh, E. (2013). Adaptation
improves neural coding efficiency despite increasing correlations in variability.
The Journal of Neuroscience, 33(5), 2108–2120.
http://doi.org/10.1523/JNEUROSCI.3449-12.2013
Auksztulewicz, R., & Friston, K. (2016). Repetition suppression and its contextual
determinants in predictive coding. *Cortex, 80*, 125–140.
http://doi.org/10.1016/j.cortex.2015.11.024
Barron, H. C., Garvert, M. M., & Behrens, T. E. J. (2016). Repetition suppression: a
means to index neural representations using BOLD? *Philosophical Transactions*
of the Royal Society of London Series B, Biological Sciences, 371(1705),
20150355. http://doi.org/10.1098/rstb.2015.0355
Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., & Naccache, L.
(2009). Neural signature of the conscious processing of auditory regularities.
Proceedings of the National Academy of Sciences, 106(5), 1672–1677.
http://doi.org/10.1073/pnas.0809667106
Benucci, A., Saleem, A. B., & Carandini, M. (2013). Adaptation maintains population
homeostasis in primary visual cortex, *16*(6), 724–729.
http://doi.org/10.1038/nn.3382
Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K.-M., & Robbins, K. A. (2015). The
PREP pipeline: standardized preprocessing for large-scale EEG analysis.
Frontiers in Neuroinformatics, 9, B153. http://doi.org/10.3389/fninf.2015.00016
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433–436.
http://doi.org/10.1163/156856897X00357
Brouwer, G. J., & Heeger, D. J. (2009). Decoding and reconstructing color from
responses in human visual cortex. *The Journal of Neuroscience, 29*(44), 13992–
14003. http://doi.org/10.1523/JNEUROSCI.3577-09.2009
Buckner, R. L., Goodman, J., Burock, M., Rotte, M., Koutstaal, W., Schacter, D., et
al. (1998). Functional-Anatomic Correlates of Object Priming in Humans
Revealed by Rapid Presentation Event-Related fMRI. *Neuron, 20*(2), 285–296.
http://doi.org/10.1016/S0896-6273(00)80456-0
Caharel, S., d'Arripe, O., Ramon, M., Jacques, C., & Rossion, B. (2009). Early
adaptation to repeated unfamiliar faces across viewpoint changes in the right
hemisphere: Evidence from the N170 ERP component. *Neuropsychologia, 47*(3),
639–643. http://doi.org/10.1016/j.neuropsychologia.2008.11.016
Chaumon, M., Bishop, D. V. M., & Busch, N. A. (2015). A practical guide to the
selection of independent components of the electroencephalogram for artifact
correction. *Journal of Neuroscience Methods, 250*, 47–63.
http://doi.org/10.1016/j.jneumeth.2015.02.025
Cichy, R. M., Pantazis, D., & Oliva, A. (2014). Resolving human object recognition in
space and time. *Nature Publishing Group*. http://doi.org/10.1038/nn.3635
Cichy, R. M., Ramirez, F. M., & Pantazis, D. (2015). Can visual information encoded
in cortical columns be decoded from magnetoencephalography data in humans?
NeuroImage, 121, 193–204. http://doi.org/10.1016/j.neuroimage.2015.07.011

Clifford, C. W. G. (2002). Perceptual adaptation: motion parallels orientation. *Trends*
in Cognitive Sciences, *6*(3), 136–143.
Clifford, C. W. G. (2014). The tilt illusion: phenomenology and functional implications.
Vision Research, *104*, 3–11. http://doi.org/10.1016/j.visres.2014.06.009
Cui, Y., Wang, Y. V., Park, S. J. H., Demb, J. B., & Butts, D. A. (2016). Divisive
suppression explains high-precision firing and contrast adaptation in retinal
ganglion cells. *eLife*, *5*. http://doi.org/10.7554/eLife.19460
Dehaene, S., Naccache, L., Cohen, L., Le Bihan, D., Mangin, J.-F., Poline, J.-B., &
Rivière, D. (2001). Cerebral mechanisms of word masking and unconscious
repetition priming. *Nature Publishing Group*, *4*(7), 752–758.
http://doi.org/10.1038/89551
Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of
single-trial EEG dynamics including independent component analysis. *Journal of*
Neuroscience Methods, *134*(1), 9–21.
http://doi.org/10.1016/j.jneumeth.2003.10.009
Dickinson, J. E., Almeida, R. A., Bell, J., & Badcock, D. R. (2010). Global shape
aftereffects have a local substrate: A tilt aftereffect field. *Journal of Vision*,
10(13), 5–5. http://doi.org/10.1167/10.13.5
Dickinson, J. E., Morgan, S. K., Tang, M. F., & Badcock, D. R. (2017). Separate
banks of information channels encode size and aspect ratio. *Journal of Vision*,
17(3), 27. http://doi.org/10.1167/17.3.27
Diederen, K. M. J., Spencer, T., Vestergaard, M. D., Fletcher, P. C., & Schultz, W.
(2016). Adaptive Prediction Error Coding in the Human Midbrain and Striatum
Facilitates Behavioral Adaptation and Learning Efficiency. *Neuron*, *90*(5), 1127–
1138. http://doi.org/10.1016/j.neuron.2016.04.019
Ester, E. F., Sutterer, D. W., Serences, J. T., & Awh, E. (2016). Feature-Selective
Attentional Modulations in Human Frontoparietal Cortex. *The Journal of*
Neuroscience, *36*(31), 8188–8199. http://doi.org/10.1523/JNEUROSCI.3935-
15.2016
Fairhall, A. L., Lewen, G. D., Bialek, W., & de Ruyter van Steveninck, R. R. (2001).
Efficiency and ambiguity in an adaptive neural code. *Nature*, *412*(6849), 787–
792. http://doi.org/10.1038/35090500
Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the*
Royal Society of London B: Biological Sciences, *360*(1456), 815–836.
http://doi.org/10.1098/rstb.2005.1622
Garcia, J. O., Srinivasan, R., & Serences, J. T. (2013). Near-real-time feature-
selective modulations in human cortex. *Current Biology*, *23*(6), 515–522.
http://doi.org/10.1016/j.cub.2013.02.013
Garrido, M. I., Kilner, J. M., Stephan, K. E., & Friston, K. J. (2009). The mismatch
negativity: A review of underlying mechanisms. *Clinical Neurophysiology*, *120*(3),
453–463. http://doi.org/10.1016/j.clinph.2008.11.029
Garrido, M. I., Sahani, M., & Dolan, R. J. (2013). Outlier Responses Reflect
Sensitivity to Statistical Structure in the Human Brain. *PLOS Comput Biol*, *9*(3),
e1002999. http://doi.org/10.1371/journal.pcbi.1002999

Gotts, S. J., Chow, C. C., & Martin, A. (2012). Repetition priming and repetition
suppression: A case for enhanced efficiency through neural synchronization.
Cognitive Neuroscience. http://doi.org/10.1080/17588928.2012.691277
Grootswagers, T., Wardle, S. G., & Carlson, T. A. (2017). Decoding Dynamic Brain
Patterns from Evoked Responses: A Tutorial on Multivariate Pattern Analysis
Applied to Time Series Neuroimaging Data. *Journal of Cognitive Neuroscience*,
29(4), 677–697. http://doi.org/10.1162/jocn_a_01068
Gross, C. G., Schiller, P. H., Wells, C., & Gerstein, G. L. (1967). Single-unit activity in
temporal association cortex of the monkey. *Journal of Neurophysiology*, *30*(4),
833–43.
Grotheer, M., & Kovács, G. (2014). Repetition probability effects depend on prior
experiences. *The Journal of Neuroscience*, *34*(19), 6640–6646.
http://doi.org/10.1523/JNEUROSCI.5326-13.2014
Henson, R. N. A., & Rugg, M. D. (2003). Neural response suppression,
haemodynamic repetition effects, and behavioural priming. *Neuropsychologia*,
41(3), 263–270. http://doi.org/10.1016/S0028-3932(02)00159-8
Kaliukhovich, D. A., & Vogels, R. (2010). Stimulus Repetition Probability Does Not
Affect Repetition Suppression in Macaque Inferior Temporal Cortex. *Cerebral
Cortex*, *21*(7), 1547–1558. http://doi.org/10.1093/cercor/bhq207
Kamitani, Y., & Tong, F. (2005). Decoding the visual and subjective contents of the
human brain, *8*(5), 679–685. http://doi.org/10.1038/nn1444
Kaposvari, P., Kumar, S., & Vogels, R. (2018). Statistical Learning Signals in
Macaque Inferior Temporal Cortex. *Cerebral Cortex*, *28*(1), 250–266.
http://doi.org/10.1093/cercor/bhw374
Kay, K. N., Naselaris, T., Prenger, R. J., & Gallant, J. L. (2008). Identifying natural
images from human brain activity. *Nature*, *452*(7185), 352–355.
http://doi.org/10.1038/nature06713
Keil, A., Debener, S., Gratton, G., Junghöfer, M., Kappenman, E. S., Luck, S. J., et
al. (2014). Committee report: Publication guidelines and recommendations for
studies using electroencephalography and magnetoencephalography.
Psychophysiology, *51*(1), 1–21. http://doi.org/10.1111/psyp.12147
Keller, A. J., Houlton, R., Kampa, B. M., Lesica, N. A., Mrsic-Flogel, T. D., Keller, G.
B., & Helmchen, F. (2017). Stimulus relevance modulates contrast adaptation in
visual cortex. *eLife*, *6*, e21589. http://doi.org/10.7554/eLife.21589
King, J. R., & Dehaene, S. (2014). Characterizing the dynamics of mental
representations: the temporal generalization method. *Trends in Cognitive
Sciences*, *18*(4), 203–210. http://doi.org/10.1016/j.tics.2014.01.002
King, J.-R., Gramfort, A., Schurger, A., Naccache, L., & Dehaene, S. (2014). Two
Distinct Dynamic Modes Subtend the Detection of Unexpected Sounds. *PloS
One*, *9*(1), e85791. http://doi.org/10.1371/journal.pone.0085791
King, J.-R., Pescetelli, N., & Dehaene, S. (2016). Brain Mechanisms Underlying the
Brief Maintenance of Seen and Unseen Sensory Information. *Neuron*, *92*(5),
1122–1134. http://doi.org/10.1016/j.neuron.2016.10.051
Kohn, A., & Movshon, J. A. (2004). Adaptation changes the direction tuning of
macaque MT neurons., *7*(7), 764–772. http://doi.org/10.1038/nn1267

Kok, P., Jehee, J. F. M., & de Lange, F. P. (2012). Less Is More: Expectation
Sharpens Representations in the Primary Visual Cortex. *Neuron*, *75*(2), 265–
270. http://doi.org/10.1016/j.neuron.2012.04.034
Kok, P., Mostert, P., & de Lange, F. P. (2017). Prior expectations induce prestimulus
sensory templates. *Proceedings of the National Academy of Sciences of the*
United States of America, *26*, 201705652.
http://doi.org/10.1073/pnas.1705652114
Kourtzi, Z., & Kanwisher, N. (2001). Representation of Perceived Object Shape by
the Human Lateral Occipital Complex. *Science*, *293*(5534), 1506–1509.
http://doi.org/10.1126/science.1061133
Kovács, G., Kaiser, D., Kaliukhovich, D. A., Vidnyánszky, Z., & Vogels, R. (2013).
Repetition probability does not affect fMRI repetition suppression for objects. *The*
Journal of Neuroscience, *33*(23), 9805–9812.
http://doi.org/10.1523/JNEUROSCI.3423-12.2013
Kumar, S., Kaposvari, P., & Vogels, R. (2017). Encoding of Predictable and
Unpredictable Stimuli by Inferior Temporal Cortical Neurons. *Journal of Cognitive*
Neuroscience, *29*(8), 1445–1454. http://doi.org/10.1162/jocn_a_01135
Kvale, M. N., & Schreiner, C. E. (2004). Short-Term Adaptation of Auditory
Receptive Fields to Dynamic Stimuli. *Journal of Neurophysiology*, *91*(2), 604–
612. http://doi.org/10.1152/jn.00484.2003
Larsson, J., & Smith, A. T. (2012). fMRI repetition suppression: neuronal adaptation
or stimulus expectation? *Cerebral Cortex*, *22*(3), 567–576.
http://doi.org/10.1093/cercor/bhr119
Liu, T., Larsson, J., & Carrasco, M. (2007). Feature-Based Attention Modulates
Orientation-Selective Responses in Human Visual Cortex. *Neuron*, *55*(2), 313–
323. http://doi.org/10.1016/j.neuron.2007.06.030
Maravall, M., Petersen, R. S., Fairhall, A. L., Arabzadeh, E., & Diamond, M. E.
(2007). Shifts in Coding Properties and Maintenance of Information Transmission
during Adaptation in Barrel Cortex. *PLoS Biology*, *5*(2), e19.
http://doi.org/10.1371/journal.pbio.0050019
Marti, S., King, J. R., & Dehaene, S. (2015). Time-Resolved Decoding of Two
Processing Chains during Dual-Task Interference. *Neuron*, *88*(6), 1297–1307.
http://doi.org/10.1016/j.neuron.2015.10.040
Maunsell, J. (2015). Neuronal Mechanisms of Visual Attention. *Annual Review of*
Vision Science, *1*(1), 373–391. http://doi.org/10.1146/annurev-vision-082114-
035431
Maunsell, J. H. R., & Treue, S. (2006). Feature-based attention in visual cortex.
Trends in Neurosciences, *29*(6), 317–322.
http://doi.org/10.1016/j.tins.2006.04.001
Movshon, J. A., & Lennie, P. (1979). Pattern-selective adaptation in visual cortical
neurones. *Nature*, *278*(5707), 850–852.
Myers, N. E., Rohenkohl, G., Wyart, V., Woolrich, M. W., Nobre, A. C., Stokes, M.
G., & Frank, M. J. (2015). Testing sensory evidence against mnemonic
templates. *eLife*, *4*, e09000. http://doi.org/10.7554/eLife.09000
Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity
(MMN) in basic research of central auditory processing: A review. *Clinical*

Neurophysiology, *118*(12), 2544–2590.
http://doi.org/10.1016/j.clinph.2007.04.026
Oostenveld, R., & Praamstra, P. (2001). The five percent electrode system for high-
resolution EEG and ERP measurements. *Clinical Neurophysiology*, *112*(4), 713–
719.
Paavilainen, P. (2013). The mismatch-negativity (MMN) component of the auditory
event-related potential to violations of abstract regularities: A review.
International Journal of Psychophysiology, *88*(2), 109–123.
http://doi.org/10.1016/j.ijpsycho.2013.03.015
Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics:
Transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442.
Rao, R. P. N., & Ballard, D. H. (1999). Predictive coding in the visual cortex: a
functional interpretation of some extra-classical receptive-field effects, *2*(1), 79–
87. http://doi.org/10.1038/4580
Rasmussen, R. G., Schwartz, A., & Chase, S. M. (2017). Dynamic range adaptation
in primary motor cortical populations. *eLife*, *6*, e21409.
http://doi.org/10.7554/eLife.21409
Rentzeperis, I., Nikolaev, A. R., Kiper, D. C., & Van Leeuwen, C. (2012).
Relationship between neural response and adaptation selectivity to form and
color: an ERP study. *Frontiers in Human Neuroscience*, *6*.
http://doi.org/10.3389/fnhum.2012.00089
Saarinen, J., Paavilainen, P., Schöger, E., Tervaniemi, M., & Näätänen, R. (1992).
Representation of abstract attributes of auditory stimuli in the human brain.
NeuroReport, *3*(12), 1149–1151.
Salti, M., Monto, S., Charles, L., King, J.-R., Parkkonen, L., Dehaene, S., &
Johansen-Berg, H. (2015). Distinct cortical codes and temporal dynamics for
conscious and unconscious percepts. *eLife*, *4*, e05652.
http://doi.org/10.7554/eLife.05652
Schacter, D. L., & Buckner, R. L. (1998). Priming and the Brain. *Neuron*, *20*(2), 185–
195.
Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-Positive Psychology:
Undisclosed Flexibility in Data Collection and Analysis Allows Presenting
Anything as Significant. *Psychological Science*, *22*(11), 1359–1366.
http://doi.org/10.1177/0956797611417632
Smirnakis, S. M., Berry, M. J., Warland, D. K., Bialek, W., & Meister, M. (1997).
Adaptation of retinal processing to image contrast and spatial scale. *Nature*,
386(6620), 69–73. http://doi.org/10.1038/386069a0
Spaak, E., Watanabe, K., Funahashi, S., & Stokes, M. G. (2017). Stable and
Dynamic Coding for Working Memory in Primate Prefrontal Cortex. *The Journal*
of Neuroscience, *37*(27), 6503–6516. http://doi.org/10.1523/JNEUROSCI.3364-
16.2017
Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in
the human occipital, parietal and frontal cortices, *16*(12), 1879–1887.
http://doi.org/10.1038/nn.3574

Repetition suppression and prediction error

Stokes, M. G., Kusunoki, M., Sigala, N., Nili, H., Gaffan, D., & Duncan, J. (2013).
Dynamic Coding for Cognitive Control in Prefrontal Cortex. *Neuron*, *78*(2), 364–
375. http://doi.org/10.1016/j.neuron.2013.01.039
Stokes, M. G., Wolff, M. J., & Spaak, E. (2015). Decoding Rich Spatial Information
with High Temporal Resolution. *Trends in Cognitive Sciences*, *19*(11), 636–638.
http://doi.org/10.1016/j.tics.2015.08.016
Summerfield, C., Trittschuh, E. H., Monti, J. M., Mesulam, M. M., & Egner, T. (2008).
Neural repetition suppression reflects fulfilled perceptual expectations, *11*(9),
1004–1006. http://doi.org/10.1038/nn.2163
Summerfield, C., Wyart, V., Mareike Johnen, V., & de Gardelle, V. (2011). Human
Scalp Electroencephalography Reveals that Repetition Suppression Varies with
Expectation. *Frontiers in Human Neuroscience*, *5*.
http://doi.org/10.3389/fnhum.2011.00067
Tang, M. F., Dickinson, J. E., Visser, T. A. W., & Badcock, D. R. (2015). The broad
orientation dependence of the motion streak aftereffect reveals interactions
between form and motion neurons. *Journal of Vision*, *15*(13), 4.
http://doi.org/10.1167/15.13.4
Todorovic, A., & de Lange, F. P. (2012). Repetition suppression and expectation
suppression are dissociable in time in early auditory evoked fields. *The Journal*
of Neuroscience, *32*(39), 13389–13395.
http://doi.org/10.1523/JNEUROSCI.2227-12.2012
Todorovic, A., van Ede, F., Maris, E., & de Lange, F. P. (2011). Prior Expectation
Mediates Neural Adaptation to Repeated Sounds in the Auditory Cortex: An
MEG Study. *The Journal of Neuroscience*, *31*(25), 9118–9123.
http://doi.org/10.1523/JNEUROSCI.1425-11.2011
Tootell, R. B., Hadjikhani, N. K., Vanduffel, W., Liu, A. K., Mendola, J. D., Sereno, M.
I., & Dale, A. M. (1998). Functional analysis of primary visual cortex (V1) in
humans. *Proceedings of the National Academy of Sciences of the United States*
of America, *95*(3), 811–817. http://doi.org/10.1038/340386a0
Tootell, R., Reppas, J. B., Dale, A. M., & Look, R. B. (1995). Visual motion aftereffect
in human cortical area MT revealed by functional magnetic resonance imaging.
Nature.
Treue, S., & Trujillo, J. (1999). Feature-based attention influences motion processing
gain in macaque visual cortex. *Nature*. http://doi.org/10.1038/21176
Ulanovsky, N., Las, L., & Nelken, I. (2003). Processing of low-probability sounds by
cortical neurons, *6*(4), 391–398. http://doi.org/10.1038/nn1032
Vinken, K., Vogels, R., & Op de Beeck, H. (2017). Recent Visual Experience Shapes
Visual Processing in Rats through Stimulus-Specific Adaptation and Response
Enhancement. *Current Biology*, *27*(6), 914–919.
http://doi.org/10.1016/j.cub.2017.02.024
Wacongne, C., Labyt, E., van Wassenhove, V., Bekinschtein, T., Naccache, L., &
Dehaene, S. (2011). Evidence for a hierarchy of predictions and prediction errors
in human cortex. *Proceedings of the National Academy of Sciences of the United*
States of America, *108*(51), 20754–20759.
http://doi.org/10.1073/pnas.1117807108

Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017a). Dynamic hidden
states underlying working-memory-guided behavior, *20*(6), 864–871.
http://doi.org/10.1038/nn.4546
Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017b). Dynamic hidden
states underlying working-memory-guided behavior, *20*(6), 864–871.
http://doi.org/10.1038/nn.4546
1112