Running head: Repetition suppression and prediction error

Prediction Error and Repetition Suppression Have Distinct Effects

on Neural Representations of Visual Information

Matthew F. Tang[1,2], Cooper A. Smout[1,2], Ehsan Arabzadeh[2,3] & Jason B.

Mattingley[1,2,4]

1. Queensland Brain Institute, The University of Queensland, St Lucia, QLD,

Australia

2. Australian Research Council Centre of Excellence for Integrative Brain

Function, Victoria, Australia

3. Eccles Institute of Neuroscience, John Curtin School of Medical Research,

The Australian National University, Canberra, ACT, Australia

4. School of Psychology, The University of Queensland, St Lucia, QLD,

Australia

Corresponding author

Matthew F. Tang

Queensland Brain Institute

The University of Queensland

St Lucia, QLD, Australia

Email: m.tang1@uq.edu.au

Abstract

Predictive coding theories argue recent experience establishes expectations in the

brain that generate *prediction errors* when violated. Prediction errors provide a

possible explanation for *repetition suppression*, in which evoked neural activity is

attenuated across repeated presentations of the same stimulus. According to the

predictive coding account, repetition suppression arises because repeated stimuli

are expected, non-repeated stimuli are unexpected and thus elicit larger neural

responses. Here we employed electroencephalography in humans to test the

predictive coding account of repetition suppression by presenting sequences of

gratings with orientations that were expected to repeat or change in separate

blocks. We applied multivariate forward modelling to determine how orientation

selectivity was affected by repetition and prediction. Unexpected stimuli were

associated with significantly enhanced orientation selectivity, whereas there was no

such influence on selectivity for repeated stimuli. Our results suggest that repetition

suppression and expectation have separable effects on neural representations of

visual feature information.

41   **Introduction**

At any moment in time, the brain receives more sensory information than can

be responded to, creating the need for selection and efficient processing of

incoming signals. One mechanism by which the brain reduces its information

processing load is to encode successive presentations of the same stimulus in a

more efficient form, a process known as neural adaptation (Fairhall, Lewen, Bialek,

& de Ruyter van Steveninck, 2001; Kvale & Schreiner, 2004; Smirnakis, Berry,

Warland, Bialek, & Meister, 1997). Such adaptation has been observed across

different sensory modalities and species, and has been suggested as a potential

mechanism for enhancing the coding efficiency of individual neurons and neuronal

populations (Adibi, McDonald, Clifford, & Arabzadeh, 2013; Benucci, Saleem, &

Carandini, 2013; Maravall, Petersen, Fairhall, Arabzadeh, & Diamond, 2007). A

particular form of neuronal adaptation, known as *repetition suppression*, is

characterised by attenuated neural responses to repeated presentations of the

same stimulus (Diederen, Spencer, Vestergaard, Fletcher, & Schultz, 2016; Gross,

Schiller, Wells, & Gerstein, 1967; Keller et al., 2017; Movshon & Lennie, 1979;

Rasmussen, Schwartz, & Chase, 2017). Here we asked whether predictive coding

theory, which assumes that sensory processing is influenced by prior exposure, can

account for changes in neural representations observed with repetition suppression.

The phenomenon of repetition suppression has been widely exploited to

investigate neural representations of sensory information. Repeated exposures allow

for more efficient representation of subsequent stimuli, as manifested in improved

behavioural performance despite a significant reduction in neural activity (Henson &

Rugg, 2003; Schacter & Buckner, 1998). Repetition suppression paradigms have

been used extensively in human neuroimaging because they are commonly

considered to be analogous to the single-cell adaptation effects observed in animal

models (see Barron, Garvert, & Behrens, 2016 for review). The exact relationship

between the effects seen in human neuroimaging studies and animal

neurophysiology has, however, yet to be fully established.

The view that repetition suppression observed in human neuroimaging

studies reflects neuronal adaptation has recently been challenged by hierarchical

predictive coding theories (Auksztulewicz & Friston, 2016; Summerfield, Trittschuh,

Monti, Mesulam, & Egner, 2008). These theories argue that the brain interprets

incoming sensory events based on what would be expected from the recent history

of exposure to such stimuli (Friston, 2005; Rao & Ballard, 1999). Predictions are

generated within each cortical area, and are bi-directionally propagated from higher

to lower areas, including to primary sensory regions, allowing for more efficient

representation of expected stimuli. When there is a precise expectation, incoming

information can be efficiently represented by recruiting a small pool of relevant

neurons (Friston, 2005). When there is a mismatch between an expectation and the

stimulus presented, i.e., when there is a *prediction error*, the stimulus is less

efficiently represented and thus elicits a larger neural response.

Some of the most compelling evidence for predictive coding comes from

human neuroimaging experiments in which the presentation of an unexpected

stimulus generates a larger response than the presentation of an expected stimulus.

In studies employing electroencephalography (EEG) and magnetoencephalography

(MEG), this effect is known as the *mismatch negativity* (Garrido, Kilner, Stephan, &

Friston, 2009; Näätänen, Paavilainen, Rinne, & Alho, 2007; Wacongne et al., 2011)

To test the hypothesis that prediction error can account for repetition

suppression effects, Summerfield and colleagues (2008) introduced an experimental

paradigm in which the identity of a face stimulus was either repeated in 80% of

trials (making the repetition *expected*) or was changed in 80% of trials (making the

repetition *unexpected*). There was a greater attenuation of the BOLD response in

the fusiform face area when a face repetition was expected, relative to when it was

unexpected, suggesting that repetition suppression is reduced by unexpected

stimuli. This attenuation of repetition suppression by failures of expectation has also

been replicated using fMRI (Larsson & Smith, 2012) and M/EEG, using high-level

stimuli such as faces (Summerfield, Wyart, Mareike Johnen, & de Gardelle, 2011),

and simple stimuli such as tones (Todorovic & de Lange, 2012; Todorovic, van Ede,

Maris, & de Lange, 2011)

A potential reconciliation of the relationship between expectation and

repetition suppression comes from work showing that while expectations decrease

the overall amount of neural activity, they can also yield sharper representations of

sensory stimuli (Kok, Jehee, & de Lange, 2012). This work goes beyond

conventional neuroimaging approaches, which typically only measure overall levels

of neural activity (Buckner et al., 1998; Kourtzi & Kanwisher, 2001; Tootell, Reppas,

Dale, & Look, 1995). Such amplitude changes could in principle be produced by one

or more different types of change in the underlying neural representations. For

instance, both sharpening, where the response to only unpredicted features is

suppressed, and gain reduction, where a multiplicative suppression occurs for all

features, could be associated with decreased population activity, even though the

amount of information carried by the representations will be markedly different.

Recently introduced multivariate pattern analytic approaches to human neuroimaging

– specifically forward encoding modelling – allow for the quantification of stimulus-

selective information contained within patterns of neural activity in human observers

(Brouwer & Heeger, 2009; Garcia, Srinivasan, & Serences, 2013; King, Pescetelli, &

Dehaene, 2016; Myers et al., 2015; Salti et al., 2015; Wolff, Jochim, Akyürek, &

Stokes, 2017). This goes beyond typical multivariate pattern analyses, which

normally produce only accuracy scores, by quantifying neural representations

evoked by sensory stimuli to reveal both the accuracy and *tuning fidelity* for the

specific feature dimension of interest.

Here we used these multivariate forward encoding methods to determine

how repetition suppression and expectation affect the way the brain represents

visual orientation information. To anticipate the results, we found that repetition

suppression had no effect on visual orientation selectivity, but that violated

expectations were associated with a significantly increased orientation-selective

response through gain modulation soon after stimulus onset. This representation

appeared to be transiently re-activated at around 200 ms, suggesting that feedback

influences the initial sensory representation of the unexpected stimulus which

allows for the updating of the sensory prior.

131   **Results**

We used a modified version of the paradigm introduced by Summerfield and

colleagues (2008), replacing the face stimuli used in that study with oriented

Gabors. These low-level stimuli allowed us to quantify the degree of orientation

selectivity in EEG activity to determine how the representations of orientation is

affected by prediction error and repetition suppression. Each of fifteen observers

participated in two EEG sessions. On each trial, two Gabors were presented

sequentially (100 ms presentation, 600 ms stimulus onset asynchrony), and these

stimulus pairs either repeated or alternated in their orientation (Figure 1A, Movie 1).

The predictability of the repeated and alternating pairs was varied in a block-wise

manner to manipulate expectation. In a *repeating* block, the orientations of the two

Gabors in a pair repeated in 80% of trials, and alternated for the remaining 20%.

These contingencies were reversed in the *alternating* block (Figure 1B). The

orientations of successive stimuli across a block were randomized to limit any

accumulated effects of adaptation and prediction. As repetition suppression and

expectation form orthogonal dimensions of the task, the design allowed us to

isolate their respective contributions to neural responses. Participants completed an

unrelated task of discriminating (red vs blue) rare coloured Gabors (which occurred

on 10% of trials).

Repetition suppression and prediction error

A

100 ms 500 ms 100 ms

Repetition

Alternation

Target

B

	Repeating block	Alternating block
Expected	Repeat (80%)	Alternating (80%)
Unexpected	Alternate (20%)	Repeat (20%)

Figure 1. Example stimulus displays and task design. **(A)** Schematic of the stimuli and timing used in the experiment. Participants viewed a rapid stream of pairs of Gabors and monitored for an infrequent coloured target (10% of trials). The stimulus orientations varied pseudorandomly across trials between 0° and 160° (in 20° steps), allowing estimation of orientation-selective information contained within patterns of EEG activity. **(B)** The orientation of the pairs of Gabors could either repeat or alternate. In one type of block, 80% of trials were orientation repeats and the remaining 20% alternated (Repeating blocks); in the other type of block these contingencies were reversed (Alternating blocks).

Movie 1. Example of a stimulus sequence of Gabors in a typical repetition block.

Repetition suppression and prediction error affect the overall level of neural

activity

The Gabors elicited a large response over occipital-parietal areas (Figure 2A).

Consistent with previous work (Cui, Wang, Park, Demb, & Butts, 2016; Keller et al.,

2017; Rentzeperis, Nikolaev, Kiper, & Van Leeuwen, 2012; Summerfield et al., 2011;

Todorovic et al., 2011; Todorovic & de Lange, 2012; Tootell et al., 1998), there was

a significant repetition suppression effect (Repeat < Alternating), such that the

response to repeated stimuli was significantly reduced compared with the response

to alternating stimuli (Figure 2A). The repetition suppression effect was evident over

a large cluster of occipital-parietal electrodes at two time intervals: an early effect

from 79 to 230 ms, and a later effect at 250 to 540 ms after the onset of the second

stimulus (cluster $p < .025$; Figure 2B and caption). A large cluster of frontal

electrodes mirrored the repetition suppression effect with a similar time course: the

ERP over these frontal sites had the same pattern, but was reversed in sign,

suggesting it originated from the same dipole as the occipital response.

Repetition suppression and prediction error

Figure 2. Univariate EEG results for the effect of repetition suppression and
prediction error on the second stimulus in a pair (T2). Grand average ERP time
courses for repetition suppression **(A)** and expectation **(C)** averaged over occipital-
parietal electrodes (O1, O2, Oz, POz, PO7, PO3, PO8, PO4). The shaded region
indicates significant differences between the conditions (two-tailed cluster-
permutation, alpha $p < .05$, cluster alpha $p < .05$, N permutations = 20,000). The
main effects of repetition suppression **(B)** and expectation **(D)** over three time
periods and across all electrodes. The main effect of repetition suppression is
displayed as Repeating minus Alternating trials. The main effect of expectation is
displayed as Expected minus Unexpected trials. Circles indicate clusters of
electrodes with significantly reduced activity, and crosses indicate clusters of
electrodes with significantly increased activity (alpha $p < .05$, cluster $p < .025$, N
permutations = 1500).
194

Also consistent with previous results (Garrido et al., 2009; Summerfield et al.,

2011; Todorovic et al., 2011; Todorovic & de Lange, 2012), there was a significant

prediction error effect (Expected < Unexpected), with a larger neural response over

a cluster of frontal electrodes 75-150 ms after stimulus presentation (Figure 2C). As

with the repetition suppression result described above, there was a prediction error

effect of opposite polarity over occipital-parietal electrodes. This effect was

significant at an early time point post-stimulus (79-130 ms), but not at later time

points (320-390 ms; Figure 2D). Finally, there was no interaction between repetition

suppression and prediction error (i.e., no significant positive or negative clusters, all

p > .05). Taken together, these results reveal both repetition suppression and

prediction error effects in the neural data, which were indexed separately as shown

in Figure 2.

Prediction errors increase the amount of orientation-selective information

contained within patterns of EEG activity

We next examined the key question of whether repetition suppression and

expectation differentially affect the neural representation of orientation information.

To do this, we used a forward encoding approach to reconstruct orientation-

selective information contained within the multivariate pattern of EEG activity

distributed across the scalp (see Methods for details). Briefly, this technique

transforms sensor-level responses into tuned 'feature' channels (Brouwer & Heeger,

2009; Garcia et al., 2013; Kay, Naselaris, Prenger, & Gallant, 2008; Myers et al.,

2015), in this case, orientation-selective features. For each trial, the presented

orientation was convolved with a canonical, orientation-selective tuning function

and regressed against the pattern of EEG activity across all sensors at each time

point. This created a spatial filter of the multivariate EEG activity that differentiated

orientations (Figure 3D). These weights were then inverted to reconstruct the model,

and multiplied against an independent set of test trials to produce responses in the

modelled orientation channels. These sets of responses were then used to evaluate

the degree of orientation selectivity in those trials. The procedure was repeated for

all time points in the trial, and a cross-validated approach was used until all trials

had been used for both training and testing.

As shown in Figure 3, the forward encoding revealed a strong, orientation-

Repetition suppression and prediction error

227   selective response derived from the multivariate pattern of EEG activity. This

228   orientation-tuned response was evident from ~50 to 470 ms after stimulus onset,

229   and peaked between ~120-250 ms (Figure 3C). Examination of the regression

230   weights revealed that this response was largely driven by activity centred over

231   occipital-parietal areas (Figure 3D).

Figure 3. Results of the forward encoding modelling for orientation-selectivity. **(A)** Time-resolved orientation tuning curve across all participants and conditions in response to the second Gabor in the pair. The forward encoding approach resulted in a tuning curve for each of the nine presented orientations. These tuning curves were then centred at each presented orientation (here labelled as 0°) to combine across all orientations. The orientation-selective response is contained within the overall pattern of EEG; activity begins soon after stimulus onset and peaks at around 250 ms before declining. **(B)** Population tuning curve of the stimulus reconstruction across participants, averaged between 50-100 ms and 150-250 ms after stimulus presentation. Each line is a fitted Gaussian response with a variable offset used to quantify orientation selectivity. Error bars indicate ±1 standard error of mean across participants. **(C)** Amplitude of the channel response over time, averaged across all conditions (black line). The thick black line indicates significant encoding of orientation based on a cluster-permutation test across participants (cluster $p < .05$, N permutations = 20,000). Encoding accuracy was reliable from 52 to 470 ms post-stimulus onset. The error shading (in grey) indicates bootstrapped 95% confidence intervals of the mean. **(D)** Topographic plots of the weights (averaged across the 9 orientation channels across all participants) derived from forward encoding at the corresponding time points shown in panel B. (a.u. = arbitrary units).

Repetition suppression and prediction error

To examine our central question of whether repetition suppression and

prediction error have differential effects on neural representations of orientation, we

split and averaged the results of the forward encoding by trial type, and fitted these

with Gaussians (see Methods) to quantify orientation selectivity (Figure 4).

Repetition suppression did not affect the amount of orientation selectivity contained

within the EEG data, with similar selectivity for repeated and alternating trials. This

was the case even though the repeated trials had a markedly smaller EEG response

over occipital and parietal electrodes, where the forward encoding model was

maximally sensitive (Figure 2A). This result is consistent with the 'efficient

representation' hypothesis of repetition suppression, which argues that the overall

neural response is smaller with repetition suppression due to more efficient coding

of stimulus information (Gotts, Chow, & Martin, 2012).

Figure 4. The effect of repetition suppression and expectation on orientation selectivity measured using forward encoding modelling. **(A).** Amount of orientation-selective information (given by the amplitude of the fitted gaussian) from the EEG signal in response to the second Gabor in a pair, shown separately for repetition suppression (upper panel) and prediction error (lower panel). The thick black line indicates significant differences between the conditions (two-tailed cluster-permutation, alpha $p < .05$, cluster alpha $p < .05$, N permutations = 20,000). Error shading indicates bootstrapped 95% confidence intervals of the mean. **(B)** The population tuning curve averaged over the significant time period (79 – 185 ms) to show how the overall stimulus representation is affected with a fitted Gaussian (line). For prediction error, the amplitude increased and the baseline decreased. There was no effect for the repetition suppression condition. Error bars indicate ±1 standard error.

Examining the effect of expectation revealed a markedly different pattern of

results. At 79 - 185 ms after the onset of the second stimulus in the pair,

orientation-selectivity increased significantly ($p < .0001$) when the stimulus was

unexpected relative to when it was expected, i.e., the prediction error was

associated with an increased representation of the stimulus at the earliest stages of

the brain's response to that stimulus. Moreover, the prediction error signal

contained enhanced information about the specific features of the stimulus that

violated the expectation, in this case the orientation of the second grating. We

conducted the same statistical tests on the three other parameters defining the

Gaussian function (namely, the width, centre orientation and baseline) to determine

how repetition suppression and expectation might affect other properties of the

neural representation. There was no reliable influence of repetition suppression on

any of these Gaussian parameters (all p > .32). For prediction error, there was a

significant decrease in baseline activity over the same time period as observed for

the increase in amplitude (79-185 ms, p = .001), but there were no significant effects

for the other parameters (all ps > .30).

We followed up this initial analysis to ensure we did not miss any small

effects of repetition suppression or expectation on all aspects of stimulus

representation. We increased the signal-to-noise by averaging the stimulus

reconstruction over this early time period (79-185 ms after stimulus presentation),

and fitted Gaussians to each participant's data individually (Figure 4B). This again

showed that the amplitude of the response was significantly ($t(14)$ = 3.34, p = .005)

higher for unexpected (M = 0.67, SE = 0.06) than for expected (M = 0.41, SE = 0.03)

stimuli. By contrast, the width of the representations was similar for unexpected (M

= 29.62°, SE = 4.72°) and expected (M = 26.72°, SE = 2.74°) stimuli, ($t(14)$ = 0.78,

p = .45). There was also a small, but non-significant ($t(14)$ = 1.94, p = .06) trend for a

smaller baseline response (i.e., non-orientation tuned activity) in the unexpected

(M = -0.01, SE = 0.07) than in the expected (M = 0.13, SE = 0.02) condition. For

comparison, we also averaged the same time period for the repetition suppression

conditions, and found similar curves for the repeated and alternating trials (*all ps >*

.18). This analysis confirms the previous result, which employed more conservative

nonparametric cluster-based testing.

We used a number of approaches to determine whether repetition

suppression and expectation interacted to affect orientation selectivity. First, we

took the difference scores between the combination of factors (e.g., expected

repetition minus unexpected repetition, and expected alternation minus unexpected

alternation) and compared these using the same cluster-based permutation testing

outlined above. This analysis revealed no significant interactions between the

factors for any parameter (all *ps* > .10). Second, we found the largest orientation-

selectivity, defined by the amplitude of the fitted Gaussian, across the 600 ms

following stimulus presentation. For each participant, this resulted in a single value

for the four conditions. Each of these values was subjected to a two-way repeated-

measures ANOVA, which again revealed no significant interaction between the

factors (all *ps* > .30)

To further examine whether orientation-selectivity contained within the overall

pattern of EEG activity differed for unexpected and expected stimuli, we used

multivariate discriminant analysis to determine whether more traditional backward

decoding produces the same pattern of results as that yielded by the forward

encoding approach described above. The same cross-validation procedure was

used as in the forward encoding approach, but accuracy was now defined as the

proportion of trials labelled with the correct orientation. To facilitate comparison

with the results with Kok et al., (2013), we took the peak classification accuracy

within a 600 ms window after presentation of the second grating of the pair. This

analysis confirmed the results of the forward encoding: orientations shown in

unexpected trials were classified better than orientations shown in expected trials

($F(1,14)$ 76.42, $p <. 00001$). Again, there was no effect of repetition on classification

accuracy ($F(1,14) = 0.027$, p = .87) nor a significant interaction ($F(1,14) = 2.52$, $p =$

.13). This suggests the finding is not specific to the analysis method but rather

reflects how expectation affects the representation of sensory information in

general.

Figure 5. Peak (naive Bayes) classification accuracy of the presented grating
orientation for expected and unexpected conditions. The dotted line indicates
chance performance (1/9 orientations). The error bars indicate ±1 standard error of
the mean.
347
Prediction error affects the temporal stability of stimulus representations

Next, we examined whether repetition suppression and prediction error

affected dynamic, ongoing stimulus representations by using cross-temporal

generalization (King & Dehaene, 2014; King, Gramfort, Schurger, Naccache, &

Dehaene, 2014; Myers et al., 2015; Spaak, Watanabe, Funahashi, & Stokes, 2017;

Stokes et al., 2013). To do this, we used the same forward encoding approach as in

the previous analysis, but now the weights were derived from one time-point on one

set of trials, and then applied at every time point in the test trials. Again, a cross-

validation approach was used, with all trials serving as both training and test. This

analysis examined whether the same spatial pattern of EEG activity that allows for

orientation selectivity generalizes to other time points, thus revealing whether the

pattern of orientation-selective activity is stable or dynamic over time.

As shown in Figure 6, the best orientation selectivity was on-axis (training

time equals test time) between 100 ms and 300 ms after stimulus presentation,

suggesting that the stimulus representation changed dynamically over time (King &

Dehaene, 2014). There was also significant off-axis orientation-selectivity from 100-

500 ms after stimulus presentation, suggesting that some aspects of the neural

representation of orientation were stable over time.

Repetition suppression and prediction error

367
Figure 6. Cross-temporal generalization of the forward encoding model based on
grating orientations for the main effects of repetition suppression (upper panels) and
prediction error (lower panels). The maps have been thresholded (indicated by
opacity) to show clusters (black outlines) of significant orientation selectivity
(permutation testing, cluster threshold p < .05, corrected cluster statistic *p* < .05,
5,000 permutations). The difference between the conditions is shown in the right-
hand column (permutation testing, cluster threshold *p* < .05, corrected cluster
statistic *p* < .05). Opacity and outlines indicate significant differences.
376
There was no effect of repetition suppression on temporal generalization of

orientation information (upper panels of Figure 6), suggesting that repetition

suppression did not affect the temporal stability of neural representations of

orientation. Examining the effect of expectation on cross-temporal generalization

confirmed that there was significantly more on-axis orientation selectivity when the

stimulus was unexpected than when it was expected (cluster *p* = .02). This

increased on-axis orientation selectivity generalized off-axis at around 300-400 ms

after stimulus onset (cluster *p* = .01). This finding suggests that the same

representation that is activated to process the prediction error is reactivated later as

the stimulus continues to be processed. Such a signal could constitute the prior of

the prediction, as this should be updated on the basis of incoming sensory

evidence, which in turn would likely require reactivation of the unexpected stimulus.

Representation of the expected feature is activated even when it is not

presented

In a final set of analyses, we investigated how expectation affects the

interpretation of incoming sensory information, one of the key questions of

predictive coding theory (Friston, 2005; Rao & Ballard, 1999). To do this, we

focused on trials in which the orientation of the first Gabor within a pair did not

match the orientation of the second Gabor (i.e., the alternating trials), and examined

EEG responses to the second stimulus (see Figure 7). Here, we trained the forward

encoding model on the orientation of the first Gabor in the pair to determine how

the interaction between the expectation and the incoming sensory information was

compared. We anticipated a lower expectation that the orientation would repeat for

an expected alternating trial, relative to trials in the unexpected alternation

condition, where the repeat should have been expected. We did not combine data

across the repeat and alternating conditions because the orientations were now

different upon presentation of the second Gabor in the pair.

Figure 7. Cross-temporal generalization of the forward encoding model for the

orientations presented in first grating relative to the onset of second grating (first

grating was presented at -600 ms). The orientation in the second grating period is

randomized but there are still periods for the unexpected alternation in which there

is significant orientation selectivity. These maps have been thresholded (indicated

by opacity) to show clusters (black outlines) of significant orientation selectivity

(permutation testing, cluster threshold $p < .05$, corrected cluster statistic $p < .05$,

5,000 permutations). The opacity in the right-hand column indicates areas of

statistically significant difference between the two conditions (permutation testing (n

= 5,000), cluster threshold p < .05, corrected cluster statistic $p < .05$).

415
416
Inspecting the plot for the difference between expected and unexpected

trials provides clues as to how this process might occur. As expected, for the time

that first grating was presented there was significant on- and off-axis encoding for

first grating orientation, with little difference between expectation conditions. More

interestingly, in the unexpected alternation condition there was significantly better

on-axis orientation selectivity for the first grating orientation between 150 and 300

ms after the onset of second grating, relative to the expected alternation condition

(cluster p = .03; upper right quadrant of right panel in Figure 7). Furthermore, over

the same time period, there was significantly better off-axis generalization of the

first grating representation to the second grating epoch (cluster p = .03; lower right

quadrant of right panel in Figure 7). This is confirmed by the occurrence of

significantly increased off-axis orientation selectivity between training time of -500

ms and +300 ms. The time course of this effect is consistent with our earlier

findings, as the unexpected condition yielded better orientation selectivity for the

second grating orientation between 79 and 150 ms after the second Gabor

presentation. An intriguing possibility is that the relevant neural circuits re-activate

the representation of the expected stimulus when faced with unexpected sensory

signals.

435   **Discussion**

Our findings demonstrate that repetition suppression and expectation have

distinct effects on neural representations of simple visual stimuli. We found that

repetition suppression had no effect on orientation selectivity, even though the

neural response to repeated stimuli was significantly decreased over occipital-

parietal areas. Unexpected stimuli, on the other hand, showed significantly

increased orientation-selectivity at an earlier time-point after stimulus onset than

expected stimuli. This same early representation of the unexpected stimulus

appeared to be reactivated at 200-300 ms after the initial neural response,

supporting the idea that sensory expectations may be updated through comparison

with incoming sensory evidence. These results suggest that repetition suppression

and prediction error are separable and independent neural computations.

The present work provides a significant advance in our understanding of how

predictive coding allows the brain to process incoming sensory information by

comparing what is expected with what actually occurs. How expectations affect

neural responses has been extensively investigated using mismatch negativity

paradigms in which an unexpected stimulus causes a significantly larger neural

response than an expected stimulus (Bekinschtein et al., 2009; Garrido et al., 2009;

Repetition suppression and prediction error

Näätänen et al., 2007). This increased response to an unexpected stimulus has

often been attributed to the generation of a prediction error that updates the

expectation based on a conflict between the sensory evidence and the prior

(Garrido et al., 2009). By contrast, our results suggest that the increased response

may reflect two distinct processes that occur at separate times. Incoming sensory

information is first evaluated against the prior (which occurs very early after stimulus

presentation). When an unexpected stimulus is detected and generates a prediction

error, the representation is amplified through gain enhancement. Later, around 300

ms after stimulus presentation, this same representation is reactivated to update the

expectation against the initially predicted representation. Our findings thus provide

a novel insight into how predictive coding might change neural representations of

sensory information.

Multivariate pattern classification methods have been widely used to examine

BOLD activity (Haxby, Connolly, & Guntupalli, 2014; Kamitani & Tong, 2005; 2006;

Kriegeskorte et al., 2008), but have only recently been applied to EEG and MEG

data (Cichy, Ramirez, & Pantazis, 2015; Garcia et al., 2013; King & Dehaene, 2014;

King et al., 2013; 2014; 2016; Myers et al., 2015; Wolff et al., 2017). These kinds of

analyses permit non-invasive imaging approaches to move beyond the

measurement of overall levels of neural activity, and instead to uncover subtle

patterns of neural activity associated with specific stimuli or task demands, which

are more directly comparable with those obtained from invasive neurophysiological

recordings of neuronal activity (Constantinescu, O'Reilly, & Behrens, 2016; Doeller,

Barry, & Burgess, 2010; Kamitani & Tong, 2006; Sprague & Serences, 2013). Such

model-based approaches are especially appealing since changes in overall activity

levels could be consistent with many different variations in the nature of the

underlying neural representations. For instance, sharpening and gain reduction

might both reduce the magnitude of an overall neural response but yield very

different effects on the amount of information conveyed by those representations.

The current work applied these model-based approaches to EEG data to

determine how prediction and repetition suppression affect neural representations

of perceptual information. We chose to use EEG so we could recover the temporal

dynamics of these effects – something that would not be possible with the BOLD

signal used in fMRI – and because EEG is the most widely-used tool for measuring

expectation effects in human participants (see Garrido:2009dp and Paavilainen,

2013 for review), thus facilitating comparison of our findings with those of other

studies. We estimated orientation-selectivity using all EEG electrodes distributed

across the scalp for two principal reasons. First, we wanted to limit experimenter

degrees of freedom (Simmons, Nelson, & Simonsohn, 2011) potentially introduced

through the post-hoc selection of subsets of electrodes. Second, given the broad

spatial resolution of EEG, we reasoned that activity recorded from electrodes at any

given scalp location could potentially carry important feature-selective information

from a number of neural sources. The results revealed that orientation-selective

information appears largely driven by electrodes over occipital-parietal regions

(Figure 3D), consistent with a number of previous studies that employed visual

decoding of M/EEG data (Cichy et al., 2015; Cichy, Pantazis, & Oliva, 2014; Stokes,

Wolff, & Spaak, 2015). Having said this, and as noted above, it is entirely possible

that the effects we observed here arose from sources well beyond the occipital and

parietal regions, potentially even including outside the visual cortical hierarchy.

Future studies using invasive methods will be needed to directly measure neuronal

responses across the brain to pinpoint the precise locations of these effects.

Surprisingly few studies have used invasive recording methods to examine

how predictive coding affects stimulus representations at the neuronal level. One

study in macaques (Kaliukhovich & Vogels, 2010) used the same design as the one

introduced by Summerfield and colleagues, but using high-level objects (fractals

and real-world objects) as stimuli and found that expectation did not attenuate

repetition suppression in either spiking activity or local field potentials within the

inferior temporal cortex. A later fMRI study in humans (Kovács, Kaiser, Kaliukhovich,

Vidnyánszky, & Vogels, 2013) used a similar stimulus set, and also found no

attenuation of repetition suppression by expectation in the same cortical region. A

follow-up study provided a potential explanation for these findings by showing that

the attenuation of neural responses associated with repetition suppression are

found with familiar stimuli, but not with unfamiliar stimuli (Grotheer & Kovács, 2014).

Viewed in this light, the stimulus sets used by Kaliukhovich and Vogels (2010) might

not have been sufficiently familiar to yield effects of expectation in their animals. In

any case, further work will need to be undertaken to determine whether expectation

effects dependent on stimulus familiarity.

Other work has shown that context plays an important role in determining

the magnitude of neuronal responses to sensory events. Thus, for example,

(Ulanovsky, Las, & Nelken, 2003) found that rare auditory stimuli generate

significantly larger responses in primary auditory cortical neurons than more

commonly occurring stimuli. This result has been interpreted as a single-neuron

analogue of the mismatch negativity, but the design used in the study did not

control for adaptation effects, thus making it difficult to draw an unambiguous

comparison with the current work. In the visual domain, oddball stimuli have also

been found to modulate neuronal activity in rats, characterised by an enhancement

of responses in the higher-order latero-intermediate area (Vinken, Vogels, & Op de

Beeck, 2017). Morever, Fiser et al. (2016) found that neurons in mouse primary

visual cortex show a greater calcium response when task-irrelevant visual stimuli

that had been presented during training were omitted, suggesting that an

established expectation had been violated. This result is consistent with the

literature on the mismatch negativity, in which the omission of an expected stimulus

results in a large prediction error (Garrido et al., 2009; Wacongne et al., 2011). In

non-human primates, neurons in the inferior temporal cortex show an enhanced

response to unexpected relative to expected stimuli (Kaposvari, Kumar, & Vogels,

2018), and population decoding accuracy is higher for unexpected compared with

expected stimuli (Kumar, Kaposvari, & Vogels, 2017).

Unlike the effects of expectation, there is a large body of electrophysiological

work showing that sensory adaptation influences neuronal activity (Adibi et al.,

2013b; Adibi, Clifford, & Arabzadeh, 2013a; Felsen et al., 2002; Kohn & Movshon,

2004; Patterson, Wissig, & Kohn, 2013) . For instance, there is a sharpening of

stimulus selectivity in MT neurons following 40 s of adaptation to a drifting grating

(Kohn & Movshon, 2004). As we have highlighted, however, prolonged adaptation is

Repetition suppression and prediction error

likely also associated with a significant prediction that the next stimulus will be the

same as the previous one. Perhaps more relevant to the current results, Patterson

et al. (2013) found that the width of orientation tuning in V1 is only marginally

sharpened following brief (400 ms) periods of adaptation. Again, however, their

study did not control for expectation, so it is impossible to determine the role of

predictive coding in their observations. Our finding that repetition suppression did

not affect the bandwidth of orientation selectivity measured using EEG is also

consistent with models of orientation adaptation based on human psychophysical

data, which suggest that adaptation does not affect the tuning width of the adapted

neural populations (Clifford, 2002; 2014; Dickinson, Almeida, Bell, & Badcock, 2010;

Dickinson, Morgan, Tang, & Badcock, 2017; Tang, Dickinson, Visser, & Badcock,

2015).

We found that prediction errors were associated with increased gain in early

stimulus representations, but did not affect the sharpness of their tuning. This lack

of sharpening is in contrast to the findings of a previous study (Kok et al., 2012), in

which a high-level prediction error led to 'sharper' multivariate decoding for

expected versus unexpected visual stimuli. In their study, Kok et al. (2012) used an

auditory tone to cue the orientation of a subsequent visual stimulus, and found

significantly less off-label classification accuracy for predicted than for unpredicted

stimuli. They concluded that predictions cause sharpening of stimulus

representations. More recently, using the same task combined with a forward

encoding approach, Kok et al. (2017) showed that response gain is increased for a

predicted stimulus (Kok, Mostert, & de Lange, 2017).

Repetition suppression and prediction error

It is natural to ask why the results of the current study differ from those of

Kok and colleagues outlined above. One possible explanation lies in the different

approaches used to generate expectations across the studies. Specifically, whereas

Kok et al. manipulated expectations by pairing an auditory cue with a visual

stimulus, we exploited the properties of the visual stimuli themselves (i.e., their

orientation) to generate expectations within blocks of trials. An intriguing possibility

is that predictions requiring integration of sensory events from two or more

modalities lead to increased gain, whereas predictions made within a single

sensory modality lead to decreased gain. This might in turn relate to the noted

differences between simple 'local' and higher-order 'global' type predictions

(Bekinschtein et al., 2009; King et al., 2014), which lead to distinct patterns of

stimulus-selective decoding. A similar discrepancy relating to the effects of

attention on sensory representations has been widely discussed, with some studies

finding sharpening of stimulus representations with attention, and others showing

gain enhancement (Liu, Larsson, & Carrasco, 2007; Maunsell, 2015; Maunsell &

Treue, 2006; Treue & Trujillo, 1999). *The differences between these results may*

potentially have arisen because the tasks relied upon different types of attention

(e.g., spatial versus feature-based). Future studies could determine whether this

same divergence occurs for prediction effects.

In summary, we have shown that repetition suppression and expectation

differentially affect the neural representation of simple, but fundamental, sensory

features. Our results further highlight how the context in which a stimulus occurs,

not just its features, affect the way it is represented by the brain. Our findings

suggest encoding priority through increased gain may be given to unexpected

events which could potentially speed responses. This prioritized representation is

then re-activated at a later time period, supporting the idea that feedback from

higher cortical areas reactivates an initial sensory representation in early cortical

areas.

596   **Method**

Participants

A group of 15 healthy adult volunteers (9 females, median age = 20.5 yr,

range = 18 to 37 yr) participated in exchange for partial course credit or financial

reimbursement (AUD$20/hr). We based our sample size on work that investigated

the interaction between repetition suppression and prediction error (Summerfield et

al., 2008), and that used forward encoding modelling to investigate orientation

selectivity using MEG with a comparable number of trials as the current study

(Myers et al., 2015). Each person provided written informed consent prior to

participation, and had normal or corrected-to-normal vision. The study was

approved by The University of Queensland Human Research Ethics Committee and

was in accordance with the Declaration of Helsinki.

Experimental setup

The experiment was conducted inside a dimly illuminated room with the

participants seated in a comfortable chair. The stimuli were displayed on a 22-inch

LED monitor (resolution 1920 x 1080 pixels, refresh rate 120 Hz) using the

PsychToolbox presentation software (Brainard, 1997; Pelli, 1997) for MATLAB

(v7.3). Viewing distance was maintained at 45 cm using a chinrest, meaning the

screen subtended 61.18° x 36.87° (each pixel 2.4' x 2.4').

Task

The stimuli were Gabors (diameter: 5°, spatial frequency: 2 c/°, 100%

contrast) presented centrally in pairs for 100 ms, separated by 500 ms (600 ms

stimulus onset asynchrony) with a variable (650 to 750 ms) inter-stimulus interval

between trials. Across the trials, the orientations of the Gabors were evenly spaced

between 0° and 160° (in 20° steps) so we could reconstruct orientation selectivity

contained within the EEG response using forward encoding modelling. The

relationship of the orientations of the pairs Gabors was also used to construct the

different repetition suppression and prediction conditions. The orientation presented

in the second Gabor in the pair could either repeat or alternate with respect to the

orientation of the first Gabor. In the alternation trials, the orientation of the first

Gabor was drawn randomly, without replacement, from an even distribution of

orientations that was different to the orientation of the second Gabor. To vary the

degree of prediction, in half of the blocks 80% of the trials had repeated

orientations and 20% of the trials had alternating orientations, whereas in the other

half of the blocks these contingencies were reversed. This design allowed us to

separately examine the effects of repetition suppression and prediction because of

the orthogonal nature of the blocked design. The blocks of 135 trials (~3 mins)

switched between the expectation of a repeating or alternating pattern, with the

starting condition counterbalanced across participants.

The participants' task was to monitor the visual streams for rare, faintly

coloured (red or green) Gabors and to discriminate the colour as quickly and

accurately as possible. Any trial with a coloured target was excluded from analysis.

The orientation match between the pairs was made to be consistent with the

dominant contingency (i.e., repeated or alternating) within that block. Pilot testing

was used prior to the main experiment to set the task at approximately threshold, to

ensure that participants focused exclusively on the colour-discrimination task rather

than the orientation contingencies associated with prediction and repetition. Only

one participant reported being aware of the changing stimulus contingencies across

the blocks when asked at the end of the experiment, and excluding this

participant's data had no effect on the key results reported here. Self-paced breaks

were provided between each of the 20 blocks within a session, at which time

feedback was provided on performance in the preceding block. Each participant

completed two sessions of 2700 trials each (5400 trials in total), with each session

lasting around 70 mins of experimental time and 45 mins of EEG setup.

EEG acquisition and pre-processing

Continuous EEG data were recorded using a BioSemi Active Two system

(BioSemi, Amsterdam, Netherlands). The signal was digitised at 1024 Hz sampling

rate with a 24-bit A/D conversion. The 64 active scalp Ag/AgCl electrodes were

arranged according to the international standard 10–20 system for electrode

placement (Oostenveld & Praamstra, 2001) using a nylon head cap. As per BioSemi

system design, the common mode sense and driven right leg electrodes served as

the ground, and all scalp electrodes were referenced to the common mode sense

during recording.

Offline EEG pre-processing was performed using EEGLAB in accordance

with best practice procedures (Bigdely-Shamlo, Mullen, Kothe, Su, & Robbins,

2015; Keil et al., 2014). The data were initially down-sampled to 256 Hz and

subjected to a 0.5 Hz high-pass filter to remove slow baseline drifts. Electrical line

noise was removed using *clean_line.m,* and *clean_rawdata.m* in EEGLAB (Delorme

& Makeig, 2004) was used to remove bad channels (identified using Artifact

Subspace Reconstruction), which were then interpolated from the neighbouring

electrodes. Data were then re-referenced to the common average before being

epoched into segments around each stimulus pair (-0.5 s to 1.25 s from the first

stimulus in the pair). Systematic artefacts from eye blinks, movements and muscle

activity were identified using semi-automated procedures in the SASICA toolbox

(Chaumon, Bishop, & Busch, 2015) and regressed out of the signal. After this stage,

any trial with a peak voltage exceeding ± 100 *uV* was excluded from the analysis.

The data were then baseline corrected to the mean EEG activity from -100 to 0 ms

before the presentation of the second Gabor in the pair.

Design

We used a modified version of a factorial design that has previously been

used to separately examine the effects of repetition suppression and prediction

error (Kaliukhovich & Vogels, 2010; Kovács et al., 2013; Summerfield et al., 2008;

2011; Todorovic et al., 2011; Todorovic & de Lange, 2012). By comparing the two

repeat conditions with the two alternating conditions, we could examine repetition

suppression while controlling for different levels of expectation. Conversely, by

comparing across the expected and unexpected trials, we could examine prediction

error while controlling for repetition suppression.

Forward encoding modelling

We used a forward encoding approach to estimate the amount of orientation-

selective information contained in the EEG data at each time point of the trial. This

approach differs from standard decoding approaches by modelling each presented

orientation as a continuous variable of a set of tuned orientation-selective channels.

The forward-encoding technique has been successfully used to reconstruct colour

(Brouwer & Heeger, 2009), spatial (Sprague & Serences, 2013) and orientation

(Ester, Sutterer, Serences, & Awh, 2016) selectivity in fMRI data. More recently it

has been extended to orientation (Garcia et al., 2013; Kok et al., 2017; Myers et al.,

2015; Wolff et al., 2017) and space (Foster, Sutterer, Serences, Vogel, & Awh, 2016)

encoding of MEG and EEG data, which allows for far greater temporal resolution

than fMRI.

Repetition suppression and prediction error

Figure 8. A schematic of the forward-encoding approach applied to EEG activity.
(A) Participants viewed individual gratings at fixation, each with a specific
orientation. **(B)** Neural activity evoked by each grating was measured over the entire
scalp. **(C)** Evoked neural responses were convolved with canonical orientation-
selective functions (grey lines in C) to determine coefficients for the different
orientations (coloured dots and lines, which match the colours of the outlined
gratings in A). These coefficients were then use to generate a regression matrix. **(D)**
General linear modelling was used on a subset of training trials to generate weights
for each channel. These weights were inverted and simultaneously applied to an
independent test set of data to recover orientation selectivity in the EEG activity. As
EEG activity has high temporal resolution we can apply the procedure to many
epochs following stimulus presentation to determine the temporal dynamics of
orientation processing (see Figure 3).

We applied forward encoding modelling to determine how repetition

suppression and prediction error affected orientation selectivity. To do this, the

second orientation (Figure 8A) in the Gabor pair in each trial was used to construct a

regression matrix, with separate regressors for the 9 orientations used across the

experiment. This regression matrix was convolved with a set of basis functions (half

cosines raised to the 8^{th} power (Figure 8C), which allowed complete and unbiased

coverage of orientation space) to allow us to pool similar information patterns

across nearby orientations (Brouwer & Heeger, 2009). We used this tuned

regression matrix to estimate time-resolved orientation selectivity contained within

the EEG activity in a 16 ms sliding window, in 4 ms steps (Figure 8B; Myers et al.,

2015). To avoid overfitting, we used a leave-one-out cross-validation procedure

722   where the regression weights were estimated for a training set and applied to an

723   independent test set (Figure 8D). All trial types (including target trials) were used in

724   training and test sets. This was done by solving the linear equation:

$$B_1 = WC_1 \tag{1}$$

726   Where B_1 (64 sensors x N training trials) is the electrode data for the training set, C_1

727   (9 channels x N training trials) is the tuned channel response across the training

728   trials, and W is the weight matrix for the sensors we want to estimate (64 sensors x

729   9 channels). W can be estimated using least square regression to solve equation (2):

$$W = (C_1 \, C_1^T)^{-1} \, C_1^T \, B_1 \tag{2}$$

731   The channel response in the test set C_2 (9 channels x N test trials) was estimated

732   using the weights in (2) and applied to activity in B_2 (64 sensors x N test trials).

$$C^2 = (W \, W^T) \, W^T \, B^2 \tag{3}$$

We repeated this process by holding one trial out as test, and training on the

remaining trials until all trials had been used in test and training. The procedure was

repeated for each trial within the trial epoch. We then shifted all trials to a common

orientation, meaning that 0° corresponded to the orientation presented on each trial.

The reconstructed channel activations were separated into the four conditions, and

averaged over trials. These responses were then smoothed with a Gaussian kernel

with a 16 ms window, and fitted with a Gaussian function (4) using non-linear least

square regression to quantify the amount of orientation selective activity.

$$G(x) = A \, exp(-\frac{(x-\varphi)^2}{2\sigma^2}) + C \tag{4}$$

Where *A* is the amplitude representing the amount of orientation selective activity,

is the orientation the function is centred on (in degrees), is the width (degrees) and

C is a constant used to account for non-orientation selective baseline shifts.

Multivariate pattern analysis

We conducted a multivariate pattern analysis to build upon the initial forward

encoding results which showed that unexpected stimuli elicit greater orientation

selectivity than expected stimuli. This analysis used the same data as the forward

encoding analysis. We used the *classify* function from *Matlab 2017a* with the

'diaglinear' option to implement a Naive Bayes classifier. For each time point, we

used the same cross-validation procedure as the forward encoding modelling with

the same averaging procedure to select train and test sets of data. The classifier

was given the orientations of the training data and predicted the orientation of the

test data. A trial was labelled correct if the presented orientation was produced. To

facilitate comparison of the results with those of (Kok et al., 2012), we found the

peak classification accuracy for each participant in the 600 ms following stimulus

presentation.

Statistical testing

A non-parametric sign permutation test was used to determine the null

distribution for testing (Wolff et al., 2017). This method makes no assumptions

about the underlying shape of the null distribution. This was done by randomly

flipping the sign of the data for the participants with equal probability. Fifty

thousand (50,000) permutations were used for the time-series data, whereas only

5000 were used for the temporal generalization plots because of the significantly

greater computational demands.

Cluster-based non-parametric correction (50,000 permutations for timeseries

768 and 5,000 for temporal generalization) was used to account for multiple

769 comparisons, and determined whether there were statistical differences between

770 the contrasting conditions. Paired-samples t-tests were used to follow up the

771 analysis in Figure 4 within a specified time window, and no correction was applied.

772 A two-way repeated measures ANOVA (implemented using GraphPad Prism 7.0c,

773 La Jolla California, USA) was used to analyse the multivariate pattern analysis

774 results shown in Figure 5.

775 **Acknowledgements**

776  This work was supported by the Australian Research Council (ARC) Centre of

777 Excellence for Integrative Brain Function (ARC Centre Grant CE140100007) to JBM

778 and EA, and by an ARC Discovery Project (DP170100908) to EA. JBM was

779 supported by ARC Australian Laureate Fellowship (FL110100103).

References

Adibi, M., McDonald, J. S., Clifford, C. W. G., & Arabzadeh, E. (2013). Adaptation
improves neural coding efficiency despite increasing correlations in variability.
The Journal of Neuroscience, 33(5), 2108–2120.
http://doi.org/10.1523/JNEUROSCI.3449-12.2013
Auksztulewicz, R., & Friston, K. (2016). Repetition suppression and its contextual
determinants in predictive coding. *Cortex, 80*, 125–140.
http://doi.org/10.1016/j.cortex.2015.11.024
Barron, H. C., Garvert, M. M., & Behrens, T. E. J. (2016). Repetition suppression: a
means to index neural representations using BOLD? *Philosophical Transactions
of the Royal Society of London Series B, Biological Sciences, 371*(1705),
20150355. http://doi.org/10.1098/rstb.2015.0355
Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., & Naccache, L.
(2009). Neural signature of the conscious processing of auditory regularities.
Proceedings of the National Academy of Sciences, 106(5), 1672–1677.
http://doi.org/10.1073/pnas.0809667106
Benucci, A., Saleem, A. B., & Carandini, M. (2013). Adaptation maintains population
homeostasis in primary visual cortex. *Nature Neuroscience, 16*(6), 724–729.
http://doi.org/10.1038/nn.3382
Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K.-M., & Robbins, K. A. (2015). The
PREP pipeline: standardized preprocessing for large-scale EEG analysis.
Frontiers in Neuroinformatics, 9, B153. http://doi.org/10.3389/fninf.2015.00016
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433–436.
http://doi.org/10.1163/156856897X00357
Brouwer, G. J., & Heeger, D. J. (2009). Decoding and reconstructing color from
responses in human visual cortex. *The Journal of Neuroscience, 29*(44), 13992–
14003. http://doi.org/10.1523/JNEUROSCI.3577-09.2009
Buckner, R. L., Goodman, J., Burock, M., Rotte, M., Koutstaal, W., Schacter, D., et
al. (1998). Functional-Anatomic Correlates of Object Priming in Humans
Revealed by Rapid Presentation Event-Related fMRI. *Neuron, 20*(2), 285–296.
http://doi.org/10.1016/S0896-6273(00)80456-0
Chaumon, M., Bishop, D. V. M., & Busch, N. A. (2015). A practical guide to the
selection of independent components of the electroencephalogram for artifact
correction. *Journal of Neuroscience Methods, 250*, 47–63.
http://doi.org/10.1016/j.jneumeth.2015.02.025
Cichy, R. M., Pantazis, D., & Oliva, A. (2014). Resolving human object recognition in
space and time. *Nature Publishing Group*. http://doi.org/10.1038/nn.3635
Cichy, R. M., Ramirez, F. M., & Pantazis, D. (2015). Can visual information encoded
in cortical columns be decoded from magnetoencephalography data in humans?
NeuroImage, 121, 193–204. http://doi.org/10.1016/j.neuroimage.2015.07.011
Clifford, C. W. G. (2002). Perceptual adaptation: motion parallels orientation. *Trends
in Cognitive Sciences, 6*(3), 136–143.
Clifford, C. W. G. (2014). The tilt illusion: phenomenology and functional implications.
Vision Research, 104, 3–11. http://doi.org/10.1016/j.visres.2014.06.009

Constantinescu, A. O., O'Reilly, J. X., & Behrens, T. E. J. (2016). Organizing
conceptual knowledge in humans with a gridlike code. *Science*, *352*(6292),
1464–1468. http://doi.org/10.1126/science.aaf0941
Cui, Y., Wang, Y. V., Park, S. J. H., Demb, J. B., & Butts, D. A. (2016). Divisive
suppression explains high-precision firing and contrast adaptation in retinal
ganglion cells. *eLife*, *5*. http://doi.org/10.7554/eLife.19460
Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of
single-trial EEG dynamics including independent component analysis. *Journal of
Neuroscience Methods*, *134*(1), 9–21.
http://doi.org/10.1016/j.jneumeth.2003.10.009
Dickinson, J. E., Almeida, R. A., Bell, J., & Badcock, D. R. (2010). Global shape
aftereffects have a local substrate: A tilt aftereffect field. *Journal of Vision*,
10(13), 5–5. http://doi.org/10.1167/10.13.5
Dickinson, J. E., Morgan, S. K., Tang, M. F., & Badcock, D. R. (2017). Separate
banks of information channels encode size and aspect ratio. *Journal of Vision*,
17(3), 27. http://doi.org/10.1167/17.3.27
Diederen, K. M. J., Spencer, T., Vestergaard, M. D., Fletcher, P. C., & Schultz, W.
(2016). Adaptive Prediction Error Coding in the Human Midbrain and Striatum
Facilitates Behavioral Adaptation and Learning Efficiency. *Neuron*, *90*(5), 1127–
1138. http://doi.org/10.1016/j.neuron.2016.04.019
Doeller, C. F., Barry, C., & Burgess, N. (2010). Evidence for grid cells in a human
memory network. *Nature*, *463*(7281), 657–661.
http://doi.org/10.1038/nature08704
Ester, E. F., Sutterer, D. W., Serences, J. T., & Awh, E. (2016). Feature-Selective
Attentional Modulations in Human Frontoparietal Cortex. *The Journal of
Neuroscience*, *36*(31), 8188–8199. http://doi.org/10.1523/JNEUROSCI.3935-
15.2016
Fairhall, A. L., Lewen, G. D., Bialek, W., & de Ruyter van Steveninck, R. R. (2001).
Efficiency and ambiguity in an adaptive neural code. *Nature*, *412*(6849), 787–
792. http://doi.org/10.1038/35090500
Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K., & Awh, E. (2016). The
topography of alpha-band activity tracks the content of spatial working memory.
Journal of Neurophysiology, *115*(1), 168–177.
http://doi.org/10.1152/jn.00860.2015
Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the
Royal Society of London B: Biological Sciences*, *360*(1456), 815–836.
http://doi.org/10.1098/rstb.2005.1622
Garcia, J. O., Srinivasan, R., & Serences, J. T. (2013). Near-real-time feature-
selective modulations in human cortex. *Current Biology*, *23*(6), 515–522.
http://doi.org/10.1016/j.cub.2013.02.013
Garrido, M. I., Kilner, J. M., Stephan, K. E., & Friston, K. J. (2009). The mismatch
negativity: A review of underlying mechanisms. *Clinical Neurophysiology*, *120*(3),
453–463. http://doi.org/10.1016/j.clinph.2008.11.029
Gotts, S. J., Chow, C. C., & Martin, A. (2012). Repetition priming and repetition
suppression: A case for enhanced efficiency through neural synchronization.
Cognitive Neuroscience. http://doi.org/10.1080/17588928.2012.691277

Gross, C. G., Schiller, P. H., Wells, C., & Gerstein, G. L. (1967). Single-unit activity in
temporal association cortex of the monkey. *Journal of Neurophysiology*, *30*(4),
833–43.
Grotheer, M., & Kovács, G. (2014). Repetition probability effects depend on prior
experiences. *The Journal of Neuroscience*, *34*(19), 6640–6646.
http://doi.org/10.1523/JNEUROSCI.5326-13.2014
Haxby, J. V., Connolly, A. C., & Guntupalli, J. S. (2014). Decoding neural
representational spaces using multivariate pattern analysis. *Annual Review of
Neuroscience*, *37*(1), 435–456. http://doi.org/10.1146/annurev-neuro-062012-
170325
Henson, R. N. A., & Rugg, M. D. (2003). Neural response suppression,
haemodynamic repetition effects, and behavioural priming. *Neuropsychologia*,
41(3), 263–270. http://doi.org/10.1016/S0028-3932(02)00159-8
Kaliukhovich, D. A., & Vogels, R. (2010). Stimulus Repetition Probability Does Not
Affect Repetition Suppression in Macaque Inferior Temporal Cortex. *Cerebral
Cortex*, *21*(7), 1547–1558. http://doi.org/10.1093/cercor/bhq207
Kamitani, Y., & Tong, F. (2005). Decoding the visual and subjective contents of the
human brain. *Nature Neuroscience*, *8*(5), 679–685. http://doi.org/10.1038/nn1444
Kamitani, Y., & Tong, F. (2006). Decoding seen and attended motion directions from
activity in the human visual cortex. *Current Biology*, *16*(11), 1096–1102.
http://doi.org/10.1016/j.cub.2006.04.003
Kaposvari, P., Kumar, S., & Vogels, R. (2018). Statistical Learning Signals in
Macaque Inferior Temporal Cortex. *Cerebral Cortex*, *28*(1), 250–266.
http://doi.org/10.1093/cercor/bhw374
Kay, K. N., Naselaris, T., Prenger, R. J., & Gallant, J. L. (2008). Identifying natural
images from human brain activity. *Nature*, *452*(7185), 352–355.
http://doi.org/10.1038/nature06713
Keil, A., Debener, S., Gratton, G., Junghöfer, M., Kappenman, E. S., Luck, S. J., et
al. (2014). Committee report: Publication guidelines and recommendations for
studies using electroencephalography and magnetoencephalography.
Psychophysiology, *51*(1), 1–21. http://doi.org/10.1111/psyp.12147
Keller, A. J., Houlton, R., Kampa, B. M., Lesica, N. A., Mrsic-Flogel, T. D., Keller, G.
B., & Helmchen, F. (2017). Stimulus relevance modulates contrast adaptation in
visual cortex. *eLife*, *6*, e21589. http://doi.org/10.7554/eLife.21589
King, J. R., & Dehaene, S. (2014). Characterizing the dynamics of mental
representations: the temporal generalization method. *Trends in Cognitive
Sciences*, *18*(4), 203–210. http://doi.org/10.1016/j.tics.2014.01.002
King, J. R., Faugeras, F., Gramfort, A., Schurger, A., Karoui, El, I., Sitt, J. D., et al.
(2013). Single-trial decoding of auditory novelty responses facilitates the
detection of residual consciousness. *NeuroImage*, *83*, 726–738.
http://doi.org/10.1016/j.neuroimage.2013.07.013
King, J.-R., Gramfort, A., Schurger, A., Naccache, L., & Dehaene, S. (2014). Two
Distinct Dynamic Modes Subtend the Detection of Unexpected Sounds. *PloS
One*, *9*(1), e85791. http://doi.org/10.1371/journal.pone.0085791

King, J.-R., Pescetelli, N., & Dehaene, S. (2016). Brain Mechanisms Underlying the
Brief Maintenance of Seen and Unseen Sensory Information. *Neuron*, *92*(5),
1122–1134. http://doi.org/10.1016/j.neuron.2016.10.051
Kohn, A., & Movshon, J. A. (2004). Adaptation changes the direction tuning of
macaque MT neurons. *Nature Neuroscience*, *7*(7), 764–772.
http://doi.org/10.1038/nn1267
Kok, P., Jehee, J. F. M., & de Lange, F. P. (2012). Less Is More: Expectation
Sharpens Representations in the Primary Visual Cortex. *Neuron*, *75*(2), 265–
270. http://doi.org/10.1016/j.neuron.2012.04.034
Kok, P., Mostert, P., & de Lange, F. P. (2017). Prior expectations induce prestimulus
sensory templates. *Proceedings of the National Academy of Sciences of the
United States of America*, *26*, 201705652.
http://doi.org/10.1073/pnas.1705652114
Kourtzi, Z., & Kanwisher, N. (2001). Representation of Perceived Object Shape by
the Human Lateral Occipital Complex. *Science*, *293*(5534), 1506–1509.
http://doi.org/10.1126/science.1061133
Kovács, G., Kaiser, D., Kaliukhovich, D. A., Vidnyánszky, Z., & Vogels, R. (2013).
Repetition probability does not affect fMRI repetition suppression for objects. *The
Journal of Neuroscience*, *33*(23), 9805–9812.
http://doi.org/10.1523/JNEUROSCI.3423-12.2013
Kriegeskorte, N., Mur, M., Ruff, D. A., Kiani, R., Bodurka, J., Esteky, H., et al. (2008).
Matching Categorical Object Representations in Inferior Temporal Cortex of Man
and Monkey. *Neuron*, *60*(6), 1126–1141.
http://doi.org/10.1016/j.neuron.2008.10.043
Kumar, S., Kaposvari, P., & Vogels, R. (2017). Encoding of Predictable and
Unpredictable Stimuli by Inferior Temporal Cortical Neurons. *Journal of Cognitive
Neuroscience*, *29*(8), 1445–1454. http://doi.org/10.1162/jocn_a_01135
Kvale, M. N., & Schreiner, C. E. (2004). Short-Term Adaptation of Auditory
Receptive Fields to Dynamic Stimuli. *Journal of Neurophysiology*, *91*(2), 604–
612. http://doi.org/10.1152/jn.00484.2003
Larsson, J., & Smith, A. T. (2012). fMRI repetition suppression: neuronal adaptation
or stimulus expectation? *Cerebral Cortex*, *22*(3), 567–576.
http://doi.org/10.1093/cercor/bhr119
Liu, T., Larsson, J., & Carrasco, M. (2007). Feature-Based Attention Modulates
Orientation-Selective Responses in Human Visual Cortex. *Neuron*, *55*(2), 313–
323. http://doi.org/10.1016/j.neuron.2007.06.030
Maravall, M., Petersen, R. S., Fairhall, A. L., Arabzadeh, E., & Diamond, M. E.
(2007). Shifts in Coding Properties and Maintenance of Information Transmission
during Adaptation in Barrel Cortex. *PLoS Biology*, *5*(2), e19.
http://doi.org/10.1371/journal.pbio.0050019
Maunsell, J. H. R. (2015). Neuronal Mechanisms of Visual Attention. *Annual Review
of Vision Science*, *1*(1), 373–391. http://doi.org/10.1146/annurev-vision-082114-
035431
Maunsell, J. H. R., & Treue, S. (2006). Feature-based attention in visual cortex.
Trends in Neurosciences, *29*(6), 317–322.
http://doi.org/10.1016/j.tins.2006.04.001

Movshon, J. A., & Lennie, P. (1979). Pattern-selective adaptation in visual cortical
neurones. *Nature*, *278*(5707), 850–852.
Myers, N. E., Rohenkohl, G., Wyart, V., Woolrich, M. W., Nobre, A. C., Stokes, M.
G., & Frank, M. J. (2015). Testing sensory evidence against mnemonic
templates. *eLife*, *4*, e09000. http://doi.org/10.7554/eLife.09000
Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity
(MMN) in basic research of central auditory processing: A review. *Clinical*
Neurophysiology, *118*(12), 2544–2590.
http://doi.org/10.1016/j.clinph.2007.04.026
Oostenveld, R., & Praamstra, P. (2001). The five percent electrode system for high-
resolution EEG and ERP measurements. *Clinical Neurophysiology*, *112*(4), 713–
719.
Paavilainen, P. (2013). The mismatch-negativity (MMN) component of the auditory
event-related potential to violations of abstract regularities: A review.
International Journal of Psychophysiology, *88*(2), 109–123.
http://doi.org/10.1016/j.ijpsycho.2013.03.015
Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics:
Transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442.
Rao, R. P. N., & Ballard, D. H. (1999). Predictive coding in the visual cortex: a
functional interpretation of some extra-classical receptive-field effects. *Nature*
Neuroscience, *2*(1), 79–87. http://doi.org/10.1038/4580
Rasmussen, R. G., Schwartz, A., & Chase, S. M. (2017). Dynamic range adaptation
in primary motor cortical populations. *eLife*, *6*, e21409.
http://doi.org/10.7554/eLife.21409
Rentzeperis, I., Nikolaev, A. R., Kiper, D. C., & Van Leeuwen, C. (2012).
Relationship between neural response and adaptation selectivity to form and
color: an ERP study. *Frontiers in Human Neuroscience*, *6*.
http://doi.org/10.3389/fnhum.2012.00089
Salti, M., Monto, S., Charles, L., King, J.-R., Parkkonen, L., Dehaene, S., &
Johansen-Berg, H. (2015). Distinct cortical codes and temporal dynamics for
conscious and unconscious percepts. *eLife*, *4*, e05652.
http://doi.org/10.7554/eLife.05652
Schacter, D. L., & Buckner, R. L. (1998). Priming and the Brain. *Neuron*, *20*(2), 185–
195.
Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-Positive Psychology:
Undisclosed Flexibility in Data Collection and Analysis Allows Presenting
Anything as Significant. *Psychological Science*, *22*(11), 1359–1366.
http://doi.org/10.1177/0956797611417632
Smirnakis, S. M., Berry, M. J., Warland, D. K., Bialek, W., & Meister, M. (1997).
Adaptation of retinal processing to image contrast and spatial scale. *Nature*,
386(6620), 69–73. http://doi.org/10.1038/386069a0
Spaak, E., Watanabe, K., Funahashi, S., & Stokes, M. G. (2017). Stable and
Dynamic Coding for Working Memory in Primate Prefrontal Cortex. *The Journal*
of Neuroscience, *37*(27), 6503–6516. http://doi.org/10.1523/JNEUROSCI.3364-
16.2017

Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in
the human occipital, parietal and frontal cortices. *Nature Neuroscience*, *16*(12),
1879–1887. http://doi.org/10.1038/nn.3574
Stokes, M. G., Kusunoki, M., Sigala, N., Nili, H., Gaffan, D., & Duncan, J. (2013).
Dynamic Coding for Cognitive Control in Prefrontal Cortex. *Neuron*, *78*(2), 364–
375. http://doi.org/10.1016/j.neuron.2013.01.039
Stokes, M. G., Wolff, M. J., & Spaak, E. (2015). Decoding Rich Spatial Information
with High Temporal Resolution. *Trends in Cognitive Sciences*, *19*(11), 636–638.
http://doi.org/10.1016/j.tics.2015.08.016
Summerfield, C., Trittschuh, E. H., Monti, J. M., Mesulam, M. M., & Egner, T. (2008).
Neural repetition suppression reflects fulfilled perceptual expectations. *Nature*
Neuroscience, *11*(9), 1004–1006. http://doi.org/10.1038/nn.2163
Summerfield, C., Wyart, V., Mareike Johnen, V., & de Gardelle, V. (2011). Human
Scalp Electroencephalography Reveals that Repetition Suppression Varies with
Expectation. *Frontiers in Human Neuroscience*, *5*.
http://doi.org/10.3389/fnhum.2011.00067
Tang, M. F., Dickinson, J. E., Visser, T. A. W., & Badcock, D. R. (2015). The broad
orientation dependence of the motion streak aftereffect reveals interactions
between form and motion neurons. *Journal of Vision*, *15*(13), 4.
http://doi.org/10.1167/15.13.4
Todorovic, A., & de Lange, F. P. (2012). Repetition suppression and expectation
suppression are dissociable in time in early auditory evoked fields. *The Journal*
of Neuroscience, *32*(39), 13389–13395.
http://doi.org/10.1523/JNEUROSCI.2227-12.2012
Todorovic, A., van Ede, F., Maris, E., & de Lange, F. P. (2011). Prior Expectation
Mediates Neural Adaptation to Repeated Sounds in the Auditory Cortex: An
MEG Study. *The Journal of Neuroscience*, *31*(25), 9118–9123.
http://doi.org/10.1523/JNEUROSCI.1425-11.2011
Tootell, R. B., Hadjikhani, N. K., Vanduffel, W., Liu, A. K., Mendola, J. D., Sereno, M.
I., & Dale, A. M. (1998). Functional analysis of primary visual cortex (V1) in
humans. *Proceedings of the National Academy of Sciences of the United States*
of America, *95*(3), 811–817. http://doi.org/10.1038/340386a0
Tootell, R., Reppas, J. B., Dale, A. M., & Look, R. B. (1995). Visual motion aftereffect
in human cortical area MT revealed by functional magnetic resonance imaging.
Nature.
Treue, S., & Trujillo, J. (1999). Feature-based attention influences motion processing
gain in macaque visual cortex. *Nature*. http://doi.org/10.1038/21176
Ulanovsky, N., Las, L., & Nelken, I. (2003). Processing of low-probability sounds by
cortical neurons. *Nature Neuroscience*, *6*(4), 391–398.
http://doi.org/10.1038/nn1032
Vinken, K., Vogels, R., & Op de Beeck, H. (2017). Recent Visual Experience Shapes
Visual Processing in Rats through Stimulus-Specific Adaptation and Response
Enhancement. *Current Biology*, *27*(6), 914–919.
http://doi.org/10.1016/j.cub.2017.02.024
Wacongne, C., Labyt, E., van Wassenhove, V., Bekinschtein, T., Naccache, L., &
Dehaene, S. (2011). Evidence for a hierarchy of predictions and prediction errors

in human cortex. *Proceedings of the National Academy of Sciences of the United*
States of America, 108(51), 20754–20759.
http://doi.org/10.1073/pnas.1117807108
Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic hidden
states underlying working-memory-guided behavior. *Nature Neuroscience, 20*(6),
864–871. http://doi.org/10.1038/nn.4546
1058