Sporulation-specific cell division defects in *ylmE* mutants of *Streptomyces coelicolor* are

rescued by additional deletion of *ylmD*

Le Zhang[1], Joost Willemse[1], Paul A. Hoskisson[2] and Gilles P. van Wezel[1, *]

[1] *Molecular Biotechnology, Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE, The*

Netherlands

[2] *Strathclyde Institute of Pharmacy and Biomedical Science, University of Strathclyde, 161*

Cathedral Street, Glasgow, G4 0RE, United Kingdom

* Author for correspondence. Tel: +31 71 5274310; email: g.wezel@biology.leidenuniv.nl.

ABSTRACT

Cell division during the reproductive phase of the *Streptomyces* life-cycle requires tight

coordination between synchronous formation of multiple septa and DNA segregation. One

remarkable difference with most other bacterial systems is that cell division in *Streptomyces* is

positively controlled by the recruitment of FtsZ by SsgB. Here we show that deletion of *ylmD*

(SCO2081) or *ylmE* (SCO2080), which lie in operon with *ftsZ* in the *dcw* cluster of actinomycetes,

has major consequences for sporulation-specific cell division in *Streptomyces coelicolor*.

Electron and fluorescence microscopy demonstrated that *ylmE* mutants have a highly aberrant

phenotype with defective septum synthesis, and produce very few spores with low viability and

high heat sensitivity. FtsZ-ring formation was also highly disturbed in *ylmE* mutants. Deletion of

ylmD had a far less severe effect on sporulation. Interestingly, the additional deletion of *ylmD*

restored sporulation to the *ylmE* null mutant. YlmD and YlmE are not part of the divisome, but

instead localize diffusely in aerial hyphae, with differential intensity throughout the sporogenic

part of the hyphae. Taken together, our work reveals a function for YlmD and YlmE in the

control of sporulation-specific cell division in *S. coelicolor*, whereby the presence of YlmD alone

results in major developmental defects.

INTRODUCTION

INTRODUCTION

In unicellular bacteria, cell division divides a mother cell in two identical daughter cells, each

containing a single copy of the chromosome. The control of cell division thereby revolves

around finding the mid-cell position, and chromosome segregation and septum synthesis are

closely coordinated in time and space to avoid DNA damage by the nascent septum. The cell

division scaffold is formed by FtsZ, which is a homologue of tubulin [1] and forms a contractile ring

(or Z-ring) that mediates the recruitment of the cell division machinery to the division site

(reviewed in [2,3]). Septum-site selection and Z-ring stabilization are mediated by proteins like

FtsA and ZipA [4-6], ZapA [7] and SepF [8,9]. Z-ring (dis-)assembly is thereby actively controlled

(reviewed in [10]).

Streptomycetes are filamentous Gram-positive bacteria that belong to the phylum of

Actinobacteria. These bacteria produce over 60% of all known antibiotics and many other

bioactive natural products [11,12]. Exponential growth of the vegetative hyphae is achieved by

apical growth and branching. At this stage of the life cycle, cell division does not affect physical

separation of the cells, but instead long syncytial cells are formed that are separated by cross-

walls [13]. Hence, streptomycetes are model organisms for the study of multicellularity and

bacterial morphogenesis [14,15].

Most divisome components except FtsZ are not required for vegetative division,

presumably reflecting the lack of constriction [16]. Spacing between the cross-walls is highly

variable, and little is known of the way septum-site selection is controlled. Recently, using cryo-

electron tomography, our lab and others showed that intracellular membrane assemblies or

cross-membranes are involved in DNA protection during septum synthesis in young vegetative

hyphae, suggesting a novel way of cell-division control [17,18]. In addition, these multicellular

bacteria have a complex cytoskeleton, which among others plays a role in the organization of

the tip growth machinery [19,20].

Canonical division resulting in cell fission occurs during sporulation-specific cell division,

which requires all components of the divisome [16,21]. At this stage of the life cycle up to a

hundred septa are formed in a short time span, following a highly complex process of

coordinated cell division and DNA segregation, and visualized as long ladders of Z-rings [22].

Eventually, chains of unigenomic spores are formed, which have a thick protective spore wall

facilitating long-term survival in the environment. Though *ftsZ* null mutants are viable, they fail

to produce septa and hence do not sporulate, cell division is not essential for growth of

Streptomyces, which provides a unique system for the study of this process [23,24].

Sporulation-specific cell division is controlled by the SsgA-like proteins (SALPs), which are

exclusively found in morphologically complex actinobacteria [25,26]. The canonical view is that cell

division is negatively controlled by the action of the Min system that inhibits division away from

midcell, and by nucleoid occlusion (Noc) to avoid septum synthesis near the chromosome to

avoid DNA damage, as seen in *B. subtilis* and *E. coli*. In contrast, cell division in streptomycetes is

positively controlled by the recruitment of FtsZ to future septum sites by SsgB, in an SsgA-

dependent manner [27]. As a consequence, both SsgA and SsgB are required for sporulation [28,29].

We recently showed that SepG (formerly called YlmG) assists in docking of SsgB to the

membrane, and also plays a major role in maintaining the nucleoid shape in the spores [30].

Many of the genes for the components of the divisome and the cell-wall biosynthetic

machinery are located in the so-called *dcw* cluster (division and cell-wall synthesis; [31]). Most of

these genes have been studied extensively and their functions have been well characterized.

However, little is known of the genes *ylmD* and *ylmE* that lie immediately downstream of, and

likely form an operon with, *ftsZ* on the genome of streptomycetes and many other bacteria,

including firmicutes. Earlier work on a mutant of *S. venezuelae* lacking both *ylmD* and *ylmE*

showed that the double mutant had no obvious phenotypic defects [32].

In this study, we show that deletion of *ylmE* alone results in severe cell division defects,

while deletion of *ylmD* only slightly affects sporulation. Interestingly, the cell division defects of

ylmE mutants were rescued by the additional deletion of *ylmD*, and *ylmDE* mutants had no

obvious sporulation defects. These data strongly suggest that expression of YlmD alone is

detrimental for sporulation-specific cell division in streptomycetes, which is counteracted by

YlmE. This is consistent with the phylogenetic evidence that some bacteria only harbor an

ortholog of *ylmE,* whereas *ylmD* never occurs without *ylmE*.

RESULTS & DISCUSSION

Phylogenetic analysis of YlmE (SCO2080) and YlmD (SCO2081)

Many of the genes in the *dcw* gene cluster have been extensively studied and their functions are

well established. However, this is not the case for *ylmD* and *ylmE*, which lie immediately

downstream of *ftsZ* in many bacteria. In all *Streptomyces* genomes analyzed, *ftsZ* (SCO2082),

ylmD (SCO2081) and *ylmE* (SCO2080) form an operon, an observation that is supported by high-

resolution transcript mapping [33]; moreover there is apparent translational fusion between *ftsZ*

and *ylmD* (overlapping start and stop codons) and only 6 nt spacing between *ylmD* and *ylmE*.

This transcriptional coupling to *ftsZ* suggests that these genes may play a prominent role in the

cell division process. Transcript levels of *ftsZ* are similar to those of *ylmD* and *ylmE* during

vegetative growth on MM agar plates; however, transcription of *ftsZ* is enhanced during

sporulation, while that of *ylmD* and *ylmE* is not significantly altered [34]. YlmD and YlmE are wide-

spread in Gram-positive bacteria, and particularly in actinobacteria. SCO2081 and SCO2080

share 32% aa identity with YlmD and YlmE of *B. subtilis*, respectively. Orthologues of these

proteins are also found in several genera of Gram-negative bacteria. Phylogenetic analysis of

the YlmD and YlmE proteins in actinobacteria shows that while YlmE is widespread, YlmD is

often absent in actinobacteria, such as *Stackebrandtia, Catenulispora, Salinispora,*

Micromonospora, Amycolatopsis and *Mycobacterium*, several of which are spore-forming

actinobacteria (Fig. 1). Remarkably, there are no examples of *ylmD* being present in the absence

of *ylmE* and thus it would appear that loss of YlmD (SCO2081) has occurred on multiple

occasions, given that there is wide but patchy distribution of this gene across distinct

actinobacteria lineages and the sequences clade tightly within the accepted actinomycete

phylogenies.

Analysis of *ylmD* using the EMBL String engine [35] shows functional linkage in two groups

to cell division associated genes (*sepF* (SCO2079)*,* SCO2085 and *ftsZ*) along with a group of

mainly hypothetical proteins (STRING Data link: http://bit.ly/2gc5kCB). YlmD has a domain that

has homology to the multiple-copper polyphenol oxidoreductase laccases, which are

oxidoreductases that are widely distributed in both prokaryotes and eukaryotes [36]. A similar

analysis of YlmE also indicates a functional linkage to cell-wall biosynthesis and cell division

(STRING Data link: http://bit.ly/2gbPyHK). YlmE is a member of the family of YBL036c-like

proteins, which generally contain pyridoxal 5-phosphate dependent enzymes. The structure of

YBL036c from *Saccharomyces cerevisiae* was resolved at 2.0 Å resolution (PDB 1CT5;[37]). The

protein has homology to the N-terminal domains of alanine racemases but lacks of the β-

sandwich domain which would likely limit the activity of YBL036c as alanine or non-specific

amino acid racemase [38]. To test the hypothesis that YlmE may have alanine racemase activity

and thus may play a role in determining the amino acid composition of the peptidoglycan, we

tested purified YlmE for alanine racemase assays as described previously [39], using alanine

racemase Alr as positive control. Whereas purified Alr successfully catalyzed the conversion of L-

alanine to D-alanine, YlmE could not perform this reaction under the same conditions, and over-

expression of YlmE failed to restore a D-Ala prototrophy to *alr* null mutants (data not shown).

These observations coupled with protein structure homology data make it highly unlikely that

YlmE functions as an alanine racemase *in vivo*.

143 ***ylmD* and y*lmE* are required for proper sporulation**

144 To analyze the role of *ylmD* and *ylmE*, deletion mutants were created in *S. coelicolor* M145 as

145 detailed in the Materials and Methods section. The +25 to +696 region of *ylmE* (SCO2080) or the

146 +25 to +705 region of *ylmD* (SCO2081) were replaced by the apramycin resistance cassette,

147 followed by deletion of the cassette using the Cre recombinase so as to avoid polar effects. For

148 each mutant, four independent mutants were selected and all had the same phenotype.

149 Therefore, one was selected for more detailed analysis, designated GAL47 (*S. coelicolor* M145

150 Δ*ylmD*) and GAL48 (*S. coelicolor* M145 Δ*ylmE*). The M145 *ylmDE* mutant (GAL130) was created

151 by replacing *ylmD* in *ylmE* null mutant GAL48 by the apramycin resistance cassette.

152 The *ylmD* null mutant GAL47 had a wild-type-like appearance, while *ylmE* null mutant

153 GAL48 hardly produced any grey pigment after 5 days incubation on SFM agar plates, indicative

154 of a failure to complete full development. Surprisingly, *ylmDE* mutant GAL130 had a grey

155 appearance similar to that of the parental strain M145 (Fig. 2A). Phase-contrast light microscopy

156 of impression prints of the top of the colonies demonstrated that while the parent produced

157 typical long spore chains, the *ylmD* null mutant produced abundant but aberrantly sized spores,

158 the *ylmE* null mutant produced fewer spores and those produced were unusually large. The

159 *ylmDE* double mutant produced abundant spores, though some were irregularly sized (Fig. 2B

160 and Fig. S1). The same defect in sporulation was observed on R5 and MM mannitol agar plates.

161 Antibiotic production was not affected, and the mutants produced normal levels of the

162 pigmented antibiotics actinorhodin and undecylprodigiosin.

163 To further ascertain that the sporulation defects were indeed solely due to the deletion

164 of the respective genes, we introduced plasmids expressing *ylmD* or *ylmD-egfp* in the in the

ylmD null mutant and *ylmE* or *ylmE-egfp* in the *ylmE* mutant, in all cases with transcription

directed from the native *ftsZ* promoter region. While the majority of the spores of the

complemented *ylmD* mutant had a regular appearance, those of the complemented *ylmE*

mutants still showed variable lengths (Fig. 2B). This partial complementation suggests that the

deletion of *ylmE* may have polar effect on the expression of its downstream genes. This is

supported by the fact that introduction of a plasmid harboring the entire region from *murD* to

divIVA fully restored sporulation to *ylmE* mutants (Fig 2B & Fig. S2). Promoter probing using the

redD reporter system [40] showed that besides the intergenic promoter between *ylmE* and *sepF*,

an additional promoter is located within *ylmE*, suggesting that deletion of the entire *ylmE* gene

negatively affects the transcription of *sepF* (data not shown). This may explain why *ylmE* alone

failed to fully complement the *ylmE* null mutant.

To quantify the spore lengths and their size distribution (*i.e.* variability), approximately 300

spores from the parental strain *S. coelicolor* M145, its *ylmD* and *ylmE* mutants and the

genetically complemented strains were measured from light images from phase-contrast

microscopy (100x magnification). Spore sizes were compared using a boxplot analysis (Fig. 2C).

Wild-type spores typically had lengths between 0.8-1.2 µm, with an average length of 1.01 ±

0.15 µm, while *ylmD* mutant spores showed a slightly broader distribution (Fig. 2C), with an

average length of 1.12 ± 0.27 µm. The average length was reduced to wild-type levels by

complementation, namely to 1.03 µm and 1.05 µm by introduction of *ylmD* and *ylmD-egfp*,

respectively. Spores of the *ylmE* null mutant were much larger and with a significantly wider

variation, showing an average length of 2.23 ± 0.86 µm (Fig. 2C). Genetic complementation of

the *ylmE* mutant restored spore length variation of *ylmE* mutant to wild-type levels, with an

average size of 1.03 ± 0.19 μm. Partial complementation of the *ylmE* null mutant was seen when

constructs expressing *ylmE* or *ylmE-egfp* were introduced into the mutant, with spore lengths of

1.62 ± 0.50 μm and 1.85 ± 0.59 μm, respectively. The average spore length of the combined

ylmDE mutant was much closer to that of the parental strain, namely 1.23 ± 0.34 μm, thereby

showing significantly smaller size variation than *ylmE* null mutants (Fig. 2C). The statistical

validity of the variations in spore sizes between mutants and the parental strain was validated

by a Kolgomorov-Smirnov test and a Mann-Whitney U test, which confirmed that the sizes of

mutant spores deviated significantly from those of wild-type spores.

To study the spores of *ylmD* and *ylmE* null mutants at high resolution, cryo-scanning

electron microscopy (SEM) was performed, which again demonstrated the sporulation defects.

The parental strain *S. coelicolor* M145 produced abundant spore chains, with nearly all hyphae

fully developed into mature spore chains (Fig. 3AB). The *ylmD* null mutant GAL47 frequently

produced aberrantly sized spores (Fig. 3CD), while in *ylmE* null mutant GAL48, precious few and

often aberrant spores were identified (Fig. 3EF). The same approach was used to create *ylmD*

and *ylmE* mutants in *Streptomyces lividans* 66, and these again showed the same morphology,

strongly supporting the notion that the observed phenotypes were solely due to the mutations

in *ylmD* or *ylmE* (data not shown). The highly similar phenotypes of the *S. coelicolor* and *S.*

lividans mutants support the notion that the genes play a key role in sporulation-specific cell

division and are consistent with the conserved function of the genes in sporulation-specific cell

division in *Streptomyces*.

Deletion of *ylmD* and especially *ylmE* results in reduced robustness of spores

Viability of the spores was tested by plating around 1000 spores - counted in a hemocytometer -

of *S. coelicolor* M145 and its mutants GAL47 (Δ*ylm*D) and GAL48 (Δ*ylmE*) onto SFM agar plates.

While the wild-type strain had close to 100% viability, spores of the *ylmD* and *ylmE* mutants had

reduced viability, with 50% and 60% viable spores, respectively. Spores of in particular the *ylmE*

mutant were heat sensitive, with only 11 ± 1% viability after 15 min incubation at 58^0C, as

compared to 47 ± 6% survival for *ylmD* null mutant spores and 55 ± 5% for the parental strain.

Genetic complementation of the *ylmE* null mutant restored wild-type survival to heat shock (58

± 3%). The same was true when *ylmD* was deleted in the *ylmE* null mutant (56 ± 4% survival).

To analyze the possible cell-wall defects in more detail, all strains were grown on SFM

agar plates for 5 days and analyzed by Transmission Electron Microscopy (TEM) (Fig. 4). The

parent produced typical spore chains and thick-walled spores. Conversely, *ylmD* and *ylmE*

mutant spores were deformed and highly irregular. In particular the spores of the *ylmE* mutant

had a thin wall similar to that of hyphae, suggesting that spore-wall synthesis was compromised.

YlmD and YlmE are required for peptidoglycan synthesis at the septum

To analyze cell wall and membrane distribution in *ylmD* and *ylmE* mutants, fluorescence

microscopy was performed on five-days old SFM surface-grown cultures of mutants GAL47

(M145 Δ*ylmD*) and GAL48 (M145 Δ*ylmE*) and compared to the parental strain M145.

Peptidoglycan precursors were stained with FITC-WGA or Oregon-WGA and membranes

visualized by staining with FM5-95. In wild-type pre-division aerial hyphae, long symmetrical

septal ladders were observed when stained for cell wall or membranes (Fig. 5A). In contrast,

aerial hyphae of the mutants showed highly disturbed cell wall and membrane distribution,

which was more pronounced in *ylmE* than in *ylmD* mutants (Fig. 5A). Septation of *ylmD* mutants

was complemented via re-introduction of *ylmD*; similarly, septation was restored to the *ylmE*

mutant by re-introduction of *ylmE*, although peptidoglycan synthesis and septum spacing was

still irregular (Fig. 5B). These data again show that the consequences of deleting *ylmE* are much

more severe for sporulation-specific cell division then when *ylmD* is deleted.

Cellular localization of YlmD and YlmE

Given the impact of *ylmDE* on sporulation, we analyzed how YlmD and YlmE were localized in

the hyphae of *S. coelicolor*. To this end, constructs based on the low-copy number vector

pHJL401 were prepared to allow the expression of YlmD-eGFP and YlmE-eGFP fusion proteins,

which were expressed from the natural *ftsZ* promoter region (see Materials and Methods

section for details). These constructs were then introduced into *S. coelicolor* M145 and (as a

control for functionality) in the mutants. The constructs partially restored development to the

respective mutants, with significant restoration of the sporulation defects in *ylmD* mutants, and

partial restoration of sporulation to *ylmE* mutants (Fig. 2B).

In vegetative hyphae, only very weak signals were obtained for YlmD-eGFP and YlmE-

eGFP, indicative of low protein expression (Fig. 6A). In early and late aerial hyphae, both YlmD-

eGFP and YlmE-eGFP became more visible and showed diffuse localization along the wall of the

aerial hyphae (Fig. 6A). During early sporulation, YlmD-eGFP and YlmE-eGFP showed an irregular

pattern of varying intensity, with some spores showing bright fluorescence indicative of high

concentrations of YlmD or YlmE, while others hardly showed any fluorescence (Fig 6A). This

suggests that YlmD and YlmE are differentially expressed throughout the nascent spore chains,

and do not exclusively co-localize with FtsZ and the divisome. To rule out possible proteolysis of

YlmD- or YlmE-eGFP fusion proteins, Western analysis was performed on extracts of surface-

grown *S. coelicolor* mycelia, using antibodies against GFP. *S. coelicolor* M145 with empty vector

and *S coelicolor* M145 expressing freely mobile eGFP [41] were included as controls. Only a single

band was identified in strains GAL50 and GAL49, corresponding to the expected lengths for

YlmD-eGFP and YlmE-eGFP, respectively (Fig. S3). In both cases, the predicted mass of the fusion

protein is around 52 kDa, which conforms well to the apparent mobility seen by Western

analysis. The strain expressing only eGFP showed the expected band of around 27 kDa for the

non-fused protein. Therefore, we conclude that the observed localizations can indeed be

ascribed to intact YlmD- or YlmE-eGFP.

The fluorescence intensity of YlmD-eGFP and YlmE-eGFP was measured on images

representing different developmental stages (Fig. 6B). As development progressed, the

fluorescence intensities increased for both YlmD-eGFP and YlmE-eGFP and reached peak levels

during the phase corresponding to late aerial growth and early sporulation. When spores

matured, YlmD-eGFP and YlmE-eGFP signals decreased to the level as in vegetative growth.

These results suggest YlmD and YlmE proteins are active primarily during the phase of

sporulation-specific cell division.

Localization of FtsZ is disturbed in *ylmD* and *ylmE* null mutants

To investigate the localization of FtsZ in the mutants, integrative plasmid pKF41 that expresses

FtsZ-eGFP from the native *ftsZ* promoter region [42] was introduced into the *ylmD* and *ylmE*

mutants, to generate strain GAL52 and GAL53, respectively. Prior to sporulation, typical ladders

of Z-rings were observed in the parental strain *S. coelicolor* M145, while *ylmD* and *ylmE* null

mutants showed abnormal Z-ladders (Fig. 7). In the absence of YlmD, ladders were still observed,

but the intensity varied and spacing between the individual rings was less regular, with many

neighboring Z-rings either close together or widely spaced. Consistent with the sporulation

defect, *ylmE* null mutants produced very few Z-rings, and the few Z-ladders that where formed

were highly irregular or unfinished, and significantly shorter than in the parental strain. Thus,

FtsZ localization is irregular in *ylmD* mutants and highly compromised in *ylmE* mutants.

A model for sporulation control by YlmDE in streptomycetes

Our data show that mutants of *S. coelicolor* lacking *ylmE* have severe developmental defects,

with highly compromised cell-wall synthesis and aberrant cell division. Mutants which lack *ylmD*

sporulated well, though many spores exhibited aberrant shapes and variable lengths.

Surprisingly, the additional deletion of *ylmD* greatly rescued the sporulation-deficient

phenotype of *ylmE* mutants, with the *ylmDE* null mutant producing abundant and mostly

regularly sized spores.

We propose that *YLMD* AND *YLME* Form a two-part system, reminiscent of toxin-

antitoxin systems [43]. Similarly dysfunction of one protein in the presence of another has also

been reported for the sporulation control protein WhiJ, following the observation that media-

dependent sporulation defects caused by a point mutation in *whiJ* (SCO4543) could be rescued

by the full deletion of *whiJ* [44]. The authors proposed a model wherein repression of

developmental genes by WhiJ could be released when WhiJ interacts with a WhiJ-associated

protein encoded by the adjacent gene SCO4542 [44]. It was suggested that the aberrant protein

WhiJ* still binds to operator sequences of developmental genes, leading to their transcriptional

repression, but fails to associate with SCO4542, with a permanent block of transcription of

developmental genes as a result [44]. Similarly, the expression of YlmD alone is detrimental for

sporulation, suggesting that YlmD acts in a deleterious manner towards sporulation, yet this

effect can be relieved by the presence of YlmE. Supporting evidence for this is that deletion of

both genes allows *Streptomyces* to develop normally, and by the observation that *ylmD* is never

found alone in bacterial genomes, while the occurrence of an orphan *ylmE* is seen frequently.

Further support comes from the report that *ylmDE* double mutants of *S. venezuelae* also

sporulate relatively normally [32]. This also indicates that this two-part system functions in a

similar manner in a range of *Streptomyces* species. This apparently rules out major divergence

between the two morphologically distinct *Streptomyces* clades, namely those that sporulate in

submerged cultures and those that do not [45,46]. However, it will be very interesting to see if

submerged sporulation of *S. venezuelae* is prevented by the deletion of just *ylmE*. Currently, we

are performing detailed structural and functional analysis of YlmD and YlmE, to elucidate their

biochemical function and their precise role in bacterial cell division.

METHODS

Phylogenetic analysis of *ylmD* and *ylmE*

The amino acid sequence of YlmD and YlmE were extracted from StrepDB

(http://strepdb.streptomyces.org.uk) and used to search the NCBI database

(www.ncbi.nlm.nih.gov) using BLASTP against the non-redundant protein sequence database.

Alignment of YlmD and YlmE was generated using ClustalW [47] followed by manual editing in

MEGA v. 4.0. The neighbour-joining trees [48] were generated with default parameters settings as

implemented in MEGA v. 4.0 [49]. The maximum-likelihood trees were made using the best fit

models predicted by MEGA. Tree reliability was estimated by bootstrapping with 1000

replicates. Trees were drawn with either N-J or ML algorithms gave trees of similar topologies

indicating that the phylogenies are likely to reliable.

Bacterial strains and media

All bacterial strains used in this study are listed in Table S1. *E. coli* JM109 was used for routine

cloning and ET12567 [50] to prepare nonmethylated DNA to bypass the methyl-specific restriction

system of *S. coelicolor*. *E. coli* strains were propagated in Luria broth, where appropriate

supplemented with antibiotics for selection, namely ampicillin (100 µg/ml end concentration),

apramycin (50 µg/ml) and/or chloramphenicol (25 µg/ml). *S. coelicolor* A3(2) M145 [51] and *S.*

lividans 66 [52] were obtained from the John Innes Centre strain collection. *S. coelicolor* strains

were grown on soya flour medium (SFM) or minimal media mannitol (MM) agar plates for

phenotypic characterization and on R5 agar plates for regeneration of protoplasts [53]. Antibiotics

used for screening Streptomyces were apramycin (20 µg/ml end concentration) and

thiostrepton (10 µg/ml).

Plasmids and constructs

All plasmids and constructs described in this study are summarized in Table S2. The

oligonucleotides used for PCR are listed in Table S3. PCR reactions were performed using Pfu

DNA polymerase as described [54]. All inserts of the constructs were verified by DNA sequencing,

which was performed at BaseClear (Leiden, The Netherlands).

Constructs for the deletion of ylmD and ylmE

The strategy for creating knock-out mutants is based on the unstable multi-copy vector pWHM3

[55] as described previously [56]. For each knock-out construct roughly 1.5 kb of upstream and

downstream region of the respective genes were amplified by PCR from cosmid St4A10 that

contains the *dcw* cluster of *S. coelicolor*. The upstream region was thereby cloned as an EcoRI-

XbaI fragment, and the downstream part as an XbaI-BamHI fragment, and these were ligated

into EcoRI-BamHI-digested pWHM3 (for the precise location of the oligonucleotides see Table

S3). In this way, an XbaI site was engineered in-between the flanking regions of the gene of

interest. This was then used to insert the apramycin resistance cassette *aac(3)IV* flanked by *loxP*

sites, using engineered XbaI sites. The presence of the *loxP* recognition sites allows the efficient

removal of the apramycin resistance cassette following the introduction of a plasmid pUWL-Cre

expressing the Cre recombinase [57,58]. Knock-out plasmids pGWS728 and pGWS729 were created

for the deletion of nucleotide positions +25/+696 of *ylmE* (SCO2080) and +25/+705 of *ylmD*

(SCO2081), whereby +1 refers to the translational start site of the respective genes. This

allowed first the replacement by the apramycin resistance cassette. Subsequently the

apramycin resistance cassette was removed using expression of pUWLCre. To create a *ylmDE*

double deletion mutant, construct pGWS1044 was created which contains the upstream region

of *ylmD* and downstream region of *ylmE* flanked by *loxP* sites, and with the apramycin

resistance cassette *aac(3)IV* inserted in between.

For complementation of the *ylmE* and *ylmD* null mutants, the entire coding regions of

SCO2080 and SCO2081 (with stop codons) were amplified from the *S. coelicolor* M145

chromosome using primer pairs ylmE_F+1 and ylmE_R+723 and ylmD_F+1 and ylmD_R+732,

respectively. The PCR products were digested with StuI/XbaI, and inserted downstream of the

native *ftsZ* promoter region in pHJL401, respectively. Thus constructs pGWS1042 and

pGWS1043 were generated that express *ylmE* and *ylmD*, respectively, under control of the *S.*

coelicolor ftsZ promoter region. Alternatively, construct pKR8 was used for complementation;

pKR8 is based on integrative vector pIJ8600 [59] and contains the 2227742-2241015 region of the

S. coelicolor genome, encompassing the end of *murX* (SCO2087), the entire coding sequences of

murD, ftsW, murG, ftsQ, ftsZ, ylmD, ylmE, sepF, sepG, divIVA and a large part of SCO2076

(encoding Ile-tRNA synthetase).

Constructs for the localization of YlmD and YlmE

The entire coding regions of SCO2080 and SCO2081 (without stop codons) were amplified from

the *S. coelicolor* M145 chromosome using primer pairs ylmE_F+1 and ylmE_R+717 and

ylmD_F+1 and ylmD_R+726, respectively. The PCR products were digested with StuI/BamHI, and

inserted downstream of the native *ftsZ* promoter region and immediately upstream of *egfp* in

pHJL401. The latter is a highly stable vector with low copy number that generally results in wild-

type transcription levels and is well suited for among others complementation experiments [40].

Thus constructs pGWS757 and pGWS758 were generated that express YlmE-eGFP and YlmD-

eGFP, respectively, from the native *S. coelicolor ftsZ* promoter region. Plasmid pKF41 expresses

FtsZ-eGFP from its own promoter region [42]. Constructs pGWS757 and pGWS758 were also used

to complement *ylmE* and *ylmD* null mutants, respectively.

eGFP detection using western blot analysis

Strains were grown on SFM agar plates containing the appropriate antibiotics overlaid with

cellophane disks. Biomass (mycelium and spores) was collected after 6 days of growth at 30 °C

and suspended in urea lysis buffer (8M urea, 2M thiourea, 4% CHAPS, 1mM PMSF and 65mM

DTT), followed by cell disruption via sonication. Western blots were performed as described [60].

Briefly, cell lysates were obtained after centrifugation at 30,000x g for 30 min at 4°C. Proteins

were separated on a 12.5% SDS-PAGE gel, and then transferred to PVDF membrane using Mini

Trans-Blot (Bio-Rad). As primary antibody a 1:2000 dilution of GFP antibody (ThermoFisher, REF

A11122) was used and incubated for 16 h at 4°C in the presence of 3% BSA. After washing, the

membrane was incubated with alkaline phosphatase conjugated anti-rabbit IgG (Sigma) as

secondary antibody for 1 h at room temperature. Visualization was done using BCIP/NBT color

development substrate.

Microscopy

Fluorescence and light microscopy were performed as described previously [61]. For

peptidoglycan staining we used FITC-labeled wheat germ agglutinin (FITC-WGA) or Oregon

Green 488 conjugated WGA (Oregon-WGA); for membrane staining, we used FM5-95 (all

obtained from Molecular Probes). All images were background-corrected, setting the signal

outside the hyphae to 0 to obtain a sufficiently dark background. These corrections were

made using Adobe Photoshop CS4. Cryo-scanning electron microscopy (cryo-SEM) and

transmission electron microscopy (TEM) were performed as described [62]. For quantification of

fluorescence intensity of YlmD-eGFP and YlmE-eGFP, the arbitrary average intensity values of

manually selected areas of interest were determined as relative fluorescence intensity. All green

fluorescent images were made with equal exposure time.

Data analysis

For the Kolmogorov–Smirnov (KS) test, the critical value D_{crit} was calculated for the number of

measured spores (n) and significance level (α) of 0.05: $D_{crit} = \frac{\sqrt{-0.5*\ln(\frac{\alpha}{2})}}{\sqrt{n}}$. The difference ($D_n$)

between the observed cumulative distribution function (F_{obs}) and the expected cumulative

distribution function (F_{exp}) was calculated as: $D_n = |F_{exp}(x) - F_{obs}(x)|$. The maximal value was

identified as D_{max}. The null hypothesis that the data comes from the specified distribution was

rejected if $D_{max} > D_{crit.}$ The Mann-Whitney U test was done using the SPSS software release 24.0.

Computer analysis

DNA   and   protein   database   searches   were   completed   by   using   StrepDB

(http://strepdb.streptomyces.org.uk/). Phylogenetic analysis was done using the STRING engine

at EMBL (www.string.EMBL).

DATA AVAILABILITY

All strains, materials and data will be made available upon request.

ACKNOWLEDGEMENTS

430 This work was supported by the Netherlands Organization for Scientific Research (NWO), via

431 VICI grant 10379 to GPvW.

COMPETING FINANCIAL INTERESTS

434 The authors declare no conflict of interests.

AUTHOR CONTRIBUTIONS

437 LZ performed the molecular biology experiments and created the mutants and expression

438 constructs, JW performed the imaging and PAH the phylogenetic analysis. GVW conceived the

439 experiments together with the other authors. All authors wrote and approved the manuscript.

REFERENCES

Bi, E. F. & Lutkenhaus, J. FtsZ ring structure associated with division in *Escherichia coli*. *Nature* **354**, 161-164 (1991).

Goehring, N. W. & Beckwith, J. Diverse paths to midcell: assembly of the bacterial cell division machinery. *Curr Biol* **15**, R514-526 (2005).

Adams, D. W. & Errington, J. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. *Nat Rev Microbiol* **7**, 642-653 (2009).

Hale, C. A. & de Boer, P. A. J. Direct Binding of FtsZ to ZipA, an essential component of the septal ring structure that mediates cell division in *E. coli. Cell* **88**, 175-185 (1997).

RayChaudhuri, D. ZipA is a MAP-Tau homolog and is essential for structural integrity of the cytokinetic FtsZ ring during bacterial cell division. *EMBO J* **18**, 2372-2383 (1999).

Sekine, H., Mimura, J., Yamamoto, M. & Fujii-Kuriyama, Y. Unique and overlapping transcriptional roles of arylhydrocarbon receptor nuclear translocator (Arnt) and Arnt2 in xenobiotic and hypoxic responses. *J Biol Chem* **281**, 37507-37516 (2006).

Gueiros-Filho, F. J. & Losick, R. A widely conserved bacterial cell division protein that promotes assembly of the tubulin-like protein FtsZ. *Genes Dev* **16**, 2544-2556 (2002).

Hamoen, L. W., Meile, J. C., de Jong, W., Noirot, P. & Errington, J. SepF, a novel FtsZ-interacting protein required for a late step in cell division. *Mol Microbiol* **59**, 989-999 (2006).

Ishikawa, S., Kawai, Y., Hiramatsu, K., Kuwano, M. & Ogasawara, N. A new FtsZ-interacting protein, YlmF, complements the activity of FtsA during progression of cell division in *Bacillus subtilis*. *Mol Microbiol* **60**, 1364-1380 (2006).

Romberg, L. & Levin, P. A. Assembly dynamics of the bacterial cell division protein FTSZ: poised at the edge of stability. *Annu Rev Microbiol* **57**, 125-154 (2003).

Barka, E. A. *et al.* Taxonomy, physiology, and natural products of the *Actinobacteria*. *Microbiol Mol Biol Rev* **80**, 1-43 (2016).

Hopwood, D. A. *Streptomyces in nature and medicine: the antibiotic makers*. (Oxford University Press, 2007).

Wildermuth, H. & Hopwood, D. Septation during sporulation in *Streptomyces coelicolor. J Gen Microbiol* **60**, 51-59 (1970).

Claessen, D., Rozen, D. E., Kuipers, O. P., Sogaard-Andersen, L. & van Wezel, G. P. Bacterial solutions to multicellularity: a tale of biofilms, filaments and fruiting bodies. *Nat Rev Microbiol* **12**, 115-124 (2014).

Flärdh, K. & Buttner, M. J. *Streptomyces* morphogenetics: dissecting differentiation in a filamentous bacterium. *Nat Rev Microbiol* **7**, 36-49 (2009).

McCormick, J. R. Cell division is dispensable but not irrelevant in *Streptomyces*. *Curr Opin Microbiol* **12**, 689-698 (2009).

Celler, K., Koning, R. I., Willemse, J., Koster, A. J. & van Wezel, G. P. Cross-membranes orchestrate compartmentalization and morphogenesis in *Streptomyces*. *Nat Comms* **7**, 11836 (2016).

Yague, P. *et al.* Subcompartmentalization by cross-membranes during early growth of *Streptomyces* hyphae. *Nat Comms* **7**, 12467 (2016).

19   Holmes, N. A. *et al.* Coiled-coil protein Scy is a key component of a multiprotein
assembly controlling polarized growth in *Streptomyces*. *Proc Natl Acad Sci U S A* **110**,
E397-406 (2013).
20   Celler, K., Koning, R. I., Koster, A. J. & van Wezel, G. P. Multidimensional view of the
bacterial cytoskeleton. *J Bacteriol* **195**, 1627-1636 (2013).
21   Jakimowicz, D. & van Wezel, G. P. Cell division and DNA segregation in *Streptomyces*:
how to build a septum in the middle of nowhere? *Mol Microbiol* **85**, 393-404 (2012).
22   Schwedock, J., McCormick, J. R., Angert, E. R., Nodwell, J. R. & Losick, R. Assembly of the
cell division protein FtsZ into ladder like structures in the aerial hyphae of *Streptomyces*
coelicolor. *Mol Microbiol* **25**, 847-858 (1997).
23   McCormick, J. R., Su, E. P., Driks, a. & Losick, R. Growth and viability of *Streptomyces*
coelicolor mutant for the cell division gene *ftsZ*. *Mol Microbiol* **14**, 243-254 (1994).
24   McCormick, J. R. & Losick, R. Cell division gene *ftsQ* is required for efficient sporulation
but not growth and viability in *Streptomyces coelicolor* A3(2). *J Bacteriol* **178**, 5295-5301
(1996).
25   Traag, B. A. & van Wezel, G. P. The SsgA-like proteins in actinomycetes: small proteins up
to a big task. *Antonie Van Leeuwenhoek* **94**, 85-97 (2008).
26   Girard, G. *et al.* A novel taxonomic marker that discriminates between morphologically
complex actinomycetes. *Open Biol* **3**, 130073 (2013).
27   Willemse, J., Borst, J. W., de Waal, E., Bisseling, T. & van Wezel, G. P. Positive control of
cell division: FtsZ is recruited by SsgB during sporulation of *Streptomyces*. *Genes Dev* **25**,
89-99 (2011).
28   Keijser, B. J., Noens, E. E., Kraal, B., Koerten, H. K. & van Wezel, G. P. The *Streptomyces*
coelicolor ssgB gene is required for early stages of sporulation. *FEMS Microbiol Lett* **225**,
59-67 (2003).
29   van Wezel, G. P. *et al. ssgA* is essential for sporulation of *Streptomyces coelicolor* A3(2)
and affects hyphal development by stimulating septum formation. *J Bacteriol* **182**, 5653-
5662 (2000).
30   Zhang, L., Willemse, J., Claessen, D. & van Wezel, G. P. SepG coordinates sporulation-
specific cell division and nucleoid organization in *Streptomyces coelicolor*. *Open Biol* **6**,
150164 (2016).
31   Mingorance, J., Tamames, J. & Vicente, M. Genomic channeling in bacterial cell division. *J*
Mol Recognit **17**, 481-487 (2004).
32   Santos-Beneit, F., Gu, J. Y., Stimming, U. & Errington, J. *ylmD* and *ylmE* genes are
dispensable for growth, cross-wall formation and sporulation in *Streptomyces*
venezuelae. *Heliyon* **3**, e00459 (2017).
33   Romero, D. A. *et al.* A comparison of key aspects of gene regulation in *Streptomyces*
coelicolor and *Escherichia coli* using nucleotide-resolution transcription maps produced
in parallel by global and differential RNA sequencing. *Mol Microbiol* **94**, 963-987 (2014).
34   Swiatek, M. A. *et al.* The ROK family regulator Rok7B7 pleiotropically affects xylose
utilization, carbon catabolite repression, and antibiotic production in *Streptomyces*
coelicolor. *J Bacteriol* **195**, 1236-1248 (2013).
35   Szklarczyk, D. *et al.* The STRING database in 2011: functional interaction networks of
proteins, globally integrated and scored. *Nucleic Acids Res* **39**, D561-D568 (2011).

36 Claus, H. Laccases and their occurrence in prokaryotes. *Arch Microbiol* **179**, 145-150
(2003).

37 Eswaramoorthy, S. *et al.* Structure of a yeast hypothetical protein selected by a
structural genomics approach. *Acta Crystallogr D Biol Crystallogr* **59**, 127-135 (2002).

38 Percudani, R. & Peracchi, A. A genomic overview of pyridoxal-phosphate-dependent
enzymes. *EMBO R* **4**, 850-854 (2003).

39 Tassoni, R., van der Aart, L. T., Ubbink, M., van Wezel, G. P. & Pannu, N. S. Structural and
functional characterization of the alanine racemase from *Streptomyces coelicolor* A3(2).
Biochem Biophys Res Comm **483**, 122-128 (2017).

40 van Wezel, G. P., White, J., Hoogvliet, G. & Bibb, M. J. Application of *redD*, the
transcriptional activator gene of the undecylprodigiosin biosynthetic pathway, as a
reporter for transcriptional activity in *Streptomyces coelicolor* A3(2) and *Streptomyces*
lividans. *J Mol Microbiol Biotechnol* **2**, 551-556. (2000).

41 Zacchetti, B. *et al.* Aggregation of germlings is a major contributing factor towards
mycelial heterogeneity of *Streptomyces. Sci Rep* **6**, 27045 (2016).

42 Grantcharova, N., Lustig, U. & Flärdh, K. Dynamics of FtsZ assembly during sporulation in
Streptomyces coelicolor A3(2). *J Bacteriol* **187**, 3227-3237 (2005).

43 Yamaguchi, Y., Park, J. H. & Inouye, M. Toxin-antitoxin systems in bacteria and archaea.
Ann Rev Genet **45**, 61-79 (2011).

44 Ainsa, J. A., Bird, N., Ryding, N. J., Findlay, K. C. & Chater, K. F. The complex *whiJ* locus
mediates environmentally sensitive repression of development of *Streptomyces*
coelicolor A3(2). *Antonie Van Leeuwenhoek* **98**, 225-236 (2010).

45 Girard, G. *et al.* A novel taxonomic marker that discriminates between morphologically
complex actinomycetes. *Open Biol* **3**, 130073 (2013).

46 van Dissel, D., Claessen, D. & Van Wezel, G. P. Morphogenesis of *Streptomyces* in
submerged cultures. *Adv Appl Microbiol* **89**, 1-45 (2014).

47 Thompson, J. D., Higgins, D. G. & Gibson, T. J. Clustal-W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific
gap penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673-4680 (1994).

48 Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Mol Biol Evol* **4**, 406-425 (1987).

49 Tamura, K., Dudley, J., Nei, M. & Kumar, S. MEGA4: Molecular Evolutionary Genetics
Analysis (MEGA) software version 4.0. *Mol Biol Evol* **24**, 1596-1599 (2007).

50 MacNeil, D. J. *et al.* Analysis of *Streptomyces avermitilis* genes required for avermectin
biosynthesis utilizing a novel integration vector. *Gene* **111**, 61-68 (1992).

51 Bentley, S. D. *et al.* Complete genome sequence of the model actinomycete
Streptomyces coelicolor A3(2). *Nature* **417**, 141-147 (2002).

52 Cruz-Morales, P. *et al.* The genome sequence of *Streptomyces lividans* 66 reveals a novel
tRNA-dependent peptide biosynthetic system within a metal-related genomic island.
Genome Biol Evol **5**, 1165-1175 (2013).

53 Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F. & Hopwood, D. A. *Practical*
Streptomyces genetics. (John Innes Foundation, 2000).

54 Colson, S. *et al.* Conserved *cis*-acting elements upstream of genes composing the
chitinolytic system of streptomycetes are DasR-responsive elements. *J Mol Microbiol*
Biotechnol **12**, 60-66 (2007).
55 Vara, J., Lewandowska-Skarbek, M., Wang, Y. G., Donadio, S. & Hutchinson, C. R. Cloning
of genes governing the deoxysugar portion of the erythromycin biosynthesis pathway in
Saccharopolyspora erythraea (*Streptomyces erythreus*). *J Bacteriol* **171**, 5872-5881
(1989).
56 Świątek, M. A., Tenconi, E., Rigali, S. & van Wezel, G. P. Functional analysis of the N-
acetylglucosamine metabolic genes of *Streptomyces coelicolor* and role in control of
development and antibiotic production. *J Bacteriol* **194**, 1136-1144 (2012).
57 Fedoryshyn, M., Welle, E., Bechthold, A. & Luzhetskyy, A. Functional expression of the
Cre recombinase in actinomycetes. *App Microbiol Biotechnol* **78**, 1065-1070 (2008).
58 Khodakaramian, G. *et al.* Expression of Cre recombinase during transient phage infection
permits efficient marker removal in *Streptomyces*. *Nucleic Acids Res* **34**, e20 (2006).
59 Sun, J., Kelemen, G. H., Fernandez-Abalos, J. M. & Bibb, M. J. Green fluorescent protein
as a reporter for spatial and temporal gene expression in *Streptomyces coelicolor* A3(2).
Microbiology **145**, 2221-2227 (1999).
60 Vijgenboom, E. *et al.* Three *tuf*-like genes in the kirromycin producer *Streptomyces*
ramocissimus. *Microbiology* **140**, 983-998 (1994).
61 Willemse, J. & van Wezel, G. P. Imaging of *Streptomyces coelicolor* A3(2) with Reduced
Autofluorescence Reveals a Novel Stage of FtsZ Localization. *PLoS ONE* **4**, e4242 (2009).
62 Piette, A. *et al.* From dormant to germinating spores of *Streptomyces coelicolor* A3(2):
new perspectives from the *crp* null mutant. *J Proteome Res* **4**, 1699-1708 (2005).
592
593

FIGURE LEGENDS

Figure 1. Phylogenetic analysis of YlmE in actinobacteria and its genetic linkage to *ylmD*. A phylogenetic tree is shown of YlmE in actinobacteria (left) and genetic linkage of *ylmE* (grey) to *ftsZ* (black) and *ylmD* (white) (right).

Figure 2. Phenotype of the *S. coelicolor ylmD* and *ylmE* mutants. (A) sporulation of *S. coelicolor* M145 and its *ylmD*, *ylmE* and *ylmDE* mutants on SFM agar after 5 days incubation. Note the lack of grey pigmentation of the *ylmE* mutant, indicative of a sporulation defect. (B) phase-contrast micrographs of impression prints of the strains shown in (A) as well as the complemented mutants. Arrows point at irregularly shaped spores in the *ylmD* mutant. The sporulation defect of the *ylmD* mutant could be complemented by introduction of wild-type *ylmD* and *ylmD-eGFP*, while complementation of the *ylmE* mutant by wild-type *ylmE* or *ylmE-egfp* restored sporulation, although irregularly sized spores were often produced. Full complementation of *ylmE* mutants was obtained by introduction of pKR8, which contains part of the *S. coelicolor dcw* cluster. Bar, 5 µm. (C) Size distributions of wild-type spores and those of *ylmD, ylmE* and *ylmDE* mutants presented as boxplots. Data are presented as median and interquartile range in boxplots, with whiskers spread to the maximal and minimal values. The numbers (n) of spores measured for each strain are indicated. Assessment of normality of spore length data was performed by a Kolgomorov-Smirnov test, the results showing that spore length data for all mutants followed non-normal distribution ($D_{max} > D_{crit}$). The shown two-tailed P values between each mutant and the parental strain were calculated using a Mann-Whitney U test. The two-tailed P values were

far below 0.05, which shows that the sizes of the mutant spores deviated significantly from those of wild-type spores.

Figure 3. Cryo-scanning electron micrographs of aerial hyphae of *S. coelicolor* M145 and its *ylmD* and *ylmE* mutants. The parental strain produced wild-type spores, the *ylmD* mutant produced abundant but often irregular spores, while the *ylmE* null mutant produced occasional spores with highly irregular sizes. Cultures were grown on SFM agar plates for 5 days at 30^0C. Bars: top row, 1 µm; bottom row, 5 µm.

Figure 4. Transmission electron micrographs of spores of *S. coelicolor* M145 and its *ylmD* and *ylmE* mutants. Wild-type spores (M145) show regular sizes and appearance (A). In contrast, spores of the *ylmD* (B) and *ylmE* (C) null mutants have an irregular appearance. Note the lighter appearance of the spore wall in the *ylmD* null mutant and the lack of the typical thick spore wall in *ylmE* mutants. Cultures were grown on SFM agar plates for 5 days at 30^0C. Bar, 500 nm.

Figure 5. Fluorescence microscopy of cell wall, DNA and membranes.

(A) Fluorescent micrographs of hyphae stained for cell-wall synthesis (FITC-WGA or Oregon-WGA) or membranes (FM5-95). An overlay of these images is presented in the third column, and the corresponding light image in the last column. Bar, 5 µm. (B) Fluorescence micrographs showing DNA and cell-wall distribution in the complemented *ylmD* and *ylmE* mutants. While ladders of septa were formed in both strains, indicative that sporulation was restored to the

mutants, in particular the complemented *ylmE* mutant formed imperfect septa. Cultures were grown on SFM agar plates for 5 days at 30^{0}C. Bar, 10 µm.

Figure 6. Localization of YlmD-GFP and YlmE-GFP.

(A) Sporogenic aerial hyphae of *S. coelicolor* M145 at different stages of development were imaged by fluorescence microscopy visualizing YlmD-eGFP and YlmE-eGFP. Stages were: vegetative growth, early aerial development, late aerial development, early sporulation and mature spores. During spore maturation, YlmD and YlmE had a 'patchy' localization, suggesting that at this stage YlmD and YlmE may co-localize with the cell-wall synthetic machinery. Bar, 5 µm. (B) Relative fluorescence intensity of YlmD-eGFP and YlmE-eGFP during *Streptomyces* development. VEG, vegetative growth; EA, early aerial growth; LA, late aerial growth; ES, early sporulation; MS, mature spores.

Figure 7. Localization of FtsZ-eGFP in *S. coelicolor* M145 and its *ylmD* and *ylmE* mutants. FtsZ-eGFP formed typical ladders in wild-type cells (M145). In contrast, YlmE is required for the formation of ladders of FtsZ, while the absence of YlmD caused irregular spacing between the septa. Cultures were grown on SFM agar plates for 5 days at 30^{0}C. Bar, 5 µm.

tr|C9YVD1|C9YVD1_STRSW Putative uncharacterized protein Streptomyces scabies
tr|Q82AD3|Q82AD3_STRAW Uncharacterized protein Streptomyces avermitilis
tr|Q9S2X1|Q9S2X1_STRCO Putative uncharacterized protein SCO2080 Streptomyces coelicolor
tr|G0PZI7|G0PZI7_STRGR Putative uncharacterized protein Streptomyces griseus
tr|E4N9R1|E4N9R1_KITSK Putative uncharacterized protein Kitasatospora setae
tr|E2S959|E2S959_9ACTO YggS family pyridoxal phosphate enzyme Aeromicrobium marinum
tr|D2B842|D2B842_STRRD Alanine racemase domain protein Streptosporangium roseum
tr|A3TJT1|A3TJT1_9MICO Putative uncharacterized protein Janibacter sp. HTCC2649
tr|Q0RNN5|Q0RNN5_FRAAA Putative enzyme with PLP-binding domain Frankia alni
tr|D3PZX2|D3PZX2_STANL Alanine racemase domain protein Stackebrandtia nassauensis
tr|Q6AE68|Q6AE68_LEIXX Putative uncharacterized protein Leifsonia xyli
tr|B0RIJ7|B0RIJ7_CLAMS Putative uncharacterized protein Clavibacter michiganensis subsp. sepedonicus
tr|D1A7U2|D1A7U2_THECD Alanine racemase domain protein Thermomonospora curvata
tr|A0LTM7|A0LTM7_ACIC1 Putative uncharacterized protein Acidothermus cellulolyticus
tr|C7QGS7|C7QGS7_CATAD Alanine racemase domain protein Catenulispora acidiphila
tr|J7L1G1|J7L1G1_NOCAA Alanine racemase N-terminal domain protein Nocardiopsis alba
tr|D2Q0G4|D2Q0G4_KRIFD Alanine racemase domain protein Kribbella flavida
tr|B8HGX5|B8HGX5_ARTCA Alanine racemase domain protein Arthrobacter chlorophenolicus
tr|B2GJP3|B2GJP3_KOCRD Putative uncharacterized protein Kocuria rhizophila
tr|C5CA25|C5CA25_MICLC Putative uncharacterized protein Micrococcus luteus
tr|A6WCW9|A6WCW9_KINRD Alanine racemase domain protein Kineococcus radiotolerans
tr|I0H160|I0H160_ACTM4 Uncharacterized protein Actinoplanes missouriensis
tr|A4X9R2|A4X9R2_SALTO Alanine racemase domain protein Salinispora tropica
tr|D9T677|D9T677_MICAI Alanine racemase domain protein Micromonospora aurantiaca
tr|F4FHR0|F4FHR0_VERMA Alanine racemase domain-containing protein Verrucosispora maris (strain AB-18-032)
tr|D8I163|D8I163_AMYMU Pyridoxal-5-phosphate dependent enzyme class III Amycolatopsis mediterranei
tr|G4J3W6|G4J3W6_9PSEU Putative uncharacterized protein Saccharomonospora paurometabolica
tr|D6ZC34|D6ZC34_SEGRD Alanine racemase domain protein Segniliparus rotundus
tr|F9VQ15|F9VQ15_9ACTO Putative uncharacterized protein Gordonia alkanivorans
tr|E4WIN5|E4WIN5_RHOE1 Putative alanine racemase Rhodococcus equi
tr|D5USH0|D5USH0_TSUPD Alanine racemase domain protein Tsukamurella paurometabola
tr|A0R009|A0R009_MYCS2 Putative uncharacterized protein Mycobacterium smegmatis
sp|P67083|Y2148_MYCTU UPF0001 protein Rv2148c/MT2207 Mycobacterium tuberculosis

0.05

ftsZ *ylmD* *ylmE*

M145	**Δ*ylmD***	**Δ*ylmE***

M145   ΔylmD   ΔylmE

A

	Cell wall	Membrane	Merge	Light
M145				
Δ*ylmD*				
Δ*ylmE*				

B

	Cell wall	Membrane	Merge	Light
Δ*ylmD* + *ylmD*				
Δ*ylmE* + *ylmE*				

A

	YlmD-eGFP	Light	YlmE-eGFP	Light

Vegetative hyphae

Early aerial hyphae

Late aerial hyphae

Early spore chains

Mature spores

B

	FtsZ-eGFP	Light
M145		
ΔylmD		
ΔylmE		