1 **Using Biogeography to Assess Key Adaptation Strength in Two Bird Families**
2
3

4   Abdel H. Halloway[1], Christopher J. Whelan[1,2], Çağan H. Şekercioğlu[3,4], and Joel S. Brown[1,5]
5
6
7
8   [1]Department of Biological Sciences, University of Illinois at Chicago
9   845 W. Taylor St. (M/C 066) Chicago, IL 60607
10
11   [2]Department of Cancer Physiology, Moffitt Cancer Center
12   SRB-4, 12902 USF Magnolia Drive Tampa, FL 33612
13
14   [3]Department of Biology, University of Utah
15   257 South 1400 East, Rm. 201, Salt Lake City, UT 84112-0840
16
17   [4]College of Sciences, Koç University, Rumelifeneri, Istanbul 34450, Turkey
18   [5]Integrated Mathematical Oncology, Moffitt Cancer Center
19   SRB-4, 12902 USF Magnolia Drive Tampa, FL 33612
20
<u>Corresponding Author</u>
Abdel Halloway
Department of Biological Sciences, University of Illinois at Chicago
845 W. Taylor St. (M/C 066) Chicago, IL 60607
ahallo2@uic.edu

Abstract

Adaptations can be thought of as evolutionary technologies which allow an organism to

exploit environments. Among convergent taxa, adaptations may be largely equivalent with the

taxa operating in a similar set of environmental conditions, divergent with the taxa operating in

different sets of environmental conditions, or superior with one taxon operating within an

extended range of environmental conditions than the other. With this framework in mind, we

sought to characterize the adaptations of two convergent nectarivorous bird families, the New

World hummingbirds (Trochilidae) and Old World sunbirds (Nectariniidae), by comparing their

biogeography. Looking at their elevational and latitudinal gradients, hummingbirds not only

extend into but also maintain species richness in more extreme environments. We suspect that

hummingbirds have a superior key adaptation that sunbirds lack, namely a musculoskeletal

architecture that allows for hovering. Through biogeographic comparisons, we have been able to

assess and understand adaptations as evolutionary technologies among two convergent bird

families, a process that should work for most taxa.

Keywords

Biogeography, Key Adaptations, Sunbird, Hummingbird, Convergence, Species Gradients

Introduction

Convergent evolution provides startling examples of how natural selection shapes the traits of species to optimize fitness (1). Species' morphologies, physiologies, and behaviors become fine-tuned to their shared ecologies somewhat irrespective of evolutionary history (2). Besides the species level, convergence can happen between higher taxa (e.g., families and orders). One striking example is the convergence between the New World nectarivorous hummingbirds (order Apodiformes, family Trochilidae) and the Old World passerine nectarivores including the Hawaiian honeycreepers (order Passeriformes, family Fringillidae), Australian honeyeaters (order Passeriformes, family Meliphagidae) and the Asian and African sunbirds (order Passeriformes, family Nectariniidae). Some or all members of these families show convergent adaptations for nectarivory, particularly elongated bills and extensile tongues. Their remarkable convergence, especially between hummingbirds and sunbirds, makes them ripe for analysis of adaptations as evolutionary technologies.

Adaptations can be thought of as evolutionary technologies that allow an organism to operate within an environment. Among evolutionary convergent taxa, adaptations might be equivalent leading to similar fitness in similar environmental conditions, e.g. the convergent snake constrictor families Boidae and Pythonidae (Fig. 1a) (3). In this case, both clades operate under similar fundamental niches. Differences in adaptations, though, can change the fundamental niches of the convergent clades and open new ecological opportunities (4). Such adaptations are known as as key adaptations. Key adaptations may be divergent evolutionary technologies with the taxa occupying different fundamental niches, e.g. the ankle bones of the grandorder Euarchonta (four orders of mammals including primates) that promote arboreal living (Fig. 1b). Hummingbirds and hawkmoths (order Lepidoptera, family Sphingidae) are instructive

examples of convergent families displaying divergent evolutionary technologies. As vertebrates

and invertebrates respectively, they strongly differ in virtually all aspects of ontogeny,

morphology, and physiology. Yet, their coexistence in the New World suggests that one set of

evolutionary technologies is not superior to the other under all circumstances leading to a

partitioning of environmental conditions (5).

Key adaptations may also be superior evolutionary technologies allowing a taxon to

expand beyond its original range of environments and have a greater fundamental niche (Fig.

1c). Examples of superior key adaptations include the retractable necks among turtles of the

suborder Cryptodira, which protect them from predation, and the infrared-sensing pits among

vipers of the subfamily Crotalinae, which allows them to "see" mammals at night. When a key

adaptation creates a superior evolutionary technology, we might see the replacement of another

clade – typically but not always the ancestral clade – by the new one (6; 7; 8). Or the ancestral

clade may persist where the derived clade has yet to colonize as is the case with the Pleurodira

turtles of the Southern hemisphere. Since hummingbirds and sunbirds do not occur sympatrically

and therefore do not interact, it is hard to discern whether they represent equivalent, divergent, or

superior evolutionary technologies. We hypothesize that hummingbirds represent the latter

compared to sunbirds and other nectarivorous passerines with hummingbirds possessing a

superior key adaptation making them an example of "progressive evolution" (9; 6).

Hummingbirds display a stronger mutualistic co-adaptation with flowers compared to

sunbirds (10; 11). All hummingbirds feed almost exclusively on nectar, only supplementing

protein intake by eating small insects (12). As such, they have evolved distinct anatomical and

morphological features suited to nectar foraging. In addition to an elongated bill and extensile

tongue, the hummingbird's tongue acts as a micro-pump for reaching and gathering nectar (13;

14). They possess large breast muscles (30% of body weight), skeletal architecture common to

Apodiformes, and dense erythrocyte counts for delivering a steady supply of oxygen to feed

extremely active muscles (10). Specialized wings allow hummingbirds to hover and fly

backwards. Sunbirds, on the other hand, are not as tightly adapted to nectar feeding with many

species supplementing their diet with insects, seeds, fruit, and flower heads, and others being

largely insectivorous (15). They also show large variation in bill and flight morphology with the

flowerpeckers and the *Hedydipna* and *Hypogramma* sunbirds having broad, flat tongues. In

addition, all sunbirds lack the musculoskeletal architecture to hover and must perch to feed (11).

These anatomical differences along with differences in species richness (364 hummingbird

species vs. 147 sunbird species) suggest that hummingbirds have a superior key adaptation not

found in sunbirds (16). Furthermore, the geographic isolation between the taxa allowed for

independent diversification, making them ideal convergent clades to assess adaptations.

Testing for a key adaptation requires two things: elucidating a mechanistic hypothesis for

its ecological and functional role and a comparison between clades (7). When comparing clades,

species richness and diversification rates have typically been used (4). Besides these properties,

we also surmise that a greater geographical range would be seen with a superior key adaptation.

By increasing net fitness overall, a superior key adaptation should increase the fitness of a clade

at the margins of its range (17); therefore, clades with a superior key adaptation will be better

able to handle abiotic stress and live under harsher climatic regimes. Looking at the convergent

mice genera *Peromyscus* and *Apodemus*, *Peromyscus* inhabits colder, more arid, and higher

habitats compared to *Apodemus* due to its more efficient and widely used torpor state (18; 19;

20). Between species richness and biogeography, comparing the latter may be more useful to

assess adaptations as evolutionary technologies since biogeographical extent explicitly depends

upon a clades' overall net fitness and directly tests its ecological role.

To characterize the adaptations of hummingbirds and sunbirds, we compared their

biogeography by analyzing each family's latitudinal and elevational distribution. We demonstrate

that hummingbirds as a clade inhabit more extreme latitudes and maintain their species richness

at higher elevations. We hypothesize these differences in biogeography reveal a superior key

adaptation present in hummingbirds but absent in sunbirds. We speculate that the key adaptation

may be either the unique tongue of the hummingbirds or the unique wing architecture that allows

for hovering. We further speculate on the role adaptations as evolutionary technologies play in

influencing an organism's ability to exploit the environment.

<u>Materials and Methods</u>

To assess the adaptations of hummingbirds and sunbirds, we gathered each family's

latitudinal and elevational gradient of species richness. These gradients are robust geographic

patterns that generally show species richness declining towards higher altitudes and more

extreme latitudes (21; 22; 23; 24; 25; 26; 27). Numerous environmental properties change along

both gradients. Aridity declines significantly around 30 to 40 degrees latitude; a thinner

atmosphere and more variable daily temperatures occur with increased elevation; and more

variable seasonal temperatures, less productivity, and colder temperatures occur with both. Taxa

with superior evolutionary technologies should be better able to deal with these challenges (28).

To compare the biogeography of hummingbirds and sunbirds, we gathered the latitudinal

and elevational range of all species from each family. Elevational ranges came from a global bird

ecology database covering all the bird species of the world (29) while latitudinal ranges of the

families were taken from shapefiles downloaded from BirdLife International and NatureServe

with data extracted using R packages "sp", "raster", "rasterVis", "maptools", and "rgeos" (30).

All latitudinal extremes located in the Southern hemisphere were converted to negative values,

and latitudinal maxima and minima were rounded up and down to the nearest integer

respectively. For example, the hummingbird species *Amazilia amabilis* which ranges from

14.17N to 3.98S would have its range taken as 15 to -4. An additional measure of distance from

the equator, hereafter referred to as "polewardness", was created. If a species' range crossed the

equator, then the poleward range was taken to be from 0 to the maximum distance from the

equator. For *A. amabilis*, its poleward range would be 0 to 15 degrees. The poleward range of an

only Northern or Southern hemispheric species would simply be the absolute value of its

latitudinal range.

With ranges in hand, we compared the families in two ways. First, we compared several

empirical cumulative distribution functions (ECDFs) based upon the three geographical

properties (elevation, latitude, and polewardness) for each family. Each ECDF started from sea

level, the South Pole, and the equator and traced to higher altitudes, northward, and more

extreme poles. For each geographical property, three ECDFs were created with a species'

presence based on the minimum, the midpoint, and the maximum of its range. Since species

which cross the equator are not necessarily symmetric about it, the midpoint of a species

poleward range may not accurately reflect its bias towards the equator or poles. Therefore, we

created another measurement of species presence for polewardness, its expected value (see SI).

This led to ten different ECDFs for each family: minimum, maximum, and midpoint for

elevation, latitude, and polewardness and the additional measure of expected polewardness. Each

type of ECDF was then compared between families using the Kolmogorov-Smirnov and

Anderson-Darling minimum difference estimation (MDE) tests with the assumption that the

hummingbird ECDF is less than the sunbird ECDF (one-tailed tests).

The ECDF analysis tells us whether the distributions differ, not necessarily how they

differ. Therefore, we additionally sought to characterize each family's distribution by measuring

changes in species richness with polewardness and elevation. To do so, we first counted the

number of species in poleward and elevation intervals of 5 degrees and 500 meters for each

family. If the edge of a species' range was at the cutoff point of the interval, it would be

considered present in the lower interval but not in the upper interval due to previous rounding. In

the example with *A. amabilis*, this would mean that the species is counted in the 10 to 15 degree

interval but not the 15 to 20 degree interval. The frequency data were then normalized such that

the interval with the highest number of species became 1 to remove any effect of total species

richness. This gave us four sets of data based on a 2x2 factorial: sunbird and hummingbird

polewardness and elevation. A logistic curve (eq. 1) was then fitted onto each of the four sets of

data – the normalized species richness, S_N, per interval vs. the midpoint of each interval – with

variables a and b determining position and steepness of the curve respectively.

$$S_N = \frac{1}{1 + ae^{bx}} \tag{1}$$

We then found the inflection point and the two points of the maximum magnitude of

curvature (MMC points) for each curve. Inflection points indicate how well each family

maintains species richness while MMC points give us the start and end of the decline in species

richness. The functions and their key points characterize the shape of each family's gradient.

<u>Results</u>

Broadly, our results show that hummingbirds extend farther poleward and higher in

elevation than do sunbirds. Hummingbirds extend from 62 degrees north to 56 degrees south and

up to 5000 m in elevation (SI Table 1,2; Fig. 2,3). Sunbirds, on the other hand, extend only from

36 degrees north to 40 degrees south and up to 4880 m in elevation (SI Table 1,2; Fig. 2,3). Both

families show the same general pattern of an initial increase in species richness followed by a

decline moving poleward and to higher altitudes (Fig. 4a, b). In addition, hummingbirds maintain

their species richness at higher elevations and more extreme latitudes than sunbirds. ECDF

results confirm this difference in biogeography between hummingbirds and sunbirds with

elevation constituting the greatest difference (SI Table 3, SI Fig. 1).

Both hummingbirds and sunbirds reach approximately the same maximum elevation,

around 5000m (SI Table 1, SI Fig. 1a). Even though both hummingbirds and sunbirds extend to

roughly the same elevation, hummingbirds have a higher normalized species richness at higher

elevations compared to sunbirds. The inflection point for sunbirds occurs at 2087m and

hummingbirds at 2533m (SI Table 4, Fig. 4c). Sunbirds and hummingbird species richness

values both start to decline around the same elevation –1764 and 1898m respectively – but

sunbirds plateau at a lower elevation compared to hummingbirds –2410m vs. 3458m respectively

– indicating a more gradual decline in the normalized species richness of hummingbirds (SI

Table 4, Fig. 4c).

Regarding latitude, hummingbirds occur farther from the equator than do sunbirds, 60-65

degrees vs. 35-40 degrees respectively (SI Table 2, SI Fig. 1b). Also, hummingbird normalized

species richness is at its greatest divergence from sunbird normalized species richness at mid-

latitudinal ranges. The hummingbirds' inflection point is 22.14 degrees latitude versus 18.92

degrees for sunbirds (SI Table 4, Fig. 4d). Hummingbirds also begin their declines further from

the equator than do sunbirds –14.99 and 9.44 degrees respectively. Both plateau around the same

latitude – 29.29 vs. 28.39 degrees respectively (SI Table 4, Fig. 4d).

Discussion

Sunbirds and hummingbirds are two convergent nectarivorous bird families with different

evolutionary technologies. While hummingbirds are extremely specialized to nectar feeding,

sunbirds vary, ranging from the highly specialized sugarbirds to the passerine-like flowerpeckers

(10; 11). These differences in evolutionary technologies should reflect differences in the families'

distribution and biogeography. As one moves higher in elevation and towards the poles,

hummingbirds maintain their species richness more than sunbirds. Though extending to roughly

the same elevational maximum, normalized hummingbird species richness declines at a much

slower rate than sunbirds. The same is true for latitude; in addition, hummingbirds extend into

more extreme latitudes (farther north and south) than sunbirds. Clear from our results is that

hummingbirds have a greater biogeographical extent than sunbirds, likely reflecting a superior

key adaptation.

One potential hypothesis for the biogeographical differences of hummingbirds and

sunbirds could be dispersal limitation. Firstly, there is a lack of suitable land below $40°$ S

Secondly, Old World mountain ranges may form a barrier to sunbird dispersal as they primarily

run along the east-west axis in contrast to New World mountain ranges which primarily run

along a north-south axis. We reject this hypothesis on the grounds that hummingbirds are

frequently found in montane habitats. Not only do hummingbirds maintain species richness at

higher elevations as our study showed, they have higher species richness in the mountains of

western North and South America compared to the flat-lying eastern regions and frequently

undertake migrations in mountainous areas. Even if sunbirds were dispersal limited,

hummingbirds are still more speciose than sunbirds even when taking latitudinal range into

account. Of the 364 species, only 15 hummingbirds are found at latitudes where sunbirds are

absent. Even if we assume that expansion into the northern latitudes led to the evolution of these

15 species, it still only accounts for approximately 4% of hummingbird species. The difference

in species richness between the families cannot solely be due to dispersal limitation. Instead, we

feel that the combined evidence of species richness and biogeography is highly suggestive of one

or more key adaptations in hummingbirds.

Our spatial analyses cannot tell us what the key adaptations are, but we can speculate on

what they may be. Though hummingbirds and highly specialized sunbirds show many

similarities, they do differ in specific areas. Likely, the key adaptation deals with the differences

in their foraging, specifically how they feed and how they fly. With feeding, one possibility for

hummingbirds' key adaptation may be their unique tongues. The tongues of hummingbirds have

recently been shown to act as micropumps, a way of quickly and efficiently gathering nectar

from flowers, in contrast to the previously assumed capillary action (13; 14). This unusual

feeding method may allow hummingbirds to more efficiently gather nectar compared to sunbirds.

Not enough is known about sunbird tongues, however, to see how the two taxa compare in nectar

gathering abilities. Studies indicate that hummingbirds and sunbirds gather nectar at seemingly

comparable rates suggesting that the amount gathered is not the key difference (31; 32; 33; 14

[personal calculation]). If the tongue is the key adaptation, it will be for the fact that

micropumping requires no energy expenditure on the part of hummingbirds, which removes a

cost, while sunbirds apparently intake nectar through suction, a potentially energetically

expensive system (34; 35). More research needs to be done on the tongues of sunbirds to see how

they compare with the tongues of hummingbirds.

Another possibility of the key adaptation that separates hummingbirds and sunbirds is

hummingbirds' ability to hover and fly in all directions (10). Adaptations for hovering include

shortened arm bones, longer hand bones, a relatively fixed V-shaped arm position, a shallow

ball-and-cup joint between the coracoid and sternum, a large sternum with a deep keel onto

which large breast muscles – pectoralis and supracoracoideus – attach, and red blood cells and

hemoglobin adapted for higher-oxygen affinity and carrying capability (36; 37; 39; 38). All these

anatomical features are adaptations to stiff-winged flight and are seen to a lesser extreme within

other bird families of the order Apodiformes (36; 37; 38). What truly differentiates the flight of

hummingbirds is the axial rotation of the humerus and wrist bones during flight (38).

Hummingbirds can create lift on the upstroke – in addition to the downstroke seen in all birds –

due to wing inversion caused by axial rotation of the wrist (39). Wrist flexibility comes from

changes in carpal structure and the deletion of key ligaments and is seen in birds outside of

Apodiformes (40; 38; 41). Additional power for each downstroke and upstroke also comes from

the axial rotation of humerus, driven by the pectoralis, supracoracoideus, and other muscles (42;

39; 38). The humerus can rotate up to 180º due to a unique humeroscapular joint (43; 36). In

hummingbirds, the humeral head (condyle) is placed along the axis of the shaft instead of the

terminal position, a feature unique to them (44; 45). Together, this suite of adaptations allows

hummingbirds to hover effectively when foraging (46).

Other evolutionary technologies may also benefit hummingbirds in secondary ways. For

example, hummingbirds sustain flight more efficiently at higher altitudes, likely due to their

denser erythrocyte count, expanding their fundamental niche to higher elevations (47). We feel

though that hovering remains the likeliest candidate for a hummingbird key adaptation. Many of

the musculoskeletal changes are seen only in Apodiformes with the shifting of the condyle seen

only in Trochilidae. Such efficient hovering is likely an evolutionarily implastic and ancestral

trait that arose only once among Aves. Through this adaptation, hummingbirds have

fundamentally changed the rules of their nectarivory; they exist as a new type of bauplan while

sunbirds are still effectively a derived passerine (6; 48).

We speculate three possible reasons for the evolution of hovering. Firstly, hummingbirds

can exploit the nectar of plants without perches, potentially opening a new resource for them. As

other nectarivorous birds need to perch while feeding, flowers without perches may represent a

relatively abundant and constant resource without competition from other bird species. Evolution

of hovering in this scenario may be a virtuous cycle as hovering is more efficient at high nectar

volumes which occur in the absence of competition (49). Secondly, hummingbirds may be better

able to escape predation due to their unique flying abilities. With the ability to fly in all

directions, hummingbirds may more easily avoid predators, a useful ability especially when

feeding at a flower with blocked sightlines (50). Furthermore, the musculoskeletal changes in the

hummingbirds are shown to make them extraordinarily agile (51). Finally, while hovering is

energetically costly, it is also time efficient (52). Hovering birds spend less time gathering

resources at flowers than birds which rely on perches. This means that hovering becomes more

energetically efficient compared to perching when birds feed within clustered flower patches (53;

54).

There could be many reasons why hummingbirds developed their key adaptation.

Hummingbirds underwent an expansive radiation during the uplift of the Andes beginning

around 10 mya (55). Living in such rapidly changing conditions could have necessitated the

evolution of a more efficient foraging system. As mentioned earlier, greater oxygen capacity is

beneficial to both hovering and living in low oxygen conditions. There is also the possibility that

the rise of the Andes freed up niche space that would have otherwise been taken up by a

competing family like hawkmoths (Sphingidae), a sort of ecological and evolutionary constraint

(5). These factors, along with hummingbirds' evolutionary history, may combine to lead to the

evolution of hovering (46). Furthermore, sunbirds may face their own internal constraints,

genetic or otherwise, preventing them from evolving a key adaptation (56). Whatever the case

may be, our results suggest the evolution of hovering (or some other adaptation) allowed

hummingbirds to more efficiently take advantage of a resource and expand their fundamental

niche.

The real test of evolutionary technologies would come from seeing what happens when

the two clades meet. Deliberately shifting species across the globe would obviously be unethical

but previous or accidental species invasions may offer such a test. For example, European

Lumbricid earthworms have colonized parts of North America that are farther north than their

American counterparts (57). Both sets of earthworms are ecological equivalents and have

convergent features to fill the role of soil turners. The invasive European earthworms, though,

are known to tolerate environmental stress through protective cocoons during times of drought

and high glucose and glycogen content in cells to prevent freezing during winter (58; 59). These

adaptations may have allowed European earthworms to colonize the colder climes of Canada and

expand their range beyond the North American species.

Through biogeographic analysis, we show that hummingbirds inhabit more hostile climes

than sunbirds, likely due to the possession of a superior evolutionary technology. Going forward,

biogeographic comparison between clades may reveal itself to be a powerful tool to reveal

differences in evolutionary technologies and illuminate the interaction between adaptation and

environment.

Acknowledgements

Abdel Halloway wishes to thank the NSF for funding his graduate studies. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant Nos. DGE-0907994 and DGE-1444315. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors(s) and do not necessarily reflect the views of the National Science Foundation.

References

1. Mooney HA, Dunn EL. Convergent evolution of Mediterranean☐climate evergreen

sclerophyll shrubs. Evolution. 1970 Jun;24(2):292-303.

2. Darwin C. On the origin of species. London, UK: John Murray; 1859

3. Simpson GG. History of the fauna of Latin America. American Scientist. 1950 Jul

1;38(3):361-89.

4. Galis, F. Key innovations and radiations. In: Wagner GP (eds.) The character concept in

evolutionary biology. San Diego, California, USA: Academic Press; 2001. p.581-605

5. Halloway A, Whelan CJ, Brown JS. The hummingbird and the hawk-moth: Species

distribution, geographical partitioning, and macrocompetition across the United States.

bioRxiv [Preprint]. 2018 Jan 1. Available from https://doi.org/10.1101/212894

6. Rosenzweig ML, McCord RD. Incumbent replacement: evidence for long-term

evolutionary progress. Paleobiology. 1991;17(3):202-13.

7. Heard SB, Hauser DL. Key evolutionary innovations and their ecological mechanisms.

Historical biology. 1995 Aug 1;10(2):151-73.

8. Silvestro D, Antonelli A, Salamin N, Quental TB. The role of clade competition in the

diversification of North American canids. Proceedings of the National Academy of

Sciences. 2015 Jul 14;112(28):8684-9.

9. Dawkins R. Human chauvinism. Evolution. 1997 Jun 1;51(3):1015-21.

10. Johnsgard PA. The hummingbirds of North America. Smithsonian Institution; 1997.

11. Mann CF, Cheke RA. Sunbirds: a guide to the sunbirds, flowerpeckers, spiderhunters and

sugarbirds of the world. Bloomsbury Publishing; 2001

12. Yanega GM, Rubega MA. Feeding mechanisms: Hummingbird jaw bends to aid insect

capture. Nature. 2004 Apr;428(6983):615.

13. Rico-Guevara A, Rubega MA. The hummingbird tongue is a fluid trap, not a capillary

tube. Proceedings of the National Academy of Sciences. 2011 Jun 7;108(23):9356-60.

14. Rico-Guevara A, Fan TH, Rubega MA. Hummingbird tongues are elastic micropumps.

Proceedings of the Royal Society B: Biological Sciences. 2015 Aug

22;282(1813):20151014.

15. Oyugi JO, Brown JS, Whelan CJ. Foraging behavior and coexistence of two sunbird

species in a Kenyan woodland. Biotropica. 2012 Mar;44(2):262-9.

16. BirdLife International. Bird species distribution maps of the world. [Accessed 2018]

17. Holt RD, Lawton JH, Gaston KJ, Blackburn TM. On the relationship between range size

and local abundance: back to basics. Oikos. 1997 Feb 1;78(1):183-90.

18. Morhardt JE. Body temperatures of white-footed mice (Peromyscus sp.) during daily

torpor. Comparative biochemistry and physiology. 1970 Mar 15;33(2):423-39.

19. Walton JB, Andrews JF. Torpor induced by food deprivation in the wood mouse

Apodemus sylvaticus. Journal of Zoology. 1981 Jun 1;194(2):260-3.

20. Montgomery, WI. *Peromyscus* and *Apodemus*: patterns of similarity in ecological

equivalents. In: Kirkland Jr GL, Layne JN. (eds.) Advances in the study of *Peromyscus*

(Rodentia). Lubbock, Texas, USA: Texas Tech University; 1989. p.293-366,

21. Terborgh J. Bird species diversity on an Andean elevational gradient. Ecology. 1977

Sep;58(5):1007-19.

22. Graham GL. Changes in bat species diversity along an elevational gradient up the

Peruvian Andes. Journal of mammalogy. 1983 Nov 25;64(4):559-71.

23. Rosenzweig ML. Species diversity in space and time. Cambridge University Press; 1995

May 4.

24. Kaufman DM. Diversity of New World mammals: universality of the latitudinal gradients

of species and bauplans. Journal of Mammalogy. 1995 May 19;76(2):322-34.

25. Sánchez Cordero V. Elevation gradients of diversity for rodents and bats in Oaxaca,

Mexico. Global Ecology and Biogeography. 2001 Jan 1;10(1):63-76.

26. Sanders NJ. Elevational gradients in ant species richness: area, geometry, and Rapoport's

rule. Ecography. 2002 Feb;25(1):25-32.

27. Hillebrand H. On the generality of the latitudinal diversity gradient. The American

Naturalist. 2004 Jan 15;163(2):192-211.

28. Buckley LB, Hurlbert AH, Jetz W. Broad scale ecological implications of ectothermy and

endothermy in changing environments. Global Ecology and Biogeography. 2012

Sep;21(9):873-85.

29. Şekercioğlu CH. Bird functional diversity and ecosystem services in tropical forests,

agroforests and agricultural areas. Journal of Ornithology. 2012 Aug 1;153(1):153-61.

30. BirdLife International and NatureServe. Bird species distribution maps of the world.

[Accessed 2015]

31. Hainsworth FR. On the tongue of a hummingbird: its role in the rate and energetics of

feeding. Comparative Biochemistry and Physiology Part A: Physiology. 1973 Sep

1;46(1):65-78.

32. Schlamowitz R, Hainsworth FR, Wolf LL. On the tongues of sunbirds. The Condor. 1976

Jan 1;78(1):104-7.

33. Paton DC, Collins BG. Bills and tongues of nectar☐feeding birds: A review of

morphology, function and performance, with intercontinental comparisons. Australian

Journal of Ecology. 1989 Dec;14(4):473-506.

34. Liversidge R. The tongues and feeding methods of sunbirds, white-eyes, and sugarbirds.

In: Skead CJ. (eds.) The Sunbirds of Southern Africa, also Sugarbirds, White-eyes and the

Spotted Creeper. Cape Town, South Africa: Balkema; 1967. p.27-32

35. Downs CT. Some preliminary results of studies on the bill and tongue morphology of

Gurney's Sugarbird and some southern African sunbirds. Ostrich-Journal of African

Ornithology. 2004 Sep 1;75(3):169-75.

36. Schuchmann KL. Family Trochilidae (Hummingbirds). In: del Hoyo J, Elliot A, Sargatal J.

(eds.) Handbook of the Birds of the World, Vol.5. Barcelona, Spain: Lynx Edicions; 1999.

p.468-535

37. Chantler P. Family Apodidae (Swifts). In: del Hoyo J, Elliot A, and Sargatal J, (eds.)

Handbook of the Birds of the World, Vol.5. Barcelona, Spain: Lynx Edicions; 1999. p.388-

417

38. Zusi RL. Introduction to the skeleton of hummingbirds (Aves: Apodiformes, Trochilidae)

in functional and phylogenetic contexts. Ornithological Monographs. 2013 Sep 1;77(1):1-

94.

39. Warrick D, Hedrick T, Fernández MJ, Tobalske B, Biewener A. Hummingbird flight.

Current Biology. 2012 Jun 19;22(12):R472-7.

40. Ros IG, Bassman LC, Badger MA, Pierson AN, Biewener AA. Pigeons steer like

helicopters and generate down-and upstroke lift during low speed turns. Proceedings of

the National Academy of Sciences. 2011 Dec 13;108(50):19990-5.

41. Parslew B. Predicting power-optimal kinematics of avian wings. Journal of The Royal
Society Interface. 2015 Jan 6;12(102):20140953.

42. Hedrick TL, Tobalske BW, Ros IG, Warrick DR, Biewener AA. Morphological and
kinematic basis of the hummingbird flight stroke: scaling of flight muscle transmission
ratio. Proceedings of the Royal Society B: Biological Sciences. 2011 Dec
14;279(1735):1986-92.

43. Ingram K. Hummingbirds and miscellaneous orders (Apodiformes, Caprimulgiformes,
Coliiformes, Trogoniformes, Musophagiformes, and Cuculiformes). In: Fowler ME. (eds.)
Zoo and Wild Animal Medicine. Philadelphia, Pennsylvania, USA: WB Saunder
Company; 1978. p.335-346

44. Karhu AA. A new genus and species of the family Jungornithidae (Apodiformes) from the
Late Eocene of the Northern Caucasus, with comments on the ancestry of hummingbirds.
In: Olson SL, Wellnhofer P, Mourer-Chauviré C, Steadman DW, and Martin LD (eds.)
Avian Paleontology at the Close of the 20[th] Century: Proceedings of the 4[th] International
Meeting of the Society of Avian Paleontology and Evolution, Washington, D.C, 4-7 June
1996. Washington, DC, USA: Smithsonian Institution Press; 1999. p.207-216

45. Videler JJ. Avian flight. Oxford, UK: Oxford University Press; 2006 Aug 10.

46. Donoghue MJ, Sanderson MJ. Confluence, synnovation, and depauperons in plant
diversification. New Phytologist. 2015 Jul 1;207(2):260-74.

47. Berger M. Energiewechsel von kolibris beim schwirrflug unter höhenbedingungen.
Journal fuer Ornithologie. 1974 Jul 1;115(3):273-88.

48. Vincent TL, Brown JS. Evolutionary game theory, natural selection, and Darwinian
dynamics. Cambridge University Press; 2005 May 23.

49. Dreisig H. Why do some nectar foragers perch and others hover while probing flowers?.

Evolutionary Ecology. 1997 Sep 1;11(5):543-55.

50. Lima SL. Ecological and evolutionary perspectives on escape from predatory attack: a

survey of North American birds. The Wilson Bulletin. 1993 Mar 1:1-47.

51. Dakin R, Segre PS, Straw AD, Altshuler DL. Morphology, muscle capacity, skill, and

maneuvering ability in hummingbirds. Science. 2018 Feb 9;359(6376):653-7.

52. Hainsworth FR. Why hummingbirds hover: A commentary. The Auk. 1986 Oct

1;103(4):832-3.

53. Pyke GH. Why hummingbirds hover and honeyeaters perch. Animal Behaviour. 1981 Aug

1;29(3):861-7.

54. Wolf LL, Hainsworth FR. Economics of foraging strategies in sunbirds and

hummingbirds. In: Behavioral energetics: the cost of survival in vertebrates. Aspey WP,

Lustick SI. Columbus, Ohio, USA: Ohio State University Press; 1983. p.223-264

55. McGuire JA, Witt CC, Remsen Jr JV, Corl A, Rabosky DL, Altshuler DL, Dudley R.

Molecular phylogenetics and the diversification of hummingbirds. Current Biology. 2014

Apr 14;24(8):910-6.

56. Arnold SJ. Constraints on phenotypic evolution. The American Naturalist. 1992 Nov

1;140:S85-107.

57. Frelich LE, Hale CM, Reich PB, Holdsworth AR, Scheu S, Heneghan L, Bohlen PJ.

Earthworm invasion into previously earthworm-free temperate and boreal forests.

Biological Invasions. 2006; 8(6):1235-45.

58. James SW, Hendrix PF. Invasion of exotic earthworms into North America and other

regions. Earthworm Ecology. 2004 Mar 29;441:75-88.

59. Holmstrup M, Overgaard J, Bindesbøl AM, Pertoldi C, Bayley M. Adaptations to

overwintering in the earthworm Dendrobaena octaedra: genetic differences in glucose

mobilisation and freeze tolerance. Soil Biology and Biochemistry. 2007 Oct

1;39(10):2640-50.

Fig. 1 How differences in adaptations between two convergent taxa (purple and green) may lead

may alter the set of environmental variables (its fundamental niche) under which a taxon has

positive fitness (colored region). (a) Adaptations are largely equivalent, and the taxa survive

under similar conditions. (b) Adaptations are divergent, and there is little overlap in

environmental conditions in which both taxa survive. (c) The superior adaptations of the purple

clade mean it can survive under a greater set of environmental variables (i.e., it has a larger

fundamental niche) compared to the green clade.

Fig. 2 A map of species density of hummingbirds (purple) and sunbirds (green). Richer colors

represent greater species density. Scales are chosen to reflect the difference in overall species

richness of each taxon. Hummingbirds not only have higher species density but also extend

farther.

Fig. 3 Changes in species density with elevation for hummingbirds (purple) and sunbirds

(green). Though both clades extend to similar altitudes, hummingbirds maintain species richness

at higher elevations as denoted by the richer colors.

Fig. 4 A plot of the normalized species richness S_N of hummingbirds and sunbirds, along with

the fitted line, for elevation (a, c) and "polewardness" (b, d). Triangles and purple lines denote

hummingbirds, and circles and green lines denote sunbirds. Hummingbirds maintain species

richness at higher elevations and mid-latitudinal ranges and extend farther latitudinally than

sunbirds. Inflection (cross) and MMC points (asterisks) also are shown (c, d). Inflection points

come later in hummingbirds than sunbirds. With regard to elevation, hummingbird S_N and

sunbird S_N start their decline at a similar spot but hummingbird S_N declines more slowly. With

latitude, sunbird S_N declines earlier than hummingbird S_N.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

489