Effects of single cage housing in the rat and mouse pilocarpine models of epilepsy

Keeping social contact in experimental epilepsy

Manouze H[1,2], Ghestem A[2], Poillerat V[3], Bennis M[1], Ba-M'hamed S[1], Benoliel JJ[3,4], Becker C[3*], Bernard C[2*]

[1]Lab of Pharmacology, Neurobiology & Behavior (URAC-37), Cadi Ayyad University, Marrakech, Morocco

[2]Aix Marseille Univ, Inserm, INS, Institut de Neurosciences des Systèmes, Marseille, France

[3]Université Paris Descartes, Sorbonne Paris Cité, Faculté de Médecine, F-75006, Paris, France

[4]AP-HP, Hôpital de la Pitié-Salpêtrière, Service de Biochimie Endocrinienne et Oncologique, F-75013, Paris, France

CBer, CBec, AG and JJB Designed Research; HM, AG, VP and MB Performed Research and Analyzed data; CBer, CBec, SBM and JJB Wrote the paper

Corresponding author: Christophe.bernard@univ-amu.fr
* Equally contributing last author

2 Figures, 1 Table

150 words in the abstract

106 words in the Significance Statement

399 words in the Introduction

494 words in the Discussion

Acknowledgments

This work was supported by the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement no602102 (EPITARGET), and Fondation pour la Recherche sur le Cerveau. HM was supported by NEUREN Project (PIRSES-GA-2012-318997).

Conflict of Interest

The authors declare no conflicts of interest.

ABSTRACT

Many experimental approaches require housing rodents in individual cages, including in epilepsy research. However, rats and mice are social animals; and individual housing constitutes a stressful situation. The goal of the present study was to determine the effects of individual housing as compared to conditions maintaining social contact on stress markers and epilepsy. Control male mice housed individually for 6 weeks displayed anhedonia, anxiety and biological markers of stress as compared to mice housed 3 per cage. Pilocarpine-treated mice housed together showed increased levels anhedonia, anxiety and stress markers as compared to the control group. The differences were larger in pilocarpine treated mice housed individually. In the male rat pilocarpine model, we also found increased stress markers in isolated animals as compared to rats kept in pairs. Seizures were 16 times more frequent in singly housed animals as compared to animals kept in pairs. Daily interactions with an experimenter in otherwise singly housed animals was sufficient to produce results identical to those found in animals kept in pairs. We propose that social isolation produces a severe phenotype in terms of stress and seizure frequency as compared to animals maintaining social contact (at least in these two models), a factor that needs to be taken into account for data interpretation. This may be particularly relevant for preclinical studies.

Significance Statement

Many experimental approaches require housing rodents in individual cages, a stressful condition for social animals, even in an enriched environment context. Using the pilocarpine model of epilepsy in rats and mice, we report that singly housing animals develop a more severe phenotype in terms of stress and epilepsy as compared to animals maintaining social contact. We propose that social isolation adds a degree of complexity for the interpretation of data. This may be particularly relevant for preclinical studies.

Introduction

Social isolation (single housing) and barren environment are stressful to rodents; for a recent review cf. (Mumtaz et al., 2018). In fact, social isolation of rodents after weaning is often used as an experimental manipulation to model early life stress in humans. Social isolation for 8 weeks after weaning produces anxiety/depressive-like behaviour and cognitive deficits and favours the emergence of pathological traits such as addictive behaviour (Filipovic et al., 2017). These behavioural alterations are associated with biological modifications including oxidative stress, the production of stress hormones and inflammatory cytokines (Krugel et al., 2014; Shao et al., 2015; Butler et al., 2016; Filipovic et al., 2017). Therefore, many regulation agencies recommend social housing and environmental enrichment to improve welfare of rats and mice in animal facilities (Lidster et al., 2016). However, single housing of rodents may be imposed by the experimental protocol to obtain accurate measurements of biological parameters in individual animals, such as in the fields of toxicology, drug addiction, and neurological disorders. Experimental epilepsy is a typical example. In order to precisely assess seizures with continuous video-EEG recordings, instrumented animals are singly housed to prevent the severing of EEG wires and aggressive behaviour from others, to clearly identify individuals on the video, etc. Several studies have shown that environment enrichment has positive effects on seizure frequency or severity in experimental models (Morelli et al., 2014; Kotloski and Sutula, 2015; Dezsi et al., 2016; Vrinda et al., 2017), but the effect of social isolation starting from the beginning of the experimental procedure has not been assessed.

There exist many types of experimental models of epilepsy in different rodent species and strains (Levesque et al., 2016; Loscher, 2017; Becker, 2018). The effects of social isolation should be studied for each model, as results may be model-, strain- and species-

specific. For example, Wistar and Sprague Dawley rats display different phenotypes in terms

of depression-like profile and cognitive deficits in the kainic acid and pilocarpine models

(Inostroza et al., 2011; Inostroza et al., 2012). Here we focus on the widely used pilocarpine

model of experimental epilepsy in adult male Swiss mice and Wistar rats (Levesque et al.,

2016). Numerous studies show a dysregulation of the HPA axis in patients and in

experimental epilepsy, which is model-dependent (Maguire and Salpekar, 2013; Wulsin et al.,

2016; Mumtaz et al., 2018). Since stress also results in the activation of the HPA axis, we

reasoned that social isolation in experimental models of epilepsy may produce a strong HPA

axis response via the contributions of both -isolation and epilepsy-induced stress (Gunn and

Baram, 2017). We also hypothesized that the combination of both factors may exacerbate the

epilepsy phenotype as compared to animals maintaining social interactions. Clinical studies

support this hypothesis. Stressful life events are associated with increased risk of seizure

occurrence in patients with epilepsy (Baldin et al., 2017; Kotwas et al., 2017). In some

patients, the fear to have a seizure and social isolation due to stigmatization may increase the

allostatic load, with a direct impact on seizure frequency (Kotwas et al., 2017). We thus

assessed the consequences of animal housing on stress response and epilepsy severity, acting

on a single experimental variable: social interaction (environmental enrichment was

provided). We also propose an alternate solution to maintain social contact in singly housed

animals, to decrease their stress level. We hope to foster discussions on how to interpret

results (including negative preclinical studies) obtained in different housing conditions.

METHODS
Animals

We used male Wistar rats (200 to 250g; Charles Rivers Laboratories, Les Oncins, France),

aged of 9 weeks (at their arrival in laboratory). Swiss male mice (8 weeks old) were obtained

from the animal husbandry of the Faculty of Sciences, Cadi Ayyad University, Marrakech

(Morocco). The animals were kept under controlled environmental conditions (23±1 °C;

night-day cycle (12 h-12 h)) with *ad libitum* access to food and water. Zeitgeber (ZT) 0 was at

7:30 am (time when the light was switched on in the animal facility). All procedures were

conducted in accordance with approved institutional protocols, and with the provisions for

animal care and use prescribed in the scientific procedures on living animals, European

Council Directive: EU2010/63and by the Council Committee of Research Laboratories of the

Faculty of Sciences, Cadi Ayyad University, Marrakech. All efforts were made to minimize

any animal suffering.

Experimental design

In mice

The experimental protocol is shown in Figure 1A. Swiss male mice were housed in groups

of 3 per cage. In order to validate the stress state of same mouse throughout experimental

protocol, one mouse from each litter was randomly picked, ear-marked and returned to its

own cage. They were left together during one week without experimental intervention to

allow adaptation to a novel housing environment. After one week, we performed different

behavioral tests for each ear-marked mouse in order to obtain reference levels. Sucrose

preference was assessed every day during 1 week. At the end of the week, we evaluated

anxiety levels (Day-3: elevated plus maze – EPM) and performance in the novel object

recognition (NOR) test (Day -2-1). Animals were then randomly assigned to two groups to

receive a pilocarpine injection to trigger status epilepticus (pilo group– experimental

procedure described hereafter) or saline (non-pilo group). Both groups were further divided

into two groups: group-housing (social condition – SC, n =18, 3/cage) and one individual-

housing condition (isolated condition – IC, n = 6, 1/cage). Individual housing used ear marked

mice. After 4 weeks exposure to different housing conditions, ear-marked mice from social

and isolated conditions were tested for anhedonia (Day 46-53), EPM (Day 54) and NOR (55-

56). Behavioral tasks (EPM and NOR) were recorded and analyzed using

Ethovision®XTNoldus 8.5 video tracking program (Noldus, Netherlands) connected to a

video camera (JVC). At the end of each behavioral session apparatuses were cleaned with a

75% ethanol solution to remove any of odor or trace. The behavioral tests were performed

between 8:00 and 12:00 a.m. during the light cycle to avoid the circadian-related fluctuation

in the performance of the mice. Before the beginning of behavioral tests, the animals were

transferred to the testing room in their home cages and left there to habituate for 60 min.

At the end of the experimental protocol, we measured ACTH, CORT and BDNF levels as

in rats (described below).

In rats

The experimental protocol is shown in Figure 2A. Animals were received in sets of 4 from the

same litter from the vendor. Two main groups of animals were used: animals with

spontaneous seizures following pilocarpine-induced status epilepticus (pilo group –

experimental procedure described hereafter) and control animals (non-pilo group). Pilo and

non-pilo animals were further divided into three groups:

- Isolated group: rats were singly housed and were not handled during the experimental

    period except for cage cleaning and body weight measurement once a week.

- Handled group: rats were singly housed and were handled daily until the end of

    experiment (see below).

- Paired group: rats were kept in buddy pairs with social interaction upon arrival in the

    animal facility.

Handling was performed twice daily (ZT2 and ZT9) throughout the experimental period by the same experimenter. Briefly, each handling session consisted of stroking animals for 1 minute each in their cage. Then, each rat was gently handled by experimenter's hands (without wearing gloves), while being softly stroked from the head to the tail for 2 minutes. Finally, the rats were placed back in their home cage and fed by the experimenter for 2 minutes. Isolated and paired groups were left undisturbed, except for weekly cage cleaning and body weighing.

Animals from the non-pilo group were randomized to the three housing groups when received. Animals from the pilo group were randomized after pilocarpine-induced status epilepticus. Since we have no knowledge of stress history of the animals when we receive them, randomization decreases this confounding factor.

Sample size

A preliminary analysis made on isolated (n=4) and handled (n=4) pilo groups for seizure frequency allowed us to perform a power analysis. We determined that n=3 per group was enough. The experimental procedure in the pilo groups was conducted twice. The first and second group of simultaneously recorded animals being composed of (n=4 isolated, n=4 handled, n=6 paired) and (n=3 isolated, n=4 handled, n=6 paired), respectively. In the paired groups, only one animal was instrumented. Seizure analysis was thus performed in 6 animals. Results were similar in the two sets; and data was pooled together.

For the mice experiments, the number of subjects was defined on the basis of the study design. Preliminary analysis was conducted with n=3 for social condition and n=3 for isolated condition where the results were non-significant. In this case, a second study was done with a sample size of 3 per group.

Status epilepticus induction and electrode implantation

In rats, status epilepticus (SE) was induced by a single i.p injection of pilocarpine (pilo) 320 mg/kg, one week after receiving the animals from the vendor. In mice, we used repeated 100 mg/kg i.p. injections every 20 min until SE onset 17 days after receiving the animals (1 week for habituation and 10 days for baseline of behavioural tasks). To reduce peripheral effects, animals were pre-treated with Methyl-scopolamine (1mg/kg) 30 min prior pilo injection. SE was stopped by diazepam (10 mg/kg i.p., twice within a 15 min interval) after 60 min and 90 min of SE; respectively. At the end of these injections, mice and rats were hydrated with saline (2ml i.p. twice within 2 h) and fed with a porridge made of soaked pellets, until they resumed normal feeding behavior. All drugs were obtained from Sigma.

In rats, four weeks following SE, the telemetry implant was surgically inserted intraperitoneally under anesthesia (ketamine [1 mg/ kg]/xylasine [0.5 mg/kg] i.p) and connected to screws on the surface of the brain by two electrodes; one above the cortex (4.0 mm anteroposterior, 2.0 mm mediolaterally, compared with bregma), the second, above the cerebellum as reference. The EEG signal was transmitted by radio to a data acquisition and processing system (DSI). In the paired group, both animals developed epilepsy but only one rat was equipped with the telemetry system as two animals cannot be recorded simultaneously with EEG transmitters in the same cage, while the other was monitored with video only. Animals were left to recover during one week before switching on the transmitter.

Monitoring of spontaneous recurrent seizures

Continuous (24/7) video-EEG recordings started in the 6^{th} week after SE, a period sufficient to reach stability in seizure frequency (Williams et al., 2009), and were stopped at week 10. We verified that animals displayed stable seizure activity, quantifying seizure frequency during each successive week. Spontaneous recurrent seizures were detected and quantified

using both visual inspections of the EEG and a semi-automatic way usingClampfit 10.2. All

detected seizures were verified and reconfirmed using "NeuroScore" software. Video

recordings were used to assess seizure severity according to Racine's scale during the light

phase in the animal facility. Rats kept in pairs never had spontaneous seizures simultaneously,

which allowed a correct assessment of seizure severity of the EEG monitored animal. Finally,

keeping animals in pairs did not prevent/alter continuous EEG recordings in the equipped

animals (no loss of signal).

Behavioral and biological parameters

Body weights of mice and rats were measured weekly at ZT 2.

Sucrose consumption test

In mice, the baseline for sucrose consumption was assessed every day for 1 week, 7

days after their arrival in laboratory. The sucrose preference was calculated at the 5th week

following the separation and treatment. In rats, sucrose consumption was assessed at week 4

and week 8 following animal reception. Sucrose and water intakes were measured daily at ZT

2.

Briefly, animals were given a free choice between two bottles, one with 1% sucrose

solution and another with tap water. The location of the bottles was alternated every day to

prevent possible effects of side preference in drinking behaviour. The consumption of water

and sucrose solution was estimated by weighting the bottles. For the paired, the volume of

sweet water consumed by rat was taken as the total consumed volume divided by 2. For the

social groups, the volume of sweet water consumed by mouse was taken as the total

consumed volume divided by the total number of animals at each time. Sweet water

consumption corresponds to sucrose preference, which is calculated as a percentage of the

volume of sucrose intake over the total volume of fluid intake using the following equation:

Sucrose preference= V (sucrose solution)/ (V (sucrose solution) +V (water)) X100%.

Elevated plus maze test

The elevated plus maze has been described as a simple method for assessing anxiety-like

behavior in rodents (Pellow et al., 1985; Lapiz-Bluhm et al., 2008). The elevated plus maze

consisted of four arms (two open without walls and two enclosed arms by 15 cm high walls)

50 cm long and 5 cm wide, which were joined at a square central area (5×5 cm) to form a

plus sign. The maze floor and the side/end walls of the enclosed arms were made of clear

Plexiglas. The room illumination of the elevated plus maze apparatus was under an

approximate brightness of 200 lux. Briefly, the test consisted of placing gently the mouse in

the central arena of the elevated plus maze, facing the junction of an open and closed arm.

The mouse was allowed to freely explore the maze for 5 min while the duration and frequency

of entries into open arms was recorded. Anxiety index (Rao and Sadananda, 2016) was

calculated as: 1 - ((open-arm time/total time) + (open-arm entries/total entries))/2.

Novel object recognition test

The novel object recognition test is widely used to evaluate object recognition memory in

rodents (Ennaceur and Aggleton, 1994; Reger et al., 2009; Gaskin et al., 2010). The apparatus

consisted of an open field ($50 \times 50 \times 50$ cm high) made of Plexiglas with the inside painted

matt black. The objects to be discriminated were available as three plastic objects. 24h before

testing, mice were first habituated in the open field arena in the absence of any object for

10min. During the training trial, two identical objects (approximately 10 cm) were placed in

the back corner of the box. The mouse was then positioned at the midpoint of the wall

opposite to the objects and the total time spent exploring the two objects was recorded for 10

min. During the test trial (60 min after training trial), one object used during training was

replaced with a novel object and both of them were placed in the middle of the back wall. The

animal was then allowed to explore freely for 5 min and the time spent exploring of each

object was recorded. The discrimination between the novel and familiar objects was

calculated by the discrimination index [(time spent exploring the novel object - time spent

exploring the familial object) / total time x 100)].

ACTH, corticosterone and BDNF levels

Isolated, handled and paired rats in the non-pilo group were killed by decapitation after light

anesthesia with isoflurane at the beginning of the 8[th] week following the separation at ZT4

and at the 6[th] weeks after the separation and treatment at ZT4 for pilo and non-pilo mice.

Animals were decapitated in a quiet separate room, one by one, with the bench cleaned

between animals. Trunk blood was collected in less than 5 sec after decapitation in dry tube

and EDTA tube in order to obtain serum and plasma samples, respectively. The plasma was

prepared by a 15 min centrifugation at 1600g, 4°C. The ACTH concentration was determined

according the manufacturer's instructions (Clinisciences, France). For corticosterone and

BDNF levels, blood was centrifuged at 3500g for 10 min at 4°C and the serum was stored at

80°C until used. Corticosterone and BDNF concentrations were determined according to the

manufacturer's instructions (Coger, Promega, France).

Statistical analysis

Statistical analysis was performed using Sigma Plot 11.0 software. All data is presented as

mean ± SEM. The effect on body weight and anhedonia (rat) was measured using the

ANOVA with repeated measures followed by Bonferroni *post-hoc* analysis. The biological

and seizures parameters were tested according to the one-way analysis of variance (ANOVA)

followed by Bonferroni multiple comparison test. The effects of group and pilo treatment on

anxiety index, discrimination index, ACTH, BDNF and corticosterone were measured using

the two-way ANOVA followed by Bonferroni *post-hoc* analysis. The effects on anhedonia

was measured using the non-parametric *Mann–Whitney U*. The significance threshold was set

at $p<0.05$.

RESULTS

We first tested the consequences of social isolation on behavioral parameters and stress markers in adult pilo and non-pilo mice (**Figure 1A**).

Repeated measures ANOVA showed a significant effect of time ($F_{(5,100)}$=128.08, p<0.0001[a]) on body weight evolution; while the experimental group ($F_{(3,20)}$ =2.23, p=0.11[a]) and the interaction of group and age ($F_{(15,100)}$ =1.18, p=0.29[a]) had no effect (**Figure 1B**). At the 4th and 5th week post-social isolation, *post-hoc* analysis showed that both pilo-SC and pilo-IC groups gained significantly less weight than the non pilo-SC and the non pilo-IC group. There was no significant difference between all groups until the 3rd week. Thus, the pilo group gained less weight than the non-pilo group, but housing conditions had no effect in both groups.

We used the sucrose preference test to assess anhedonia (lack of interest in rewarding stimuli). The typical behaviour of animals is a bias toward the sweetened drink. Lack of preference for the sweetened drink indicates anhedonia, which signs stress-related disorders (Gold, 2015). At baseline (before changing housing conditions), animals from different groups (different cages) did not show differences in sucrose preference (**Figure 1C**). Four weeks later, our results demonstrated that mice injected by pilo consumed less of sweet water than mice injected with vehicle. In addition, mice in isolated condition had sweet water consumption lower than those in social condition (**Figure 1C**). From day 1 until day 7, the non-parametric *Mann–Whitney U* test, applied on sweet water consumption, revealed that the pilo-SC (day 1: U=0.00; Z=-10.89; p<0.01; day 7: U=0.00; Z=-9.35; p<0.01 [B-1]) and pilo-IC (day1: U=0.00; Z=-10.23; p<0.01; day 7: U=0.00; Z=-10.09; p<0.01 [B-1]) groups consumed significantly less of sweet water than the non pilo-SC and non pilo-IC groups. Additionally, the *Mann–Whitney U* showed that both non pilo-IC (day1: U=2.00, Z= -4.87, p<0.05; day 7: U=1.50; Z= -4.72; p<0.01 [B-1]) and pilo-IC (day 1: U=1.00; Z=-4.42; p<0.01; day 7: U=5.00;

Z= -5.00; p<0.05 [B-1]) groups had a significant decrease in sucrose preference as compared to

non pilo-SC and pilo-SC groups, respectively. Moreover, the Mann–Whitney U test did not

show any effect of day on sweet consumption (p>0.05). Therefore, social isolation induces

anhedonia in the control (non pilo) group. The pilo group socially housed also displayed

anhedonia as compared to the control group, but the anhedonia phenotype was increased in

the socially isolated pilo group.

We then tested animals for anxiety-like behavior. All groups tested before separation

had similar anxiety levels (**Figure 1D**). Two-way ANOVA demonstrated significant

difference in anxiety index recorded among the factors group ($F_{(3,40)}$=12.93, p<0.0001[b]) and

time ($F_{(1,40)}$=64.32, p<0.0001[b]) as well as the interaction of group $\times$ time ($F_{(3,100)}$=11.84,

p<0.0001[b]). *Post-hoc* comparisons confirmed that the pilo-treated groups showed a significant

increase in anxiety (p<0.01 and p<0.001) with respect to the non pilo groups (**Figure 1D**). In

the isolated condition, the anxiety index was significantly larger in both pilo and non pilo

mice as compared to mice in social housing (p<0.05). These results show that social isolation

readily increases anxiety levels in the control group. The pilo group maintaining social

contact displayed increased anxiety as compared to the control group (epilepsy effect), but

anxiety was further exacerbated in the pilo isolated group (combining the effects of epilepsy

and social isolation).

Next, we assessed the effect of social isolation on memory performance using the NOR

test. All groups performed similarly at baseline before separation and treatment (**Figure 1E**).

The two-way ANOVA test showed significant difference in the discrimination index among

the factors group ($F_{(3,40)}$=6.95, p=0.0007[c]) and time ($F_{(1,40)}$=43.30, p<0.0001[c])as well as the

interaction of group $\times$ time ($F_{(3,100)}$=8.89, p<0.0001[c]) (**Figure1E**).The discrimination index for

the pilo-treated group was notably lower than that of the non pilo group (p<0.0001).

However, there was no significant effect of housing conditions in NOR performance in either

pilo or non pilo groups. Hence, the cognitive deficits in the NOR test are only due to the

epilepsy condition.

Finally, we looked at the effect of social isolation on biological markers. For ACTH and

corticosterone, two stress-related hormones, two-way ANOVA analysis showed a significant

effect of treatment ($F_{(1,20)}=74.62$, $p<0.0001$; $F_{(1,20)}=117.3$, $p<0.0001$[d,e]) and social isolation

($F_{(1,20)}=20.55$, $p=0.0002$[d,e]; $F_{(1,20)}=18.56$, $p=0.0003$[d,e]) (**Figure 1F**). Our results showed a

significant increase of ACTH and corticosterone levels in pilo as compared to non pilo mice

($p<0.001$). Moreover, this increase was observed in isolated condition as compared to social

housing ($p<0.05$).

Strong stressful situations can trigger a state of vulnerability to depression in some

individuals, which can be identified with low serum BDNF levels (Blugeot et al., 2011;

Becker et al., 2015). We thus assessed serum BDNF in the different groups. Although serum

BDNF level was not affected in the pilo group as compared to the non pilo group when

animals were socially housed ($F_{(1,20)}=0.47$, $p=0.5011$[f]); social isolation had a significant effect

($F_{(1,20)}=16.96$, $p=0.0005$[f]) in both pilo and non pilo groups (**Figure 1F**). *Post-hoc* comparison

confirmed that isolated mice showed a significant decrease in BDNF ($p<0.05$) when

compared to socially housed mice. There was no significant difference between pilo and non

pilo groups.

Together, the results obtained in control (non pilo) mice demonstrate that social

isolation has a strong effect on many behavioral and biological stress indicators. They

recapitulate the main features of social isolation performed after weaning to model early life

stress (Mumtaz et al., 2018). In animals maintained in groups, the epilepsy had a direct effect

on stress indicators. These effects were exacerbated in singly housed pilo mice. We conclude

that social isolation produces a stronger stress phenotype (with the tests used here) in the pilo

mouse experimental model of epilepsy. The lack of EEG recording devices for mice

prevented us to test the effect of social isolation on epilepsy severity. This issue was investigated in the pilocarpine rat model.

Because of their larger size, it was not technically possible to maintain rats in small colonies while performing wireless 24/7 recordings. For the social group, animals were kept as pairs (only one was equipped with wireless transmitter, but both received pilocarpine injection). We added a third group (handled) of singly housed animals but which could interact with the experimenter everyday (see methods). The rationale for this group is that some EEG studies are performed with wired systems, which precludes group housing, as animals tend to severe the wires. We first analysed general stress markers in control (non pilo) animals to determine whether social isolation had similar effects to those described in mice. The ANOVA repeated measures showed a significant effect of social condition ($F_{(2,17)}$=41.58, $p<0.0001^{g}$) and time ($F_{(6,102)}$ =266.83, $p<0.0001^{g}$) on the gain of body weight (**Figure 2B**). At the 4th week, *post-hoc* analysis showed that the isolated group gained significantly less weight than the handled and the paired group. There was no significant difference between all groups from the 1st to the 3th week.

We used the sucrose preference test to assess anhedonia. ANOVA repeated measures showed a main effect of social isolation on sweet water consumption (**Figure 2C**) already during the third week ($F_{(2,12)}$=21.66, $p<0.0001^{h}$) up to the last week ($F_{(2,12)}$=97.85, $p<0.0001^{i}$) with no time effect in both weeks (first week: $F_{(6,72)}$ =0.11, $p>0.05$ and last week: $F_{(6,72)}$ = 1.09, $p>0.05$; respectively ^{h,i}). During the third and the last week, *post-hoc* analysis revealed that sweet water consumption was significantly lower in the isolated than in the handled ($p<0.01$, $p<0.001$) and paired groups ($p<0.05$, $p<0.01$ and $p<0.001$). Thus, just two weeks of social isolation is sufficient to produce a state of anhedonia.

We then tested stress-related ACTH and corticosterone hormones. One-way ANOVA revealed a significant effect of social isolation on hormone levels (ACTH: $F_{(2,17)}$=5.05,

p<0.02[j]; corticosterone: $F_{(2,17)}$=8.42, p<0.01[j]; **Figure 2D**). The *post-hoc* analysis showed that

ACTH and corticosterone levels were significantly increased in the isolated group as

compared to the handled (t=2.69, p<0.05 and t=3.57, p<0.01; respectively) and paired (t=2.93,

p<0.05; t=3.70, p<0.01; respectively) groups.

Finally, we assessed serum BDNF levels. One-way ANOVA revealed a significant

effect of social isolation on BDNF levels ($F_{(2,17)}$= 6.18, p<0.01[l]). BDNF levels were

significantly lower in the isolated group as compared to the handled (t=2.38, p<0.05) and

paired (t=3.47, p<0.01) groups (**Figure 1D**). There was no significant difference between the

handled and paired groups in all tested biological parameters (ACTH: t=0.18; corticosterone:

t=0.42 and BDNF: t=1.02, p>0.05).

We conclude that social isolation in the strains of control rats and mice used here

produces aphenotype associated with stress. We then compared how different housing

conditions influence the epilepsy phenotype.

One-way ANOVA analysis showed a significant difference between groups for seizure

parameters (frequency, duration and severity) ($F_{(2,26)}$=10.73, *p*<0.001; $F_{(2,26)}$=30.90, *p*<0.0001;

$F_{(2,26)}$=4.07, *p*<0.05; respectively [m,n,o]) (**Figure 3**). Figure 3-1 shows seizure frequency for

each week for each experimental group, and for each individual animal. *Post-hoc* analysis

showed that the seizure frequency and duration were considerably increased in the isolated

group as compared to the handled and paired groups (handled: t=4.29 and t=7.82, p<0.001;

paired: t= 3.45 and t=4.07, p<0.01; respectively). The handled group significantly

demonstrated lower Racine scores (seizure severity) as compared to the isolated group

(t=2.72, p<0.05); whereas, there was no significant difference between the paired and isolated

groups (t=1.94, p>0.05). We only assessed the severity of the seizures during the light phase

in the animal facility. Finally, there was no significant difference between the handled and

paired groups for seizure frequency and seizure severity. Thus, lack of social interaction

dramatically increased the seizure phenotype, with seizures being 16 times more frequent in

isolated animals as compared to the those socially housed or those having daily interaction

with experimenters.

DISCUSSION

Our results show that single housing exacerbates the phenotype in Sprague Dawley rats and

Swiss mice in the pilocarpine experimental model of epilepsy. It is important to note that

daily social interaction with the experimenter was sufficient to prevent the development of a

severe epilepsy phenotype in rats although they were singly housed. It was as efficient as

keeping animals in pairs. Although we could not quantify it, we noted that isolated pilo

animals were very aggressive and displayed escaping behaviour when cages were changed

every week. Their cages were close to one another, suggesting that visual contact was not

sufficient to prevent the effect of social isolation. Abnormal reaction to handling is a direct

consequence of single housing (Hatch et al., 1965). In contrast, rats and mice socially housed

(or in daily contact with an experimenter) remained calm when handled. In the laboratory, we

are also performing electrophysiological recordings with high density silicon probes in

experimental rat models of epilepsy. The equipment is protected with a copper mesh hat,

which precludes social housing. We are using large cages, divided in two, separated by a grid

allowing physical interaction. Each compartment contains one animal with epilepsy. Animals

can make nose contact and interact with their whiskers through the bars. Animals do not show

any sign of anxiety, and wire connectors are easily inserted, as in control animals (no

struggle). This type of housing allows wired recordings in rats and mice with epilepsy, whilst

maintaining a certain level of social interaction.

Our data on control animals confirms the large body of literature reporting that social

isolation constitutes a major stressful situation that can lead to a depression-like profile

(Mumtaz et al., 2018). Among the numerous markers that can be assessed, we focused on

general ones such ACTH, corticosterone and anhedonia. When rats and mice are singly

housed, ACTH and corticosterone levels are increased (Veenema et al., 2005; Djordjevic et

al., 2012). In pilocarpine experimental models of epilepsy, ACTH and corticosterone levels

are also increased (Mazarati et al., 2009; Inostroza et al., 2012; Ngoupaye et al., 2013). A

direct comparison of ACTH and corticosterone values between studies is difficult as they

depend upon the time of collection and the way blood is collected (*in vivo* or *post mortem* as

done here). Despite the differences in experimental procedures, all results are qualitatively

similar (increase due social isolation or epilepsy). To the best of our knowledge, we here

provide the first evaluation of the contribution of both factors: social isolation and epilepsy.

We demonstrate that epilepsy in socially housed mice produces an activation of the HPA axis

and anhedonia, responses which are amplified in isolated mice with epilepsy. We found the

same exacerbation in the rat experimental model between isolated and paired rats. A strong

activation of the HPA axis in isolated animals with epilepsy may contribute to the expression

of co-morbidities such as anxiety, cognitive deficits and depression-like behaviour.

In rats with epilepsy, we found that spontaneous seizures were 16 times more frequent

in isolated animals as compared to animals kept in pairs or singly housed animals but with

daily interactions with experimenter. During the 3 weeks of continuous recordings (starting 6

weeks after pilocarpine-induced status epilepticus), we did not find an increase in seizure

frequency (Figure 3-1), suggesting that a global steady state activity had been reached

(Williams et al., 2009). Interictal activity and seizures follow a multidien rhythm in the

pilocarpine experimental rat model of epilepsy (Baud et al., 2019) as in humans (Baud et al.,

2018). The fact that seizures occur at specific phases of this rhythm is not a confounding

factor as its frequency is between 5 and 7 days in this rat model (Baud et al., 2019), which means that we captured at least 3 cycles. In the present study, isolated animals had a high seizure frequency (≈16/day). It is difficult to compare this value with the existing literature as housing conditions are rarely mentioned. In addition, the beginning (how long after status epilepticus) and duration of 24/7 EEG recordings vary from one study to another. In the rat pilocarpine model, typical values for singly housed animals in Sprague Dawley rats are: 3/day during the early phase of epilepsy (Behr et al., 2017), 8/day (Paolone et al., 2019), 1-13/day (Tai et al., 2017), 10/day (Bankstahl et al., 2012); similar to values reported in Wistar rats: ≈4.5/day (Bajorat et al., 2011; Bajorat et al., 2018). Our study, although on the high end, remains in the same order of the magnitude as previously published work (we limited our analysis of the literature to the ongoing decade). We evaluated seizure severity only during the light phase, which constitutes a limitation of the study. We cannot rule out that seizures occurring during the dark phase may be less severe (including subclinical electrographic seizures).

Finding the mechanisms underlying the difference in phenotypes between the different housing conditions was beyond the scope of this study. We do not claim that the strong seizure phenotype found in the isolated group is solely due to the stronger activation of the HPA axis, although it may contribute to it. We may tentatively propose that social isolation produces a vulnerability phenotype, as suggested by the low levels of serum BDNF (Becker et al., 2015). We suggest that epileptogenesis occurring in networks "weakened" by social isolation-induced stress would result in a strong phenotype.

The field of epilepsy research makes use of many different experimental models, strains and species. Results are clearly model-, strain- and species-specific, which may explain some conflicting results. We propose that another factor to consider is housing conditions. Some patients with epilepsy may experience social isolation, which correlates

with more severe forms of epilepsy (Kotwas et al., 2017). Results obtained in experimental

animal models kept in isolation may be highly relevant to such patient situation, or any other

situation associated with high levels of stress. Results obtained in socially housed animals

may be relevant to other subsets of patients, with a less severe phenotype. Our results may

have important consequences in terms of data interpretation, in particular when testing anti-

epileptogenesis or anti-seizure strategies. When animals are kept in isolation, two

intermingled factors need to be taken into account: social isolation-induced stress and

epilepsy. Different interpretations can thus be provided to the existing body of literature in

experimental epilepsy. Some studies report a large decrease in seizure frequency when

animals are treated with a given compound. It is possible that the compound may have

targeted the isolation-dependent stress component of the phenotype and not epilepsy itself. A

reduction in seizure frequency by a factor of 16 would be equivalent to the difference we

report between paired and isolated animals. For example, inhibition of kelch like ECH

associated protein 1 (Keap1) reduces seizure frequency by a factor 20 in isolated kainic acid-

treated Sprague-Dawley animals (Shekh-Ahmad et al., 2018). Keap1 controls nuclear factor

erythroid 2-related factor 2 (Nrf2), a key regulator of antioxidant defence mechanisms. The

authors suggest that these results support "the hypothesis that reactive oxygen species

generation is a key event in the development of epilepsy" (Shekh-Ahmad et al., 2018). Since

social isolation produces reactive oxygen species (Filipovic et al., 2017), it is possible that

activating antioxidant defence mechanisms may have targeted the isolation-induced

component of the seizure phenotype. Indeed, treating animals with a strong antioxidant had

no effect on seizure frequency in the kainic acid rat model when animals interacted daily with

experimenters, whilst the treatment was very efficient when animals are characterized by high

levels of reactive oxygen species before status epilepticus (in press). These considerations

provide another possible interpretation of the Keap1 data (Shekh-Ahmad et al., 2018).

However, their conclusion that antioxidant treatment is strongly disease modifying in the

context of social isolation is not only valid but also highly clinically relevant to socially

isolated patients. Finally, many (often unpublished) preclinical studies performed in isolated

animals did not evidence any significant effect on seizure frequency. Since the social isolation

stress component could act as a confounding factor producing a strong phenotype, there may

be false negative results; i.e. the tested drugs may be very efficient in experimental animals

maintaining social interaction.

Although we cannot claim that maintaining social interaction between rodents in

laboratory conditions corresponds to "normality" (as in the wild), single housing does

produce strong biological alterations that render data interpretation more complex, in both

physiological and pathological conditions. We recommend that material and methods should

systematically report housing conditions, and that all efforts should be made to maintain

social interaction. Finally, these results were obtained in one experimental model

(pilocarpine), in two species, one strain for each species and only in males. Similar studies

should be performed in females, other strains and models such kainic acid, kindling, tetanus

toxin etc. Our results and conclusions should not be seen as a criticism of previous works.

Rather varying housing conditions provide a unique opportunity to study, with the same

experimental model, different situations found in patients with high and low stress levels.

REFERENCES

Bajorat R, Wilde M, Sellmann T, Kirschstein T, Kohling R (2011) Seizure frequency in

pilocarpine-treated rats is independent of circadian rhythm. Epilepsia 52:e118-122.

Bajorat R, Porath K, Kuhn J, Gossla E, Goerss D, Sellmann T, Kohling R, Kirschstein T

(2018) Oral administration of the casein kinase 2 inhibitor TBB leads to persistent

KCa2.2 channel up-regulation in the epileptic CA1 area and cortex, but lacks anti-

seizure efficacy in the pilocarpine epilepsy model. Epilepsy Res 147:42-50.

Baldin E, Hauser WA, Pack A, Hesdorffer DC (2017) Stress is associated with an increased

risk of recurrent seizures in adults. Epilepsia 58:1037-1046.

Bankstahl M, Bankstahl JP, Loscher W (2012) Inter-individual variation in the anticonvulsant

effect of phenobarbital in the pilocarpine rat model of temporal lobe epilepsy. Exp

Neurol 234:70-84.

Baud MO, Ghestem A, Benoliel JJ, Becker C, Bernard C (2019) Endogenous multidien

rhythm of epilepsy in rats. Exp Neurol.

Baud MO, Kleen JK, Mirro EA, Andrechak JC, King-Stephens D, Chang EF, Rao VR (2018)

Multi-day rhythms modulate seizure risk in epilepsy. Nat Commun 9:88.

Becker AJ (2018) Review: Animal models of acquired epilepsy: insights into mechanisms of

human epileptogenesis. Neuropathol Appl Neurobiol 44:112-129.

Becker C, Bouvier E, Ghestem A, Siyoucef S, Claverie D, Camus F, Bartolomei F, Benoliel

JJ, Bernard C (2015) Predicting and treating stress-induced vulnerability to epilepsy

and depression. Ann Neurol 78:128-136.

Behr C, Levesque M, Stroh T, Avoli M (2017) Time-dependent evolution of seizures in a

model of mesial temporal lobe epilepsy. Neurobiol Dis 106:205-213.

Blugeot A, Rivat C, Bouvier E, Molet J, Mouchard A, Zeau B, Bernard C, Benoliel JJ, Becker
C (2011) Vulnerability to depression: from brain neuroplasticity to identification of
biomarkers. J Neurosci 31:12889-12899.

Butler TR, Karkhanis AN, Jones SR, Weiner JL (2016) Adolescent Social Isolation as a
Model of Heightened Vulnerability to Comorbid Alcoholism and Anxiety Disorders.
Alcohol Clin Exp Res 40:1202-1214.

Dezsi G, Ozturk E, Salzberg MR, Morris M, O'Brien TJ, Jones NC (2016) Environmental
enrichment imparts disease-modifying and transgenerational effects on genetically-
determined epilepsy and anxiety. Neurobiol Dis 93:129-136.

Djordjevic J, Djordjevic A, Adzic M, Radojcic MB (2012) Effects of chronic social isolation
on Wistar rat behavior and brain plasticity markers. Neuropsychobiology 66:112-119.

Ennaceur A, Aggleton JP (1994) Spontaneous recognition of object configurations in rats:
effects of fornix lesions. Exp Brain Res 100:85-92.

Filipovic D, Todorovic N, Bernardi RE, Gass P (2017) Oxidative and nitrosative stress
pathways in the brain of socially isolated adult male rats demonstrating depressive-
and anxiety-like symptoms. Brain Struct Funct 222:1-20.

Gaskin S, Tardif M, Cole E, Piterkin P, Kayello L, Mumby DG (2010) Object familiarization
and novel-object preference in rats. Behav Processes 83:61-71.

Gold PW (2015) The organization of the stress system and its dysregulation in depressive
illness. Mol Psychiatry 20:32-47.

Gunn BG, Baram TZ (2017) Stress and Seizures: Space, Time and Hippocampal Circuits.
Trends Neurosci 40:667-679.

Hatch AM, Wiberg GS, Zawidzka Z, Cann M, Airth JM, Grice HC (1965) Isolation syndrome
in the rat. Toxicol Appl Pharmacol 7:737-745.

Inostroza M, Cid E, Menendez de la Prida L, Sandi C (2012) Different emotional disturbances

in two experimental models of temporal lobe epilepsy in rats. PLoS One 7:e38959.

Inostroza M, Cid E, Brotons-Mas J, Gal B, Aivar P, Uzcategui YG, Sandi C, Menendez de la

Prida L (2011) Hippocampal-dependent spatial memory in the water maze is

preserved in an experimental model of temporal lobe epilepsy in rats. PLoS One

6:e22372.

Kotloski RJ, Sutula TP (2015) Environmental enrichment: evidence for an unexpected

therapeutic influence. Exp Neurol 264:121-126.

Kotwas I, McGonigal A, Bastien-Toniazzo M, Bartolomei F, Micoulaud-Franchi JA (2017)

Stress regulation in drug-resistant epilepsy. Epilepsy Behav 71:39-50.

Krugel U, Fischer J, Bauer K, Sack U, Himmerich H (2014) The impact of social isolation on

immunological parameters in rats. Arch Toxicol 88:853-855.

Lapiz-Bluhm MD, Bondi CO, Doyen J, Rodriguez GA, Bedard-Arana T, Morilak DA (2008)

Behavioural assays to model cognitive and affective dimensions of depression and

anxiety in rats. J Neuroendocrinol 20:1115-1137.

Levesque M, Avoli M, Bernard C (2016) Animal models of temporal lobe epilepsy following

systemic chemoconvulsant administration. Journal of neuroscience methods 260:45-

52.

Lidster K, Jefferys JG, Blumcke I, Crunelli V, Flecknell P, Frenguelli BG, Gray WP,

Kaminski R, Pitkanen A, Ragan I, Shah M, Simonato M, Trevelyan A, Volk H,

Walker M, Yates N, Prescott MJ (2016) Opportunities for improving animal welfare

in rodent models of epilepsy and seizures. J Neurosci Methods 260:2-25.

Loscher W (2017) Animal Models of Seizures and Epilepsy: Past, Present, and Future Role

for the Discovery of Antiseizure Drugs. Neurochem Res 42:1873-1888.

Maguire J, Salpekar JA (2013) Stress, seizures, and hypothalamic-pituitary-adrenal axis

targets for the treatment of epilepsy. Epilepsy Behav 26:352-362.

Mazarati AM, Shin D, Kwon YS, Bragin A, Pineda E, Tio D, Taylor AN, Sankar R (2009)

Elevated plasma corticosterone level and depressive behavior in experimental

temporal lobe epilepsy. Neurobiol Dis 34:457-461.

Morelli E, Ghiglieri V, Pendolino V, Bagetta V, Pignataro A, Fejtova A, Costa C,

Ammassari-Teule M, Gundelfinger ED, Picconi B, Calabresi P (2014) Environmental

enrichment restores CA1 hippocampal LTP and reduces severity of seizures in

epileptic mice. Exp Neurol 261:320-327.

Mumtaz F, Khan MI, Zubair M, Dehpour AR (2018) Neurobiology and consequences of

social isolation stress in animal model-A comprehensive review. Biomed

Pharmacother 105:1205-1222.

Ngoupaye GT, Bum EN, Daniels WM (2013) Antidepressant-like effects of the aqueous

macerate of the bulb of Gladiolus dalenii Van Geel (Iridaceae) in a rat model of

epilepsy-associated depression. BMC Complement Altern Med 13:272.

Paolone G, Falcicchia C, Lovisari F, Kokaia M, Bell WJ, Fradet T, Barbieri M, Wahlberg LU,

Emerich DF, Simonato M (2019) Long-term, targeted delivery of GDNF from

encapsulated cells is neuroprotective and reduces seizures in the pilocarpine model of

epilepsy. J Neurosci.

Pellow S, Chopin P, File SE, Briley M (1985) Validation of open:closed arm entries in an

elevated plus-maze as a measure of anxiety in the rat. J Neurosci Methods 14:149-167.

Rao RM, Sadananda M (2016) Influence of State and/or Trait Anxieties of Wistar Rats in an

Anxiety Paradigm. Ann Neurosci 23:44-50.

Reger ML, Hovda DA, Giza CC (2009) Ontogeny of Rat Recognition Memory measured by

the novel object recognition task. Dev Psychobiol 51:672-678.

Shao Y, Yan G, Xuan Y, Peng H, Huang QJ, Wu R, Xu H (2015) Chronic social isolation

decreases glutamate and glutamine levels and induces oxidative stress in the rat

hippocampus. Behav Brain Res 282:201-208.

Shekh-Ahmad T, Eckel R, Dayalan Naidu S, Higgins M, Yamamoto M, Dinkova-Kostova

AT, Kovac S, Abramov AY, Walker MC (2018) KEAP1 inhibition is neuroprotective

and suppresses the development of epilepsy. Brain 141:1390-1403.

Tai TY, Warner LN, Jones TD, Jung S, Concepcion FA, Skyrud DW, Fender J, Liu Y,

Williams AD, Neumaier JF, D'Ambrosio R, Poolos NP (2017) Antiepileptic action of

c-Jun N-terminal kinase (JNK) inhibition in an animal model of temporal lobe

epilepsy. Neuroscience 349:35-47.

Veenema AH, Sijtsma B, Koolhaas JM, de Kloet ER (2005) The stress response to sensory

contact in mice: genotype effect of the stimulus animal. Psychoneuroendocrinology

30:550-557.

Vrinda M, Sasidharan A, Aparna S, Srikumar BN, Kutty BM, Shankaranarayana Rao BS

(2017) Enriched environment attenuates behavioral seizures and depression in chronic

temporal lobe epilepsy. Epilepsia 58:1148-1158.

Williams PA, White AM, Clark S, Ferraro DJ, Swiercz W, Staley KJ, Dudek FE (2009)

Development of spontaneous recurrent seizures after kainate-induced status

epilepticus. JNeurosci 29:2103-2112.

Wulsin AC, Solomon MB, Privitera MD, Danzer SC, Herman JP (2016) Hypothalamic-

pituitary-adrenocortical axis dysfunction in epilepsy. Physiol Behav 166:22-31.

Figure Legends

Figure 1: Effect of social isolation conditions on behavioral and biological parameters in mice injected or not with pilocarpine. (**A**) Experimental protocol in non-pilo and pilo mice. (**B**) The average body weight over 6 weeks, (**C**) the sweet water consumption measured during the pre and post-social isolation. (**D**) Anxiety-like behavior and memory function (**E**) are shown for isolated (IC) and social mice (SC) of the pilo and non-pilo group. (**F**) The levels of ACTH concentration, serum corticosterone and serum BDNF are shown for isolated (n= 6) and social group (n=6) of the non-pilo and pilo mice group. Data are mean ± SEM. *$p<0.05$, **$p<0.01$ and ***$p< 0.001$ in comparison with the vehicle group; [#]$p<0.05$, [##]$p<0.01$ and [###]$p<0.001$ in comparison with social condition group. [§§]$p<0.01$ and [§§§]$p<0.001$ compared to the baseline.

Figure 2: Effect of social isolation conditions on behavioral and biological parameters in rats. (**A**) Experimental protocol in non-pilo and pilo rat groups. (**B**) The average body weight over seven weeks, (**C**) the sweet water consumption measured during the first and last week, and (**D**) the levels of ACTH concentration, serum corticosterone and serum BDNF levels are shown for isolated (n= 5), interaction (handled, n= 5) and paired (n= 10 [5 couples]) rats of the non-pilo group. The isolated group displayed strong behavioral and biological alterations as compared to the two other groups. Data are mean ± SEM. *$p<0.05$, **$p<0.01$ and ***$p<0.001$ in comparison with the isolated group; [#]$p<0.05$, [##]$p<0.01$ and [###]$p<0.001$ in comparison with the handled group.

Figure 3: **Effect of social isolation conditions on spontaneous seizures in rats**. The average seizure frequency (number of seizures per hour), duration and severity are shown for isolated, interaction (handled) and paired rats (n=11, n=12 and n=6) of the pilo group. Isolated rats

displayed a very severe epileptic phenotype as compared to the other groups. Values are

expressed as mean ± SEM. *$p<0.05$, **$p<0.01$ and ***$p<0.001$ in comparison with the

isolated group.

Figure 3-1: Distribution of seizures per week and for each animal. Top left panel. The

average seizure frequency (seizures per hour) over the three weeks of EEG recording is

shown for isolated (n=11), handled (n=12) and paired rats (n=6). No significant difference

was shown between weeks in seizure frequency for all experimental groups. Other panels

show the average seizure frequency per week for each animal. Two animals (out of 11) in the

isolated group had a seizure frequency similar to that found in paired or handled animals

(0.03/h).

Data are mean ± SEM. ANOVA two way followed by Bonferroni *post-hoc,* *$p<0.05$ and

**$p<0.01$ refers to handled or paired groups vs. isolated group comparison.

FIGURE 1

A

B **C**

D

FIGURE 2

FIGURE 3

A- Statistical analysis of the effects of mice groups (grouped/isolated) and treatment (Veh and pilo) on behavioral, biological and biochemical parameters:

DATA	DATA STRUCTURE (Normality test)	TYPE OF TEST	POWER
Body weight (a)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F(3,20) = 2.23$, $P = 0.11$ Time effect: $F(5,100) = 128.08$, $P < 0.0001$ Interaction: $F(15,100) = 1.18$, $P = 0.29$ (CI = 95%)
EPM (b)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F(3,40) = 12.93$, $P < 0.0001$ Time effect: $F(1,40) = 64.32$, $P < 0.0001$ Interaction: $F(3,40) = 11.84$, $P < 0.0001$ (CI = 95%)
NOR (c)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F(3,40) = 6.95$, $P = 0.0007$ Time effect: $F(1,40) = 43.30$, $P < 0.0001$ Interaction: $F(3,40) = 8.89$, $P < 0.0001$ (CI = 95%)

ACTH (d)	yes	Two way ANOVA Post hoc test : Bonferroni test	Group effect: $F(1,20) = 20.55$, $P = 0.0002$ Treatment effect: $F(1,20) = 74.62$, $P < 0.0001$ Interaction: $F(1,20) = 0.09$, $P = 0.75$ (CI = 95%)
Corticosterone (e)	yes	Two way ANOVA Post hoc test :	Group effect: $F(1,20) = 18.56$,

		Bonferroni test	P = 0.0003 Treatment effect: $F_{(1,20)} = 117.3$, P < 0.0001 Interaction: $F_{(1,20)} = 0.006$, P = 0.93 (CI = 95%)
BDNF (f)	yes	Two way ANOVA Post hoc test : Tukey test	Group effect: $F_{(1,20)} = 16.96$, P = 0.0005 Treatment effect: $F_{(1,20)} = 0.4696$, P = 0.5011 Interaction: $F_{(1,20)} = 0.0002$, P = 0.9889 (CI = 95%)

B-Statistical analysis of the group (1) and day effects (2) on sucrose preference in mice using non-parametric *Mann–Whitney U* test after 4 weeks of isolation.

B-1	Non pilo-IC vs Non pilo-SC	Pilo-IC vs Pilo-SC	Pilo-SC vs Non pilo-SC	Pilo-IC vs Non pilo-IC
Day 1	U= 2.00; Z= -4.87; P= 0.0108 *	U= 1.00; Z=-4.42; P=0.0043**	U= 0.00 ; Z=-10.89; P= 0.0022**	U=0.00 ; Z=-10.23; p=0.0022**
Day 2	U= 4.00; Z= -3.96; P= 0.0238*	U= 2.50; Z= -5.56; P=0.0108*	U= 0.00 ; Z= -10.63 P= 0.0022**	U=0.00; Z=- 12.49 p= 0.0022**
Day 3	U= 3.00; Z=- 4.75 P=0.0152*	U= 4.5; Z= -4.98; P= 0.0303*	U= 0.00 ; Z= -10.17; P= 0.0022**	U= 0.00; Z= -10.20 p=0.002**
Day 4	U= 2.50; Z= -3.31 P= 0.0108*	U= 2.00 ; Z=-4.60; P=0.0087**	U= 0.00; Z=- 12.37; P= 0.002**	U= 0.00; Z=-11.77 p=0.002**
Day 5	U= 4.00; Z=-3.91 P=0.0238*	U= 2.00;Z=-4.44; P=0.0108 *	U=0.00; Z=-10.84; P= 0.0022**	U= 0.00; Z=-11.74; p= 0.0022**
Day 6	U= 0.00; Z=-5.84 P= 0.0022**	U= 5.00 ; Z=-4.06; P=0.0390*	U= 0.00 ; Z=-11.21 P= 0.0022**	U= 0.00; Z=-9.82; p= 0.0022**

Day 7	U= 1.50; Z= -4.72 P= 0.0065**	U=5.00;Z=-5.00; P=0.0390*	U= 0.00; Z=-9.35 P= 0.0022**	U= 0.00; Z=-10.09 p= 0.0022**

B-2	Day 2 vs day 1	Day 3 vs day 1	Day 4 vs day 1	Day 5 vs day 1	Day 6 vs day 1	Day 7 vs day 1
Non-pilo-SC	U= 14.50 ; Z=0.11 ; P=0.6147	U=16.50 : Z=-0.26 ; P=0.8528	U=15.00 ; Z=-0.27 ; P=0.6861	U=16.50 ; Z=-0.13 ; P=0.8355	U=15.00 ; Z=-0.27 ; P=0.6688	U=16.50 ; Z=-0.16 ; P=0.8550
Non-Pilo-IC	U=14.00 ; Z=1.18 ; P=0.5714	U=16.00 ; Z=-0.34 ; p=0.7900	U=13.00 ; Z= 0.90 ; P=0.4567	U=14.50 ; Z=0.72 ; P= 0.6234	U=16.50 ; Z=-0.40 ; P=0.8506	U=17.50 ; Z=0.09 ; P=0.9848
Pilo-SC	U= 14.50 ; Z= 0.53 P= 0.6212	U=16.50 ; Z= 0.37 ; P=0.8550	U=14.00 ; Z=-1.19 ; P= 0.5584	U=16.00 ; Z= -0.22 ; P= 0.7944	U=17.50 ; Z=0.00 ; P= 0.9784	U=13.00 ; Z= 0.84 : P= 0.4632
Pilo-IC	U=16.00 ; Z= -0.45 ; P= 0.7965	U=16.50 ; Z= 0.20 ; P= 0.8528	U=12.00 ; Z=-0.92 P= 0.3896	U=15.50 ; Z=-0.11 ; P=0.7403	U=15.00 ; Z= 0.33 ; P= 0.6732	U=15.00 ; Z=0.32 ; P=0.6753

C: Statistical analysis of the effects of rat groups (isolated/handled/paired) on behavioral, biological, biochemical and parameters

DATA	DATA STRUCTURE (Normality test)	TYPE OF TEST	POWER
Body weight (g)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F_{(2,17)} = 41.58$, $P < 0.0001$ Time effect: $F_{(6,102)} = 266.83$, $P < 0.0001$ Interaction: $F_{(12,102)} = 2.10$, $P = 0.02$ (CI = 95%)
Anhedonia First week (h)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F_{(2,12)} = 21.66$, $P < 0.0001$ Time effect: $F_{(6,72)} = 0.11$, $P = 0.99$ Interaction: $F_{(12,102)} = 0.16$, $P = 0.99$ (CI = 95%)
Anhedonia Last week (i)	yes	ANOVA with repeated measures Post hoc test :Bonferroni test	Group effect: $F_{(2,12)} = 97.85$, $P < 0.0001$ Time effect: $F_{(6,72)} = 1.09$,

			P = 0.37 Interaction: F(12,72) = 0.13, P = 0.99 (CI = 95%)
ACTH (j)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,17) = 5.05, P= 0.019 (CI = 95%)
Corticosterone (k)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,17) = 8.42, P = 0.0029 (CI = 95%)
BDNF (l)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,17) = 6.18, P = 0.0096 (CI = 95%)
Seizure frequency (m)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,26) = 10.73, P = 0.0004 (CI = 95%)
Seizure duration (n)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,26) = 30.90, P < 0.0001 (CI = 95%)
Seizure severity (o)	yes	One way ANOVA Post hoc test : Bonferroni test	F(2,26) = 4.07, P = 0.028 (CI = 95%)