1

3   High genomic diversity of multi-drug resistant wastewater *Escherichia coli*

5   Norhan Mahfouz[1,*], Serena Caucci[2,3,*], Eric Achatz[1], Torsten Semmler[4], Sebastian Guenther[4],

6   Thomas U. Berendonk[2,*], and Michael Schroeder[1,*,#]

[1] Biotec, TU Dresden

[2] Institute for Hydrobiology, TU Dresden

[3] United Nations University Institute for Integrated Management of Material Fluxes and of

Resources

[4] Institute of Microbiology und Epizootics, FU Berlin

* These authors contributed equally

Correspondence: Michael Schroeder, ms@biotec.tu-dresden.de

Keywords: Antibiotic Resistance, Wastewater Treatment, Pan-Core genome, Environment

Conflict of interest statement: The authors declare no conflict of interest.

Abstract

Wastewater treatment plants play an important role in the release of antibiotic resistance into the

environment. It has been shown that wastewater contains multi-drug resistant *Escherichia coli*,

but information on strain diversity is surprisingly scarce. Here we present an exceptionally large

dataset on multidrug resistant *Escherichia coli*, originating from wastewater, over a thousand

isolates were phenotypically characterized for twenty antibiotics and for 103 isolates whole

genomes were sequenced. To our knowledge this is the first study documenting such a

comprehensive diversity of multi-drug resistant *Escherichia coli* in wastewater. The genomic

diversity of the isolates was unexpectedly high and contained a high number of resistance and

virulence genes. To illustrate the genomic diversity of the isolates we calculated the pan genome

of the wastewater *Escherichia coli* and found it to contain over sixteen thousand genes. To

analyse this diverse dataset, we devised a computational approach correlating genotypic variation

and resistance phenotype, this way we were able to identify not only known, but also candidate

resistance genes. Finally, we could verify that the effluent of a wastewater treatment plant will

contain multi-drug resistant *Escherichia coli* belonging to clinically important clonal groups.

Introduction

In 1945, Alexander Fleming, the discoverer of Penicillin, warned of antibiotic resistance. Today, the WHO echoes this warning, calling antibiotic resistance a global threat to human health. Humans are at the center of the modern rise of resistance. The human gut [1], clinical samples [2,3], soil [4,5], and wastewater [6] all harbor resistant bacteria and resistance genes. At the heart of modern resistance development is a human-centred network of clinics, industry, private homes, farming, and wastewater. Recent studies suggest that wastewater contains a significant amount of antibiotic resistant *Escherichia coli*, specifically extended-spectrum beta-lactamase-producing *Escherichia coli* [7]. Particularly, multidrug-resistant (MDR) clones (normally defined as those resistant to three or more drug classes) are of great concern. Past studies have documented the presence of MDR *Escherichia coli* isolates in wastewater on the basis of phenotypic resistance testing [8], but a comprehensive analysis of the clonal composition of MDR *Escherichia coli* in wastewater employing whole genome analysis is largely lacking. Therefore the current information on the genomic diversity of antibiotic resistant *Escherichia coli* in wastewater is very limited. Recent metagenomic studies have documented that human-associated bacteria are strongly reduced in the wastewater and its treatment process [9]. To investigate the genomic diversity as well as virulence genes and resistance determinants for wastewater *Escherichia coli*, we proceeded as sketched in Fig. 1: We collected 1178 *Escherichia coli* isolates from a waste treatment plant's inflow and outflow in the city of Dresden, Germany. We selected 20 antibiotics, which are the most prescribed ones in the area from which the wastewater inflow originates (data provided by the public health insurer AOK). We analyzed the isolates' resistance to these 20 antibiotics and selected 103 isolates for whole genome sequencing. Our analysis reveals a surprisingly high genomic diversity of MDR *Escherichia coli* in the wastewater with very flexible genomes harboring a high variation of virulence genes and resistance determinants. Using this diversity we developed a computational approach to identify not only known, but also novel candidate resistance genes.

Figure 1: Wastewater plays an important role in antibiotic resistance development. Wastewater *Escherichia coli* isolates are tested for antibiotic resistance and sequenced. Many isolates are multi-drug resistant and potentially pathogenic. Their large pan-genome is a source of potentially novel resistance genes.

68
69

Results

The wastewater pan-genome. The concept of evolution implies that genomes of organisms of

the same species differ. Differences range from small single nucleotide polymorphisms to large

genome rearrangements. As a consequence, *Escherichia coli* possesses a core of genes present

in all genomes, as well as genes only present in some genomes, or even just in one. The union of

all of these genes is called the pan-genome. It is believed, that the *Escherichia coli* core genome

comprises around 1400-1500 genes, while the pan-genome may be of infinite size [10].

To assess the degree of genomic flexibility of the wastewater isolates, we relate the wastewater

pan-genome and the wastewater core genome. At 16582 genes, the wastewater pan-genome is

nearly six times larger than the wastewater core genome of 2783 genes, a reservoir of some

14000 genes. Despite this large reservoir, the size difference of nearly 1000 genes between the

wastewater *Escherichia coli* core genome and the whole species core genome suggests that the

full diversity of *Escherichia coli* is still not covered in our wastewater sample.

The balance between maintaining the core genome and spending energy on acquisition of new

genetic material can be captured by the ratio of the core genome size and the average genome

size, which is 4700 genes in our sample. This means that only 1400/4700 = 30% of genes in our

wastewater *Escherichia coli* are core genes. Most of the non-core genes are very unique and

appear only in one or two isolates each. More precisely, 50% of the pan-genome genes appear in

only one or two isolates each. This implies that the investigated wastewater *Escherichia coli* are

highly individual.

This high diversity is also illustrated in Fig. 2, which compares the wastewater *Escherichia coli* to

a clinical dataset of *Escherichia coli.* The figure clearly shows that the *Escherichia coli* of clinical

origin are more homogeneous and hence their pan-genome is smaller. In contrast, the diversity of

the wastewater *Escherichia coli* match other datasets comprising mixtures of commensal and

pathogenic *Escherichia coli*, as well as *Shigella* genomes (see Table 1). This underlines the great

diversity of *Escherichia coli* genomes in the wastewater. Interestingly, the variation of the

wastewater genomes after the treatment plant was not reduced.

Figure 2: The pan-genome at the outflow has the same size as at the inflow, suggesting that highly flexible
Escherichia coli emerge from a treatment plant. The wastewater pan-genome is larger than a clinical pan-
genome one and of similar size to (see Table 1) highly diverse samples comprising pathogenic,
commensal, and lab *Escherichia coli*, as well as *Shigella*.

Ref	Pan	Core	Strains	Path.	Comm.	Lab	*Shig.*
This study	16582	2783	92	28	62	0	0
Kaas et al., 2012[11]	16373	1702	186		171		15
Vieira et al., 2011[12]	14986	1957	29	21	8	0	6
Gordienko et al., 2013[13]	12000	2000	32	16	6	3	7
Lukjancenko et al., 2010[14]	13000	1472	53	35	11	7	0
Rasko et al., 2008 [15]	13000	2344	17	14	1	2	0
Touchon et al., 2009[16]	11432	1976	20	10	3	0	7

108   **Table 1 :** Highly diverse samples comprising pathogenic, commensal, and lab *Escherichia coli*, as well as
109   *Shigella.*

Resistance genes in the wastewater pan-genome. Wastewater *Escherichia coli* are known to
host antibiotic resistance genes. While there are many known resistance genes (see e.g. CARD
[17]), they fall mostly into a few groups, such as beta-lactamases. Here, we seek to confirm and
expand the space for candidate resistance genes. Firstly, we measured antibiotic resistance in all
1178 isolates to the 20 antibiotics. As a positive control we included also two antibiotics to which
at least clinical *E. coli* are reported to be inherently resistant (kanamycin and cephalotin). Fig. 3
reveals a high degree of resistance and big differences between different antibiotics, including a
general trend indicating greater resistance to antibiotics that have been available for longer.
Specifically, antibiotics from the 50s and 60s have a significantly different number of resistances
than the more recent antibiotics (Welch test, p-value < 0.0025, also significant without including
kanamycin and cephalotin). However, there is no significant difference in the number of
resistances between isolates from the inflow and the outflow (p-value 0.0001), suggesting that
wastewater treatment is not affecting resistance.

Figure 3: 1178 Wastewater *Escherichia coli* isolates are tested for antibiotic resistance to 20 antibiotics. The antibiotics kanamycin and cephalotin were included as a positive control as *E. coli* is reported to be inherently resistant to those antibiotics. Nearly all isolates are multi-drug resistant. Generally, isolates are more susceptible to betalactams and fluoroquinolones than to tetracyclins and aminoglycosides. Surprisingly, the outflow isolates show similar resistance as inflow (p-value 0.0001), suggesting that wastewater treatment is not reducing resistance development.

Next, we correlated the presence of each gene in the sequenced isolates with their phenotypic

antibiotic resistance profiles. We excluded meropenem and imipenem, since nearly all isolates

are susceptible. For each of the 18 remaining antibiotics, we list the top ten candidate resistance

genes in Table 2. These 180 genes comprise 88 unique confirmed genes, including many well-

known resistance genes, such as efflux pumps (MT1297 and *emr*E), membrane and transport

proteins (*aida*-I*, yia*V*, yij*K*, pit*A*, ics*A, and *pag*N), tetracycline (*tet*A, *tet*R, and *tet*C),

chloramphenicol (*cat*), and piperacillin (the beta lactamase *bla*2) resistance genes. However, the

180 genes also comprise a large number of open reading frames encoding hypothetical proteins

(41) and genes not yet linked to antibiotic resistance (116). These genes have to be studied

further to determine whether they are novel resistance genes or just correlating (e.g. because

they are on the same genetic element with a resistance gene). Nearly all of the identified genes

are found both in inflow and outflow genomes suggesting that the waste water treatment does not

impact on the presence or absence of known and candidate resistance genes.

#	Amikacin	Gentamicin	Kanamycin	Tobramycin	Doxycycline	Tetracycline	Cefepime	Cefotaxime	Ceftazidime	Cefuroxime Sod.	Cephalotin	Piperacillin	Ciprofloxacin	Levofloxacin	Nalidixic Acid	Norfloxacin	Chloramphenicol	Fosfomycin
1	Hypothetical Protein	4-hydroxyacetophenone monooxygenase hapE	Transposase IS200 like protein	Autotransporter precursor aida-I	Tetracycline resistance protein, class B tetA	Oxygen-dependent choline dehydrogenase betA	Ash protein family protein	Hypothetical Protein	cell division protein	Type-1 restriction enzyme R protein hsdR	GTPase era	Beta-lactamase TEM precursor bla	Virulence regulon transcriptional activator virB	Transposon Tn10 protein tetD	Mercuric resistance operon regulatory protein merR	Transposon Tn10 protein tetD	Chloramphenicol acetyltransferase cat	Invasin
2	Caudovirales tail fiber assembly protein	Phosphoadenosine phosphosulfate reductases	putative multidrug-efflux transporter/MT1297	putative protease yhbU precursor	Tetracycline repressor protein class B tetR	NAD/NADP-dependent betaine aldehyde dehydrogenase betB	Fibronectin type III protein	Hypothetical Protein	Plasmid stability protein	Type I restriction enzyme EcoKI M protein hsdM	Prophage CP4-57 regulatory protein alpA	Transposon Tn3 resolvase tnpR	Sporulation initiation inhibitor protein Soj	Tetracycline resistance protein, class B tetA_1	Mercuric resistance protein merC	Tetracycline resistance protein, class B tetA_1	Streptomycin 3''-adenylyltransferase ant1	Putative DNA-invertase Rac pinR
3	Swarming motility protein ybiA	putative multidrug-efflux transporter/MT1297	Phosphotransferase enzyme family protein	Chaperone protein dnaK	Transposon Tn10 TetC protein tetC	HTH-type transcriptional regulator betI	Transcriptional activator perC	Transcriptional activator perC	HTH-type transcriptional regulator cmtR	mrr restriction system protein	Hypothetical Protein	Tyrosine recombinase xerD	putative HTH-type transcriptional regulator	Tetracycline repressor protein class B from transposon Tn10 tetR	mercuric transport protein merT	Tetracycline repressor protein class B from transposon Tn10 tetR	Chromosome-partitioning ATPase soj	Transcriptional repressor dicA
4	Phospholipase ytpA	Phosphotransferase enzyme family protein	Hypothetical Protein	putative ABC transporter ATP-binding protein yjjK	HTH-type transcriptional regulator cmtR	Tetracycline resistance protein, class B tetA	Hypothetical Protein	Hypothetical Protein	Phage-related minor tail protein	Outer membrane protein lcsA precursor	Hypothetical Protein	Acetyltransferase (GNAT) family protein	DNA-binding transcriptional regulator dicC	Transposon Tn10 protein tetC	Mercuric transport protein periplasmic component precursor merP	Transposon Tn10 protein tetC	parG	Hypothetical Protein
5	Carbonic anhydrase 1 cynT	Hypothetical Protein	Streptomycin 3''-adenylyltransferase ant1	cell envelope integrity inner membrane protein tolA	Tetracycline resistance protein, class C tetA	Tetracycline repressor protein class B tetR	Hypothetical Protein	Hypothetical Protein	Phage tail protein E	Hypothetical Protein	Hypothetical Protein	Virulence regulon transcriptional activator virB	Hypothetical protein	putative HTH-type transcriptional regulator	Anti-adapter protein iraM	CAAX amino terminal protease self-immunity	Hypothetical Protein	Hypothetical Protein
6	Hypothetical Protein	Hypothetical Protein	Hypothetical Protein	Inner membrane protein yiaV precursor	putative inner membrane transporter yedA	Transposon Tn10 TetC protein tetC	Chromosome partition protein smc	Hypothetical Protein	Hypothetical Protein	Fibronectin type III protein	Transposon Tn10 tetD protein	Transposase	Lysine--tRNA ligase lysS	DNA-binding transcriptional regulator dicC	Hypothetical protein	mRNA interferase pemK	Hypothetical Protein	Hypothetical Protein
7	Xanthine dehydrogenase molybdenum-binding subunit xdhA	Hypothetical Protein	Zinc-responsive transcriptional regulator	Entericidin B membrane lipoprotein	Tetracycline repressor protein class A from transposon 1721 tetR	High-affinity choline transport protein betT	Hypothetical Protein	Invasin	Hypothetical Protein	Hypothetical Protein	putative multidrug-efflux transporter/MT1297	Tetracycline resistance protein, class B tetA	Transposon Tn10 protein tetD	Hypothetical protein	Mercuric reductase merA_1	Antitoxin pemI	Acetyltransferase (GNAT) family protein	Molybdenum cofactor biosynthesis protein A
8	Nicotinate dehydrogenase FAD-subunit ndhF	Hypothetical Protein	merE protein	Low-affinity inorganic phosphate transporter 1 pitA	Hypothetical Protein	Formate dehydrogenase H fdhF	Aldehyde-alcohol dehydrogenase adhE	Hypothetical Protein	Tyrosine recombinase xerC	Hypothetical Protein	Phosphotransferase enzyme family protein	Tetracycline repressor protein class B tetR	Tetracycline resistance protein, class B tetA_1	CAAX amino terminal protease self-immunity	Hypothetical protein	putative HTH-type transcriptional regulator	putative multidrug-efflux transporter/MT1297	ATP-dependent zinc metalloprotease ftsH4
9	Nicotinate dehydrogenase small FeS subunit ndhS	Phage polarity suppression protein psu	Phosphoadenosine phosphosulfate reductases	Methyl-accepting chemotaxis protein II tar	Transposon Tn10 tetD protein	S-fimbrial protein subunit sfaH	Aldehyde-alcohol dehydrogenase adhE	Hypothetical Protein	Hypothetical Protein	Hypothetical Protein	Outer membrane protein pagN precursor	Transposon Tn10 tetC protein	Tetracycline repressor protein class B transposon Tn10 tetR	mRNA interferase pemK	zinc-responsive transcriptional regulator	DNA-binding transcriptional regulator dicC	Phosphotransferase enzyme family protein	Molybdenum cofactor biosynthesis protein A
10	putative fimbrial-like protein ElfG precursor elfG	DNA primase traC	Caudovirales tail fiber assembly protein	Leucine-specific-binding protein precursor livK	putative multidrug-efflux transporter/MT1297	Beta-lactamase TEM precursor bla	Cob(I)yrinic acid a,c-diamide adenosyltransferase yvqK	Type-1 restriction enzyme R protein hsdR	Hypothetical Protein	Hypothetical Protein	Tetracycline resistance protein, class B tetA	Multidrug transporter emrE	Transposon Tn10 protein TetC	Antitoxin PemI	MerE protein	Caudovirales tail fiber assembly protein	Leucine-specific-binding protein precursor livK	Hypothetical Protein

Table 2: Known and candidate resistance genes from correlation of genomes to resistance phenotype. Top 10 genes for 18 antibiotics.

Virulence genes Generally, *Escherichia coli* strains exhibit great variation. Many exist as

harmless commensals in the human gut, but some are classified as intra- (InPEC) or extra-

intestinal pathogenic *Escherichia coli* (ExPEC [18]). Based on their virulence genes profile the

pathogenic potential of Escherichia coli isolates can be determined[7]. The sequenced isolates

contain some 700 of the 2000 *Escherichia coli* virulence factors in the virulence factor database

[19], averaging to 153 and to 155 virulence factors per isolate for inflow and outflow, respectively.

Hence, there is no significant difference (Welch test, CI 95%) between inflow and outflow. In

particular, we found combinations of virulence factors for 16 isolates (see methods), which are

indicative of ExPEC. Eight of these 16 isolates were obtained from the outflow of the treatment

plant (see Fig. 4).

Besides the presence of known virulence factors, the pathogenic potential can be assessed using

genotyping with multi-locus sequence types [20] and phylogroups [21]. Broadly, *Escherichia coli* has,

among other s, four phylogroups, A, B1, B2 and D. Commensal *Escherichia coli* fall mostly into

groups A and B1 and ExPEC into B2 and D [21]. Fig. 4 shows a phylogenetic tree of the sequenced

wastewater *Escherichia coli* isolates along with the commensal phylogroups A (red) and B1 (blue)

and the pathogenicity-associated groups B2 (yellow) and D (green), as well as the finer-grained

multi-locus sequence types. The tree is based on genomic variations compared to the reference

genome of *Escherichia coli* K12 MG1655. Fig. 4 reveals that nearly one third of isolates belong to

group B2 and D, in which ExPEC are usually found. In particular, B2 and D include 14 of the 16

potential ExPEC isolates. Remarkably, half of the B2 and D isolates are from the wastewater

treatment plant's outflow.

Figure 4: Phylogeny and pathogenic potential of wastewater *Escherichia coli*. Phylogenetic tree, multi-locus sequence types, and phylogroups of 92 sequenced wastewater *Escherichia coli* isolates reveal 16 potential ExPEC isolates (marked with a black star) in phylogroups B2 (yellow) and D (green), which are associated with pathogenicity. Half of the potentially pathogenic isolates stem from the outflow of the treatment plant.

Discussion

Pan and core genome.

It is well known that wastewater treatment reduces the bacterial abundance, in addition a recent metagenomic study has shown that the bacterial community in wastewater is very different to the human gut community and that the number of detected genera is reduced in the wastewater[9]. Consequently, our expectation was that the genomic diversity of *Escherichia coli* should be reduced. We were very surprised to find an unexpectedly high genomic diversity, which is illustrated in the large pangenome. A possible explanation for this high genomic diversity is that the *Escherichia coli* cells within the wastewater originate not only from human faeces, but also from a multitude of different animal faeces collected via the surface runoff into the sewers. This would also explain why the pangenome of the wastewater *Escherichia coli* is considerably larger than the clinical pangenome reported by Land et al.[22]. Generally, many authors have pointed out that *Escherichia coli* has a large and flexible pan genome. Lapierre *et al.* argue that *Escherichia coli* appears to have unlimited ability to absorb genetic material and hence its pan genome is open [10]. In a recent study comprising over 2000 genomes Land *et al.* put this into numbers and arrive at a pan genome of 60000-89000 gene families for over 2000 sequenced *Escherichia coli* genomes [22]. The study by Land *et al.* (24) is based on clinical isolates, in contrast our study is the first, which has calculated the pangenome of *Escherichia coli* for wastewater. Interestingly, our results seem to be in concordance and suggest that within our study we still have not reached the saturation of the detected diversity (Fig. 2), indicating that the full genomic diversity of *Escherichia coli* in the wastewater is probably even larger than what we report here. Worryingly, this is also reflected in a high diversity of resistance and virulence genes. This documents that the wastewater contains a significant amount of multi-drug resistant (MDR) *Escherichia coli,* which also carry a suit of virulence genes suggesting that some of those MDR have a pathogenic potential. Furthermore, we did not find a significant difference in genomic diversity between inflow and outflow of the wastewater treatment plant, suggesting that selection against genome diversity and resistance determinants does not seem to occur.

Pathogenic potential and resistance. Resistant bacteria may or may not be pathogenic. While ultimate proof for pathogenicity can only be obtained from in vivo studies, we wanted to understand the pathogenic potential of the isolates by analysing the genome for suitable markers. Here we chose to consider three independent approaches: classification by phylogenetic groups, by multi-locus sequence tags, and by identification of specific virulence factors (see methods). While the three approaches showed consistent results, they are by no means proof for pathogenicity, since there can be exceptions to these classification rules. As an example, consider the strain ED1a (O81), which was isolated from a healthy man, but belongs to the phylogenetic group B2 [16]. Similarly, pathogenicity may not only arise from the acquisition of genes, but also from the loss [23].

Regarding resistance there are similar confounding factors. *Escherichia coli* is inherently resistant to kanamycin and cephalotin, which is also clearly shown in Fig. 3. This supports the notion that, generally antibiotic resistance is ancient [24] and naturally occurring in the environment. Nonetheless, there are pronounced differences between pristine and human environments [25]. This is also supported by Fig. 3, which shows that antibiotics introduced in the 60s have more resistances than those introduced later (p-value < 0.0025), which suggests, that the naturally occurring resistances do not play a major role in the emergence of observed resistances.

From clinic to river. We have shown that there are *Escherichia coli* at the wastewater outflow, which are multi-drug resistant and have pathogenic potential. But are they abundant enough to have an impact in the aquatic system they are released into? They do. The percentage of possibly pathogenic *Escherichia coli* in the outflow is considerable and may correspond to a large absolute amount. If an average of 100 *Escherichia coli* colony forming units (CFU) are released per ml, then 10^{13} CFUs per day are released (assuming a release of $10^5\,m^3$ per day). This is in accordance with Manaia *et al.*, who showed that 10^{10}-10^{14} CFU of ciprofloxacin-resistant bacteria are released by a mid-sized wastewater treatment plant [26]. Supporting these results, a study in a Japanese river shows the presence of pathogenic *Escherichia coli* [27]. In this study they sequenced over 500 samples from the Yamato river and most of their prevalent multi-drug resistant and clinical strains are also present in our samples. In a related study, Czekalski *et al.*

found that particle-associated wastewater bacteria are the responsible source for antibiotic

resistance genes in the sediments of lake Geneva in Switzerland [28]. Assuming that the river Elbe

is comparable to these aquatic systems, it suggests, that the urban environment (including clinics)

and river are connected with wastewater treatment plants in between.

Composition of phylogroups. It is interesting to compare the breakdown into phylogenetic

groups of wastewater *Escherichia coli* to compare samples from human and animal

environments. It is, e.g., known that the phylogenetic group B2 is more abundant among

commensal *Escherichia coli* from human faeces (43%) than from farm animals (11%) [29].

Therefore, the composition of wastewater *Escherichia coli* as shown in Fig. 4 resembles

commensal *Escherichia coli* from farm animals more closely. Similarly, Tenaillon *et al.* find that

groups A and B1 make up one third in human faeces [29], whereas we find two thirds. This

suggests that animal feces play an important role for resistance also of urban wastewater

treatment plants and this is probably part of the explanation for the high observed genomic

diversity.

Random sampling and novel resistance mechanisms. The initial 1178 isolates were sampled

randomly over different times of the year, from two different inflows and the outflow of the

wastewater treatment plant. In contrast, the 103 sequenced isolates were chosen in such way

that all of the phenotypes encountered were represented (see methods). Within a phenotype

group isolates were chosen randomly. This random, but representative choice and the

subsequent link from genotype to phenotype is an example of high-throughput hypothesis-free

analysis. And although, there was no pre-defined resistance mechanism, which we aimed to hit,

many of the well-known resistance genes were ranked high. This supports the hope that high-

throughput, hypothesis-free methods such as deep sequencing will help to uncover novel

resistance mechanisms and in particular that some of the candidate resistance genes will prove

to have a causal link to resistance. The results show that the here outlined computational

approach to correlate genomic and phenotypic information for wastewater *Escherichia coli*

significantly assists to identify a larger part of the existing resistome of *Escherichia coli*.

Conclusion

Overall, we have shown for the first time that *Escherichia coli* isolates from wastewater have a surprisingly large pan-genome, which harbors virulence genes, known and novel candidate resistance genes. We developed a computational approach based on genomic and phenotypic correlation for *Escherichia coli* and show that applying this to wastewater will discover novel parts of the resistome in *Escherichia coli*. Finally, together with the estimates on absolute *Escherichia coli* abundance, we could demonstrate that there is a considerable pathogenic potential in the outflow of a wastewater treatment plant. Using *Escherichia coli* as an example, this study demonstrates the importance of investigating wastewater with modern bioinformatics and strain specific genomic analysis in order to estimate the extent of genomic variation and resistance determinants for bacteria with clinical relevance present in the environment.

Methods

Collection. 1178 samples were collected from the municipal wastewater treatment plant

Dresden, Germany. Samples were collected on 11/4/2012 (Spring 2012), 30/7/2012 (Summer

2012), 21/1/2013 (Winter 2012), 27/3/2013 (Spring 2013), 6/8/2013 (Summer 2013), 14/10/2013

(Autumn 2013), and 17/12/2013 (Winter 2013). Samples were collected either at the outflow

(OUT) or at one of two inflow locations (Altstadt ALT and Neutstadt NEU), representing the area

south and north of the river Elbe).

Isolation. *Escherichia coli* and total coliforms bacteria were enumerated via serial fold dilution

plating of the original wastewater (triplicate samples). Wastewaters were diluted in double distilled

water, until the enumeration of bacterial colonies was possible. *Escherichia coli* and coliform

counts were always performed in triplicates. The *Escherichia coli* colonies were selected and

picked after overnight growth at 37°C on a selective chromogenic media (OXOID Brilliance

Escherichia coli/Coliform Selective Agar, Basingstoke, England). To minimize the risk of colony

contamination, picked colonies were spiked a second time on the same selective media and pure

single colonies were grown overnight on LB media at 37°C and stored on glycerol stock at -80° C.

Resistance phenotyping. Antibiotic resistance phenotypes were determined by the agar

diffusion method using 20 antibiotic discs (OXOID, England) according to EUCAST (or CLSI

when EUCAST was not available) [7,8]. The selected drugs belong to the most commonly

prescribed antibiotics for diseases caused by bacteria according to the German health insurance

AOK Plus: piperacillin ($100\mu g$), nalidixic acid ($30\mu g$), chloramphenicol ($30\mu g$), imipenem ($10\mu g$),

cefotaxime ($30\mu g$), cephalotin ($30\mu g$), kanamycin ($30\mu g$), tetracycline ($30\mu g$), gentamicin ($10\mu g$),

amikacin ($30\mu g$), ciprofloxacin ($5\mu g$), fosfomycin ($50\mu g$), doxycycline ($30\mu g$), cefepime ($30\mu g$),

ceftazidime ($10\mu g$), levofloxacin ($5\mu g$), meropenem ($10\mu g$), norfloxacin ($10\mu g$), cefuroxime sod.

($30\mu g$), tobramycin ($10\mu g$) [30]. After 24 hours of incubation at 37°C, the resistance diameters were

measured. Clustering of antibiotics and of isolates was performed using the R function heatmap.2

from the R library [31] Heatplus and hierarchical clustering of matrices based on Euclidean

distances between isolates and between antibiotics.

Sequencing. To select isolates representative of phenotype, we clustered isolates according to

the diameters of inhibition zone against the 20 antibiotics using k-means clustering based on

Euclidean distances between isolates (vectors of 20 inhibition zone diameters). The analysis and

graphs were produced using R version 3.2.4 [31]. As clusters may be highly skewed in number of

cluster members, we tested all cluster numbers from 1 to 100 and plotted within class sum of

squares against k. At $k = 47$, the sum of squares tails off and there is a steep local decrease, so

that $k = 47$ was fixed as k-means parameter. We obtained 103 isolates, which were subsequently

used for sequencing and further analysis. To further validate the choice, we plotted the average

number of resistances against number of isolates and antibiotics vs. number of isolates for the

total 1178 and the selected 103 isolates (see Supp Fig. 1) and concluded that both distributions

are roughly similar. 3000ng DNA were extracted from each of the 103 selected isolates using

MasterPure extraction kit (Epicentre) according to the manufacturer's instructions. Sequencing

was performed using Illumina Flex GL.

Assembly. Genomes were assembled with Abyss (version 1.5.2) [32]. In order to optimize k for the

best assembly, k-mer values had to be empirically selected from the range of 20-48 (see Supp.

Fig. 2) on a per sample basis to maximize contiguity [3]. To determine the k-mer length that

achieved highest contiguity, the 28 assemblies per draft genome/isolate were compared based on

N50 values. 11 assemblies with an *N50* statistic of less than 5×10^4 bp were excluded [33].

Genes. Reference gene clusters were computed from 58 complete *Escherichia coli* genomes

(see Table 2) available in June 2015 from NCBI. Genes were identified in wastewater and

reference genomes using Prokka (version 1.11) [34]. Genes were clustered at 80% using CD-HIT [35]

(version 4.6.3, arguments -n 4 -c 0.8 -G 1 -aL 0.8 –aS 0.8 -B 1). Genes with over 90% sequence

identity, but only 30% coverage, as well as genes with 80% or greater identity and covered to

phage and virus sequences [36] were discarded. A gene cluster is defined to be present in an

isolate if there is a Prokka gene in the genome, which is longer than 100 amino acids and has

over 80% sequence identity and coverage against the gene cluster representative.

Pan- and core-genome. To generate the pan- and core-genome size graph we followed the

procedure in [3,16]. We had 92 genomes available. We varied *i* from one to 92. At each subset size

i, we randomly selected *i* genomes and computed the sizes of the union (pan) and intersection

(core) of gene clusters. This random selection was carried out 2000 times in each step.

Gene clusters to rank genes by correlation to phenotype. Prokka genes were identified in all

isolate genomes and then clustered with CD-HIT at 60% sequence identity and 50% coverage

(arguments -n 4 -c 0.6 -G 1 -aL 0.8 -aS 0.5 -B 1). A 80% identity cutoff was also tried but

dismissed, because the 60% threshold yielded 25% less clusters while adequately clustering

homologous gene sequences with lower sequence similarity. This threshold value is also

supported by the widespread default use of the BLOSUM62 matrix, the basis of which is

sequences clustered by 62% sequence identity.

Tree. The phylogenetic tree of 92 isolates was built following the procedure of [37,38] using FastTree

version 2.1 [39]. Sequence reads were aligned to *Escherichia coli* K12 MG 1665 and single

nucleotide variant calling was carried out using GATK [40]. Quality control for variant calling was

performed; variants supported by more than ten reads or likelihood score greater than 200 were

always in the range of 84 – 99% of variants called per isolate with the exception of 2 isolates

where only 59% and 60% of the variants were above the threshold for quality and supporting

reads. FastTree 2.1 [39] was then used to build the maximum likelihood tree based on the

sequences derived from variant calling.

Phylogrouping. For phylogrouping, the classification system established by Clermont *et al.* [21]

based on the genes chuA and yjaA and the DNA fragment TspE4.C2 was used. Blast was

performed to check each genome assembly for presence or absence of the aforementioned

elements with an identity cutoff ≥ 90%.

MLST. Concerning epidemiology and Multi-Locus Sequence Typing, we used the webserver at

https://cge.cbs.dtu.dk/services/MLST/ that follows the MLST scheme in [41] for predicting MLSTs

from whole genome sequence data [42]. 92 Draft genome assemblies were submitted and results

were obtained; 5 isolates were unidentified demonstrating novel sequence types.

Virulence factors. Virulence factors protein sequences were downloaded from VFDB: Virulence

Factors database [19,43]. 2000 sequences, which are *Escherichia coli* related, were chosen.

Sequences were then clustered at 80% sequence identity using CD-HIT (version 4.6.3,

arguments -n 4 -c 0.8 -G 1 -aL 0.8 -aS 0.8 –B 1). A virulence factor was considered present in an

isolate's genome if there is a Prokka gene in the genome that has over 80% sequence identity

and coverage against the virulence factor cluster representative.

ExPEC classification. There are intra- and extra-intestinal pathogenic *Escherichia coli*, which

can be classified from the presence of virulence factors [44-47]. InPEC are characterised by the

virulence factors stx1, stx2, escV, and bfpB. They are ExPEC if they contain over 20 of 58

virulence factors afa/draBC, bmaE, gafD, iha cds, mat, papEF, papGII, III, sfa/foc, etsB, etsC, sitD

ep, sitD ch, cvaC MPIII, colV MPIX, eitA, eitC, iss, neuC, kpsMTII, ompA, ompT, traT, hlyF, GimB,

malX, puvA, yqi, stx1, stx2, escV, bfp, feob, aatA, csgA, fimC, focG, nfaE, papAH, papC, sfaS,

tsh, chuA, fyuA, ireA, iroN, irp2, iucD, iutA, sitA, astA, cnf1, sat, vat, hlyA, hlyC, ibeA, tia, and pic.

Data availability statement

Genome assemblies of the analyzed isolates that support the findings of the study will be made

available on the NCBI upon paper publication.

Bioproject	Biosample	Accession	strain
PRJNA380388	SAMN06641941	NBBP00000000	Escherichia coli Win2013_WWKa_OUT_3
PRJNA380388	SAMN06641940	NBBQ00000000	Escherichia coli Win2013_WWKa_OUT_29
PRJNA380388	SAMN06641933	NBBR00000000	Escherichia coli Win2013_WWKa_OUT_18
PRJNA380388	SAMN06641932	NBBS00000000	Escherichia coli Win2013_WWKa_OUT_24
PRJNA380388	SAMN06641931	NBBT00000000	Escherichia coli Win2013_WWKa_OUT_1
PRJNA380388	SAMN06641928	NBBU00000000	Escherichia coli Win2013_WWKa_NEU_65
PRJNA380388	SAMN06641927	NBBV00000000	Escherichia coli Win2013_WWKa_NEU_20
PRJNA380388	SAMN06641926	NBBW00000000	Escherichia coli Win2013_WWKa_NEU_60
PRJNA380388	SAMN06641901	NBBX00000000	Escherichia coli Win2013_WWKa_ALT_23
PRJNA380388	SAMN06641884	NBBY00000000	Escherichia coli Win2012_WWKa_OUT_49
PRJNA380388	SAMN06641883	NBBZ00000000	Escherichia coli Win2012_WWKa_OUT_8
PRJNA380388	SAMN06641882	NBCA00000000	Escherichia coli Win2012_WWKa_OUT_34
PRJNA380388	SAMN06641881	NBCB00000000	Escherichia coli Win2012_WWKa_OUT_35
PRJNA380388	SAMN06641880	NBCC00000000	Escherichia coli Win2012_WWKa_OUT_29
PRJNA380388	SAMN06641879	NBCD00000000	Escherichia coli Win2012_WWKa_OUT_26
PRJNA380388	SAMN06641878	NBCE00000000	Escherichia coli Win2012_WWKa_OUT_33
PRJNA380388	SAMN06641877	NBCF00000000	Escherichia coli Win2012_WWKa_OUT_21
PRJNA380388	SAMN06641876	NBCG00000000	Escherichia coli Win2012_WWKa_OUT_2
PRJNA380388	SAMN06641875	NBCH00000000	Escherichia coli Win2012_WWKa_NEU_7
PRJNA380388	SAMN06641874	NBCI00000000	Escherichia coli Win2012_WWKa_OUT_14
PRJNA380388	SAMN06641873	NBCJ00000000	Escherichia coli Win2012_WWKa_NEU_51
PRJNA380388	SAMN06641872	NBCK00000000	Escherichia coli Win2012_WWKa_NEU_31
PRJNA380388	SAMN06641871	NBCQ00000000	Escherichia coli Win2012_WWKa_NEU_37
PRJNA380388	SAMN06641870	NBCR00000000	Escherichia coli Win2012_WWKa_NEU_16
PRJNA380388	SAMN06641869	NBCS00000000	Escherichia coli Win2012_WWKa_NEU_19
PRJNA380388	SAMN06641868	NBCT00000000	Escherichia coli Win2012_WWKa_NEU_12
PRJNA380388	SAMN06641867	NBCU00000000	Escherichia coli Win2012_WWKa_ALT_65
PRJNA380388	SAMN06641866	NBCV00000000	Escherichia coli Win2012_WWKa_NEU_1
PRJNA380388	SAMN06641865	NBCW00000000	Escherichia coli Win2012_WWKa_ALT_49
PRJNA380388	SAMN06641864	NBCX00000000	Escherichia coli Win2012_WWKa_ALT_54
PRJNA380388	SAMN06641863	NBCY00000000	Escherichia coli Sum2013_WWKa_OUT_5
PRJNA380388	SAMN06641862	NBCZ00000000	Escherichia coli Sum2013_WWKa_OUT_39
PRJNA380388	SAMN06641861	NBDA00000000	Escherichia coli Sum2013_WWKa_OUT_49
PRJNA380388	SAMN06641860	NBDB00000000	Escherichia coli Sum2013_WWKa_OUT_3
PRJNA380388	SAMN06641859	NBDC00000000	Escherichia coli Sum2013_WWKa_OUT_31
PRJNA380388	SAMN06641858	NBDD00000000	Escherichia coli Sum2013_WWKa_OUT_2
PRJNA380388	SAMN06641857	NBDE00000000	Escherichia coli Sum2013_WWKa_OUT_21
PRJNA380388	SAMN06641856	NBDF00000000	Escherichia coli Sum2013_WWKa_NEU_53
PRJNA380388	SAMN06641855	NBDG00000000	Escherichia coli Sum2013_WWKa_NEU_46
PRJNA380388	SAMN06641854	NBDH00000000	Escherichia coli Sum2013_WWKa_NEU_39
PRJNA380388	SAMN06641853	NBDI00000000	Escherichia coli Sum2013_WWKa_ALT_44
PRJNA380388	SAMN06641852	NBDJ00000000	Escherichia coli Sum2013_WWKa_NEU_29
PRJNA380388	SAMN06641851	NBDK00000000	Escherichia coli Spr2013_WWKa_OUT_27
PRJNA380388	SAMN06641844	NBDL00000000	Escherichia coli Sum2013_WWKa_ALT_41
PRJNA380388	SAMN06641843	NBDM00000000	Escherichia coli Sum2013_WWKa_ALT_27
PRJNA380388	SAMN06641842	NBDN00000000	Escherichia coli Spr2013_WWKa_OUT_56
PRJNA380388	SAMN06641841	NBDO00000000	Escherichia coli Sum2013_WWKa_ALT_20

PRJNA380388	SAMN06641840	NBJM00000000	Escherichia coli Spr2013_WWKa_OUT_5
PRJNA380388	SAMN06641839	NBJN00000000	Escherichia coli Spr2013_WWKa_OUT_55
PRJNA380388	SAMN06641838	NBJO00000000	Escherichia coli Spr2013_WWKa_OUT_32
PRJNA380388	SAMN06641837	NBJP00000000	Escherichia coli Spr2013_WWKa_OUT_45
PRJNA380388	SAMN06641822	NBJQ00000000	Escherichia coli Spr2013_WWKa_OUT_15
PRJNA380388	SAMN06641821	NBJR00000000	Escherichia coli Spr2013_WWKa_OUT_29
PRJNA380388	SAMN06641820	NBJS00000000	Escherichia coli Spr2013_WWKa_NEU_6
PRJNA380388	SAMN06641819	NBJT00000000	Escherichia coli Spr2013_WWKa_OUT_11
PRJNA380388	SAMN06641818	NBJU00000000	Escherichia coli Spr2013_WWKa_NEU_15
PRJNA380388	SAMN06641817	NBJV00000000	Escherichia coli Spr2013_WWKa_NEU_37
PRJNA380388	SAMN06641816	NBJW00000000	Escherichia coli Spr2013_WWKa_ALT_63
PRJNA380388	SAMN06641815	NBJX00000000	Escherichia coli Spr2013_WWKa_ALT_71
PRJNA380388	SAMN06641814	NBJY00000000	Escherichia coli Spr2013_WWKa_ALT_51
PRJNA380388	SAMN06641813	NBJZ00000000	Escherichia coli Spr2013_WWKa_ALT_55
PRJNA380388	SAMN06641812	NBKA00000000	Escherichia coli Spr2013_WWKa_ALT_43
PRJNA380388	SAMN06641811	NBKB00000000	Escherichia coli Spr2013_WWKa_ALT_27
PRJNA380388	SAMN06641810	NBKC00000000	Escherichia coli Spr2013_WWKa_ALT_41
PRJNA380388	SAMN06641809	NBKD00000000	Escherichia coli Spr2012_WWKa_OUT_37
PRJNA380388	SAMN06641808	NBKE00000000	Escherichia coli Spr2012_WWKa_OUT_54
PRJNA380388	SAMN06641807	NBKF00000000	Escherichia coli Spr2012_WWKa_OUT_25
PRJNA380388	SAMN06641806	NBKG00000000	Escherichia coli Spr2012_WWKa_OUT_3
PRJNA380388	SAMN06641805	NBKH00000000	Escherichia coli Spr2012_WWKa_OUT_16
PRJNA380388	SAMN06641804	NBKI00000000	Escherichia coli Spr2012_WWKa_OUT_13
PRJNA380388	SAMN06641803	NBKJ00000000	Escherichia coli Spr2012_WWKa_NEU_74
PRJNA380388	SAMN06641802	NBKK00000000	Escherichia coli Spr2012_WWKa_OUT_12
PRJNA380388	SAMN06641801	NBKL00000000	Escherichia coli Spr2012_WWKa_NEU_31
PRJNA380388	SAMN06641800	NBKM00000000	Escherichia coli Spr2012_WWKa_NEU_51
PRJNA380388	SAMN06641799	NBKN00000000	Escherichia coli Spr2012_WWKa_NEU_24
PRJNA380388	SAMN06641798	NBKO00000000	Escherichia coli Spr2012_WWKa_ALT_27
PRJNA380388	SAMN06641797	NBKP00000000	Escherichia coli Spr2012_WWKa_ALT_35
PRJNA380388	SAMN06641796	NBKQ00000000	Escherichia coli Aut2013_WWKa_OUT_3
PRJNA380388	SAMN06641793	NBKR00000000	Escherichia coli Aut2013_WWKa_OUT_10
PRJNA380388	SAMN06641792	NBKS00000000	Escherichia coli Aut2013_WWKa_OUT_20
PRJNA380388	SAMN06641791	NBKT00000000	Escherichia coli Aut2013_WWKa_NEU_51
PRJNA380388	SAMN06641789	NBKU00000000	Escherichia coli Aut2013_WWKa_NEU_53
PRJNA380388	SAMN06641788	NBKV00000000	Escherichia coli Aut2013_WWKa_NEU_44
PRJNA380388	SAMN06641786	NBKW00000000	Escherichia coli Aut2013_WWKa_ALT_65
PRJNA380388	SAMN06641785	NBKX00000000	Escherichia coli Aut2013_WWKa_NEU_28
PRJNA380388	SAMN06641784	NBKY00000000	Escherichia coli Aut2013_WWKa_ALT_59
PRJNA380388	SAMN06641782	NBKZ00000000	Escherichia coli Aut2013_WWKa_ALT_48
PRJNA380388	SAMN06641780	NBLA00000000	Escherichia coli Aut2013_WWKa_ALT_45
PRJNA380388	SAMN06641779	NBLB00000000	Escherichia coli Aut2013_WWKa_ALT_30
PRJNA380388	SAMN06641778	NBLC00000000	Escherichia coli Aut2013_WWKa_ALT_17
PRJNA380388	SAMN06641777	NBLD00000000	Escherichia coli Aut2013_WWKa_ALT_13
PRJNA380388	SAMN06670745	NBNO00000000	Escherichia coli Win2012_WWKa_OUT_19

Table 3: Accession numbers of 92 de novo assembled wastewater *Escherichia coli* genomes.

References

1   Hu, Y. *et al.* Metagenome-wide analysis of antibiotic resistance genes in a large cohort of human gut
microbiota. *Nature communications* **4**, 2151, doi:10.1038/ncomms3151 (2013).
2   Sommer, M. O., Dantas, G. & Church, G. M. Functional characterization of the antibiotic resistance reservoir
in the human microflora. *Science* **325**, 1128-1131, doi:10.1126/science.1176950 (2009).
3   Salipante, S. J. *et al.* Large-scale genomic sequencing of extraintestinal pathogenic Escherichia coli strains.
Genome research **25**, 119-128, doi:10.1101/gr.180190.114 (2015).
4   Forsberg, K. J. *et al.* The shared antibiotic resistome of soil bacteria and human pathogens. *Science* **337**,
1107-1111, doi:10.1126/science.1220761 (2012).
5   Riesenfeld, C. S., Goodman, R. M. & Handelsman, J. Uncultured soil bacteria are a reservoir of new
antibiotic resistance genes. *Environmental microbiology* **6**, 981-989, doi:10.1111/j.1462-2920.2004.00664.x
(2004).
6   Rizzo, L. *et al.* Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes
spread into the environment: a review. *The Science of the total environment* **447**, 345-360,
doi:10.1016/j.scitotenv.2013.01.032 (2013).
7   Gomi, R. *et al.* Occurrence of Clinically Important Lineages, Including the Sequence Type 131 C1-M27
Subclone, among Extended-Spectrum-beta-Lactamase-Producing Escherichia coli in Wastewater.
Antimicrobial agents and chemotherapy **61**, doi:10.1128/AAC.00564-17 (2017).
8   Kappell, A. D. *et al.* Detection of multi-drug resistant Escherichia coli in the urban waterways of Milwaukee,
WI. *Frontiers in microbiology* **6**, 336, doi:10.3389/fmicb.2015.00336 (2015).
9   Bengtsson-Palme, J. *et al.* Elucidating selection processes for antibiotic resistance in sewage treatment
plants using metagenomics. *The Science of the total environment* **572**, 697-712,
doi:10.1016/j.scitotenv.2016.06.228 (2016).
10   Lapierre, P. & Gogarten, J. P. Estimating the size of the bacterial pan-genome. *Trends Genet* **25**, 107-110,
doi:10.1016/j.tig.2008.12.004 (2009).
11   Kaas, R. S., Friis, C., Ussery, D. W. & Aarestrup, F. M. Estimating variation within the genes and inferring the
phylogeny of 186 sequenced diverse Escherichia coli genomes. *BMC Genomics* **13**, 577, doi:10.1186/1471-
2164-13-577 (2012).
12   Vieira, G. *et al.* Core and panmetabolism in Escherichia coli. *J Bacteriol* **193**, 1461-1472,
doi:10.1128/JB.01192-10 (2011).
13   Gordienko, E. N., Kazanov, M. D. & Gelfand, M. S. Evolution of pan-genomes of Escherichia coli, Shigella
spp., and Salmonella enterica. *J Bacteriol* **195**, 2786-2792, doi:10.1128/JB.02285-12 (2013).
14   Lukjancenko, O., Wassenaar, T. M. & Ussery, D. W. Comparison of 61 sequenced Escherichia coli genomes.
Microb Ecol **60**, 708-720, doi:10.1007/s00248-010-9717-3 (2010).
15   Rasko, D. A. *et al.* The pangenome structure of Escherichia coli: comparative genomic analysis of E. coli
commensal and pathogenic isolates. *J Bacteriol* **190**, 6881-6893, doi:10.1128/JB.00619-08 (2008).
16   Touchon, M. *et al.* Organised genome dynamics in the Escherichia coli species results in highly diverse
adaptive paths. *PLoS Genet* **5**, e1000344, doi:10.1371/journal.pgen.1000344 (2009).
17   McArthur, A. G. *et al.* The comprehensive antibiotic resistance database. *Antimicrobial agents and
chemotherapy* **57**, 3348-3357, doi:10.1128/AAC.00419-13 (2013).
18   Kaper, J. B., Nataro, J. P. & Mobley, H. L. Pathogenic Escherichia coli. *Nature reviews. Microbiology* **2**, 123-
140, doi:10.1038/nrmicro818 (2004).
19   Yang, J., Chen, L., Sun, L., Yu, J. & Jin, Q. VFDB 2008 release: an enhanced web-based resource for
comparative pathogenomics. *Nucleic acids research* **36**, D539-542, doi:10.1093/nar/gkm951 (2008).
20   Jaureguy, F. *et al.* Phylogenetic and genomic diversity of human bacteremic Escherichia coli strains. *BMC
genomics* **9**, 560, doi:10.1186/1471-2164-9-560 (2008).
21   Clermont, O., Bonacorsi, S. & Bingen, E. Rapid and simple determination of the Escherichia coli phylogenetic
group. *Applied and environmental microbiology* **66**, 4555-4558 (2000).
22   Land, M. *et al.* Insights from 20 years of bacterial genome sequencing. *Funct Integr Genomics* **15**, 141-161,
doi:10.1007/s10142-015-0433-4 (2015).
23   Maurelli, A. T., Fernandez, R. E., Bloch, C. A., Rode, C. K. & Fasano, A. "Black holes" and bacterial
pathogenicity: a large genomic deletion that enhances the virulence of Shigella spp. and enteroinvasive
Escherichia coli. *Proc Natl Acad Sci U S A* **95**, 3943-3948 (1998).
24   D'Costa, V. M. *et al.* Antibiotic resistance is ancient. *Nature* **477**, 457-461, doi:10.1038/nature10388 (2011).
25   Durso, L. M., Miller, D. N. & Wienhold, B. J. Distribution and quantification of antibiotic resistant genes and
bacteria across agricultural and non-agricultural metagenomes. *PLoS One* **7**, e48325,
doi:10.1371/journal.pone.0048325 (2012).

26   Manaia, C. M., Novo, A., Coelho, B. & Nunes, O. C. Ciprofloxacin Resistance in Domestic Wastewater
Treatment Plants. *Water Air Soil Poll* **208**, 335-343, doi:10.1007/s11270-009-0171-0 (2010).
27   Gomi, R. *et al.* Whole-Genome Analysis of Antimicrobial-Resistant and Extraintestinal Pathogenic
Escherichia coli in River Water. *Appl Environ Microbiol* **83**, doi:10.1128/AEM.02703-16 (2017).
28   Czekalski, N., Berthold, T., Caucci, S., Egli, A. & Burgmann, H. Increased levels of multiresistant bacteria and
resistance genes after wastewater treatment and their dissemination into lake geneva, Switzerland. *Front*
Microbiol **3**, 106, doi:10.3389/fmicb.2012.00106 (2012).
29   Tenaillon, O., Skurnik, D., Picard, B. & Denamur, E. The population genetics of commensal Escherichia coli.
Nat Rev Microbiol **8**, 207-217, doi:10.1038/nrmicro2298 (2010).
30   Caucci, S. *et al.* Seasonality of antibiotic prescriptions for outpatients and resistance genes in sewers and
wastewater treatment plant outflow. *FEMS microbiology ecology* **92**, fiw060, doi:10.1093/femsec/fiw060
(2016).
31   R: A language and environment for statistical computing (R Foundation for Statistical Computing, Vienna,
Austria, 2010).
32   Simpson, J. T. *et al.* ABySS: a parallel assembler for short read sequence data. *Genome research* **19**, 1117-
1123, doi:10.1101/gr.089532.108 (2009).
33   Hashimoto, M. *et al.* Cell size and nucleoid organization of engineered Escherichia coli cells with a reduced
genome. *Molecular microbiology* **55**, 137-149, doi:10.1111/j.1365-2958.2004.04386.x (2005).
34   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-2069,
doi:10.1093/bioinformatics/btu153 (2014).
35   Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide
sequences. *Bioinformatics* **22**, 1658-1659, doi:10.1093/bioinformatics/btl158 (2006).
36   Zhou, Y., Liang, Y., Lynch, K. H., Dennis, J. J. & Wishart, D. S. PHAST: a fast phage search tool. *Nucleic acids*
research **39**, W347-352, doi:10.1093/nar/gkr485 (2011).
37   Delsuc, F., Brinkmann, H. & Philippe, H. Phylogenomics and the reconstruction of the tree of life. *Nature*
reviews. Genetics **6**, 361-375, doi:10.1038/nrg1603 (2005).
38   Kumar, S., Filipski, A. J., Battistuzzi, F. U., Kosakovsky Pond, S. L. & Tamura, K. Statistics and truth in
phylogenomics. *Molecular biology and evolution* **29**, 457-472, doi:10.1093/molbev/msr202 (2012).
39   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood trees for large
alignments. *PloS one* **5**, e9490, doi:10.1371/journal.pone.0009490 (2010).
40   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation
DNA sequencing data. *Genome research* **20**, 1297-1303, doi:10.1101/gr.107524.110 (2010).
41   Wirth, T. *et al.* Sex and virulence in Escherichia coli: an evolutionary perspective. *Molecular microbiology* **60**,
1136-1151, doi:10.1111/j.1365-2958.2006.05172.x (2006).
42   Larsen, M. V. *et al.* Multilocus sequence typing of total-genome-sequenced bacteria. *Journal of clinical*
microbiology **50**, 1355-1361, doi:10.1128/JCM.06094-11 (2012).
43   Chen, L. *et al.* VFDB: a reference database for bacterial virulence factors. *Nucleic acids research* **33**, D325-
328, doi:10.1093/nar/gki008 (2005).
44   Antikainen, J. *et al.* New 16-plex PCR method for rapid detection of diarrheagenic Escherichia coli directly
from stool samples. *European journal of clinical microbiology & infectious diseases : official publication of*
the European Society of Clinical Microbiology **28**, 899-908, doi:10.1007/s10096-009-0720-x (2009).
45   Johnson, J. R. & Russo, T. A. Molecular epidemiology of extraintestinal pathogenic (uropathogenic)
Escherichia coli. *International journal of medical microbiology : IJMM* **295**, 383-404,
doi:10.1016/j.ijmm.2005.07.005 (2005).
46   Johnson, J. R. & Stell, A. L. Extended virulence genotypes of Escherichia coli strains from patients with
urosepsis in relation to phylogeny and host compromise. *The Journal of infectious diseases* **181**, 261-272,
doi:10.1086/315217 (2000).
47   Pitout, J. D. Extraintestinal Pathogenic Escherichia coli: A Combination of Virulence with Antibiotic
Resistance. *Frontiers in microbiology* **3**, 9, doi:10.3389/fmicb.2012.00009 (2012).