Regulation of Eag by calcium/calmodulin controls presynaptic excitability in *Drosophila*

Running title: Eag's CaMBD regulates excitability

Peter Bronk, Elena A. Kuklin, Chang Liu, Timothy D. Wiggin, Srinivas Gorur-Shandilya, Eve Marder and Leslie C. Griffith*

Department of Biology, National Center for Behavioral Genomics and Volen Center for Complex Systems, Brandeis University, Waltham, MA, 02454-9110 USA

*Correspondence:

Leslie C. Griffith
Brandeis University
Department of Biology MS008
415 South Street,
Waltham, MA, 02454-9110, USA
Tel: 781 736 3125
griffith@brandeis.edu

Acknowledgements

We would like to thank Rob Reenan (Brown University) for generously sharing *eag* constructs and for his advice on techniques for homologous recombination, and Ed Dougherty for assistance with confocal imaging.

Grants
This work was funded by R01 GM54408 to LCG, R35 NS097343 (to EM) and F32 NS098624 (to TDW).
.

Disclosures
The authors declare no competing financial interests

51
52 **Author Contributions**
53
54 PB and LCG conceived and designed the experiments. PB created mutants, performed all
55 biochemical and electrophysiology experiments and analyzed the data. CL assayed sleep and
56 memory and analyzed behavioral data. EAK performed morphological analyses of the NMJ.
57 SGS and EM conceptualized and built the model. TDW performed and analyzed the imaging
58 experiments. PB and LCG wrote the paper with editing from the other authors.
59

Abstract

Drosophila ether-à-go-go (eag) is the founding member of a large family of voltage-gated potassium channels, the KCNH family, which includes Kv10, 11 and 12, (Ganetzky et al. 1999). Mammalian EAG1 channels are inhibited by binding of calcium/calmodulin (Ca^{+2}/CaM) at sub-micromolar Ca^{2+} concentrations (Schonherr et al. 2000) as a result of the concurrent binding of Ca^{2+}/CaM to N- and C-terminal sites causing pore constriction (Whicher and MacKinnon 2016). Although the *Drosophila* EAG channel was believed to be Ca^{2+}-insensitive (Schonherr et al. 2000), both the N- and C-terminal sites are conserved. Here we show that *Drosophila* EAG is inhibited by high Ca^{2+} concentrations that are likely only present at plasma membrane Ca^{2+} channel microdomains. To test the role of this regulation *in vivo*, we engineered mutations that block CaM-binding to the major C-terminal site of the endogenous *eag* locus, disrupting Ca^{2+}-dependent gating. *eag* CaMBD mutants have reduced evoked release from larval motor neuron presynaptic terminals and show decreased Ca^{2+} influx in stimulated adult projection neuron presynaptic terminals. These results are consistent with a conductance-based multi-compartment model of the presynaptic terminal. The reduction of release in the larval neuromuscular junction drives a homeostatic increase in motor neuron somatic excitability. This misregulation of synaptic and somatic excitability has consequences for systems-level processes and leads to defects in associative memory formation in adults.

Significance Statement

Regulation of excitability is critical to tuning the nervous system for complex behaviors. We demonstrate here that the EAG family of voltage-gated potassium channel exhibits conserved gating by Ca^{2+}/CaM. Disruption of this regulation in *Drosophila* results in a decrease in evoked neurotransmitter release due to truncated Ca^{2+} influx in presynaptic terminals. In the larval motor system the blunting of evoked responses triggers a homeostatic increase in somatic excitability, which would act to compensate for the decrease in muscle stimulation. In adults, disrupted Ca^{2+} dynamics lead to defects in memory formation. These data demonstrate that the biophysical details of channels can have widespread effects on behavior at the level of cellular compartments and circuits.

Introduction

The repolarization of neurons and muscle cells by potassium channels plays an

important role in limiting the excitability of cells and returning them to a baseline membrane

potential to be ready for the next action potential. Potassium channels also play an indirect role

in regulating Ca^{2+} levels inside the cell, preventing excitotoxicity and shaping the temporal

profile of cellular responses to activity. Several families of Ca^{2+}-activated potassium channels

(e.g. small and large conductance, SK and BK channels), appear to act in a feedback mode,

with internal Ca^{2+} activating the channels to hyperpolarize the cell and limit additional Ca^{2+} entry.

However, there is one Ca^{2+}-regulated potassium channel family that appears to act in a

counterintuitive manner. The mammalian Ether-à-go-go (EAG) potassium channel is completely

inhibited at all membrane voltages in the presence of 100-300 nM Ca^{2+}/calmodulin (Ca^{2+}/CaM;

Schonherr et al. 2000; Stansfeld et al. 1996; Whicher and MacKinnon 2016; Ziechner et al.

2006). We sought to understand the role of this seemingly maladaptive regulation by disrupting

the CaM interaction with EAG in *Drosophila* where the presence of only one gene coding for

EAG simplifies the analysis.

The *Drosophila ether-à-go-go* (*eag*) gene encodes a voltage-gated delayed rectifier

potassium channel subunit, EAG. The first *eag* mutant was discovered in *Drosophila* and was

identified by its ether-induced leg shaking (Kaplan 1969). Subsequently, *Drosophila* EAG was

found to define a family of potassium channels with multiple homologs in mammals (Warmke

and Ganetzky 1994). EAG is a member of the KCNH family of potassium channels and is also

referred to as Kv10 (Gutman et al. 2003). Although the transmembrane domains of EAG are

similar in structure to Shaker-type voltage-gated channels (Warmke and Ganetzky 1994), recent

structural data has shown that EAG has a novel gating mechanism. This alternate gating

mechanism allows cytoplasmic factors like CaM to act on the pore region through the domain

linking the C-terminus to the S6 transmembrane domain (Whicher and MacKinnon 2016).

Channel closure is achieved by the simultaneous binding of Ca^{2+}/CaM to both an N-terminal and

a C-terminal site. This requirement for binding at two sites means that disruption of a single

CaM binding domain (CaMBD) blocks Ca^{2+}-dependent gating entirely (Schonherr et al. 2000).

The biophysical properties of EAG have been determined in heterologous expression

systems, and little is known about its function in the nervous system. Fly loss of function *eag*

mutants have learning deficits (Griffith et al. 1994) in addition to a robust hyperexcitability in

larval motor neurons that also causes spontaneous neuronal firing (Drysdale et al. 1991b;

Ganetzky and Wu 1983; Griffith et al. 1994; Srinivasan et al. 2012). EAG1 knock-out mice have

normal learning and memory, sensorimotor function, social behavior, anxiety and only display a

mild hyperactivity (Ufartes et al. 2013). However, the existence of multiple EAG family members

in mammals raises the possibility of compensation or redundancy and complicates *in vivo*

analysis. Cellular physiology in EAG1 knock-out mice has revealed enhanced synaptic

facilitation during high frequency stimulation (>= 50 Hz) at the parallel fiber-Purkinje cell

synapse in the cerebellum, accompanied by elevated presynaptic Ca^{2+} (Mortensen et al. 2015).

In both insects and mammals, the dominant phenotype of loss of EAG channels appears to be

increased presynaptic release.

The role of gating by Ca^{2+}/CaM *in vivo* is likely to be more complicated, and

understanding the role of Ca^{2+} inhibition requires generating an EAG channel that is voltage-

gated, but lacks Ca^{2+}/CaM inhibition. In this study we show that fly EAG is inhibited by

Ca^{2+}/CaM, and we generate mutant alleles disrupting the CaM interaction with EAG to

interrogate its function. We find that without Ca^{2+}/CaM inhibition of EAG, evoked synaptic

currents are reduced, and presynaptic Ca^{2+} is reduced during high frequency stimulation. These

defects lead to homeostatic changes in somatic excitability that go in the opposite direction. The

complex misregulation of excitability in these mutants disrupts higher level behavior,

demonstrating that the biophysical details of channel regulation can have profound effects at the

organismal level.

Materials and Methods

***Drosophila* crosses for recording and protein analysis**

Crosses to produce larvae for electrophysiology were set up in vials with 5-6 females and 5

males for each genotype to have consistent numbers of larvae between genotypes and kept at

around 22° C in the lab. Crosses were flipped every 3-5 days, but no more than 4 times. Bottle

crosses were used for protein extraction. The single alleles analyzed were hemizygous males

resulting from a cross of homozygous control or CaMBD mutant females to *w* males. The

transheterozygous alleles were females resulting from crossing 2 homozygous alleles of control

(WT12 and WT8) or CaMBD mutants (CaMBD8 and CaMBD17). Whenever possible the

transheterozygous combinations were used to reduce the effect of potential off target mutations

not eliminated during outcrossing of stocks.

Generation of mutant strains

Control and Calmodulin binding domain (CaMBD) mutant knock-in strains were generated by

targeted ends-out homologous recombination using methods described in detail previously

(Staber et al. 2011). The homologous arms were amplified by PCR from genomic DNA and

each arm was cloned into a Topo vector and were a kind gift from the laboratory of Robert

Reenan at Brown University. Mutations to the arm spanning the C2 CaMBD were made by site

directed mutagenesis using primers

CGTCCGGAAGATATCCTCCAAATCTCGTCGCACTCCGC (forward) and

GCGGAGTGCGACGAGATTTGGAGGATATCTTCCGGACG (reverse). The arms were cloned

sequentially into a P[w25.2] target vector. The full *eag* deletion mutant was generated using

piggyBack insertions containing FRT sites (Thibault et al. 2004) as described previously (Parks

et al. 2004). Briefly, Exelixis stocks PBac{RB}e03618 and PBac{WH}f02960 (p-element on X

chromosome) were used to delete the entire *eag* gene. PBac{RB}e03618/FM7a flies were

crossed to hs-FLP/Bal and male progeny were crossed to PBac{WH}f02960 females. After 48

hours, the bottle crosses were heat shocked in a 37º C water bath for 1 hour to activate the FLP

recombinase. The parents were removed after 72 hours and the bottles were heat shocked daily

(1 hour at 37º C) for 4 more days. The candidate flies were balanced and screened by PCR

using paired element-specific and genome-specific primers:

PB08014: GATCATTAAAACGTGGCCAACTAC   (5' *eag* flanking region e03618)

PB08013: CCTCGATATACAGACCGATAAAAC   (element specific primer)

PB08015: GTGGGGGTTCTTATTCTTCAGTT   (3' eag flanking region f02960).

Electrophysiology

Solutions: Nominally Ca^{2+}-free HL3.1 (Feng et al. 2004) (in mM): 70 NaCl, 5 KCl, 4

MgCl$_2$, 10 NaHCO$_3$, 5 Trehalose, 115 sucrose, 5 HEPES; pH adjusted to 7.1-7.2 with NaOH. *0*

Ca^{2+} Modified A solution (in mM): NaCl, 118; KCl, 2; MgCl$_2$, 4; Trehalose, 5; Sucrose, 45.5;

HEPES, 5; EGTA, 0.5; pH adjusted to pH 7.3-7.4 with NaOH. *Nominally Ca^{2+}-free Modified A*

solution (in mM): NaCl, 118; KCl, 2; MgCl$_2$, 4; Trehalose, 5; Sucrose, 45.5; HEPES, 5; pH

adjusted to pH 7.3-7.4 with NaOH. *HEK cell low Ca^{2+} internal patch solution* (in mM): NaCl, 2; K-

Gluconate, 130; CaCl$_2$, 0.1; MgCl$_2$, 2; EGTA, 1; HEPES, 10; pH adjusted to 7.3-7.4 with KOH.

Motor neuron low Ca^{2+} internal patch solution (in mM): 2 NaCl, 130 K-Gluconate, 0.1 CaCl$_2$, 1

EGTA, 0.5 Na-GTP, 4 Mg-ATP, 10 HEPES) adjusted to 285 mOsm with glucose, and pH 7.3-

7.4 with KOH.

HEK cell recordings by whole-cell patch were made 48 to 60 hours after transfection.

Cells were recorded at 21-22° C in 0 Ca^{2+} modified A solution (see solutions section) and a low

Ca^{2+} internal patch solution (see above). The appropriate amount of Ca^{2+} was added to the

internal solution to yield the approximate calculated free Ca^{2+} concentrations shown in the

figure. The Maxchelator program (Chris Patton, Stanford University) was used to calculate the

free Ca^{2+} concentrations. The HEK cells were voltage clamped with a Multiclamp 700A amplifier

(Axon Instruments, Molecular Devices, Sunnyvale, CA) and currents were filtered at 2.6 kHz

and recorded using pClamp 8 software (Axon Instruments, Molecular Devices). Software

sampling at 10 kHz. Leak currents were calculated offline by fitting the steady state current vs.

voltage relationship from -80 mV to -60 mV. The slope and intercept of the linear fit was used to

calculate the leak current at each membrane potential and subtracted from the recorded

sustained currents. EAG transfected cells were rejected if the initial resting potential was < 20

mV. The recording for the different EAG isoforms (Figure 1E) were performed as above except

that the internal pipette solution was changed to increase EGTA to 5 mM and 4 mM Mg-ATP

was added.

Potentials from the neuromuscular junction (NMJ) were recorded in current clamp mode

from muscle 6 of wandering third instar (gender specified in fly cross methods) larvae with

intracellular electrodes filled with 3 M KCl (20-30 MΩ) in 0.4 mM or 0.2 mM Ca^{2+} HL3.1. Signals

were amplified with an Axoclamp 2B amplifier (Molecular Devices, Sunnyvale, CA), digitized

with a Digidata 1322A (Molecular Devices), and recorded at 10 kHz with pClamp8 software

(Molecular Devices). Muscles with resting potentials more depolarized than -45 mV were not

used. The mEJP and nerve evoked recordings were done in 0.2 mM Ca^{2+} HL3.1 except with 2

mM KCl to increase the potassium driving force. The motor nerves were stimulated with a glass

pipette ~10 μm in diameter at 2-3 times the threshold voltage to first trigger a postsynaptic

response.

Synaptic currents at the NMJ were recorded from larval muscle 6 of wandering 3[rd] instar

larvae in two-electrode voltage clamp (TEVC) mode with an Axoclamp 2B amplifier and digitized

as above. We used 0.2 mM Ca^{2+} HL3.1 solution except that the KCl was adjusted to 2 mM. The

3 M KCl filled voltage sensing electrodes had resistances of 15-18 MΩ, and the current passing

electrodes filled with 3 M KCl had resistances of 10-15 MΩ. The muscles were held at -70 mV.

Cells were rejected if, after impaling the muscle with both electrodes, the average of the 2

resting potentials recorded at each electrode was more depolarized than -30 mV.

Type Is motor neuron voltages were recorded by whole-cell patch clamp in current

clamp mode with a Multiclamp 700A amplifier (Molecular Devices, Sunnyvale, CA). Filtering and

sampling as for HEK cells. Larvae were filleted in Ca^{2+}-free modified A solution and the brain left

intact. Dorsal motor neurons directly below the sheath of the ventral nerve cord were visualized

with a 40x water immersion objective and exposed by applying 0.2 % w/v protease (type XIV,

MilliporeSigma, St. Louis, MO) in Ca^{2+}-free modified A solution with a patch pipette directly

around the cluster of neurons. The patch pipettes for recording had resistances of 2-3 MΩ

containing internal solution (see solutions section). Recordings were made first in nominally

Ca^{2+}-free modified A solution and then the solution was changed to a 1.8 mM Ca^{2+} modified A

solution. Current was injected to keep the motor neurons at -60 mV. Cells were rejected if the

initial resting potential was more depolarized than -40 mV, or if the initial access resistance

exceeded 50 MΩ.

All physiological preparations were superfused with the indicated saline solutions by

gravity feed and vacuum suction at 1-2 ml/min. Electrophysiology trace plotting and analysis of

membrane current and voltage signals were performed in MATLAB (Mathworks, Natick, MA)

using scripts written by PB. Table 1. shows cell properties recorded for each of the data sets in

this study.

Protein analysis

For immunoblots, HEK cells were washed 2x with 1x PBS and then cells were scraped

in 500 μl 2x Laemmli sample buffer. The cell suspension was passed through a syringe needle

2-3x to break up the membranes. The homogenate was separated by 7.5% SDS-PAGE, and

immunoblotted with an anti-Eag rabbit polyclonal antibody raised against amino acids 1032-

1174 of *Drosophila* Eag.

For CaM-agarose purification of EAG, 1 ml of flies of each genotype were collected in a

15 ml conical centrifuge tube and frozen on dry ice. The frozen flies were vortexed and dumped

onto a sieve to collect the heads that pass through on a fine mesh cooled on dry ice. The heads

were put into 1.5 ml eppendorph tubes on dry ice, and then transferred to a Dounce

homogenizer (2 ml) on ice. 100 ml of homogenization solution (20 mM HEPES-NaOH pH 7.4, 1

mM EGTA) was added and the heads were homogenized. The homogenate was transferred to

a fresh 1.5 ml eppendorph tube and 100 ml additional homogenization solution was added to

the homogenizer to wash and collect the remaining homogenate. 200 ml of 2x solubilization

buffer (20 mM HEPES-NaOH pH 7.4, 200 mM NaCl, 2% Triton X-100, 1 mM EGTA) was added

to the homogenate. Both buffers contained a broad spectrum protease inhibitor, Complete Mini

(Roche Diagnostics, Germany). The tubes of homogenate were rotated at 4° C for 1 hour and

centrifuged for 15 minutes (4° C, 14,000 rpm). The supernatant was added to 100 μl of

sepharose beads conjugated to Calmodulin (equilibrated in same buffers used in solubilization

step) along with Ca^{2+} to a final concentration of 2 mM and rotated 1 hour at 4° C. 100 μl of the

supernatant (flow through) was removed from each sample before washing the beads in Ca^{2+}

containing buffer 3x. 100 μl of 2x Laemmli sample buffer was added to the beads and then the

samples were boiled for 10 min. Samples were separated by 7.5% SDS-PAGE, and

immunoblotted with an anti-Eag rabbit polyclonal antibody (1:500) raised against amino acids

1032-1174 of *Drosophila* Eag, and Actin antibody, 1:5000 (EMD Millipore).

Immunohistochemistry

Larval brain: Wandering 3[rd] instar larvae were filleted and pinned out in Sylgard coated

35 x 15 mm petri dishes (3 larvae/dish) containing ice cold Ca^{2+}-free HL3.1 solution. The HL3.1

solution was removed and replaced by fresh ice cold 4% paraformaldehyde in PBS and

incubated at 4° C on an orbital shaker for 4 hours. Fixed preparations were washed 3 x 5

minutes with PBS-TX (0.2% Triton-X-100 in PBS) and then blocked overnight at 4° C in PBS-

TX-BSA (0.2% Triton-X-100 and 5% bovine serum albumin in PBS). The head and first few

abdominal segments were cut from the rest of the larval body (with the brain attached) and

placed into wells of a 60 well minitray (Nunc, Denmark) (1 larva/well) containing rabbit EAG

antibody (1:500) in PBS-TX-BSA and mouse Actin antibody, 1:1500 (EMD Millipore) in PBS-TX-

BSA. The tissue was incubated at room temperature for 2 hours on an orbital shaker and

washed 3 x 5 minutes with PBS-TX. 1:200 Alexa Fluor 488 goat anti-rabbit (Invitrogen,

Molecular Probes) and 1:200 Alexa Fluor 680 goat anti-mouse (Invitrogen, Molecular Probes) in

PBS-TX-BSA were added to the well and incubated at room temperature for 1.5 to 2 hours. The

brain preparations were then washed 3 x 5 minutes and mounted on slides in Vectashield

(Vector Laboratories, Burlingame, CA) still attached to the head to keep the brain oriented.

Sequential images of the green and red emission signals were taken using the 63x objective on

a Leica TCS SP2 confocal microscope. Ten optical sections (1.2 μm each) were taken for each

brain using identical settings making sure none of the signals were saturated. The pixel

intensities were analyzed using MATLAB (Mathworks, Natick, MA) by averaging the pixel

intensity in the same region of the neuropil for each optical section and subtracting the average

pixel intensity in the background. The ratio of Green:Red pixel intensities was calculated for

each brain.

Larval NMJ: Wandering 3[rd] instar larvae were filleted and pinned out in Sylgard coated

35 x 15 mm petri dishes containing ice cold Ca^{2+}-free HL3.1 solution. The motor nerves were

cut carefully and the brain was removed before replacing the solution with ice cold 4%

paraformaldehyde in PBS and incubating at room temperature on an orbital shaker for 20

minutes. After a quick rinse in PBS, the prep was washed in PBS for 20 minutes at room

temperature on an orbital shaker. Then the larval preparations were washed 3 x 10 minutes at

room temperature in PBT (PBS + 0.05% Triton X-100). The larval tissue was then blocked for 1

hour at room temperature in blocking solution (5% heat inactivated goat serum + PBT). Mouse

4F3 anti-DLG antibody (Developmental Studies Hybridoma Bank) was diluted 1:50 in blocking

solution and added to the larval preparations and they were incubated overnight at 4º C on an

orbital shaker. After 3 x 10 minute washes in PBT, the larval preparations were incubated with

goat anti-mouse Alexa-Fluor 488 (Invitrogen) diluted 1:200 in PBS and Alexa-Fluor 594-

conjugated goat anti-HRP (1:200, Jackson ImmunoResearch Laboratories) for 2-3 hours at

room temperature. Final washes in PBT (5 x 5 minutes). The larval preparations were then

mounted on slides in Vectashield (Vector Laboratories, Burlingame, CA). Images of the NMJs

were taken sequentially at the appropriate emission spectra on a Leica TCS SP5 confocal

microscope. Only boutons associated with DLG staining were counted from muscles 6/7 in

segments A2 or A3.

Cell culture and plasmids

HEK cells were grown in DMEM (Gibco/Life Technologies) supplemented with heat inactivated

bovine serum (Invitrogen). Cells were transfected with EAG plasmids (4 μg/100 mm plate) using

the Ca^{2+}-phosphate method, and co-transfected with GFP (1 μg/100 mm plate). After 6 hours,

transfected cells were resuspended and plated onto 35 mm plates coated with poly-L-lysine. 24

hours after transfection, cells were transferred to a 26° C incubator. This was necessary for the

expression of functional *Drosophila* Eag channels in HEK cells (Li et al. 2011). HEK cells were

transfected with a pcDNA3 vector backbone containing wild type or mutant *eag* cDNA (PA). The

C2 CaM binding domain mutations, F232S,F235S were generated by site-directed mutagenesis

(Stratagene) using primers: CGTCCGGAAGATATCCTCCAAATCTCGTCGCACTCCGC

(forward) and GCGGAGTGCGACGAGATTTGGAGGATATCTTCCGGACG (reverse). The

correct sequence of the plasmids was verified over the entire length of the cDNA. The splice

isoforms EAG-PE and EAG-PF were constructed from the EAG-PA plasmid described above

with the Gibson cloning method using 5 primers:

F10: GACTCACTATAGGGAGACCCAAGCTTGGTACCGAGCTCGGATCC

R11: GCCAATTTGGCGAATTTCGAGAGACCTCCCTTCGTGTCCTCGCTGTCG

R14:GCGAATTTCGAGAGACCAGGGAGAAACAGGTTGACGGCAAAACTCCCTTCGTGTCCT

CG

F12: GTCTCTCGAAATTCGCCAAATTGGCAAGATCAGTGACACG

R13: TTCTCGTTGTCTGTCTCCGCGGCGACATTGCCAAAGCCCACC

3 PCR reactions were performed. PCR a- F10:R11, PCR b- F10:R14, PCR c- F12:R13. Region

of editing was checked by sequencing.

Calcium Imaging of Adult *Drosophila* Neurons

Homozygous *eag* CaMBD17 mutant and *eag* WT12 control female flies were crossed with transgenic flies carrying *GH146-GAL4* (Bloomington Stock # 30026) and *UAS-GCaMP6f* (Bloomington Stock # 52869). Male progeny were screened for transgene expression by eye color and housed at 25^{o}C for 5-10 days post eclosion. Experimental animals were anesthetized with ice, and brains were dissected in Adult Hemolymph-Like (AHL) saline (Wang et al. 2003). *Ex vivo* brains were secured to a Sylgard-lined perfusion chamber filled with AHL using bent tungsten pins.

The antennal lobe (AL) of the brain was located by morphology using an Olympus BX51WI microscope and 40X/0.8NA objective. A FHC concentric bipolar electrode (CBAEB50) was positioned in the center of the AL using a manual micromanipulator. During each experimental trial, 4 electrical stimulus trains were delivered to the AL by an A.M.P.I. Master-8 pulse generator with a 20 second inter-train interval. The stimulus trains were 20 pulses with a 1ms pulse duration at a 100Hz pulse rate. The stimulus voltage was raised on each trial; 3, 5, 7, and 9 volt trials were performed on each brain.

The GCaMP6f response of the *GH146* terminals in the LH was measured using a Hammamatsu Orca-ER camera controlled by Micro-Manager. Images were acquired at a 300 ms frame interval. Stimulus delivery and image acquisition were synchronized using a National Instruments USB-6212 DAQ and a custom MATLAB program. Brains were continuously perfused with room temperature (20-22^{o}C) AHL during the experiment. All brains responded to the high voltage (9V) stimulus, no brains were excluded, and the stimulus electrode was not repositioned after the experiment began.

Imaging data were analyzed using a custom MATLAB program. The program aligned the image time series to correct drift using the Image Processing Toolbox. An analysis region of interest (ROI) and background (BG) region were manually defined, the per-frame mean fluorescence of the ROI and BG were calculated, and the BG corrected fluorescent time series was calculated. The $\Delta F/F$ was calculated for each frame i using the formula: $\Delta F_i / F = (F_i - F_{1-20}) / F_{1-20}$. The peak change in $\Delta F/F$ was calculated as the $\Delta F/F$ of the stimulus frame minus the $\Delta F/F$ of the frame before the stimulus. The sustained level of $\Delta F/F$ was calculated as the mean $\Delta F/F$ of frames 1-5 seconds after the stimulus minus the $\Delta F/F$ of the frame before the stimulus. Frame time-stamps were used to generate a mean peri-stimulus $\Delta F/F$ of each trial. The timing of each frame varied by several milliseconds due to software delays so, for display, the data was re-sampled to a uniform 10 Hz time base using shape-preserving piecewise cubic interpolation.

Modelling

Characterization of EAG channels. Time series data of currents in HEK cells expressing
EAG channels during voltage clamp were used to characterize the activation functions and
kinetics of EAG channels. Peak currents in HEK cells as a function of voltage were subtracted
from peak currents in EAG-expressing HEK cells to isolate the currents due to EAG channels
alone. Leak currents were then subtracted from this I-V curve, which was then converted to a
curve of conductance *vs.* voltage using a reversal potential $E_K = -106.17$mV. EAG channels
were modelled using the Hodgkin-Huxley framework (Hodgkin and Huxley 1952). The current
due to EAG channels is given by

$$I_{EAG} = \bar{g}m(V - E)A$$

where $\bar{g}$ is the specific conductance of the channels, m is the activation variable, E is the
reversal potential and A is the surface area of the compartment. m is a time varying variable
determined by

$$\tau_m \frac{dm}{dt} = m_\infty - m$$

where m_∞ and τ_m are functions of the membrane potential V that are determined by fitting
functions to plots of conductance *vs.* voltage and time-series of conductance at various holding
voltages. Since the activity of EAG channels also depends on the intracellular calcium
concentration, the process was repeated for multiple calcium concentrations to obtain functions
that depended on both the membrane potential and the calcium concentration. For these
channels, we used

$$m_\infty(V, Ca) = \left[\frac{0.0929}{0.0929 + [Ca^{2+}]}\right]\left[\frac{1}{1 + \exp\left(\frac{V + 23.12}{-16.94}\right)}\right]$$

for wild-type EAG channels and

$$m_\infty(V, Ca) = \left[\frac{0.8740}{0.97 + [Ca^{2+}]}\right]\left[\frac{1}{1 + \exp\left(\frac{V + 23.12}{-16.94}\right)}\right]$$

for CaMBD mutant EAG channels. Note that the mutant and wild-type channels differ in their
calcium sensitivity, but not in their voltage sensitivity (Figure 1C). For both wild-type and mutant
EAG channels, we found that the timescale of activation was not a strong function of calcium
concentration; so we used

$$\tau_m(V) = 5497 - \frac{5500}{1 + \exp\left(\frac{V + 251.5}{-16.94 - 51.5}\right)}$$

In these functions, V is in units of mV, τ_m is in ms, and $[Ca^{2+}]$ is in μM.

Model of presynaptic terminal. Because EAG is inhibited only at relatively high levels of

intracellular calcium, we reasoned that a possible zone where EAG could be inhibited is the

micro domain close to voltage-gated calcium channels (VGCCs) at the presynaptic terminal. We

modelled the presynaptic terminal using a three-compartment model: one for the invading action

potential, another for the calcium micro-domain, and a third for parts of the pre-synaptic terminal

far from VGCCs. The calcium micro-domain had voltage-gated sodium channels, potassium

channels, EAG channels, and calcium channels. The other compartment had the same channel

profile, except for calcium channels. All compartments were electrically coupled to each other

with a conductance of 0.1 nS.

The activation functions and kinetics of sodium channels were determined using fits to

data from embryonic *Drosophila* neurons (O'Dowd and Aldrich 1988); potassium channels from

Drosophila photoreceptors (Hardie 1991) and calcium channels from *Drosophila* embryonic

muscles (Hara et al. 2015).

The model was integrated using the exponential Euler method (Dayan and Abbott 2001)

using a general-purpose neuron simulator written in C++, that we have made freely available at

https://github.com/sg-s/xolotl/.

Code availability. A script to reproduce Figure 5, a toolbox to interactively vary

parameters in the model, and scripts to estimate gating functions of EAG from raw data are

available at https://github.com/marderlab/eag .

Learning and memory assays

Appetitive conditioning protocol was as described previously (Liu et al., 2012). A group

of ~50 flies in a training tube alternatively received 10% 3-octanol (OCT; Merck) and 10% 4-

methylcyclohexanol (MCH; Sigma-Aldrich) for 1 min for immediate memory or for 2 min for 24 h

memory. Flies were loaded into CS+ tubes containing 2M dried sucrose filter papers with one

odor presentation and then CS- tubes without sucrose papers with the other odor. After a

certain retention time (2 min or 24 h), the trained flies were allowed to choose between OCT

and MCH for 2 min in the T-maze. A learning index was then calculated by taking the mean

performance indices (PIs) of the two reciprocally trained groups. Each PI was calculated by the

differences of flies choosing CS+ and CS- divided by the total number of flies. To control

whether the learning deficits were due to lack of sensation of sucrose or odors, flies were

allowed to choose between sucrose+ and sucrose- tubes for 2 min or between one odor (OCT

or MCH) and air for 2 min. Sucrose preference index or odor avoidance index was calculated,

respectively.

Statistics

Electrophysiology: Wilcoxon Rank Sum, unpaired Student's t-test and ANOVA done in

MATLAB (MathWorks, Natick, MA) or Excel (Microsoft).

Memory: statistical analyses were performed using GraphPad Prism 7. The Wilks-

Shapiro test was used to determine data normality. Normally distributed data were analyzed

with unpaired T-test, and data which did not pass the normality test were analyzed with a Mann-

Whitney test. Data are presented as mean ± SEM. Difference between groups were considered

significant at $p < 0.05$.

Results

***Drosophila* EAG is inhibited by high $[Ca^{2+}]_i$**

Human EAG has three calmodulin (CaM) binding sites: one N-terminal and two C-

terminal (C1 and C2) (Ziechner et al. 2006) and is inhibited by nM levels of Ca^{2+} (Schonherr et

al. 2000). Previous work on *Drosophila* EAG, in which it was compared to hEAG1, suggested

that the fly channel was not regulated by Ca^{2+} (Schonherr et al. 2002). Intrigued by the fact that

all of these sites appear to be conserved in *Drosophila* (Figure 1A), we revisited the issue of

regulation by Ca^{2+} by expressing wildtype (WT) EAG, or a C-terminal CaMBD mutant form

(F732S,F735S, Sun et al. 2004), in HEK293 cells and recording currents with varying amounts

of Ca^{2+} in the pipette internal solution. Mutant and WT channels were expressed at comparable

levels (Figure 1B) and GFP was co-expressed to identify transfected cells.

As previously reported (Schonherr et al. 2002), *Drosophila* EAG was not inhibited at or

below 1 μM $[Ca^{2+}]_i$. Only when $[Ca^{2+}]_i$ was increased to high μM levels, did we see significant

inhibition. Figure 1C (left panel) shows currents recorded in the presence of 100 μM Ca^{2+}.

Currents from cells transfected with WT EAG are clearly reduced compared to those from cells

expressing EAG carrying mutations which block CaM-binding to the major C-terminal site. The

voltage-dependence of the channel is unaffected by this mutation (Figure 1C, right panel). The

dose-response relationship for $[Ca^{2+}]_i$ is shown in Figure 1D. WT EAG current density

approaches that of untransfected cells at high $[Ca^{2+}]_i$ while the CaMBD mutant has substantial

residual current even at mM levels of Ca^{2+} in the pipette.

The C-terminal CaMBDs likely dominates Ca^{2+} sensitivity of EAG

Examination of genomic data for the *eag* gene revealed that the N-terminal CaMBD, which is a critical partner of the C-terminal site for Ca^{2+}-dependent gating (Whicher and MacKinnon 2016), is located at an exonic junction that is alternatively spliced (Attrill et al. 2016, Figure 1A). Since CaM is known to bind amphipathic helices, variation at this region is potentially important for the affinity of CaM-binding and could potentially explain the published Ca^{2+}-insensitivity of the canonical "WT" EAG (Drysdale et al. 1991a), which corresponds to the PA splice variant curated on FlyBase, due to a dysfunctional N-terminal binding domain. The splice site found in PA is also present in the PD and PG proteins, but the PE splice form has a 2 amino acid deletion, making it more like the mammalian EAG1 protein, while the PF variant has an additional 7 amino acids inserted into the predicted amphipathic helix. To determine if splice variation at this site alters the Ca^{2+}-sensitivity of EAG, we transfected each of the splice forms into HEK293 cells and assayed their response to changes in [Ca^{2+}] (Figure 1E). Neither PE nor PF were more sensitive to Ca^{2+} than the original PA isoform and all were significantly less sensitive than mEAG1, which was completely inhibited at 500 nM [Ca^{2+}]$_i$. We conclude that the functional sensitivity of *Drosophila* EAG to inhibition by Ca^{2+} is likely set by the C-terminal CaMBDs since it is unchanged by variation in the N-terminal domain.

Construction and characterization of a CaMBD mutant at the endogenous *eag* locus

The low sensitivity of EAG to Ca^{2+}/CaM inhibition is notable. Bulk cytoplasmic Ca^{2+} is modulated in the nM to low μM range in neurons. The HEK cell results show a partial gain-of-function phenotype in the CaMBD mutant EAG channel due to inhibition at very high Ca^{2+} levels (Figure 1D). The only cytoplasmic domains that have levels of Ca^{2+} (mM) that are this high are near the mouth of open Ca^{2+} channels (Parekh 2008). The affinity of EAG for Ca^{2+}/CaM suggests that it would only be inhibited if it were localized to this type of Ca^{2+} channel microdomain *in vivo*.

To determine if there were a physiological role for Ca^{2+}-dependent inhibition of EAG, we engineered the C-terminal CaMBD mutations characterized above into the endogenous *eag* locus by ends-out homologous recombination (Figure 2A). Multiple alleles containing the F732S,F735S mutations were recovered. WT strains, made in an identical manner with WT recombination arms were also generated. Like the mutant alleles, these WT strains contain a residual 76 bp loxP site located in an intron. These strains were used in all experiments as controls due to their similar genetic background.

Biochemical analysis of adult head protein from CaMBD and WT flies by partial

purification on CaM-agarose and immunoblotting shows that EAG protein is present in

comparable amounts in both mutant and WT control alleles (Figure 2B, left panel). Quantitative

immunohistochemistry of larval brain neuropil (Figure 2B, right panel) confirms this. EAG is

normally present at high concentration in neuropil and can be seen in axons and at the

neuromuscular junction (NMJ) in larvae (see Sun et al. 2004 for immunohistochemistry of native

EAG) and the CaMBD mutation did not change this distribution (data not shown).

Third instar NMJs in CaMBD mutants are morphologically normal and have no

spontaneous activity

The NMJ of the third instar larva has been extensively used to assay excitatory synaptic

structure and function. *eag* null mutants have a characteristic constellation of phenotypes in this

preparation which include increases in the number of synaptic boutons (Budnik et al. 1990) and

a robust hyperexcitability phenotype in larval motor neurons seen both postsynaptically at

isolated NMJs (Drysdale et al. 1991b; Ganetzky and Wu 1983; Griffith et al. 1994) and at the

soma (Srinivasan et al. 2012). To determine if CaMBD mutants had morphological abnormalities

we stained third instar NMJs with antibodies to HRP and DLG to visualize presynaptic and

postsynaptic structures. NMJs from CaMBD and WT strains were indistinguishable (Figure 3A)

and there was no significant difference in bouton number.

We also performed current clamp recordings from muscle 6/7 to ask if the CaMBD

mutant was functionally normal. Baseline recordings from CaMBD and WT control animals

showed no significant level of spontaneous excitatory junctional potentials (EJPs, Figure 3B). In

contrast, recordings from NMJs of $eag^{\Delta Full}$, a molecular null mutant, had high levels of

spontaneous activity. In experiments in which we evoked release by nerve stimulation in normal

saline, we did not see any supernumerary EJPs in the CaMBD mutant (P.B., data not shown).

CaMBD mutant third instar larvae exhibit decreases in the amplitude of evoked EJPs

The EJPs seen in unstimulated *eag* null mutants are the result of spontaneous

depolarization of the presynaptic nerve terminal and synchronous Ca^{2+}-dependent release of

many vesicles. To determine if there were changes in spontaneous Ca^{2+}-independent release of

single vesicles, we recorded miniature excitatory junctional potentials (mEJPs). Neither the rate

nor the amplitude of this class of events was changed in CaMBD animals compared to WT

controls (mean mEJP frequency: WT12/8, 1.92 ± 0.21 Hz; CaMBD8/17, 1.46 ± 0.15 Hz, p =

0.09. Mean mEJP amplitude: WT12/8, 0.85 ± 0.04 mV; CaMBD8/17, 0.83 ± 0.05 mV, p = 0.8).

This indicates that the probability of release for individual vesicles and the quantal postsynaptic

response to glutamate are both unaffected by loss of regulation of EAG by Ca^{2+}.

We also assayed evoked release by stimulating the motor nerve with a suction pipette

and recording activity in the muscle. We saw a significant decrease in the amplitude of

stimulated EJPs in the CaMBD mutants and a reduced quantal content (Figure 3C). EAG is also

expressed in larval muscles (Zhong and Wu 1991; 1993), so to make sure that activation of the

EAG channels in the postsynaptic muscle did not confound the amplitude of potentials recorded

in the muscle, we voltage clamped the muscle and recorded excitatory junctional currents

(EJCs). As seen in Figure 3D, CaMBD mutants had significantly smaller EJC amplitudes than

controls. There was no significant difference in the decay of the CaMBD mutant EJCs compared

to WT controls (WT: 10.2 $\pm$ 0.5 ms, CaMBD mutant: 10.8 $\pm$ 1.8, p = 0.8, unpaired Student's T-

test.

CaMBD domain mutants have reduced presynaptic Ca^{2+} influx

The CaMBD mutant's reduction in quantal content and synaptic currents is consistent

with a gain of EAG function when Ca^{2+}/CaM is not able to inhibit EAG currents. As seen in

Figure 1C and D, the F732S,F735S mutation leads to larger EAG currents when internal Ca^{2+} is

at high concentrations. This could cause more effective repolarization of activated presynaptic

terminals and an early termination of Ca^{2+} influx, which would be predicted to decrease

neurotransmitter release. To test the effects of the mutation on Ca^{2+} influx directly, we

expressed GCaMP6 in olfactory projection neurons. These neurons have well-separated axonal

and dendritic processes that allow clear visualization of the presynaptic terminals in the lateral

horn. These neurons can be directly stimulated by applying current to the antennal lobes with a

bipolar stimulating electrode.

Figure 4 shows GCaMP responses in WT and CaMBD mutant terminals with different

levels of stimulation. Ca^{2+} influx increases with stimulation strength, with CaMBD mutant

sustained levels lower than WT at all voltages. Interestingly, the form of the WT Ca^{2+} response

changes at high stimulation levels. The peak appears to plateau and a shoulder appears that

increases the width of the response. These data are consistent with the effect of the CaMBD

mutant being a gain-of-function that terminates the Ca^{2+} influx from voltage-gated Ca^{2+}

channels.

A simple model of the presynaptic terminal recapitulates changes in the width of action

potential and duration of the calcium influx

The fact that Ca^{2+} influx appears to be blunted in the CaMBD mutant supports the idea

that this mutation produces a gain of function. However since the sensitivity of *Drosophila* EAG

to Ca^{2+} is very low, it is possible that it only matters if EAG is present close to areas of high local

Ca^{2+}. In presynaptic terminals, the voltage-gated Ca^{2+} channels that control neurotransmitter

release are highly localized to the active zone, a molecular complex where vesicle fusion occurs

(Kawasaki et al. 2004). In contrast, EAG immunohistochemistry (Gillespie and Hodge 2013; Sun

et al. 2004) suggests that EAG is widely distributed to areas where it would never see Ca^{2+}

levels high enough to cause inhibition. To determine if our electrophysiological and imaging

results were consistent with the presence of two distinct pools of EAG, one that could be

inhibited by Ca^{2+} and a second that was insensitive even in WT animals, we built a model of the

presynaptic terminal that contained pools of EAG that were both inside and outside of Ca^{2+}

microdomains (Figure 5). Using parameters obtained from our HEK cell experiments, we

determined that the behavior we observed at the NMJ and in adult brain could be obtained by

modeling EAG's behavior as compartment-specific. The model shows that both membrane

voltage and Ca^{2+} currents were regulated as predicted.

Loss of Ca^{2+}-dependent inhibition of EAG increases somatic excitability

While the decrease in presynaptic release at the NMJ appears to be due to a decrease

in excitability and Ca^{2+} influx, direct recordings from terminals are difficult, so we carried out

whole-cell patch clamp of identified MNISN-Is neurons in the ventral ganglion to more closely

examine the relationship between membrane potential and firing in the CaMBD mutant. Figure

6A shows representative traces of CaMBD and WT control neurons with increasing current

injection. CaMBD neurons fire earlier and at a higher rate than that WT controls. The complete

F/I curves are shown in Figure 6B. These data demonstrate that the CaMBD mutants are

somatically hyperexcitable.

Recordings were performed both in nominally Ca^{2+}-free external solution and in 1.8 mM

external Ca^{2+}. Interestingly, there was no difference with Ca^{2+} as a main effect between the two

sets of F/I curves (3-way ANOVA, p = 0.9). This indicates that it is unlikely that the Ca^{2+}-

dependent regulation of EAG normally plays a role in somatic excitability. This could either

mean that EAG is not present in the somatodendritic compartment, or that it is located too far

from sites of Ca^{2+} influx to be inhibited. Given previous literature on *eag* null mutants (Srinivasan

et al. 2012), it is more likely that this is an issue of differential localization.

The data from the soma present a significant contrast to those obtained at the NMJ,

where there appeared to be a pronounced hypoexcitabilty. Given that these results are from

different cellular compartments of the same motor neurons, it seem likely that there is

compartment-specific homeostatic compensation occurring that increases drive to motor neuron

terminals. The lack of effect of external Ca^{2+} suggests that the alterations in excitability do not

involve acutely Ca^{2+}-dependent processes. The increase in excitability may act to partially

compensate for the decrease in evoked amplitude.

CaMBD mutants have memory defects

The demonstration of a change in somatic excitability aimed at compensating for poor

presynaptic release suggested that homeostatic changes might be able to ameliorate the effects

of this mutation at the circuit level. To test this, we asked if associative memory, a complex

behavior with a well-described circuitry, was disrupted in the CaMBD mutant. Adult flies were

conditioned by presenting one odor in the presence of sugar, an appetitive stimulus, and a

different odor with no sugar. WT flies, when asked to choose between the two odors either

immediately (short-term memory) or 24 h after training (long-term memory) will choose the

paired odor significantly above chance. CaMBD mutants have impairment in both types of

memory (Figure 7A). Input behaviors, sugar preference and odor avoidance (Figure 7B) are not

impaired. If anything, the CaMBD mutants were better than the WT control lines for avoidance

of octanol. The level of short-term memory impairment was similar to that seen with *eag* null

mutants, which interestingly also have enhanced octanol avoidance (C.L., data not shown).

These data demonstrate that the dysregulation of excitability caused by the loss of Ca^{2+}

sensitivity of the EAG channel has widespread effects on neuronal circuits and disrupts complex

behaviors.

Discussion

An obvious role of protein localization is to limit the action of a molecule to a particular

cell compartment e.g. an anchored K^+ channel will only affect the membrane potential of a small

area surrounding it; a scaffolded kinase will preferentially phosphorylate substrates in the same

complex. Less obvious consequences of binding to a scaffold are alterations of the intrinsic

activities of the complexed molecules and generation of novel complex-specific effects on

neuronal processes. A well-known example of this is the fusion of vesicles carrying

neurotransmitters. Cytosolic bulk Ca^{2+} in neurons is typically regulated in the 10 nM to 10 μM

range and only cytoplasmic Ca^{2+} microdomains close to a voltage-dependent Ca^{2+} channel

(VDCC) ever see Ca^{2+} at mM concentrations. The SNARE machinery has very low Ca^{2+} affinity

and only functions when vesicles are docked to a complex with VDCCs (Stanley 2016). The

inhibition of *Drosophila* EAG channels by Ca^{2+} is likely to be facilitated by a similar type of

interaction. In this study we provide the first insights into its role by introducing mutations into

the endogenous *eag* locus that block regulation by Ca^{2+}.

Drosophila EAG is inhibited by levels of Ca^{2+} only seen in VDCC microdomains

The fly EAG channel had been thought to be an outlier with regard to regulation by Ca^{2+}.

In this study we show that it is regulated in the same manner as mammalian EAG1 (Figure 1).

The major difference between fly and mammalian EAGs appears to be sensitivity. Inhibition of

fly EAG is only see at high μM to mM Ca^{2+}. This is strikingly different from the regulation of

mammalian EAG1, which is fully inhibited at nM levels of Ca^{2+}. How the difference is sensitivity

is achieved in the fly protein is unknown, but we present evidence that the efficacy of regulation

of EAG channels by Ca^{2+}/CaM is controlled primarily by the C-terminal CaMBDs, as alternative

splicing in the fly gene at the site of the N-terminal CaMBD does not modulate apparent affinity.

This Ca^{2+}-dependent inhibition of EAG channels is a robust and evolutionarily conserved feature

of the channel family that has so far been without demonstrated function.

Ca^{2+}-dependent inhibition of EAG may function to tune presynaptic release

EAG channels are located in presynaptic terminals in both mammals and flies (Gillespie

and Hodge 2013; Mortensen et al. 2015) and characterization of knock outs suggest that it

participates in repolarization after action potential invasion (Wu et al. 1983). These loss of

function studies make it clear that repolarizing K^{+} current is important for presynaptic function

and that the channel has a similar overall role across species. The role of Ca^{2+}-dependent

inhibition of EAG currents is not as well understood. The finding that evoked release was

decreased in the EAG CaMBD mutant in the face of normal spontaneous release dynamics and

quantal size argued that presynaptic EAG is normally exposed to very high levels of Ca^{2+} during

action potential-mediated neurotransmitter release, consistent with localization at active zones.

In mammals it has been speculated that the inhibition of EAG1 by Ca^{2+}/CaM serves as a

feedback mechanism to reduce the electrochemical driving force for Ca^{2+} influx (Schonherr et al.

2000). The idea is that EAG channels, if they were colocalized with VDCCs could control the

local membrane potential after an action potential during the time that the Ca^{2+} channels are

open. Inhibition of EAG by Ca^{2+} would allow the membrane potential to stay close the Ca^{2+}

reversal potential and blunt the electrochemical drive for Ca^{2+} influx through the open VDCC.

Our data (Figures 3 an 4) would argue against this hypothesis for the presynaptic terminal at the

larval NMJ since loss of regulation by Ca^{2+} decreases presynaptic release and evoked Ca^{2+}

influx. Whether this might be a viable mechanism for regulation of Ca^{2+} in dendrites would

require further investigation.

Another possible role for Ca^{2+}-dependent inhibition might be maintaining

neurotransmitter release during repetitive firing- essentially providing a way to turn down local

repolarization to maintain the ability to catalyze vesicle fusion. The dynamics of EAG inhibition

and its dependence of action potential pattern might provide clues to this aspect of EAG

function.

Why is regulation by Ca^{2+} different in mammals and insects?

The conservation of both N- and C-terminal CaMBDs from insects to mammals is

striking. Why would their apparent affinities be so disparate? The reason for the difference in

sensitivity between mammals and insects might have to do with how subcellular complexes are

constituted and the size of neurons and spines. The Ca^{2+}-sensitivity required to carry out a

function depends on the localization of the effector with respect to the Ca^{2+} source and the

nature of local Ca^{2+} buffering. Channels located in very close proximity to VDCCs do not require

high Ca^{2+} sensitivity, since they are exposed to very high (mM) local Ca^{2+} concentrations

(Tadross et al. 2013). Conversely, EAG channels that are inhibited with high affinity could be

affected by Ca^{2+} distant from the mouth of the VDCC. For fly neurons whose central processes

are very small, high affinity EAG inhibition might completely abolish all EAG current and render

neurons hyperexcitable. The Ca^{2+}-dependent inhibition of EAG could therefore contribute

maintaining presynaptic activity with repetitive stimulation in both species, but be tuned

differently due to differences in the location of the channels, buffering capacity of the neurons or

the size of the cell and synapse. A better understanding of the localizations of VDCCs and EAG

channels will be required to fully understand the role Ca^{2+}-dependent inhibition.

The ion channel complexes are critical to nervous system function

In neurons, ion channel-containing signaling complexes are a key class of protein

assemblage. The co-localization of proteins to the same complex can strongly influence the

kinetics and even the energetics of their activities and produce unforeseen interactions. Most of

a neuron's important biochemistry is not well-predicted by simply considering the equilibrium

state; Ca^{2+} microdomains are an excellent example of this principle. In this study, we present

data demonstrating that blocking the regulation of EAG by Ca^{2+} can disrupt the ability to form

memory (Figure 7). This argues that relatively subtle changes in the regulation of excitability can

have profound effects. There are an increasing number of examples of proteins whose function

requires their localization to a particular microdomain (Averaimo et al. 2016; Oheim et al. 2017;

Sanchez-Alonso et al. 2016). To understand how neurons work we must understand how

protein complexes shape signaling and the fly EAG channel will provide a useful model for this

endeavor.

705
Figure Legends

Figure 1. *Drosophila* EAG channels are regulated by Ca^{2+}. (A) Cartoon of the fly EAG

subunit showing the position of the highly conserved CaM binding domains (CaMBDs, not

drawn to scale). Protein sequence alignments of the three fly CaMBDs with human EAG as well

as comparison of the different *Drosophila* splice variants in the N-terminal CaMBD. Basic

residues are bold black, hydrophobic are bold gray. Arrows indicate the two phenylalanines

mutated to serines to disrupt CaM binding in the C2 domain. (B) Western blot of HEK293 cell

extracts expressing wild type (EAG-PA) or CaMBD (F232S,F235S) mutant plasmid DNA in

combination with an eGFP plasmid. (C) Sample traces of wild type or F232S,F235S mutant

EAG currents and I-V relation of EAG current densities recorded under whole-cell patch clamp

from HEK293 cells with 100 μM internal Ca^{2+} in pipette solution. (D) Plot comparing mean wild

type and CaMBD mutant EAG current densities at various internal Ca^{2+} concentrations at a

membrane voltage of 40 mV. Wild type (PA) and CaMBD mutant mean current densities are

significantly different at 100 and 500 μM internal Ca^{2+}, p = 0.04 for each concentration, Wilcoxon

rank sum test. (E) Plot of mean wild type EAG current densities for the N-terminal CaMBD

splice variants at 40 mV with mouse EAG for comparison. n represents number of cells

recorded. All values are mean ± SEM. There were no significant differences, p = 0.4, 2-factor

ANOVA between *Drosophila* splice isoforms across Ca^{2+} concentrations.

Figure 2. Characterization of F232S,F235S CaMBD homologous recombination mutant

***Drosophila*.** (A) Schematic of the targeted region (exons 12-14) of the *Drosophila eag* locus

and the target vector used to generate the CaMBD mutants with F232S,F235S mutations. The

control line used the same target vector without the F232S,F235S mutations. (B) Western blot

(left) from a CaM bead pull-down of fly EAG from whole head extract of CaMBD mutant, WT

control and the *eag$^{\Delta Full}$* null allele. The flow through and 2nd bead elutions were run to confirm all

of EAG was bound and released in the 1st elution. Flow through shows actin levels, indicating

loading was comparable for these samples. Larval brain immunohistochemistry (right) with EAG

antibody. Plot of the mean pixel intensity of the neuropil normalized to Actin immunostaining for

control and CaMBD mutant larvae. There is no significant difference in the normalized pixel

intensities for WT and CaMBD mutants, p = 0.4, unpaired Student's t-test. n represents number

of brains. Values are mean ± SEM.

Figure 3. Characterization of the CaMBD mutant larval neuromuscular junction (NMJ). (A) Sample images from immunohistochemistry of WT and CaMBD mutant larval neuromuscular junctions (muscles 6/7) with HRP and DLG antibodies to compare morphology of motor neuron boutons. Plot of number of boutons at the muscle 6/7 NMJs for WT and CaMBD mutants. There is no significant difference in the number of CaMBD mutant boutons compared to WT, p = 0.6, unpaired Student's t-test. n represents number of NMJs from 4 larvae each for WT control and CaMBD mutant. (B) Sample traces of spontaneous excitatory junctional potentials (EJPs) in WT, CaMBD mutant and a molecular null of *eag* recorded in 0.4 mM external Ca^{2+}. Plot of the mean spontaneous EJP frequencies. The CaMBD mutants are significantly different from the *eag* full deletion, $p = 1.1 \times 10^{-7}$, Wilcoxon rank sum test. n represents number of cells from 5 WT larvae, 6 CaMBD mutant larvae, and 7 full *eag* deletion larvae. (C) Sample traces of stimulated EJPs and plot of mean quantal content in 0.2 mM external Ca^{2+}. The CaMBD mutants had a significantly smaller quantal content, p = 0.03, Wilcoxon rank sum test. n represents number of cells from 9 larvae for WT and 6 larvae for the CaMBD mutants. (D) Sample traces of stimulated excitatory junctional currents (EJCs) and plot of mean EJC amplitudes recorded in 0.2 mM external Ca^{2+} and normalized to muscle size. The mean current density of the CaMBD mutant EJCs were significantly smaller than controls, $p = 4.7 \times 10^{-4}$, unpaired Student's t-test. n represents number of larvae (1 cell recorded per larva). n.s. = not significant. Values are mean ± SEM.

Figure 4. CaMBD mutants have reduced evoked presynaptic Ca^{2+} influx (A) Mean $\Delta F/F$ (%) traces from imaging of GCaMP6 responses in the lateral horn axon terminals of *GH146-GAL4+* neurons from adult brains before, during, and after electrical stimulation of the antennal lobe. Shaded areas represent the $25^{th}/75^{th}$ percentile bounds. Arrow indicates time of stimulus (20 X 1 msec, 100 Hz). (B) Bar graphs of mean peak change in $\Delta F/F$ (top) and sustained level of $\Delta F/F$ (bottom) for each stimulus amplitude. With a 7 V stimulus, the mean peak change was significantly lower in CaMBD mutants compared to controls, p = 0.03, and the sustained level was significantly lower in CaMBD mutants at 7 V, p = 0.004, and 9V, p = 0.001; unpaired Student's t-test with Bonferroni correction for multiple comparisons. n represents the number of brains. Values are mean ± SEM.

Figure 5. Action potential changes due to the CaMBD mutation in a simple model of the presynaptic terminal. (A) Schematic of the multi-compartment model of the presynaptic terminal. The presynaptic terminal is modeled as 3 electrically coupled compartments: (1) the

invading action potential (gray box, top), (2) the Ca^{2+} microdomain with voltage-gated Ca^{2+}

channels and Ca^{2+} influx (black box), and (3) the parts of the synaptic terminal outside the Ca^{2+}

microdomain (dotted black box). (B) Evoked potentials in response to invading action potential

in the Ca^{2+} microdomain with WT EAG channels (black) and mutant EAG channels (gray). In

WT, EAG channels are inhibited by incoming Ca^{2+} and this generates a prolonged potential. The

CaMBD mutant EAG channels have reduced sensitivity to Ca^{2+} and the membrane potential

quickly repolarizes due to a gain of EAG function. (C) Comparison of evoked potentials in the

Ca^{2+} microdomain (solid black) and outside the Ca^{2+} microdomain (dotted). Since EAG channels

outside the micro-domain are not inhibited by incoming Ca^{2+}, they quickly repolarize the

membrane after an action potential. (D) Comparison of evoked Ca^{2+} influx with WT (black) and

CaMBD mutant EAG (gray). Similar to Figure 4, Ca^{2+} influx is larger and more prolonged in cells

with WT EAG than in cells with CaMBD mutant EAG. All parameters of the model are identical

between WT and mutant, and between the two compartments of the presynaptic terminal.

Figure 6. Motor neurons show somatic hyperexcitability (A) Sample traces of whole-cell

patch current clamp recordings from Type Is larval motor neuron cell bodies at selected current

stimulus steps (80 pA and 120 pA) in 0 mM or 1.8 mM external Ca^{2+}. (B) Spike frequency vs.

stimulus curves (F/I) for nominally Ca^{2+}-free and 1.8 mM external Ca^{2+}. There is no significant

difference in the mean spike frequencies with or without Ca^{2+}, $p = 0.9$, 3-factor ANOVA). The

CaMBD mutant mean spike frequencies were significantly higher than controls (0 mM Ca^{2+}: $p =$

3.1×10^{-5}; 1.8 mM Ca^{2+}: $p = 0.03$, 2-factor ANOVA). n represents number of motor neurons

recorded. Plotted values are mean ± SEM. All cells recorded in the 1.8 mM external Ca^{2+}

condition were first recorded in the nominally Ca^{2+}-free condition.

Figure 7. Adult behavioral phenotypes of control and CaMBD mutant adults. (A) Plot of the

performance index for 2 min and 24 h memory of CaMBD mutant 17.4.1 and WT control 12.6.1

adults in a *white*[+] genetic background. The CaMBD mutants had a significantly lower

performance index for appetitive conditioning, $p = 0.007$ for both 2 min and 24 h tests, 2-factor

ANOVA with Sidak's multiple comparison test. n represents the number of groups tested (see

methods). CaMBD mutant 8.8.1 showed a similar 2 min memory deficit, though 24 h memory

was not assayed (C.L., data not shown). (B) Quantification of sugar preference ($p = 0.4$) and

avoidance of 4-methylcyclohexanol (MCH, $p = 0.08$) and 3-octanol (OCT $p = 0.0002$). All values

are mean ± SEM and comparisons done with unpaired Student's T-test.

References

Attrill H, Falls K, Goodman JL, Millburn GH, Antonazzo G, Rey AJ, Marygold SJ, and
FlyBase c. FlyBase: establishing a Gene Group resource for Drosophila melanogaster. *Nucleic*
Acids Res 44: D786-792, 2016.

Averaimo S, Assali A, Ros O, Couvet S, Zagar Y, Genescu I, Rebsam A, and Nicol X. A
plasma membrane microdomain compartmentalizes ephrin-generated cAMP signals to prune
developing retinal axon arbors. *Nature communications* 7: 12896, 2016.

Budnik V, Zhong Y, and Wu C-F. Morphological plasticity of motor axons in *Drosophila*
mutants with altered excitability. *J Neurosci* 10: 3754-3768, 1990.

Dayan P, and Abbott LF. *Theoretical Neuroscience: Computational and mathematical*
modeling of neural systems. Cambridge, MA: MIT Press, 2001.

Drysdale R, Warmke J, Kreber R, and Ganetzky B. Molecular characterization of *eag*: A
gene affecting potassium channels in *Drosophila melanogaster*. *Genetics* 127: 497-505, 1991a.

Drysdale R, Warmke J, Kreber R, and Ganetzky B. Molecular characterization of eag: a gene
affecting potassium channels in Drosophila melanogaster. *Genetics* 127: 497-505, 1991b.

Feng Y, Ueda A, and Wu CF. A modified minimal hemolymph-like solution, HL3.1, for
physiological recordings at the neuromuscular junctions of normal and mutant Drosophila
larvae. *J Neurogenet* 18: 377-402, 2004.

Ganetzky B, Robertson GA, Wilson GF, Trudeau MC, and Titus SA. The eag family of K+
channels in Drosophila and mammals. *Ann N Y Acad Sci* 868: 356-369., 1999.

Ganetzky B, and Wu CF. Neurogenetic analysis of potassium currents in Drosophila:
synergistic effects on neuromuscular transmission in double mutants. *J Neurogenet* 1: 17-28,
1983.

Gillespie JM, and Hodge JJ. CASK regulates CaMKII autophosphorylation in neuronal growth,
calcium signaling, and learning. *Frontiers in molecular neuroscience* 6: 27, 2013.

Griffith LC, Wang J, Zhong Y, Wu CF, and Greenspan RJ. Calcium/calmodulin-dependent
protein kinase II and potassium channel subunit eag similarly affect plasticity in Drosophila. *Proc*
Natl Acad Sci U S A 91: 10044-10048, 1994.

Gutman GA, Chandy KG, Adelman JP, Aiyar J, Bayliss DA, Clapham DE, Covarriubias M,
Desir GV, Furuichi K, Ganetzky B, Garcia ML, Grissmer S, Jan LY, Karschin A, Kim D,
Kuperschmidt S, Kurachi Y, Lazdunski M, Lesage F, Lester HA, McKinnon D, Nichols CG,
O'Kelly I, Robbins J, Robertson GA, Rudy B, Sanguinetti M, Seino S, Stuehmer W,
Tamkun MM, Vandenberg CA, Wei A, Wulff H, Wymore RS, and International Union of P.
International Union of Pharmacology. XLI. Compendium of voltage-gated ion channels:
potassium channels. *Pharmacol Rev* 55: 583-586, 2003.

Hara Y, Koganezawa M, and Yamamoto D. The Dmca1D channel mediates Ca(2+) inward
currents in Drosophila embryonic muscles. *J Neurogenet* 29: 117-123, 2015.

Hardie RC. Voltage-sensitive potassium channels in Drosophila photoreceptors. *J Neurosci* 11:
3079-3095, 1991.

Hodgkin AL, and Huxley AF. A quantitative description of membrane current and its
application to conduction and excitation in nerve. *J Physiol* 117: 500-544, 1952.

Kaplan W, Trout WE III. The behavior of four neurological mutants of Drosophila. *Genetics* 61:
399-409, 1969.

Kawasaki F, Zou B, Xu X, and Ordway RW. Active zone localization of presynaptic calcium
channels encoded by the cacophony locus of Drosophila. *J Neurosci* 24: 282-285, 2004.

Li Y, Liu X, Wu Y, Xu Z, Li H, Griffith LC, and Zhou Y. Intracellular regions of the Eag
potassium channel play a critical role in generation of voltage-dependent currents. *J Biol Chem*
286: 1389-1399, 2011.

Mortensen LS, Schmidt H, Farsi Z, Barrantes-Freer A, Rubio ME, Ufartes R, Eilers J,
Sakaba T, Stuhmer W, and Pardo LA. KV 10.1 opposes activity-dependent increase in Ca2+
influx into the presynaptic terminal of the parallel fibre-Purkinje cell synapse. *J Physiol* 593: 181-
196, 2015.

O'Dowd DK, and Aldrich RW. Voltage-clamp analysis of sodium channels in wild-type and
mutant *Drosophila* neurons. *J Neurosci* 8: 3633-3643, 1988.

Oheim M, Schmidt E, and Hirrlinger J. Local energy on demand: Are 'spontaneous' astrocytic
Ca2+-microdomains the regulatory unit for astrocyte-neuron metabolic cooperation? *Brain Res*
Bull 2017.

Parekh AB. Ca2+ microdomains near plasma membrane Ca2+ channels: impact on cell
function. *J Physiol* 586: 3043-3054, 2008.

Parks AL, Cook KR, Belvin M, Dompe NA, Fawcett R, Huppert K, Tan LR, Winter CG,
Bogart KP, Deal JE, Deal-Herr ME, Grant D, Marcinko M, Miyazaki WY, Robertson S, Shaw
KJ, Tabios M, Vysotskaia V, Zhao L, Andrade RS, Edgar KA, Howie E, Killpack K, Milash
B, Norton A, Thao D, Whittaker K, Winner MA, Friedman L, Margolis J, Singer MA,
Kopczynski C, Curtis D, Kaufman TC, Plowman GD, Duyk G, and Francis-Lang HL.
Systematic generation of high-resolution deletion coverage of the Drosophila melanogaster
genome. *Nature genetics* 36: 288-292, 2004.

Sanchez-Alonso JL, Bhargava A, O'Hara T, Glukhov AV, Schobesberger S, Bhogal N,
Sikkel MB, Mansfield C, Korchev YE, Lyon AR, Punjabi PP, Nikolaev VO, Trayanova NA,
and Gorelik J. Microdomain-Specific Modulation of L-Type Calcium Channels Leads to
Triggered Ventricular Arrhythmia in Heart Failure. *Circ Res* 119: 944-955, 2016.

Schonherr R, Gessner G, Lober K, and Heinemann SH. Functional distinction of human
EAG1 and EAG2 potassium channels. *FEBS Lett* 514: 204-208, 2002.

Schonherr R, Lober K, and Heinemann SH. Inhibition of human ether a go-go potassium
channels by Ca(2+)/calmodulin. *Embo J* 19: 3263-3271., 2000.

Srinivasan S, Lance K, and Levine RB. Contribution of EAG to excitability and potassium
currents in Drosophila larval motoneurons. *J Neurophysiol* 107: 2660-2671, 2012.

Staber CJ, Gell S, Jepson JE, and Reenan RA. Perturbing A-to-I RNA editing using genetics
and homologous recombination. *Methods Mol Biol* 718: 41-73, 2011.

Stanley EF. The Nanophysiology of Fast Transmitter Release. *Trends Neurosci* 39: 183-197,
2016.

Stansfeld CE, Roper J, Ludwig J, Weseloh RM, Marsh SJ, Brown DA, and Pongs O.
Elevation of intracellular calcium by muscarinic receptor activation induces a block of voltage-
activated rat ether-a-go-go channels in a stably transfected cell line. *Proc Natl Acad Sci U S A*
93: 9910-9914, 1996.

Sun XX, Hodge JJ, Zhou Y, Nguyen M, and Griffith LC. The *eag* potassium channel binds
and locally activates calcium/calmodulin-dependent protein kinase II. *J Biol Chem* 279: 10206-
10214, 2004.

Tadross MR, Tsien RW, and Yue DT. Ca2+ channel nanodomains boost local Ca2+
amplitude. *Proc Natl Acad Sci U S A* 110: 15794-15799, 2013.

Thibault ST, Singer MA, Miyazaki WY, Milash B, Dompe NA, Singh CM, Buchholz R,
Demsky M, Fawcett R, Francis-Lang HL, Ryner L, Cheung LM, Chong A, Erickson C,
Fisher WW, Greer K, Hartouni SR, Howie E, Jakkula L, Joo D, Killpack K, Laufer A,
Mazzotta J, Smith RD, Stevens LM, Stuber C, Tan LR, Ventura R, Woo A, Zakrajsek I,
Zhao L, Chen F, Swimmer C, Kopczynski C, Duyk G, Winberg ML, and Margolis J. A
complementary transposon tool kit for Drosophila melanogaster using P and piggyBac. *Nature*
genetics 36: 283-287, 2004.

Ufartes R, Schneider T, Mortensen LS, de Juan Romero C, Hentrich K, Knoetgen H,
Beilinson V, Moebius W, Tarabykin V, Alves F, Pardo LA, Rawlins JN, and Stuehmer W.
Behavioural and functional characterization of Kv10.1 (Eag1) knockout mice. *Human molecular*
genetics 22: 2247-2262, 2013.

Wang JW, Wong AM, Flores J, Vosshall LB, and Axel R. Two-photon calcium imaging
reveals an odor-evoked map of activity in the fly brain. *Cell* 112: 271-282, 2003.

Warmke JW, and Ganetzky B. A family of potassium channel genes related to eag in
Drosophila and mammals. *Proc Natl Acad Sci U S A* 91: 3438-3442, 1994.

Whicher JR, and MacKinnon R. Structure of the voltage-gated K(+) channel Eag1 reveals an
alternative voltage sensing mechanism. *Science* 353: 664-669, 2016.

Wu CF, Ganetzky B, Haugland FN, and Liu AX. Potassium currents in *Drosophila:* different
components affected by mutations of two genes. *Science* 220: 1076-1078, 1983.

Zhong Y, and Wu CF. Alteration of four identified K+ currents in Drosophila muscle by
mutations in eag. *Science* 252: 1562-1564, 1991.

Zhong Y, and Wu CF. Modulation of different K+ currents in Drosophila: a hypothetical role for
the Eag subunit in multimeric K+ channels. *J Neurosci* 13: 4669-4679, 1993.

Ziechner U, Schonherr R, Born AK, Gavrilova-Ruch O, Glaser RW, Malesevic M, Kullertz
G, and Heinemann SH. Inhibition of human ether a go-go potassium channels by
Ca2+/calmodulin binding to the cytosolic N- and C-termini. *The FEBS journal* 273: 1074-1086,
2006.
924

Table 1. Conditions and cell properties for electrophysiological recordings

Cell Type	Genotype	# Cells	Ca^{2+}	RMP (mV)		Cm (pF)		Rm (GΩ)		Rs (MΩ)	
				Initial	Final	Initial	Final	Initial	Final	Initial	Final
Motor Neuron	WT	5	0 mM Ext.	-50.2±1.3	-47.6±2.9	23.5±0.4	21.9±0.3	0.91±0.1	1.01±0.1	35.0±4.3	43.4±8.7
Motor Neuron	CaMBD mut.	7	0 mM Ext	-56.3±1.5	-56.0±1.5	25.2±0.8	22.9±0.8	0.88±0.2	1.02±0.2	36.9±3.9	53.5±8.7
Motor Neuron	WT	3	1.8 mM Ext.	-48.0±1.5	-49.3±1.9	20.3±0.5	19.1±0.5	0.92±0.1	0.96±0.1	40.6±7.0	52.6±13.6
Motor Neuron	CaMBD mut.	4	1.8 mM Ext.	-55.3±2.2	-55.3±1.9	22.6±1.4	21.0±1.3	1.08±0.3	0.92±0.2	48.9±10.8	65.9±10.6
HEK	WT	13	13 nM Int.	-38.5±2.0	-41.9±3.0	16.2±2.7	15.5±2.8	1.13±0.3	0.88±0.2	8.3±1.1	10.2±1.8
HEK	CaMBD mut.	9	13 nM Int.	-37.2±2.0	-36.6±3.0	17.6±3.8	16.8±4.2	0.6±0.1	0.42±0.1	9.1±1.7	11.5±1.5
HEK	WT	4	100 nM Int.	-43.3±3.4	-37.3±4.3	17.3±3.1	16.5±3.3	1.09±0.4	0.43±0.1	4.9±0.4	7.5±1.2
HEK	CaMBD mut.	5	100 nM Int.	-45.6±1.7	-43.2±0.7	20.5±4.2	27.3±10.7	1.30±0.6	1.26±0.6	8.3±0.7	12.6±1.1
HEK	WT	6	10 μM Int.	-41.5±4.1	-37.7±5.0	19.7±2.5	19.3±2.9	0.69±0.1	0.86±0.2	10.8±3.8	13.9±5.1
HEK	CaMBD mut.	4	10 μM Int.	-37.8±5.9	-38.5±10.5	17.5±5.0	13.9±2.2	1.01±0.4	0.77±0.2	12.0±4.4	13.1±2.9
HEK	WT	7	100 μM Int.	-41.7±0.9	-28.9±4.8	14.6±2.6	14.2±2.7	1.23±0.3	0.95±0.2	8.5±1.8	15.4±3.2
HEK	CaMBD mut.	6	100 μM Int.	-40.7±3.0	-43.5±4.2	16.8±1.8	16.2±1.6	1.57±0.4	0.91±0.2	6.7±1.6	11.3±2.2
HEK	WT	9	500 μM Int.	-39.7±4.3	-18.2±4.3	21.1±3.1	15.2±2.8	0.85±0.2	0.70±0.2	11.0±4.0	8.3±1.3
HEK	CaMBD mut.	7	500 μM Int.	-43.6±2.1	-34.9±4.2	24.5±4.0	15.8±2.9	0.76±0.2	0.53±0.2	8.3±1.0	11.6±3.4
HEK	EAG-PA	4	10 nM Int.	-37.8±3.7	-35.5±3.5	25.3±7.6	23.6±7.1	0.41±0.2	0.38±0.1	14.7±3.6	16.4±3.6
HEK	EAG-PE	3	10 nM Int.	-36.7±3.3	-41.7±1.7	17.5±9.3	17.0±9.1	1.23±0.4	1.11±0.4	11.5±3.2	13.7±3.8
HEK	EAG-PF	2	10 nM Int.	-34.5±1.5	-28.5±1.5	18.7±5.9	20.2±7.9	0.20±.04	0.43±0.2	14.0±0.9	14.4±0.5
HEK	GFP	4	10 nM Int.	-16.8±5.4	-21.3±6.6	12.7±2.7	12.2±2.7	1.95±0.2	1.34±0.2	10.4±2.0	11.7±2.8
HEK	EAG-PA	5	500 nM Int.	-38.2±3.7	-35.5±3.5	14.9±2.0	14.2±2.1	1.17±0.3	0.75±0.1	9.6±0.7	18.3±5.5
HEK	EAG-PE	8	500 nM Int.	-36.8±1.3	-29.9±3.0	13.6±1.9	13.3±1.8	0.56±0.2	0.33±0.1	8.2±0.9	10.1±1.0
HEK	EAG-PF	5	500 nM Int.	-32.8±3.0	-27.8±2.7	18.0±3.5	17.5±3.4	1.0±0.5	0.53±0.2	6.7±0.3	7.9±0.6
HEK	EAG-PA	3	1 μM Int.	-48.3±3.9	-42.0±7.5	10.1±0.3	9.7±0.2	1.09±0.5	1.0±0.4	8.2±0.9	10.4±1.9
HEK	EAG-PE	4	1 μM Int.	-39.5±3.0	-36.0±2.1	16.4±3.7	16.0±3.8	0.60±0.3	0.59±0.3	10.7±1.3	13.5±1.7
HEK	EAG-PF	4	1 μM Int.	-40.0±6.1	-26.5±6.6	19.6±5.8	18.7±5.4	0.67±0.4	0.85±0.6	13.4±2.7	16.6±3.3
HEK	GFP	3	1 μM Int.	-24.7±4.9	-18.3±6.2	12.2±2.0	11.5±2.2	1.02±.03	1.0±0.1	7.5±1.5	18.1±7.7
Muscle (VC)	WT	28	0.2 mM Ext.	-46.6±1.5		4.4±0.2 (nF)		0.0038 ±0.0003		0.14 ±0.0009	
Muscle (VC)	CaMBD mut.	30	0.2 mM Ext.	-46.8±1.1		4.8±0.1 (nF)		0.0027 ±0.0003		0.14 ±0.0009	

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7