Title: Paternal grandparental exposure to crop failure or surfeit during a childhood
slow growth period and epigenetic marks on third generation´s growth-,
glucoregulatory and stress genes

Authors: Lars Olov Bygren[1,2]*, Patrick Müller[1], David Brodin[1], Gunnar Kaati[1], John G. Kral[3]

Affiliations:

[1] Department of Biosciences and Nutriotion, Karolinska Institutet, SE_14183 Huddinge,
Sweden.

[2] Department of Community Medicine and Rehabilitation, Umeå University, SE_90187,
Umeå, Sweden.

[3] SUNY Downstate Medical Center, 450 Clarkson Avenue, Box 40, Brooklyn, NY 11203,
USA.

*Correspondence to: lars.olov.bygren@ki.se

Abstract: This latest in our series of papers describes transgenerational methylation related to

mid-childhood food availability in 19[th] century Överkalix, Sweden. Failed vs.bountiful crops

differentially influenced methylation in grandchildren of exposed paternal grandparents. In

this case study of 8 tracked cases with differential ancestral exposure, adult progeny differed

in methylated CpGs in three Amigo site gene pathways: "insulin processing", "adipose tissue

growth" and "hypothalamic development", reflecting famine, excess food and reaction to

food-insecurity stress. This is the first demonstration of transgenerational epigenetic

inheritance in humans following grandparental childhood exposure, an early developmental

origin of adult disease.

Non-technical summary: Paternal grandparents availability of food during their mid-

childhood development perceiding the prepubertal growth spurt induced epigenetic marks in

three gene pathways reflecting famine, excess food and reaction to food-insecurity stress.

Our epidemiological findings of adverse transgenerational effects of ancestral overnutrition

and conversely, beneficial effects of famine during SGP[1] prompted us to study individuals

whose paternal grandparents were randomly exposed, during their Slow Growth Period (SGP)

- a sensitive period preceding the pre-pubertal growth spurt - to failed or bumper crops

potentially engendering epigenetic marks on mechanistic molecular gene pathways.

For detailed analyses of specific gene ontology (GO) pathways, we tested the three posited

structural classes of genes associated with each of 40 preselected GO terms

(http://amigo.geneontology.org) for differentially methylated CpG positions (DMRs) between

the experimental groups. We selected GO pathways with average differences in methylation

between groups >0.1 and p-values <0.05 and performed detailed analyses of those differing

> 5%. The posited pathways comprised glucoregulatory genes that were seen as immediate

sensors of decreased glycogen stores, whereas lipids and ketones were posited to reflect the

duration and magnitude of the fast or responses to overnutrition. Individual reactions to food

insecurity were assumed to have activated the HPA-axis. We focused on two putative

response lines in direct descendants from a) grandfather to son with grandson and b) paternal

grandmother to son with granddaughter, adjusted for crop size during famine or surfeit. We

found 39 DMRs in grandchildren after grandparental differential exposure to food availability

in GO-pathways associated with famine, excess food and reactions to environmental stressors.

Nine of them were large, three GOs exhibiting remarkable DMRs in grandchildren (Table 1).

Results

Exploiting a natural experiment, we studied the paternal lineage transgenerational responses

in grandsons and granddaughters and assessed their DNA-methylations. We found 39 DMRs

amoung 40 posited pathways in Islands, North and South shores, North and South shelves

related to famine, excess and stress, 9 pathways with DMRs above 5%, and 3 above 14%.

Paternal grandparents´exposure vs. the adult grandchild methylation.

The main metabolic pathways implicated in grandpaternal SGP exposure to food-insecurity

stress transferred to grandsons were "appetite", "insulin processing", "ketone body catabolic

process" and "hypothalamic development". Differential grandmaternal exposure to feast or

famine seemed to have induced marks in "adipogenesis", "insulin-like growth factor reception

(IGF-R) binding", "ketone body catabolic process" and "adrenal gland development" (Table

1).

Three sentinel metabolic pathways implying transgenerational epigenetic inheritance.

GO:0030070 "indulin processing" influences the formation of insulin by proteolysis of the

precursor preproinsulin via C-peptide.This is the first signal sequence elevated from

preproinsulin. Proinsulin is then cleaved to release the C peptide. This pathway had a DMR in

the male line found in the Southern Shore involving 2 genes and 4 CpGs. Genes PCSK2

[proprotein convertas (also called NE2 HUMAN] and CPE Carboxypeptidase E (also called

CPBE HUMAN). The pathway is related to GO 1901142 "insulin metabolic process".

GO:1904179 , "adipose development" includes gene NR1H4 (also called BAR;FXR;HRR1;

RIP14) the nuclear hormone receptor pthr 24082, "bile acid receptor". Other genes are SIRT1

(also called SIR2L1) and chromatin regulatory protein sir2. phtr 11085

GO 21854 "hypothalamus development", involves GSX1(GShomebox 1), PTX2 (Pituitary

homebox 2), UBB (Polyubiquitin-B) and CRH (Corticoliberin)

Parental age data and selection

Average age at birth of first child was 32 years, 31 years and 27 years in the three generations

when grandparents had been exposed to famine and 34 years, 27 years and 27 years when

grandparents had been exposed to feast. All seven index persons were alive at the age of 74,

one died at the age of 63. For grandparents and parents exposed to famine during SGP, the

average age of death was 61 years and 78 years whereas when correspondingly exposed to

feast it was 87 years and 71 years (Table 2).

Discussion

We present a case series of epigenomes of 8 adult grandchildren with archival data exhibiting

their paternal grand-paternal or -maternal exposure to crop failure or large crops during their

SGP linked to differential methylation of genes associated with energy balance and

hypothalamic development. The salient novel finding is that grandparental childhood

exposure was reflected in grandchild profiles of CpG methylation of key genes.

In earlier papers we described a unique cohort of descendants of grandparents exposed

randomly to these weather-dependent variations in food availability. Data emanated from

detailed archival records enabling correlations between demographic and crop yield

statistics[1,2,3,4,5,6] . Our SGP findings have been confirmed in Germany for food deprivation[7]

and in Sweden for parental loss[8]. Our current paper adds mechanistic information based on

blood sampling of eight 75-year-old grandchildren enabling determination of candidate

epigenetic markers reflecting differential methylation in gene pathways posited to be affected

by exposure of their grandparents to crop failure versus bountiful harvest during the

grandparental childhood SGP preceding the prepubertal peak stature growth.

We propose that known adaptations to energy deficiency or excess in cells, organs or

organisms are expressed transgenerationally appearing as epigenetic marks in humans.

Famine might induce marks around promoters in at least two types of pathways: one general

or nutrient-specific, related to diminished substrate stores of glycogen, lipid or protein, and/or

excess, nutritoxic exposure , the other to environmental stresses of famine through activation

of the hypothalamo-pituitary-adrenal (HPA) axis. Based on our epidemiological findings of

transgenerational inheritance we posited that changes in DNA methylation levels of cytosine

and guanine (CpG probes) differentially methylated to form 5-methylcytosine affecting genes

in one generation can explain transgenerational epigenetic inheritance in the third generation[9].

A 2015 review concluded that evidence suggesting that acquired epigenetic marks are passed

to the next generation was limited and many other have noted the absence of any mechanism

by which gene-regulatory information is transferred from somatic cells to germ cells in the

study of transgenerational epigenetic inheritance, reviewed in Nagy and Turecki[10]. Our

epidemiological and epigenetic results demonstrate sex differences, e.g. in development of

preproinsulin as has been seen in human pancreatic islets[11]. The sex-bound epigenetic

inheritance has been interpreted simply as epigenetic actions and disease manifestations being

sex specific, as is the case with many conventional genetic variants[12]. The transcriptions are

also not fully understood. Most confusion emanates from not knowing how exposed somatic

cells can communicate their exposure to the germline which induces changes lasting for

generations.

It is difficult to get an insight into epigenetic inheritance if one believes in a real rather than

just a theoretical barrier between somatic and germline cells. One route overcoming the

barrier might be extracellular nano-vesicles of neighboring to germ cells shed by somatic cells

containing the material required for transcription able to bypass the barrier[13]. In the SGP the
male primordial germ cells form active spermatozoa whereas human ovarian stem cells that
can be modified in the SGP probably are required.

Presently the preponderance of evidence suggests transgenerational cumulative effects of
exposures: diet, behavior, environmental chemicals, activity and the microbiome. For reviews
see Sales et al.[9] and Vaiserman et al.[14].

We have focused on the paternal line to discern transgenerational epigenetic inheritance
disentangled from maternal-fetal and maternal-infant influences. The pathways in the paternal
line are probably similar in the maternal line. We found three interesting gene pathways
influenced by paternal grandparents´exposure resulting in grandchildren´s DMRs, pathways
induced by famine, overnutrition and food insecurity stress. When the paternal grandfather
had been exposed to famine, the grandson exhibited DMRs of insulin processing. When the
paternal grandmother had been exposed the granddaughter had DMRs of GOs related to the
environmental stress such as "hypothalamus development"(influencing the HPA-axis) related
to increased female susceptibility to stress, yet explaining the benefit to mental health recently
found in a different setting[7].

A strength of this study is that it exploits a natural secular phenomenon viz. variable food
security, the effects of which are documented in a homogeneous well documented 19[th]
century population, allowing exceptional tracking of ancestors. Few founders colonized the
area 500 years earlier whereby the area metaphorically became an island of native speakers of
Swedish surrounded by Sami- and Finnish-speaking neighbors. Furthermore the index
persons, the grandchildren are dispersed all over Sweden[5] and have neither been exposed to
famine nor, being born before WW2, experienced current food excesses during their
childhood.

A second strength is the focus on two lines of inheritance that emerged in the epidemiological
studies, the line from the paternal grandmother´s son and his daughter and the line from the
paternal grandfather with son and grandson. Thirdly our *a priori* selection of gene ontologies,
GOs, based on our earlier epidemiological findings of cardiovascular morbidity related to
food security reduces the risk of spurious correlations. A confound of multiple testing could
be suspected having 40 GOs but the ontologies were chosen for their evidence based
responses to only three potential pathways: feast, famine and stress.

Our earlier published studies in the community and correlation of age at the birth of the first

child, number of children and variance of survival over three generations ruled out that

variable availability of food during SGP was caused selectively[2]. In human studies measures

are very crude but the figures for the eight subjects do not demonstrate any biased selection

owing to ancestors´random exposure during the SGP. The main weakness is having only 8

index cases causing low statistical power. Furthermore, these 8 were survivors aged 75 years

during which their own exposures might have induced methylations confounding those

inherited from the grandparents and parents. On the other hand epimutations analogous to

mutation most often do not appear on the clinical horizon until late in life.

Confounding could have occurred through differences in the exposure to the feast and famine

between index cases owing to social circumstances such as being member of a family with

better food resources, or genes affecting sensitivity to undernutrition, or cellular heterogeneity

differentially affecting allele-specific or other variables´ methylation[12] . However, the natural

experiment exposed all families, showing little variation in poverty in the 19[th] century

mitigating such confound, supported by our earlier research in the community[4] . The

isolation of the community during the centuries up to the present might have resulted in an in-

bred cohort and diminished assortative mating, both benefitting the epigenetic analysis.

Human transgenerational inheritance still awaits more studies in other settings. Two questions

are pressing. How many generations can the epigenetic marks in humans persist? Our own

recent unpublished studies in a different setting indicate that it persists at least for four

generations. The effects and mechanisms of interaction between genetic and epigenetic

inheritance also remain to be determined.

We describe in a small case study of 8 subjects the presence of epigenetic changes attributable

to environmental influences potentially engendering several multifactorial diseases and

conditions pontentially providing guidance for preventive and therapeutic interventions

targeting methylation.

Methods

Classification of exposure to food availability in the environment

A tradition of using a scale for crop estimates was introduced for the years 1816-1849 in

Statistics Sweden (Tabellverket) by the 19[th] century statistician Hellstenius[15]. It was a seven-

degree scale running from total failure to good or bountiful crop and has been used since in

demographic research. For the years 1865-1902, time of our 8 grandparents´ births, statistical

tables of crop yields were the primary source, validated against food price price statistics[16],

and qualitative reports from a gubernatorial office (Hushållningssällskapet) of crops and

government aid. Using these sources, we created an ordinal seven-item "Hellstenius scale".

Over the years the criteria for the items of the scale have changed apace with changes in the

Swedish language. The current translation is as follows: Total failure (0), Sparse general

growth (1), Weak or Small (2), Below average (3), Average or mediocre (4), Above average

(5) and Good or bountiful (6) crop. We defined 0-2 as famine, 3-4 as mediocre, and 5-6 as

excess food availability. We empirically used May 1 of the year following crop failure as the

time of least food availability and November 1 of the year following good harvest and

slaughtering of pigs and cattle as the time of greatest food availability (excess). Grandparent

age on those dates was used to relate food availability to his or her SGP.

SGP, The slow growth period

Two periods in our first analysis were posited to differ in demand for food during famines and

consequently to differ in transgenerational responses: the prepubertal growth peak and the

precedimg slow growth period. These two periods were derived by superimposing the stature

growth velocity curves of Prader et al[17] and the ages of 19[th] century pubarche described by

Tanner[18]. The periods for ancestors in the 19[th] century were set at the ages 8-11 years for

female ancestors and at 9-12 years of age for male ancestors. Excess food was not considered

to be an issue in the 19[th] century Överkalix but to analyze potential transgenerational

responses we were obliged to focus on the SGP were we indeed found mismatches reflecting

transgenerational responses not only to famine but also to excess food availability. Thus, we

introduced the concept of the "slow growth period"[1], the causes and mechanisms of which we

discuss here in depth.

Samples and pedigrees

Eight subjects, alive at 75 years of age and consenting to blood sampling were selected from the 1935 birth cohort in an original epidemiologic study of transgenerational responses to variable availability of food during ancestors' SGP in Överkalix, Sweden, described in Tinghög et al[5]. Two paternal grandmothers were exposed during SGP to famine (grade 0-2) in 1867 and 1900 respectively, and two were exposed to surfeit (grade 5-6) in 1871 and 1879. Two paternal grandfathers were exposed to famine (grade 0-2) in1867 and 1877 respectively and two were exposed to surfeit (grade 5-6) in 1887 and 1881.These subjects represented four pairs of grandchildren of paternal grandparents exposed to feast or famine (Fig1).

Whole blood samples in EDTA were drawn at county health services units close to the subjects´ residence, frozen at 70^o C and sent to the project center. Pedigrees were originally tracked at parish offices in Sweden and Finland and presently in the National archive, the Regional archive of north Sweden, the Research archive of Umeå University, the National tax bureau and digitalized genealogical net-based registers.

Metabolic pathways posited

The gene ontologies chosen *a priori* described processes related to cardiovascular risks and were related to hunger, excess food and reactions to environmental stress. (http://amigo.geneontology.org). Posited processes during *famine* were "glucose homeostasis" and "glucose transport", "ketone body biosynthesis and catabolic processes", "cellular ketone body metabolic pathways", and "cholesterol homeostasis". In *hunger and excess* "insulin processing and binding", insulin-like growth factor", "insulin receptor signaling and "autophagy pathways" were posited. Individuals´ reactions to disasters were assumed to influence hypothalamus and adrenal gland development. Availability and absorption of folate stimulating methyl donors was of obvious interest. Other posited GOs include appetite, ghrelin, leptin, calcium signaling, eating behavior, adipose and fat cell development, insulin secretion, and lipids.

Altogether these added up to 40 pathways, many of them overlapping and covering three kinds of pathways reflecting the three posited reactions.

Genome-wide methylation

Genomic DNA extraction from whole-blood (in one case from buffy coat) was carried out

using GeneCatcher Blood kit (Invitrogen, Carlsbad, CA, USA). 500 ng of genomic DNA was

bisulfite converted with EZ-96 DNA Methylation kit (Zymo Research, Irvine, CA, USA) and

genome wide DNA methylation analysis was carried out using the Infinium MethylationEPIC

Bead Chip (Illumina, San Diego, CA, USA). The Laboratory procedures were done according

to the manufacturers´ protocol.

The array was designed for genome wide methylation analysis with coverage across gene

regions with sites in the promoter region, 5UTR, enhancer and gene body, and interrogating

more than 850 000 methylation sites at single nucleotide resolution. This technique uses two

different probe types (Infinium 1 and 2) with different characteristics, thus requiring

normalization to reduce technical bias. For analysis, visualization and extraction of

methylation data GenomeStudio software version 2011.1 (Illumina Inc) was used.

Methylation levels (beta values) were estimated as the ratio of signal intensity of the

methylated alleles to the sum of methylated and unmethylated intensity signals. The beta-

values vary from 0 (no methylation) to 1 (100% methylation).

The chip covers CpG islands, shores, shelves and the promoted genes. GOs containing genes

and biological pathways were assigned from the literature, primarily from the Gene Ontology

project (geneontology.org) chosen from their known intra-generational relation to excess and

lack of food.

The assay protocol combined bisulfite conversion of genomic DNA and whole-genome

amplification with array-based capture and scoring of the CpG loci. Signal intensity was

measured by scanner to generate beta values, the degree of methylation at a locus. Allele-

specific single base extension of the probes incorporated a biotin nucleotide or a dinitrophenyl

labeled nucleotide. Signal amplification of the incorporated label further improved the overall

signal-to-noise ratio of the assay.

Human genomes punctuated by DNA sequences with high frequencies of CpG sites, >200

bases, with CpG content of 50% were termed "islands" (CGIs). Differential methylation

induced by environment visible in CGIs or in "northern- and southern-*shores*", within 2 kilo

base-pairs (<2kbp) of the islands, and "northern- and southern *shelves*" <2 kbp from the

shores were studied in CGIs, shores and shelves in grandchildren of grandparents having

experienced extremes of food availability during their childhood SGP.

Statistical analysis

The association between genes and GO terms was retrieved from Gene Ontology Association

(UniProt-GOA) Database, http//www.ebi.ac.uk/GOA.

Number of genes associated with a GO term and represented with at least one probe on the

EPIC chip and number of probes associated with the genes were analyzed among index cases,

the grandchildren.

Differentially methylated female index cases whose paternal grandmothers were exposed to

bumper harvests in their mid-childhood SGP and to crop failure respectively were recorded as

well as male index cases whose paternal grandfather had the same extreme exposures.

The criteria for differential methylation was a p-value < 0.05 and an average difference in

beta-value between the group averages >0.1. The share of differentially methylated probes

was given as percent.

Tables were prepared after beta-mixture quantile normalization (BMIQ) correcting probe

design bias. Tables were prepared with the BMIQ-normalised beta values and annotations.

The analyses were carried out in Bioconductor-package ChAMP package default probe

filtering based on detection p-value, minimum bead count and probes possibly confounded by

SNPs and cross hybridisation (https//www.ncbi.nim.nih.gov/pubmed/24063430) left 813007

probes for further analysis.

References

1.Bygren LO, Kaati G, Edvinsson E. Longevity determined by paternal ancestors´nutrition
during their slow growth period. Acta Biotheoretica 2001;49:53-59.

2.Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality
determined by nutrition during parents' and grandparents' slow growth period.
Eur J Hum Genet 2002;10:682–8.
283

284 3.Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K et al. Sex-specific, male-line
285 transgenerational responses in humans. Eur J Hum Genet 2006;14:159-166

286 4.Kaati G, Bygren LO, Pembrey M, Sjostrom M. Transgenerational response to
287 nutrition, early life circumstances and longevity. Eur J Hum Genet 2007;15:784–90.
288

289 5.Tinghög P, Carstensen J, Kaati G, Edvinsson S, Sjöström M, Bygren LO.
290 Migration and mortality trajectories: a study of individuals born in therural community of
291 Överkalix. Soc Sci Med. 2011;73:744-51.
292

6.Bygren LO, Tinghög P, Carstensen J, Edvinsson S, Kaati G, Pembrey ME,
Sjöström M. Change in paternal grandmothers' early food supply influenced
cardiovascular mortality of the female grandchildren. BMC Genet 2014;15:12.

7.Van den Berg GJ, Pinger PR. Transgenerational effects of childhood conditions on third
generation health and educational outcomes. EconHumBiol 2016;23:103-120.

8.Vågerö D, Rajaleid K. Does childhood trauma influence offspring´s birth
characteristics?IntJEpidemiol 2016;

9.Sales VM, Ferguson-Smith AC, Patti ME. Epigenetic mechanisms of transmission of
metabolic disease across generations. Cell Met 2017;25:559-571.
302

303 10.Nagy C, Turecki G.Transgenerational epigenetic inheritance: an open discussion.
304 Epigenomics 2015;7:781-790.

305 11.Hall E, Volkov P, Dayeh T, Esguerra JLS, Salö S et al. Sex difference in the genome-wide
306 DN A methylation pattern and impact on gene expression, microRNA levels and insulin
307 secretion in human pancreatic islets. Genome Biology 2014;15:522-544.
308

12.Schaefer S, Nadeau JH. The genetics of epigenetic inheritance: Modes, molecules and
mechanisms Volume 90, Quart Review Biology 2015;90: Pages

13.Earon SA, Jayasooriah N, Buckland ME, Buckland ME, Martin DI, Cropley JE, Suter CM.
Roll over Weismann: extracellular vesicles in the transgenerational transmisson of
environmental effects Epigenomics 2015; 7:1165-71.

14. Vaiserman AM, Koliada AK, Jirtles RL. Non-genomic transmission of longevity between
generations: potential mechanisms and evidence across species. Epigenetics Chromatin
2017;DOI 10.1186/s 13072-017-0145-1

15. Hellstenius, J. Skördarna i Sverige och deras verkningar. [Harvests in Sweden and their
repercussions]. Statistisk Tidskrift 77-119. Stockholm, Sweden 1871 (sic).

16. Jörberg, L. (1972). A history of prices in Sweden 1732–1914. CWK Gleerup, Lund,
Sweden.

17. Prader, A., R.H. Largo, L. Molinari and C. Issler (1989). Physical growth of Swiss
children from birth to 20 years of age. First Zurich Longitudinal Study of Growth and
Development. Helvetica Paediatrica Acta 43:Suppl 52: 1-125

18. Tanner, J.M. A History of the Study of Human Growth. Cambridge University Press,
Cambridge 1981

Figure 1. Flowchart

Table 1. Gene ontology pathways describing reactions to hunger, excess food and stress at paternal
grandparent´s exposure to variable food availability, followed by DMRs in grandchildren

Gene Ontology pathway	Genes	CpGs	MethylCpG	DMR%	Shore, Shelf[3]
Paternal grandfather with son and grandson					
32100 Positively regulating appetite	3	163	10	6%	Northern shore
30070 Insulin processing[1]	2	4	1	25%	Southern shore
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
21854 Hypothalamus development[2]	6	33	5	15%	Southern shelf
Paternal grandmother with son and granddaughter					
1904179 Positive adipose development	2	11	2	18%	Northern shore
70341 Fat cell proliferation[4]	5	16	3	18%	Southern shelf
51599 IGF receptor binding[5]	9	94	12	13%	Northern shelf
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
30325 Adrenal gland development	12	57	4	7%	Southern shelf

1. The formation of mature insulin by proteolysis of the precursor preproinsulin. The signal sequence
is first elevated from preproinsulin, proinsulin is then cleaved to release the Cpeptide, leaving the A
and B chain of mature insulin.
2. The progression of the hypothalamus region of the forebrain from its initial formation to its
mature state.
3. CpG sites >200 bases with CpG content of 50% are termed "islands" (CGIs). Differential
methylation induced by environment are most often seen in "northern- and southern-shores" <2kb
from the islands, and "northern- and southern shelves" <2kb apart of the shores.
4. The multiplication or reproduction of fat cells resulting in expansion of their population.
5. Interacting selectively and non-covalently with the insulin-like growth factor receptor.

Table 2. Reproductive fitness following grandparental famine and feast in Slow Growth Period

Index person	Age at first child´s birth			Number of children		Longevity		
Exposure	Grandparent	Father	Grandchild	Father	Grandchild	Grandparent	Father	Grandchild
Paternal grandfathers exposure								
Famine								
Index person No 15	39	38	NA	4	NA	60	69	74+
Index person No 53	27	31	27	7	1	49	98	74+
Feast								
Index person No 14	28	23	30	6	3	80	70	74+
Index person No 23	33	26	NA	11	NA	94	83	63
Paternal grandmothers exposure								
Famine								
Index person No 59	39	38	34	3	2	80	83	74+
Index person No 86	24	20	22	6	2	55	64	74+
Feast								
Index person No 3	36	29	23	10	4	86	65	74+
Index person No 22	35	31	29	4	2	89	68	74+

340 NA. Not available