1
2
3

Title: Paternal grandparental exposure to crop failure or surfeit during a childhood slow growth period: Epigenetic marks on grandchildren´s growth, glucoregulatory and stress genes

4
5

Authors: Lars Olov Bygren[1,2*], Patrick Müller[1], David Brodin[1], Gunnar.Kaati[1], Jan-Åke Gustafsson[1], John G. Kral[3]

Affiliations:

7
8

[1] Department of Biosciences and Nutrition, Karolinska Institutet, SE_14183 Huddinge, Sweden.

9
10

[2] Departments of Community Medicine and Rehabilitation, Umeå University, SE_90187, Umeå, Sweden.

11
12

[3] Departments of Surgery, Medicine, Cell Biology, SUNY Downstate Medical Center, 450 Clarkson Avenue, Box 40, Brooklyn, NY 11203, USA.

[*] Correspondence to: lars.olov.bygren@ki.se

ABSTRACT

This latest in our series of papers describes transgenerational methylation related to mid-

childhood food availability in 19[th] century Överkalix, Sweden. Failed vs. bountiful crops

differentially influenced methylation in grandchildren of paternal grandparents exposed to

feast or famine during their Slow Growth Period (SGP), a sensitive period preceding the pre-

pubertal growth spurt. In this case-study of 8 tracked 75-year old progeny with differential

ancestral exposure, we found, in 40 posited gene ontology pathways, 39 differentially

methylated CpG regions (DMRs) related to famine, excess food and food-insecurity stress, 9

of which with DMRs above 5%. Three gene ontology terms (GOs) "insulin processing",

"adipose development" and "hypothalamus development" were key, with DMRs >14%.

An unbiased test of known pathways revealed four nuclear transcription factors upstream of

promotors repressing the pathway following paternal grandparental famine experience, as

well as 4 upregulated GOs with average DMRs >20%. We conclude that this is the first

demonstration of human transgenerational inheritance of epigenetic marks following

ancestral childhood exposure to variable food availability inducing early developmental

origins of adult disease.

Paternal grandparent food supply, preceding their prepubertal growth spurt, induced

epigenetic marks in three gene pathways reflecting famine, excess food and food-insecurity

stress. Our epidemiological findings of adverse transgenerational effects of ancestral

overnutrition and conversely, beneficial effects of famine during SGP[1-6] prompted us to

study individuals whose paternal grandparents were randomly exposed during their Slow

Growth Period (SGP) - a sensitive period preceding the pre-pubertal growth spurt - to failed

or bumper crops potentially engendering epigenetic marks on mechanistic molecular gene

pathways.

For detailed analyses of specific gene ontology (GO) pathways, we tested three posited

structural classes of genes associated with each of 40 preselected GO terms

(http://amigo.geneontology.org) for differentially methylated CpG regions (DMRs) between

the experimental groups. We selected GO pathways with average methylation differing >

5%. The posited pathways comprised glucoregulatory genes interpreted as immediate

sensors of decreased glycogen stores, whereas those of lipids and ketones were posited to

reflect the duration and magnitude of the famine or alternatively, responses to

overnutrition. Food insecurity stress was assumed to have activated the HPA-axis. We

focused on two putative response lines in direct descendants from a) grandfather to son

with grandson and b) paternal grandmother to son with granddaughter, adjusted for crop

size during famine or surfeit. We found 39 DMRs in grandchildren after grandparental

differential exposure to food availability in GO-pathways associated with famine, excess food

and environmental stressors. Nine of them were large, three GOs exhibiting remarkable

DMRs in grandchildren (Table 1). An unbiased analysis detected transcriptional pathways,

such as CCAAT-box (AKA C/EBPbeta) affecting adipogenesis, laying down fat, the primal

tissue for energy storage.

RESULTS

Exploiting an "experiment" of nature, we studied DNA methylation in grandsons and

granddaughters in the paternal lineage. We found 39 DMRs among 40 posited pathways in

islands, North and South *shores* and North and South *shelves* related to famine, nutrient

excess and stress, among which 9 pathways had DMRs above 5%, and 3 above 14%. The

other 31 pathways had 0-5% DMRs. An unbiased test of pathways in *islands* revealed 2

pathways with DMRs 40% and 2 with DMRs 22 and 21% (Table 1b).

Paternal grandparental exposure vs. methylated genes in adult grandchildren.

The main metabolic pathways implicated in grandpaternal SGP exposure to food-insecurity

stress transferred to grandsons were "appetite", "insulin processing", "ketone body

catabolic process" and "hypothalamic development". Differential grandmaternal exposure to

feast or famine was associated with marks in "adipogenesis", "insulin-like growth factor

receptor (IGF-R) binding", "ketone body catabolic process" and "adrenal gland

development" (Table 1). The probands' 4 ketone body catabolism genes (GO 30070) were

the same for grandfathers´ as for grandmothers´ exposures , supporting the findings. The

directions of the differences were intuitive: famine decreased energy-demanding pathways

whereas overnutrition enhanced them. Altogether three pathways were key.

Three key gene ontology terms implying transgenerational epigenetic inheritance.

GO 0030070: "insulin processing" modulates insulin production downstream from

proteolysis of the precursor preproinsulin via C-peptide. Proinsulin is cleaved to release C

peptide. This pathway had a DMR in the male line in the Southern Shore involving 2 genes:

PCSK2, proprotein convertase (also called NE2 HUMAN) in chromosome 20 and CPE

Carboxypeptidase E (also called CPBE HUMAN) in chromosome 2. This pathway is related to

GO 1901142: "insulin metabolic process".

GO 1904179: "adipose development" includes gene NR1H4 (also called BAR;FXR;HRR1;

RIP14) and the nuclear hormone receptor pthr 24082, "bile acid receptor". Other genes are

SIRT1 (also called SIR2L1) and chromatin regulatory protein sir2, pthr 11085.

GO 0021854: "hypothalamus development" involves GSX1 (GShomeobox1), PTX2 (Pituitary

homeobox2), UBB (Polyubiquitin-B) and CRH (Corticoliberin; Corticotropin-releasing

hormone).

Unbiased correlations (Table 1b).

Regardless of sex, the following gene ontology terms were highly correlated with famine:

GO 0016602: A transcription complex that binds to the CCAAT-box upstream of promotors

typically consisting of three subunits, i.e. NFYa-c. Four genes were repressed: NFYA on

Chromosome 6, NFYB on chromosome 12, NFYC on chromosome 1 and ING2 on

chromosome 4.

GO 0045540: "Regulation of cholesterol biosynthesis" (formation of cholesterol) was

repressed in 13 genes, such as ERLINI, chromosome 8 and 10; SECI4L2, chromosome 22 and

SODI, chromosome 21.

Several cell differentiation pathways correlated with food surplus, notably the following two:

GO 0030854: "Positive regulation of granulocyte differentiation" available for expression of

its 9 genes, e.g. HAXI on chromosome 1, OGT on the X chromosome, ILS on chromosome 5

and HCSL1 on chromosome 3.

GO 00 45616: "Regulation of keratinocyte differentiation" with 16 genes, such as HAXI, OGT,

ILS and C1QC on chromosome 1.

Parental age data and selection

Average paternal ages at birth of first child were 32 , 31 and 27 years in the three

generations with grandparents exposed to famine and 34 , 27 and 27 years with

grandparents exposed to feast. Seven index persons were alive at 74, the eighth having died

at 63 years. For grandparents and parents exposed to famine during SGP, the average ages

of death were 61 and 78 years respectively whereas after feast they were 87 and 71 years

(Table 2). Sequential genetics findings contributing to differential methylation were not

taken into account.

DISCUSSION

We propose that known adaptations to energy deficiency or excess in cells, organs or

organisms are expressed transgenerationally, appearing as epigenetic marks in humans.

Famine might induce marks around promoters in at least two types of pathways: one

general or nutrient-specific, related to *diminished* substrate stores of glycogen, lipid or

protein, or *excess*, nutritoxic exposure (cf. eutrophication in plants), the other to

environmental stresses of famine through activation of the hypothalamo-pituitary-adrenal

(HPA) axis, as in food insecurity.

In earlier papers we described a unique cohort of descendants of grandparents exposed

randomly to these weather-dependent variations in food availability. Data emanated from

detailed archival records enabling correlations between demographic and crop yield

statistics[1-6]. Our finding of SGP sensitivity findings have been replicated in Germany and

Sweden [7,8]. Our current paper adds mechanistic information based on blood sampling of

seven 75-year-old grandchildren and one 63 years old, enabling determination of candidate

epigenetic markers reflecting differential methylation in gene pathways posited to be

affected by exposure of their grandparents to crop failure versus bountiful harvest during

the grandparental childhood SGP preceding the prepubertal peak stature growth.

Changes in DNA methylation levels of cytosine and guanine (CpGs) differentially methylated

in one generation could explain transgenerational epigenetic inheritance in the third

generation[9].

A 2015 review concluded that evidence suggesting that acquired epigenetic marks pass to

the next generation was limited; many others have noted the absence of any mechanism by

which gene-regulatory information is transferred from somatic cells to germ cells in the

study of transgenerational epigenetic inheritance (review: Nagy and Turecki[10]). Our

epidemiological and epigenetic results demonstrate sex differences, e.g. in development of

preproinsulin as has been seen in human pancreatic islets[11]. The sex-bound epigenetic

inheritance has been interpreted simply as epigenetic actions and disease manifestations

being sex specific, as is the case with many conventional genetic variants[12]. The

transcriptions are also not fully understood. Most confusion emanates from not knowing

how exposed somatic cells can communicate their exposure to the germline which induces

changes lasting for generations.

It is difficult to understand or accept epigenetic inheritance if there is a real rather than just

a theoretical barrier between somatic and germline cells. One route to overcome this barrier

might be the proposal that extracellular nano-vesicles, neighboring germ cells, shed by

somatic cells contain the material required for transcription, able to bypass the barrier[13]. In

the SGP the male primordial germ cells form active spermatozoa whereas human ovarian

stem cells that can be modified in the SGP probably are required.

Presently the preponderance of evidence suggests transgenerational cumulative effects of

exposures: diet, behavior, environmental chemicals, activity and the microbiome. For

reviews, see Sales et al.[9] and Vaiserman et al.[14].

We have focused on the paternal line to discern transgenerational epigenetic inheritance

disentangled from maternal-fetal and maternal-infant influences. The pathways in the

paternal line are probably similar in the maternal line. We found three interesting gene

pathways influenced by paternal grandparents´ exposure resulting in grandchildren´s DMRs,

pathways induced by famine, overnutrition and food insecurity stress. When the paternal

grandfather had been exposed to famine, the grandson exhibited DMRs of insulin

processing. When the paternal grandmother had been exposed the granddaughter had

DMRs of GOs related to environmental stress such as "hypothalamus

development"(influencing the HPA-axis) related to increased female susceptibility to stress,

yet explaining the benefit to mental health recently found in a different setting[7].

A strength of our study is that it exploits a natural secular phenomenon viz. variable food

security, the effects of which are documented in a homogeneous well-documented 19[th]

century population, allowing exceptional tracking of ancestors. Few founders colonized the

area 500 years earlier whereby the area metaphorically became an island of native speakers

of Swedish surrounded by Sami- and Finnish-speaking neighbors. Furthermore the index

persons, the grandchildren, are dispersed all over Sweden[5] and have neither been exposed

to famine nor, being born before WW2, experienced current food excesses during their

childhood, preceding the current obesity epidemic. Differences in blood tissue composition

or heterogeneity of blood probably were of minor importance in this very homogenous

population.

A second strength is the focus on two lines of inheritance that emerged in the

epidemiological studies, the line from the paternal grandmother's son and his daughter and

the line from the paternal grandfather with son and grandson. Thirdly our *a priori* selection

of gene ontologies, GOs, based on our earlier epidemiological findings of cardiovascular

morbidity related to food security reduces the risk of spurious correlations. A confound of

multiple testing could be suspected having 40 GOs but the ontologies were chosen for their

evidence-based responses to only three potential pathways: feast, famine and stress. In

addition, our unbiased test identified related transcriptional genes.

Our earlier published studies in the community and correlation of age at the birth of the first

child, number of children and variance of survival over three generations ruled out that

variable availability of food during SGP was caused selectively[2]. In human studies, measures

are very crude but the figures for the eight subjects do not demonstrate any biased selection

owing to ancestors' random exposure during the SGP. The main weakness is having only 8

index cases, with low statistical power. Furthermore, seven were survivors up to age 75

years (one to 63) during which their own exposures might have induced methylations

confounding those inherited from the grandparents and parents. On the other hand,

epimutations analogous to mutation most often do not appear on the clinical horizon until

late in life. The subjects were sufficiently healthy to participate and travel to the nearest

health facility to give blood.

Confounding could have occurred through differences in exposure to the feast and famine

between index cases owing to social circumstances such as belonging to a family with better

food resources, or to other mechanisms differentially affecting allele-specific or other

variables' methylation[12]. However, the natural experiment exposed all families with little

variation in poverty in the 19[th] century mitigating this confound, as supported by our earlier

research in the community[4]. The isolation of this community during the centuries up to the

present might have resulted in a relatively inbred cohort and diminished assortative mating,

both benefitting the epigenetic analysis. We present a case series of epigenomes of 8 adult

grandchildren with archival data exhibiting their paternal grand-paternal or -maternal

exposure to crop failure or large crops during their SGP linked to differential methylation of

genes associated with energy balance and hypothalamic development. The salient novel

finding is that grandparental childhood exposure was reflected in grandchild profiles of CpG

methylation of key genes. Our epidemiological data from the cohort from which these

strategic cases were drawn strengthens the molecular findings and indicates possible

mechanisms of inheritance that might be important for the field. The presence of epigenetic

changes attributable to environmental influences potentially engendering several prevalent

multifactorial chronic diseases and conditions. These findings replicated in other settings

may provide guidance for preventive and therapeutic interventions targeting methylation.

METHODS

Classification of exposure to food availability in the environment

A seven point scale for estimating annual crop sizes was employed during 1816-1849

[Statistics Sweden (Tabellverket)][15]. It spanned from "total failure" to "abundant" crops and

has been used since in demographic research. For the years 1865-1902, birth years of the 8

grandparents , statistical tables of crop yields were the primary source, validated against

food price statistics[16] and qualitative reports from a gubernatorial *Crops and National Aid*

Office (Hushållningssällskapet). Using these sources, we created the ordinal seven-item

"Hellstenius scale". Over the years the criteria for the items of the scale have changed apace

with changes in the Swedish language. The current translation is as follows: Total failure (0),

Sparse general growth (1), Weak or Small (2), Below average (3), Average or Mediocre (4),

Above average (5) and Good or Abundant (6) . We defined 0-2 as famine, 3-4 as mediocre,

and 5-6 as excess food availability. We used May 1 of the year following crop failure as the

time of least food availability in this area around the 66[th] latitude and determined November

1 of the year following good harvest with pig and cattle slaughter when the meat could be

frozen, as the time of greatest food availability (excess,eutrophic). Grandparent age on those

dates was used to relate food availability to his or her SGP.

SGP, the slow growth period

Two developmental periods in our first analysis were posited to differ in sensitivity to food

supply affecting transgenerational responses: the prepubertal growth peak and the

preceding slow growth period (SGP). The growth peak was seen as a sensitive period for a

child to be exposed to famine. The slowest growth period before the peak was posited to be

less prone to energy variation. These two periods were set in relation to puberty, taking into

account individual variation, nutrient availability shifting the timeframes, and realizing that

the age distribution of puberty was skewed to the right during the 19[th] century compared to

later. The two periods were derived by superimposing the stature growth velocity curves of

Prader et al.[17] upon the ages of 19[th] century pubarche described by Tanner[18]. The periods

for ancestors in the 19[th] century were set at the ages 8-10 years for female ancestors and at

9-12 years of age for male ancestors. Excess food was not understood to be a problem in the

19[th] century in Överkalix but to analyze potential transgenerational responses we were

obliged to focus on the SGP where we indeed found mismatches reflecting transgenerational

responses not only to famine but also to excess food availability. We introduced the concept

of the "slow growth period"[1], the causes and mechanisms of which we discuss here in depth.

Samples and pedigrees

Eight subjects, seven of which 75 years of age and one 63, consented to blood sampling.

They were selected from the 1935 birth cohort in an original epidemiologic study of

transgenerational responses to variable availability of food during ancestors' SGP in

Överkalix, Sweden, described in Tinghög et al.[5] During SGP two paternal grandmothers were

exposed to famine (grade 0-2) in 1867 and 1900 respectively, and two to surfeit (grade 5-6)

in 1871 and 1879. Two paternal grandfathers were exposed to famine (grade 0-2) in 1867

and 1877 respectively and two were exposed to surfeit (grade 5-6) in 1887 and 1881.These

subjects represented four pairs of grandchildren of paternal grandparents exposed to feast

or famine (Fig1). Individuals inheriting grandparents who had been exposed to medium

crops (grade 3-4) were not included in this strategic sample.

Whole blood samples in EDTA were drawn at county health service units close to the

subjects´ residence, frozen at 70°C and sent to the project center. Pedigrees were originally

tracked at parish offices in Sweden and Finland and presently in the National archive, the

Regional archive of north Sweden, the Research archive of Umeå University, the National tax

bureau and digitized genealogical net-based registries. Detailed possible confounders were

not identified in this natural experiment: probands, the grandchildren, did not experience

any lifetime exposure to famine. The modern food availability though was there from their
young adulthood.

Metabolic pathways posited

The gene ontologies chosen *a priori* described processes related to cardiovascular risks
related to hunger, excess food and reactions to environmental stress
(http://amigo.geneontology.org). Posited processes during *famine* were "glucose
homeostasis" and "glucose transport", "ketone body biosynthesis and catabolic processes",
"cellular ketone body metabolic pathways", and "cholesterol homeostasis". In *hunger and*
excess "insulin processing and binding", "insulin-like growth factor", "insulin receptor
signaling" and "autophagy pathways" were posited. Individuals´ emotional reactions to the
stress of famine were assumed to influence hypothalamic and adrenal gland development.
Availability and absorption of folate providing methyl donors was of obvious interest, though
not the focus of this report. Other posited GOs include appetite, ghrelin, leptin, calcium
signaling, eating behavior, adipogenesis, insulin secretion, and lipids.

Altogether, 40 pathways were tested, many of them overlapping and covering three kinds of
response pathways reflecting the 3 posited mechanisms.

Pathways undergoing unbiased testing

We tested all gene ontology terms in a public database, (UniProt-GOA) including about
40 000 gene ontologies, for grandchildren's responses to paternal grandparental exposure to
poor *versus* excess food.

Genome-wide methylation

Genomic DNA extraction from whole-blood (buffy coat in one) used the GeneCatcher Blood
kit (Invitrogen, Carlsbad, CA, USA). 500 ng of genomic DNA was bisulfite converted with EZ-
96 DNA Methylation kit (Zymo Research, Irvine, CA, USA) and genome wide DNA
methylation analysis was carried out using the Infinium MethylationEPIC Bead Chip (Illumina,
San Diego, CA, USA). The Laboratory procedures followed the manufacturers´ protocol.

The array was designed for genome wide methylation analysis with coverage across gene
regions with sites in the promoter region, 5UTR, enhancer and gene body, and interrogating

more than 850 000 methylation sites at single nucleotide resolution. This technique uses two

different probe types (Infinium 1 and 2) with different characteristics, thus requiring

normalization to reduce technical bias. GenomeStudio software, version 2011.1 (Illumina

Inc.), was used for analysis, visualization and extraction of methylation data.

Methylation levels (beta values) were estimated as the ratio of signal intensity of the

methylated alleles to the sum of methylated and unmethylated intensity signals. The beta-

values vary from 0 (no methylation) to 1 (100% methylation).

The chip covers CpG islands, shores, shelves and the promoted genes. GOs containing genes

and biological pathways were assigned from the literature, primarily from the Gene

Ontology project (geneontology.org) chosen for their known intra-generational relation to

excess and lack of food.

The assay protocol combined bisulfite conversion of genomic DNA and whole-genome

amplification with array-based capture and scoring of the CpG loci. Signal intensity was

measured by scanner to generate beta values, the degree of methylation at a locus. Allele-

specific single base extension of the probes incorporated a biotin nucleotide or a

dinitrophenyl labeled nucleotide. Signal amplification of the incorporated label further

improved the overall signal-to-noise ratio of the assay.

Human genomes punctuated by DNA sequences with high frequencies of CpG sites, >200

bases, with CpG content of 50% were termed "*islands*" (CGIs). Differential methylation

induced by environment visible in CGIs or in "northern- and southern-*shores*", within 2 kilo

base-pairs (<2kbp) of the islands or in "northern- and southern-*shelves*" within another 2 kilo

base pairs from the shores (<2kbp) were studied in CGIs in grandchildren of grandparents

exposed to extremes of food availability during their childhood SGP. They are presented as

DMRs expressed as percent.

Statistical analysis

Associations between genes and GO terms were retrieved from the Gene Ontology

Association (UniProt-GOA) Database, http//www.ebi.ac.uk/GOA.

Number of genes associated with a GO term and represented with at least one probe on the

EPIC chip and number of probes associated with the genes were analyzed among index

cases, the grandchildren.

Differentially methylated female index cases whose paternal grandmothers were exposed to

bumper harvests in their mid-childhood SGP and to crop failure, respectively, were recorded

as well as male index cases whose paternal grandfather had the same extreme exposures.

The criteria for differential methylation was a p-value < 0.05 and an average difference in

beta-value between the group averages >0.1. No adjustment for multiple comparisons was

carried out. The decreased power of genome-wide analysis, by having negative controls in

the remainder of the genome, was compensated by the strict criteria chosen.

Tables were prepared after beta-mixture quantile normalization (BMIQ) correcting probe

design bias and with BMIQ-normalized beta values and annotations. The analyses were

carried out in Bioconductor-package ChAMP package default probe filtering based on

detection p-value, minimum bead count and probes possibly confounded by SNPs

whereupon cross hybridization (https//www.ncbi.nim.nih.gov/pubmed/24063430) left

813007 probes for further analysis.

REFERENCES

1.Bygren LO, Kaati G, Edvinsson E. Longevity determined by paternal ancestors´ nutrition
during their slow growth period. Acta Biotheoretica 2001;49:53-59.

2.Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality
determined by nutrition during parents' and grandparents' slow growth period.
Eur J Hum Genet 2002;10:682–8.
332

333   3.Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K et al. Sex-specific, male-line
334   transgenerational responses in humans. Eur J Hum Genet 2006;14:159-166

335   4.Kaati G, Bygren LO, Pembrey M, Sjostrom M. Transgenerational response to
336   nutrition, early life circumstances and longevity. Eur J Hum Genet 2007;15:784–90.
337

338   5.Tinghög P, Carstensen J, Kaati G, Edvinsson S, Sjöström M, Bygren LO.
339   Migration and mortality trajectories: a study of individuals born in the rural community of
340   Överkalix. Soc Sci Med. 2011;73:744-51.
341

6.Bygren LO, Tinghög P, Carstensen J, Edvinsson S, Kaati G, Pembrey ME,

Sjöström M. Change in paternal grandmothers' early food supply influenced
cardiovascular mortality of the female grandchildren. BMC Genet 2014;15:12.

7.Van den Berg GJ, Pinger PR. Transgenerational effects of childhood conditions on third
generation health and educational outcomes. EconHumBiol 2016;23:103-120.

8. Vågerö D, Pinger PR, Aronsson V, van den Berg G. Paternal grandfather´s access to food
predicts all-cause and cancer mortality in grandsons. Nat Comm

9.Sales VM, Ferguson-Smith AC, Patti ME. Epigenetic mechanisms of transmission of
metabolic disease across generations. Cell Met 2017;25:559-571.
351
352   10.Nagy C, Turecki G. Transgenerational epigenetic inheritance: an open discussion.
353   Epigenomics 2015;7:781-790.

354   11.Hall E, Volkov P, Dayeh T, Esguerra JLS, Salö S et al. Sex difference in the genome-wide DN
355   A methylation pattern and impact on gene expression, microRNA levels and insulin secretion
356   in human pancreatic islets. Genome Biology 2014;15:522-544.
357
12. Guo YT, LUO HV, Wu YM, Magdalou J, Chen LB, Wang H. Influencing factors, underlying
mechanisms and interactions affecting hypercholesterolemia in adult offspring with caffeine
exposure during pregnancy, Reprod Toxicol 2018;79:47-56

13.Earon SA, Jayasooriah N, Buckland ME, Buckland ME, Martin DI, Cropley JE, Suter CM. Roll
over Weismann: extracellular vesicles in the transgenerational transmisson of environmental
effects Epigenomics 2015; 7:1165-71.

14.Vaiserman AM, Koliada AK, Jirtles RL. Non-genomic transmission of longevity between
generations: potential mechanisms and evidence across species. Epigenetics Chromatin
2017;DOI 10.1186/s 13072-017-0145-1

15.Hellstenius, J. Skördarna i Sverige och deras verkningar. [Harvests in Sweden and their
repercussions]. Statistisk Tidskrift 77-119. Stockholm, Sweden 1871 (sic).
369
16.Jörberg, L. (1972). A history of prices in Sweden 1732–1914. CWK Gleerup, Lund,
Sweden.

17.Tanner, J.M. A History of the Study of Human Growth. Cambridge University Press,
Cambridge 1981

18.Prader, A., R.H. Largo, L. Molinari and C. Issler (1989). Physical growth of Swiss children
from birth to 20 years of age. First Zurich Longitudinal Study of Growth and Development.
Helvetica Paediatrica Acta 43:Suppl 52: 1-125

Figure 1. Flowchart

Table 1. Gene ontology pathways related to hunger, excess food and stress associated with
paternal grandparental exposure to variable food availability with DMRs in grandchildren.

Gene Ontology pathway	Genes	CpGs	MethylC pG	DMR %	Shore, Shelf[3]
Paternal grandfather with son and grandson					
32100 Positively regulating appetite	3	163	10	6%	Northern shore
30070 Insulin processing[1]	2	4	1	25%	Southern shore
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
21854 Hypothalamus development[2]	6	33	5	15%	Southern shelf
Paternal grandmother with son and granddaughter					
1904179 Positive adipose development	2	11	2	18%	Northern shore
70341 Fat cell proliferation[4]	5	16	3	18%	Southern shelf
51599 IGF receptor binding[5]	9	94	12	13%	Northern shelf
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
30325 Adrenal gland development	12	57	4	7%	Southern shelf

380
381

[1]Mature insulin formed by proteolysis of the precursor pre-proinsulin. The sequence starts
with pre-proinsulin; cleavage of proinsulin leads to release of C-peptide, leaving the A and B
chain of mature insulin.
[2]Evolution of the hypothalamic region of the forebrain from its initial formation to its mature
state.
[3]CpG sites with >200 bases with CpG content of 50% are termed "islands" (CGIs). Differential
methylation induced by environment is most often seen in "northern- and southern-shores"
<2kb from the islands, and "northern- and southern shelves" <2kb apart from the shores.
[4]Multiplication or recruitment of fat cells leading to expansion of their population
(adipogenesis).
[5]Interacting selectively and non-covalently with insulin-like growth factor receptor.

Table 1 b

395
396 **Gene ontology (GO) pathways in 8 grandchildren (4 men, 4 women) whose paternal grandparents were exposed to famine or food excess.**

397 ____

	Gene ontology pathway %	GO name	DMR
399 400			
401			
	Exposure		
GO:0016602ˇ	Famine	CCAAT binding complex	40
GO:0045540ˇ	Famine	Regulation of cholesterol biosynthetic process	40
GO:0030854ˋ	Excess	Positive regulation of granulocyte differentiation	22
GO:0030852ˇ	Excess	Regulation of keratinocyte differentiation	21

402 **Beta cutoff 0.02, Methylation difference >15%.**

		CpGs	Methylated CpGs		p-value,	exponent
404 405						
406			observed	expected		
407	GO: 0016602[a]	47	5	0.255	6.13	-6
408	GO: 0045540[b]	407	5	0.221	3.12	-6
409						
410	GO: 0030854[c]	75	6	0.017	1.53	-14
411	GO: 0030852[d]	260	7	0.059	1.95	-13

412 ____

413
____________ Numbers of genes in GO pathways, their ID, chromosome #:

[a] n= 4 NFYA, 6; NFYB, 12; NFYC, 1; ING2, 4.

[b] n=13 ERLIN1, 10; ERLIN2, 8; SEC14L2, 22; SOD1, 21; APOE, 19; APOB, 2;

FGF1, 5; POR, 7; SREBF1, 17; ABCG1, 21; SCAP, 3; PRKAA1, 5; DHCR7, 11.

[c] n= 9 HAX1, 1; OGT, X; IL-5, 5; HCLS1, 3; RUNX1, 21; KMT2E, 7; TESC, 12;

TRIB1, 8; LEF1, 4.

[d] n=16 HAX1, 1 ; OGT, X ; C1QC, 1 ; IL-5, 5 ; RARA, 17 ; CEACAM1, 19 ; HCLS1, 3 ;

RUNX1, 21 ; CUL4A, 13 ; ADIPOQ, 3 ; ZBTB46, 20 ; KMT2E, 7 ;

INPP5D, CHR_HG2232_PATCH; TESC, 12; TRIB1, 8; LEF1, 4.

Table 2. Reproductive fitness following grandparental famine or feast in the Slow Growth
Period

Index person	Age at first child´s birth			Number of children		Longevity		
Exposure	Grandparent	Father	Grandchild	Father	Grandchild	Grandparent	Father	Grandchild
Paternal grandfathers' exposure								
Famine								
Index person No 15	39	38	NA	4	NA	60	69	74+
Index person No 53	27	31	27	7	1	49	98	74+
Feast								
Index person No 14	28	23	30	6	3	80	70	74+
Index person No 23	33	26	NA	11	NA	94	83	63
Paternal grandmothers' exposure								
Famine								
Index person No 59	39	38	34	3	2	80	83	74+
Index person No 86	24	20	22	6	2	55	64	74+
Feast								
Index person No 3	36	29	23	10	4	86	65	74+
Index person No 22	35	31	29	4	2	89	68	74+

NA. Not available

17