Title: Paternal grandparental exposure to crop failure or surfeit during a childhood slow
growth period and epigenetic marks on third generational growth-, glucoregulatory and stress
genes

Authors: Lars Olov Bygren[1,2], Patrick Müller[1], David Brodin[1], Gunnar.Kaati[1], Jan-Åke
Gustafsson[1], John G. Kral[3*]

Affiliations:

[1] Department of Biosciences and Nutrition, Karolinska Institutet, SE_14183 Huddinge,
Sweden.

[2] Departments of Community Medicine and Rehabilitation, Umeå University, SE_90187,
Umeå, Sweden.

[3] Departments of Surgery, Medicine, Cell Biology, SUNY Downstate Medical Center, 450
Clarkson Avenue, Box 40, Brooklyn, NY 11203, USA.

[*] Correspondence to: lars.olov.bygren@ki.se

ABSTRACT

This latest in our series of papers describes transgenerational methylation related to mid-

childhood food availability in 19[th] century Överkalix, Sweden. Failed vs. bountiful crops

differentially influenced methylation in grandchildren of paternal grandparents exposed to

feast or famine during their Slow Growth Period (SGP), a sensitive period preceding the pre-

pubertal growth spurt. In this case-study of 8 tracked 75-year old progeny with differential

ancestral exposure, we found in 40 posited gene ontology pathways 39 differentially

methylated CpG regions (DMRs) related to famine, excess food and food-insecurity stress, 9

of which with DMRs above 5%. Three pathways named "insulin processing", "adipose

development" and "hypothalamus development" were sentinel with DMRs >14%. This is the

first demonstration of a human transgenerational inheritance of epigenetic marks following

childhood exposure to variable food availability indicating early developmental origins of

adult disease.

Paternal grandparent food supply, preceding their prepubertal growth spurt, induced

epigenetic marks in three gene pathways reflecting famine, excess food and reaction to food-

insecurity stress. Our epidemiological findings of adverse transgenerational effects of

ancestral overnutrition and conversely, beneficial effects of famine during SGP[1] prompted us

to study individuals whose paternal grandparents were randomly exposed, during their Slow

Growth Period (SGP) - a sensitive period preceding the pre-pubertal growth spurt - to failed

or bumper crops potentially engendering epigenetic marks on mechanistic molecular gene

pathways.

For detailed analyses of specific gene ontology (GO) pathways, we tested the three posited

structural classes of genes associated with each of 40 preselected GO terms

(http://amigo.geneontology.org) for differentially methylated CpG regions (DMRs) between

the experimental groups. We selected GO pathways with average differences in methylation

between groups >0.1 and p-values <0.05 and performed detailed analyses of those differing >

5%. The posited pathways comprised glucoregulatory genes that were seen as immediate

sensors of decreased glycogen stores, whereas lipids and ketones were posited to reflect the

duration and magnitude of the famine or responses to overnutrition. Individual reactions to

food insecurity were assumed to have activated the HPA-axis. We focused on two putative

response lines in direct descendants from a) grandfather to son with grandson and b) paternal

grandmother to son with granddaughter, adjusted for crop size during famine or surfeit. We

found 39 DMRs in grandchildren after grandparental differential exposure to food availability

in GO-pathways associated with famine, excess food and reactions to environmental stressors.

Nine of them were large, three GOs exhibiting remarkable DMRs in grandchildren (Table 1).

An unbiased analysis indicated transcriptional pathways, such as the CCAAT-box (data not

shown).

RESULTS

Exploiting a natural experiment, we studied the paternal lineage transgenerational responses

in grandsons and granddaughters and assessed their DNA-methylations. We found 39 DMRs

among 40 posited pathways in *islands*, North and South *shores* and North and South *shelves*

related to famine, excess and stress, among which 9 pathways had DMRs above 5%, and 3

above 14%. The other 31 pathways had 0-5% DMRs.

Paternal grandparental exposure vs. methylated genes in adult grandchildren.

The main metabolic pathways implicated in grandpaternal SGP exposure to food-insecurity

stress transferred to grandsons were "appetite", "insulin processing", "ketone body catabolic

process" and "hypothalamic development". Differential grandmaternal exposure to feast or

famine were associated with marks in "adipogenesis", "insulin-like growth factor reception

(IGF-R) binding", "ketone body catabolic process" and "adrenal gland development" (Table

1). The 4 genes of the probands for ketone body catabolism (GO 30070) were the same for

grandfathers´ and grandmothers´ transgenerational response, strengthening the analysis. The

directions of the differences were logical, famine inducing decreased energy-demanding

pathways and feast enhancing them. Three were sentinel and the genes involved were the

following:

Three sentinel metabolic pathways implying transgenerational epigenetic inheritance.

GO 0030070: "insulin processing" influences insulin from proteolysis of the precursor

preproinsulin via C-peptide. This is the first signal sequence elevated from preproinsulin.

Proinsulin is then cleaved to release the C peptide. This pathway had a DMR in the male line

found in the Southern Shore involving 2 genes: PCSK2, proprotein convertase (also called

NE2 HUMAN) in chromosome 20 and CPE Carboxypeptidase E (also called CPBE

HUMAN) in chromosome 2. The pathway is related to GO 1901142: "insulin metabolic

process".

GO 1904179: "adipose development" includes gene NR1H4 (also called BAR;FXR;HRR1;

RIP14), the nuclear hormone receptor pthr 24082, "bile acid receptor". Other genes are SIRT1

(also called SIR2L1) and chromatin regulatory protein sir2, pthr 11085.

GO 21854: "hypothalamus development" involves GSX1 (GShomeobox1), PTX2 (Pituitary

homeobox2), UBB (Polyubiquitin-B) and CRH (Corticoliberin; Corticotropin-releasing

hormone).

Parental age data and selection

Average age at birth of first child was 32 years, 31 years and 27 years in the three generations

with grandparents exposed to famine and 34 years, 27 years and 27 years with grandparents

exposed to feast. All seven index persons were alive at age 74 , (with one up to age 63). For

grandparents and parents exposed to famine during SGP, the average age of death was 61

years and 78 years whereas when correspondingly exposed to feast it was 87 years and 71

years (Table 2). Sequential genetics findings contributing to differential methylation were not

taken into account.

DISCUSSION

We present a case series of epigenomes of 8 adult grandchildren with archival data exhibiting

their paternal grand-paternal or -maternal exposure to crop failure or large crops during their

SGP linked to differential methylation of genes associated with energy balance and

hypothalamic development. The salient novel finding is that grandparental childhood

exposure was reflected in grandchild profiles of CpG methylation of key genes. We are aware

of the low power of tests but our epidemiological knowledge of the cohort from which these

strategic cases were drawn strengthens the molecular findings and the indication of possible

mechanisms of inheritance could be important for the field.

In earlier papers we described a unique cohort of descendants of grandparents exposed

randomly to these weather-dependent variations in food availability. Data emanated from

detailed archival records enabling correlations between demographic and crop yield

statistics[1-6]. Our SGP findings have been duplicated in Germany for food deprivation[7] and in

Sweden for parental loss[8]. Our current paper adds mechanistic information based on blood

sampling of seven 75-year-old grandchildren and one 63 years old, enabling determination of

candidate epigenetic markers reflecting differential methylation in gene pathways posited to

be affected by exposure of their grandparents to crop failure versus bountiful harvest during
the grandparental childhood SGP preceding the prepubertal peak stature growth.
We propose that known adaptations to energy deficiency or excess in cells, organs or
organisms are expressed transgenerationally, appearing as epigenetic marks in humans.
Famine might induce marks around promoters in at least two types of pathways: one general
or nutrient-specific, related to diminished substrate stores of glycogen, lipid or protein, and/or
excess, nutritoxic exposure , the other to environmental stresses of famine through activation
of the hypothalamo-pituitary-adrenal (HPA) axis. Based on our epidemiological findings of
transgenerational inheritance we posited that changes in DNA methylation levels of cytosine
and guanine (CpG probes) differentially methylated to form 5-methylcytosine affecting genes
in one generation can explain transgenerational epigenetic inheritance in the third generation[9].

A 2015 review concluded that evidence suggesting that acquired epigenetic marks pass to the
next generation was limited; many others have noted the absence of any mechanism by
which gene-regulatory information is transferred from somatic cells to germ cells in the study
of transgenerational epigenetic inheritance (review: Nagy and Turecki[10]). Our
epidemiological and epigenetic results demonstrate sex differences, e.g. in development of
preproinsulin as has been seen in human pancreatic islets[11]. The sex-bound epigenetic
inheritance has been interpreted simply as epigenetic actions and disease manifestations being
sex specific, as is the case with many conventional genetic variants[12]. The transcriptions are
also not fully understood. Most confusion emanates from not knowing how exposed somatic
cells can communicate their exposure to the germline which induces changes lasting for
generations.

It is difficult to understand or accept epigenetic inheritance if one believes in a real rather than
just a theoretical barrier between somatic and germline cells. One route to overcome this
barrier might be the proposal that extracellular nano-vesicles neighboring germ cells shed by
somatic cells contain the material required for transcription, able to bypass the barrier[13]. In
the SGP the male primordial germ cells form active spermatozoa whereas human ovarian
stem cells that can be modified in the SGP probably are required.

Presently the preponderance of evidence suggests transgenerational cumulative effects of
exposures: diet, behavior, environmental chemicals, activity and the microbiome. For
reviews, see Sales et al.[9] and Vaiserman et al.[14].

We have focused on the paternal line to discern transgenerational epigenetic inheritance

disentangled from maternal-fetal and maternal-infant influences. The pathways in the paternal

line are probably similar in the maternal line. We found three interesting gene pathways

influenced by paternal grandparents´ exposure resulting in grandchildren´s DMRs, pathways

induced by famine, overnutrition and food insecurity stress. When the paternal grandfather

had been exposed to famine, the grandson exhibited DMRs of insulin processing. When the

paternal grandmother had been exposed the granddaughter had DMRs of GOs related to

environmental stress such as "hypothalamus development"(influencing the HPA-axis) related

to increased female susceptibility to stress, yet explaining the benefit to mental health recently

found in a different setting[7].

A strength of this study is that it exploits a natural secular phenomenon viz. variable food

security, the effects of which are documented in a homogeneous well-documented 19[th]

century population, allowing exceptional tracking of ancestors. Few founders colonized the

area 500 years earlier whereby the area metaphorically became an island of native speakers of

Swedish surrounded by Sami- and Finnish-speaking neighbors. Furthermore the index

persons, the grandchildren, are dispersed all over Sweden[5] and have neither been exposed to

famine nor, being born before WW2, experienced current food excesses during their

childhood. Differences in blood tissue composition or heterogeneity of blood probably were

of minor importance in this very homogenous population.

A second strength is the focus on two lines of inheritance that emerged in the epidemiological

studies, the line from the paternal grandmother´s son and his daughter and the line from the

paternal grandfather with son and grandson. Thirdly our *a priori* selection of gene ontologies,

GOs, based on our earlier epidemiological findings of cardiovascular morbidity related to

food security reduces the risk of spurious correlations. A confound of multiple testing could

be suspected having 40 GOs but the ontologies were chosen for their evidence-based

responses to only three potential pathways: feast, famine and stress.

Our earlier published studies in the community and correlation of age at the birth of the first

child, number of children and variance of survival over three generations ruled out that

variable availability of food during SGP was caused selectively[2]. In human studies, measures

are very crude but the figures for the eight subjects do not demonstrate any biased selection

owing to ancestors´ random exposure during the SGP. The main weakness is having only 8

index cases, with low statistical power. Furthermore, seven were survivors up to age 75 years

(one to 63) during which their own exposures might have induced methylations confounding

those inherited from the grandparents and parents. On the other hand, epimutations analogous

to mutation most often do not appear on the clinical horizon until late in life. The subjects

were sufficiently healthy to participate and travel to the nearest health facility to give blood.

Confounding could have occurred through differences in exposure to the feast and famine

between index cases owing to social circumstances such as belonging to a family with better

food resources, or to other mechanisms differentially affecting allele-specific or other

variables´ methylation[12]. However, the natural experiment exposed all families with little

variation in poverty in the 19th century mitigating this confound, as supported by our earlier

research in the community[4]. The isolation of this community during the centuries up to the

present might have resulted in a relatively inbred cohort and diminished assortative mating,

both benefitting the epigenetic analysis.

Human transgenerational inheritance still awaits more studies in other settings. Two questions

are pressing. For how many generations can human epigenetic marks persist? Our own recent

unpublished studies in a different setting indicate that they persist for at least four generations.

The effects and mechanisms of interaction between genetic and epigenetic inheritance also

remain to be determined.

We describe in a small case study of 8 subjects the presence of epigenetic changes attributable

to environmental influences potentially engendering several prevalent multifactorial chronic

diseases and conditions. These findings may provide future guidance for preventive and

therapeutic interventions targeting methylation.

METHODS

Classification of exposure to food availability in the environment

A tradition of using a scale for crop estimates was introduced by the 19th century statistician

Hellstenius for the years 1816-1849 in Statistics Sweden (Tabellverket)[15]. It was a seven-

degree scale running from total failure to good or bountiful crop and has been used since in

demographic research. For the years 1865-1902, the period of our 8 grandparents´ births,

statistical tables of crop yields were the primary source, validated against food price

statistics[16] and qualitative reports from a gubernatorial office (Hushållningssällskapet) of

crops and government aid. Using these sources, we created an ordinal seven-item "Hellstenius

scale". Over the years the criteria for the items of the scale have changed apace with changes

in the Swedish language. The current translation is as follows: Total failure (0), Sparse

general growth (1), Weak or Small (2), Below average (3), Average or mediocre (4), Above

average (5) and Good or bountiful (6) crop. We defined 0-2 as famine, 3-4 as mediocre, and

5-6 as excess food availability. We used May 1 of the year following crop failure as the time

of least food availability in this area around the 66th latitude and November 1 of the year

following good harvest and slaughtering of pigs and cattle when the meat could be frozen, as

the time of greatest food availability (excess). Grandparent age on those dates was used to

relate food availability to his or her SGP.

SGP, the slow growth period

Two developmental periods in our first analysis were posited to differ in sensitivity to food

supply affecting transgenerational responses: the prepubertal growth peak and the preceding

slow growth period (SGP). The growth peak was seen as a serious period for a child to be

exposed to famine. The slowest growth period before the peak was posited to be less sensitive

to energy variation. These two periods were set in relation to puberty, taking into account

individual variation, nutrient availability shifting the timeframes, and realizing that the age

distribution of puberty was skewed to the right during the 19th century compared to later

times[17]. The two periods were derived by superimposing the stature growth velocity curves of

Prader et al.[17] and the ages of 19th century pubarche described by Tanner[18]. The periods for

ancestors in the 19th century were set at the ages 8-11 years for female ancestors and at 9-12

years of age for male ancestors. Excess food was not understood to be a problem in the 19th

century in Överkalix but to analyze potential transgenerational responses we were obliged to

focus on the SGP where we indeed found mismatches reflecting transgenerational responses

not only to famine but also to excess food availability. We introduced the concept of the

"slow growth period"[1], the causes and mechanisms of which we discuss here in depth.

Samples and pedigrees

Eight subjects, seven of which alive at 75 years of age (one at 63) and consenting to blood

sampling were selected from the 1935 birth cohort in an original epidemiologic study of

transgenerational responses to variable availability of food during ancestors' SGP in

Överkalix, Sweden, described in Tinghög et al.[5] Two paternal grandmothers were exposed

during SGP to famine (grade 0-2) in 1867 and 1900 respectively, and two were exposed to

surfeit (grade 5-6) in 1871 and 1879. Two paternal grandfathers were exposed to famine

(grade 0-2) in1867 and 1877 respectively and two were exposed to surfeit (grade 5-6) in 1887

and 1881.These subjects represented four pairs of grandchildren of paternal grandparents

exposed to feast or famine (Fig1). Individuals inheriting grandparents who had been exposed

to medium crops (grade 3-4) were not included in this strategic sample.

Whole blood samples in EDTA were drawn at county health service units close to the

subjects´ residence, frozen at 70^o C and sent to the project center. Pedigrees were originally

tracked at parish offices in Sweden and Finland and presently in the National archive, the

Regional archive of north Sweden, the Research archive of Umeå University, the National tax

bureau and digitized genealogical net-based registries. Detailed possible confounders were

not identified in this natural experiment: probands, the grandchildren, did not experience any

lifetime exposure to famine.

Metabolic pathways posited

The gene ontologies chosen *a priori* described processes related to cardiovascular risks and

were related to hunger, excess food and reactions to environmental stress

(http://amigo.geneontology.org). Posited processes during *famine* were "glucose homeostasis"

and "glucose transport", "ketone body biosynthesis and catabolic processes", "cellular ketone

body metabolic pathways", and "cholesterol homeostasis". In *hunger and excess* "insulin

processing and binding", "insulin-like growth factor", "insulin receptor signaling" and

"autophagy pathways" were posited. Individuals´ emotional reactions to famine were

assumed to influence hypothalamus and adrenal gland development. Availability and

absorption of folate stimulating methyl donors was of obvious interest. Other posited GOs

include appetite, ghrelin, leptin, calcium signaling, eating behavior, adipose and fat cell

development, insulin secretion, and lipids.

Altogether, 40 pathways were tested, many of them overlapping and covering three kinds of

response pathways reflecting the 3 posited reactions.

Genome-wide methylation

Genomic DNA extraction from whole-blood (buffy coat in one) used the GeneCatcher Blood

kit (Invitrogen, Carlsbad, CA, USA). 500 ng of genomic DNA was bisulfite converted with

EZ-96 DNA Methylation kit (Zymo Research, Irvine, CA, USA) and genome wide DNA

methylation analysis was carried out using the Infinium MethylationEPIC Bead Chip

(Illumina, San Diego, CA, USA). The Laboratory procedures were done according to the

manufacturers´ protocol.

The array was designed for genome wide methylation analysis with coverage across gene

regions with sites in the promoter region, 5UTR, enhancer and gene body, and interrogating

more than 850 000 methylation sites at single nucleotide resolution. This technique uses two

different probe types (Infinium 1 and 2) with different characteristics, thus requiring

normalization to reduce technical bias. For analysis, visualization and extraction of

methylation data GenomeStudio software version 2011.1 (Illumina Inc.) was used.

Methylation levels (beta values) were estimated as the ratio of signal intensity of the

methylated alleles to the sum of methylated and unmethylated intensity signals. The beta-

values vary from 0 (no methylation) to 1 (100% methylation).

The chip covers CpG islands, shores, shelves and the promoted genes. GOs containing genes

and biological pathways were assigned from the literature, primarily from the Gene Ontology

project (geneontology.org) chosen from their known intra-generational relation to excess and

lack of food.

The assay protocol combined bisulfite conversion of genomic DNA and whole-genome

amplification with array-based capture and scoring of the CpG loci. Signal intensity was

measured by scanner to generate beta values, the degree of methylation at a locus. Allele-

specific single base extension of the probes incorporated a biotin nucleotide or a dinitrophenyl

labeled nucleotide. Signal amplification of the incorporated label further improved the overall

signal-to-noise ratio of the assay.

Human genomes punctuated by DNA sequences with high frequencies of CpG sites, >200

bases, with CpG content of 50% were termed "*islands*" (CGIs). Differential methylation

induced by environment visible in CGIs or in "northern- and southern-*shores*", within 2 kilo

base-pairs (<2kbp) of the islands or in "northern- and southern-*shelves"* within another 2 kilo

base pairs from the shores (<2kbp) were studied in CGIs in grandchildren of grandparents

exposed to extremes of food availability during their childhood SGP are presented as DMRs

expressed as percent.

Statistical analysis

The association between genes and GO terms was retrieved from Gene Ontology Association

(UniProt-GOA) Database, http//www.ebi.ac.uk/GOA.

Number of genes associated with a GO term and represented with at least one probe on the

EPIC chip and number of probes associated with the genes were analyzed among index cases,

the grandchildren.

Differentially methylated female index cases whose paternal grandmothers were exposed to

bumper harvests in their mid-childhood SGP and to crop failure respectively were recorded as

well as male index cases whose paternal grandfather had the same extreme exposures.

The criteria for differential methylation was a p-value < 0.05 and an average difference in

beta-value between the group averages >0.1. No adjustment for multiple comparisons was

carried out. The decreased power of genome-wide analysis, by having negative controls in the

remainder of the genome, was compensated by the harsh criteria chosen and the logical choice

of GOs.

Tables were prepared after beta-mixture quantile normalization (BMIQ) correcting probe

design bias. Tables were prepared with the BMIQ-normalized beta values and annotations.

The analyses were carried out in Bioconductor-package ChAMP package default probe

filtering based on detection p-value, minimum bead count and probes possibly confounded by

SNPs and cross hybridization (https//www.ncbi.nim.nih.gov/pubmed/24063430) left 813007

probes for further analysis.

REFERENCES

1.Bygren LO, Kaati G, Edvinsson E. Longevity determined by paternal ancestors´ nutrition
during their slow growth period. Acta Biotheoretica 2001;49:53-59.

2.Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality
determined by nutrition during parents' and grandparents' slow growth period.
Eur J Hum Genet 2002;10:682–8.
316
317   3.Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K et al. Sex-specific, male-line
318   transgenerational responses in humans. Eur J Hum Genet 2006;14:159-166

319   4.Kaati G, Bygren LO, Pembrey M, Sjostrom M. Transgenerational response to
320   nutrition, early life circumstances and longevity. Eur J Hum Genet 2007;15:784–90.
321
322   5.Tinghög P, Carstensen J, Kaati G, Edvinsson S, Sjöström M, Bygren LO.
323   Migration and mortality trajectories: a study of individuals born in the rural community of
324   Överkalix. Soc Sci Med. 2011;73:744-51.
325
6.Bygren LO, Tinghög P, Carstensen J, Edvinsson S, Kaati G, Pembrey ME,
Sjöström M. Change in paternal grandmothers' early food supply influenced

cardiovascular mortality of the female grandchildren. BMC Genet 2014;15:12.

7.Van den Berg GJ, Pinger PR. Transgenerational effects of childhood conditions on third
generation health and educational outcomes. EconHumBiol 2016;23:103-120.

8.Vågerö D, Rajaleid K. Does childhood trauma influence offspring´s birth
characteristics?IntJEpidemiol 2016;

9.Sales VM, Ferguson-Smith AC, Patti ME. Epigenetic mechanisms of transmission of
metabolic disease across generations. Cell Met 2017;25:559-571.
335
336   10.Nagy C, Turecki G. Transgenerational epigenetic inheritance: an open discussion.
337   Epigenomics 2015;7:781-790.

338   11.Hall E, Volkov P, Dayeh T, Esguerra JLS, Salö S et al. Sex difference in the genome-wide
339   DN A methylation pattern and impact on gene expression, microRNA levels and insulin
340   secretion in human pancreatic islets. Genome Biology 2014;15:522-544.
341
12. Guo YT, LUO HV, Wu YM, Magdalou J, Chen LB, Wang H. Influencing factors,
underlying mechanisms and interactions affecting hypercholesterolemia in adult offspring
with caffeine exposure during pregnancy, Reprod Toxicol 2018;79:47-56

13.Earon SA, Jayasooriah N, Buckland ME, Buckland ME, Martin DI, Cropley JE, Suter CM.
Roll over Weismann: extracellular vesicles in the transgenerational transmisson of
environmental effects Epigenomics 2015; 7:1165-71.

14.Vaiserman AM, Koliada AK, Jirtles RL. Non-genomic transmission of longevity between
generations: potential mechanisms and evidence across species. Epigenetics Chromatin
2017;DOI 10.1186/s 13072-017-0145-1

15.Hellstenius, J. Skördarna i Sverige och deras verkningar. [Harvests in Sweden and their
repercussions]. Statistisk Tidskrift 77-119. Stockholm, Sweden 1871 (sic).
16.Jörberg, L. (1972). A history of prices in Sweden 1732–1914. CWK Gleerup, Lund,
Sweden.

17.Tanner, J.M. A History of the Study of Human Growth. Cambridge University Press,
Cambridge 1981

18.Prader, A., R.H. Largo, L. Molinari and C. Issler (1989). Physical growth of Swiss
children from birth to 20 years of age. First Zurich Longitudinal Study of Growth and
Development. Helvetica Paediatrica Acta 43:Suppl 52: 1-125

Figure 1. Flowchart

Table 1. Gene ontology pathways describing reactions to hunger, excess food and stress at paternal

grandparent´s exposure to variable food availability, followed by DMRs in grandchildren

Gene Ontology pathway	Genes	CpGs	MethylCpG	DMR%	Shore, Shelf[3]
Paternal grandfather with son and grandson					
32100 Positively regulating appetite	3	163	10	6%	Northern shore
30070 Insulin processing[1]	2	4	1	25%	Southern shore
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
21854 Hypothalamus development[2]	6	33	5	15%	Southern shelf
Paternal grandmother with son and granddaughter					
1904179 Positive adipose development	2	11	2	18%	Northern shore
70341 Fat cell proliferation[4]	5	16	3	18%	Southern shelf
51599 IGF receptor binding[5]	9	94	12	13%	Northern shelf
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
30325 Adrenal gland development	12	57	4	7%	Southern shelf

1. The formation of mature insulin by proteolysis of the precursor preproinsulin. The signal sequence

is first elevated from pre-proinsulin; proinsulin is then cleaved to release the C-peptide, leaving the A

and B chain of mature insulin.

2. The progression of the hypothalamus region of the forebrain from its initial formation to its

mature state.

3. CpG sites >200 bases with CpG content of 50% are termed "islands" (CGIs). Differential

methylation induced by environment are most often seen in "northern- and southern-shores" <2kb

from the islands, and "northern- and southern shelves" <2kb apart of the shores.

4. The multiplication or reproduction of fat cells resulting in expansion of their population.

5. Interacting selectively and non-covalently with the insulin-like growth factor receptor.

Table 2. Reproductive fitness following grandparental famine and feast in Slow Growth Period

Index person	Age at first child´s birth			Number of children		Longevity		
Exposure	Grandparent	Father	Grandchild	Father	Grandchild	Grandparent	Father	Grandchild
Paternal grandfathers exposure								
Famine								
Index person No 15	39	38	NA	4	NA	60	69	74+
Index person No 53	27	31	27	7	1	49	98	74+
Feast								
Index person No 14	28	23	30	6	3	80	70	74+
Index person No 23	33	26	NA	11	NA	94	83	63
Paternal grandmothers exposure								
Famine								
Index person No 59	39	38	34	3	2	80	83	74+
Index person No 86	24	20	22	6	2	55	64	74+
Feast								
Index person No 3	36	29	23	10	4	86	65	74+
Index person No 22	35	31	29	4	2	89	68	74+

376   NA. Not available