# Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of great crested newt (*Triturus cristatus*)

**Lynsey R. Harper[1]*, Lori Lawson Handley[1], Christoph Hahn[1,2], Neil Boonham[3,4], Helen C. Rees[5], Kevin C. Gough[6], Erin Lewis[3], Ian P. Adams[3], Peter Brotherton[7], Susanna Phillips[7] and Bernd Hänfling[1]**

[1]*School of Environmental Sciences, University of Hull, Hull, HU6 7RX, UK*

[2]*Institute of Zoology, University of Graz, Graz, Styria, Austria*

[3]*Fera, Sand Hutton, York, YO14 1LZ, UK*

[4]*Newcastle University, Newcastle upon Tyne, NE1 7RU, UK*

[5]*ADAS, School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK*

[6]*School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK*

[7]*Natural England, Peterborough, PE1 1NG, UK*

 **\*Corresponding author:** Lynsey Harper, L.Harper@2015.hull.ac.uk

**Manuscript type:** Standard article

**Running title:** Metabarcoding and qPCR detection sensitivity

**Word count:** 8716

## Summary

**1.** Environmental DNA (eDNA) analysis is a rapid, cost-effective, non-invasive biodiversity monitoring tool which utilises DNA left behind in the environment by organisms for species detection. The method is used as a species-specific survey tool for rare or invasive species in a broad range of ecosystems. More recently, eDNA and 'metabarcoding' have been combined to describe whole communities as opposed to focusing on single target species. However, whether metabarcoding is as sensitive as targeted approaches for rare species detection remains to be evaluated.

**2.** The great crested newt *Triturus cristatus* is a flagship pond species of international conservation concern and the first in the UK to be routinely monitored using eDNA. We evaluate whether eDNA metabarcoding has comparable sensitivity to targeted real-time quantitative PCR (qPCR) for great crested newt detection, and examine the implications for the future of community eDNA monitoring. Extracted eDNA samples (N = 532) were screened for great crested newt by qPCR and analysed for all vertebrate species using High-Throughput Sequencing technology.

**3.** Of all ponds, 50% tested positive for great crested newt with qPCR when the detection threshold was set to 1/12 positive qPCR replicates. Detection rate decreased to 32% when the threshold was increased to 4/12 positive qPCR replicates. With metabarcoding, newts were detected in 34% of ponds when no detection threshold was applied, and in 28% of ponds when a threshold (0.028%) was applied. Therefore, qPCR generates a higher detection rate than metabarcoding, but the metabarcoding detection rate with no threshold is broadly equivalent to that of qPCR with the stringent detection threshold. The proportion of great crested newt sequences in each sample was positively associated with the number of positive qPCR replicates (qPCR score)

suggesting eDNA metabarcoding may be indicative of species abundance.

**4.** eDNA metabarcoding holds enormous potential for holistic biodiversity assessment and routine

freshwater monitoring. We advocate this community approach to freshwater monitoring to

guide management and conservation, whereby entire communities can be initially surveyed to

best inform use of funding and time for species-specific surveys.

**Key-words:** environmental DNA (eDNA), great crested newt, High-Throughput Sequencing

(HTS), metabarcoding, ponds, real-time quantitative PCR (qPCR), *Triturus cristatus*.

## Introduction

Species monitoring has rapidly evolved with the advent of environmental DNA (eDNA) analysis (Lawson Handley 2015). eDNA analysis allows highly sensitive detection of rare and invasive species and is increasingly being used for surveys of aquatic species (Thomsen *et al.* 2012; Biggs *et al.* 2015; Davy, Kidd & Wilson 2015; Smart *et al.* 2015; Evans *et al.* 2017b). This non-invasive approach uses intra- and extracellular DNA (mucous, skin cells, faeces/urine, gametes, hair, deceased remains) released into the environment by organisms to survey for species and assess their distribution (Rees *et al.* 2014b; Lawson Handley 2015; Goldberg *et al.* 2016). Typically for eDNA analysis, DNA is extracted from environmental samples (water, soil, air) and analysed using a targeted or passive approach. The targeted approach uses species-specific primers with conventional PCR (PCR), real-time quantitative PCR (qPCR), or droplet digital PCR (ddPCR), to determine presence-absence and estimate abundance of single species (Shaw, Weyrich & Cooper 2016b; Goldberg *et al.* 2016). Conversely, the passive approach uses conserved primers and PCR to sequence whole communities with High Throughput Sequencing (HTS) (Shaw *et al.* 2016b; Valentini *et al.* 2016), commonly termed eDNA metabarcoding (Taberlet *et al.* 2012; Deiner *et al.* 2017). Passive eDNA monitoring is particularly attractive to ecologists for biodiversity assessment as a means to detect entire species assemblages alongside rare or invasive species (Lacoursière-Roussel *et al.* 2016; Blackman *et al.* 2017). However, this gain in community understanding may come at the cost of accuracy and sensitivity. Direct comparisons of these two approaches are essential to determine whether they have comparable power and yield similar results.

Although in its relative infancy, eDNA metabarcoding has proven effective for community biodiversity assessment across a range of taxa in varying environments, particularly freshwater

herpetofauna and fish (Valentini *et al.* 2016; Hänfling *et al.* 2016; Shaw *et al.* 2016a; Civade *et al.*

2016; Lacoursière-Roussel *et al.* 2016; Lopes *et al.* 2016; Evans *et al.* 2017a; Bálint *et al.* 2017).

However, eDNA metabarcoding is confounded by potential amplification bias during PCR,

preventing capture of all species present in a given area (Kelly *et al.* 2014). Species' DNA in

community samples is also in competition to bind to metabarcoding primers during PCR, where

more common templates are more likely to be amplified. High abundance species may thus prevent

detection of low abundance species, whether by fewer individuals or less DNA shed, resulting in

'species masking' (Kelly *et al.* 2014; Brandon-Mong *et al.* 2015). eDNA metabarcoding may

therefore be less capable of identifying eDNA of rare species DNA within a community than

species-specific qPCR (Goldberg *et al.* 2016).

The sensitivity of eDNA metabarcoding has been evaluated against conventional

biodiversity monitoring methods in freshwater ecosystems (Valentini *et al.* 2016; Hänfling *et al.*

2016; Shaw *et al.* 2016a; Civade *et al.* 2016; Lacoursière-Roussel *et al.* 2016; Lopes *et al.* 2016;

Evans *et al.* 2017a; Bálint *et al.* 2017), yet specific investigations comparing the sensitivity of

eDNA metabarcoding and targeted qPCR are sparse. Schnieder *et al.* (2016) achieved improved

detection of Invasive Mosquito Species (IMS) with targeted qPCR and eDNA metabarcoding as

opposed to conventional sampling. Although qPCR provided higher detection probability for two

species, metabarcoding achieved comparable results for a third species, and allowed simultaneous

detection of IMS and other taxa in a single sequencing run without development of multiple

species-specific markers. In another study, eDNA metabarcoding failed to detect wood turtle

*Glyptemys insculpta* (LeConte, 1830) in four rivers where targeted qPCR and conventional visual

survey detected the species (Lacoursière-Roussel *et al.* 2016). Amplification of longer fragments

during metabarcoding versus qPCR could account for difference in sensitivity of the two methods,

with the shorter qPCR assay being more capable of detecting heavily degraded DNA (Lacoursière-

Roussel *et al.* 2016). Further research is clearly needed to determine whether these two approaches

are comparable.

Great crested newt *Triturus cristatus* (Laurenti, 1768), hereafter crested newt, is a model

organism for eDNA-based monitoring. Crested newt secrete mucous, breed in water, and produce

aquatic eggs and larvae - all sources of DNA deposition in ponds. The species is rare in parts of

the UK and Europe, and as such all life stages are protected by UK and European legislation (Rees

*et al.* 2014a; Buxton *et al.* 2017). eDNA analysis using targeted qPCR has been repeatedly verified

against conventional surveying (bottle trapping, torchlight counts, larval netting, egg searches) for

crested newt and found to achieve comparable or improved species detection (Thomsen *et al.* 2012;

Rees *et al.* 2014a; Biggs *et al.* 2015). eDNA sampling also requires less training, is cost-efficient

(Biggs *et al.* 2014) and can be implemented in large-scale citizen science monitoring programmes

without loss of species detection (Biggs *et al.* 2015). Crested newt is the first species to be routinely

monitored using eDNA in the UK (Natural England 2015) and targeted eDNA assays are now

offered as a commercial service by ecological consultancies. The targeted eDNA assay is highly

effective for crested newt detection; however, should metabarcoding have comparable sensitivity,

this approach would allow detection of crested newt alongside pond communities and potentially

enable more cost-effective monitoring of entire ecosystems and ecological hypothesis testing.

Here, we perform a large-scale comparison (N = 532 ponds) of eDNA metabarcoding and

targeted qPCR for crested newt detection to compare method sensitivity. A single primer pair that

is vertebrate specific for mitochondrial DNA (mtDNA) and requires no *a priori* knowledge of

species composition was employed for eDNA metabarcoding. The metabarcoding results were

then compared to results obtained using the standard crested newt qPCR assay (Biggs *et al.* 2015).

Our hypotheses are: (1) eDNA metabarcoding will give equivalent results to qPCR for crested

newt detection, (2) eDNA metabarcoding sequence read count for crested newt will increase as

qPCR score (the number of positive qPCR replicates) increases, therefore eDNA metabarcoding

may be indicative of species abundance, and (3) metabarcoding primers will amplify DNA from

all taxa equally well and no bias towards amplification of crested newt will occur (bias would be

indicated by a positive association between the proportion of crested newt sequence reads and

PCR product concentration). We also examined the cost and investigator effort required by each

approach to determine whether a trade-off between cost, time and amount of data generated exists.

**Materials and methods**

SAMPLING

Samples from 532 ponds distributed across three UK counties (Cheshire, Kent and Lincolnshire)

were analysed for this project. Of these, 508 ponds were sampled as part of crested newt surveys

through Natural England's Great Crested Newt Evidence Enhancement Programme. Crested newt

egg searches were performed once during the daytime at 506/508 ponds. Any other life stages seen

were also recorded. A further 24 ponds were sampled for eDNA by ecological consultants for

private contracts but egg searches were not undertaken at these ponds. All water samples were

collected using methodology outlined by Biggs *et al.* (2015). Briefly, twenty 30 mL water samples

were collected at even intervals around the pond margin and pooled in a sterile 1 L Whirl-Pak®

stand-up bag, which was shaken to provide a single homogenised sample from each pond. Six 15

mL subsamples were taken from the mixed sample using a sterile plastic pipette (25 mL) and added

to sample tubes containing 33.5 mL absolute ethanol and 1.5 mL sodium acetate 3 M (pH 5.2) for

ethanol precipitation. Subsamples were then sent to Fera (Natural England), and ADAS (private

contracts) for eDNA analysis according to laboratory protocols established by Biggs *et al.* (2015).

Subsamples were centrifuged at 14,000 x g for 30 minutes at 6 °C, and the supernatant discarded.

Subsamples were then pooled during the first step of DNA extraction with the Qiagen® DNeasy

Blood & Tissue Kit, where 360 µL of ATL buffer was added to the first tube, vortexed, and the

supernatant transferred to the second tube. This process was repeated for all six tubes. The

supernatant in the sixth tube, containing concentrated DNA from all six subsamples, was

transferred in a 2 mL tube and extraction continued following manufacturer's instructions to

produce one eDNA sample per pond.

TARGETED qPCR FOR CRESTED NEWT

Prior to testing for crested newts, all extracted samples were tested for PCR inhibitors and sample

degradation using methodology outlined by Biggs *et al.* (2015), where an Internal Positive Control

was included in qPCR reactions of eDNA samples and a sample considered inhibited if replicates

showed different $C_q$ values (where samples move into the exponential phase of qPCR

amplification). Targeted qPCR was carried out as part of the crested newt monitoring programmes

mentioned above in the laboratories at Fera and ADAS during 2015 using a standardised protocol

(Biggs *et al.* 2015). Extracted DNA was amplified by TaqMan probe qPCR using published

primers and probe to amplify an 81 bp fragment of the cytochrome b gene: TCCBL (5'-

CGTAAACTACGGCTGACTAGTACGAA-3'), TCCBR (5-

CCGATGTGTATGTAGATGCAAACA) and TCCB_Probe (5'-

CCACGCTAACGGAGCCTCGC-3') (Thomsen *et al.* 2012). PCR reactions were set up in a total

volume of 25 µL consisting of: 3 µL of extracted template DNA, 1 µL of each primer (0.4 µM), 1

µL of probe (0.1 µM), 1x TaqMan® Environmental Master Mix 2.0 (containing AmpliTaq GOLD

DNA polymerase, Life Technologies) and $ddH_2O$. The PCR included an initial incubation for 5

min at 50 °C, then a 10 min denaturation step at 95 °C, followed by 55 cycles of denaturation at

95 °C for 30 s and annealing at 56.3 °C for 1 min. For each sample, 12 qPCR replicates were

performed and a sample recorded as positive for crested newt if one or more qPCR replicates were

positive. Positive (crested newt DNA: 1 x $10^{-1}$ ng/µL to 1 x $10^{-4}$ ng/µL) and negative controls

($ddH_2O$) were also included on each plate in quadruplicate. Following qPCR, the eDNA samples

were placed in storage at -80 °C.

METABARCODING OF VERTEBRATE COMMUNITIES

eDNA samples were stored at -20 °C until PCR amplification. Metabarcoding was performed using

published vertebrate-specific primers F1 (5'-ACTGGGATTAGATACCCC-3') and R1 (5'-

TAGAACAGGCTCCTCTAG-3') (Riaz *et al.* 2011) which amplify a 73-110 bp fragment of 12S

ribosomal RNA gene (rRNA). The assay was first validated *in silico* using ecoPCR software

(Ficetola *et al.* 2010; Bellemain *et al.* 2010) against a custom, phylogenetically curated reference

database for UK vertebrates. Full details of reference database construction are provided in

Supporting Information (Appendix 1). The complete reference database compiled in GenBank

format has been deposited in the dedicated GitHub repository for this study:

https://github.com/HullUni-bioinformatics/Harper_et_al_2017. Parameters set allowed a 50-250

bp fragment and maximum of three mismatches between the primer pair and each sequence in the

reference database. Primers were then validated against tissue DNA extracted from UK amphibian

species (Supporting Information: Appendix 2) having been previously validated *in vitro* for UK

fish communities by Hänfling *et al.* (2016). After primer validation, a two-step PCR protocol was

used to construct metabarcoding libraries from the eDNA samples. During the first PCR, the target

region was amplified using metabarcoding primers, comprised of the aforementioned specific

locus primer, random hexamers, sequencing primer and pre-adapter (Illumina 2011). DNA from

the cichlid *Rhamphochromis esox* (Boulenger, 1908) was used for PCR positive controls (six per

PCR plate; N = 114), whilst sterile molecular grade water (Fisher Scientific) substituted template

DNA for No Template Controls, NTCs (six per PCR plate; N = 114). In the second PCR, Molecular

Identification (MID) tags (unique 8-nucleotide sequences) and Illumina MiSeq adapter sequences

were added to the amplified product. Two independent libraries were constructed, each containing

266 eDNA samples, 57 NTCs, and 57 positive controls. Sequencing was performed on an Illumina

MiSeq using 2 x 300 bp V3 chemistry at Fera. The first sequencing run revealed human

contamination across samples and in some PCR controls, therefore reactions prepared for the

second sequencing run were sealed with mineral oil to minimise PCR contamination. Full details

of the eDNA metabarcoding workflow are provided in Supporting Information (Appendix 2).

BIOINFORMATIC PROCESSING

Illumina data was converted from raw sequences to taxonomic assignment using a custom pipeline

for reproducible analysis of metabarcoding data: metaBEAT (metaBarcoding and eDNA Analysis

Tool)    v0.8    (https://github.com/HullUni-bioinformatics/metaBEAT).    Bioinformatic    data

processing/analysis largely followed the workflow outlined by Hänfling *et al.* (2016), with minor

modifications. In brief, raw reads were quality trimmed using Trimmomatic v0.32 (Bolger, Lohse

& Usadel 2014), both from the read ends (minimum per base phred score Q30), as well as across

sliding windows (window size 5bp; minimum average phred score Q30). Reads were clipped to a

maximum length of 110 bp and reads shorter than 90 bp after quality trimming were discarded. To

reliably exclude adapters and PCR primers, the first 25 bp of all remaining reads were also

removed. Sequence pairs were merged into single high quality reads using FLASH v1.2.11 (Magoč

& Salzberg 2011), if a minimum of 10 bp overlap with a maximum of 10% mismatch was detected

between pairs. For reads that were not successfully merged, only forward reads were kept. To

reflect our expectations with respect to fragment size, a final length filter was applied and only

sequences of length 80-120 bp were retained. These were screened for chimeric sequences against

our custom reference database using the uchime algorithm (Edgar *et al.* 2011), as implemented in

vsearch v1.1 (Rognes *et al.* 2016). Redundant sequences were removed by clustering at 97%

identity ('--cluster_fast' option) in vsearch v1.1 (Rognes *et al.* 2016). Clusters represented by less

than five sequences were considered sequencing error and omitted from further analyses. Non-

redundant sets of query sequences were then compared against our custom reference database

using BLAST (Zhang *et al.* 2000). For any query matching with at least 98% identity to a reference

sequence across more than 80% of its length, putative taxonomic identity was assigned using a

lowest common ancestor (LCA) approach based on the top 10% BLAST matches. Sequences that

could not be assigned (non-target sequences) were subjected to a separate BLAST search against

the complete NCBI nucleotide (nt) database at 98% identity to determine the source via LCA as

described above. To ensure reproducibility of analyses, the described workflow has been deposited

in the GitHub repository.

DATA ANALYSIS

All downstream analyses were performed in the statistical programming environment R v.3.3.2.

(R Core Team 2016). Data and R scripts have been deposited in the GitHub repository.

Manipulation of the dataset produced by metaBEAT is described in Supporting Information:

Appendix 3.

*Detection thresholds and contamination*

At present, there are no standard guidelines for eDNA analysis to indicate minimum number of

positive eDNA samples or replicates required to class sites as species positive (Goldberg *et al.*

2016). Samples analysed by qPCR in this study were previously considered crested newt positive

if one or more qPCR replicates gave a positive result (Biggs *et al.* 2015). We term this analysis

qPCR NT (No Threshold). This inference of species presence is employed across many studies but

may not be reliable or reproducible (Goldberg *et al.* 2016). More stringent qPCR thresholds

reduced detection sensitivity for palmate newt *Lissotriton vulgaris* (Smart *et al.* 2015) but may be

necessary to ensure consistency and prevent false positives (Rees *et al.* 2014b). To facilitate

comparison with current qPCR scoring (our NT interpretation) and eDNA metabarcoding, we also

applied a stringent qPCR threshold of four or more positive qPCR replicates (out of 12) to infer

species presence, and termed the new analysis qPCR TA (Threshold Applied).

The raw eDNA metabarcoding dataset with no detection thresholds applied was termed

metabarcoding NT (No Threshold). A second dataset was constructed to reduce the potential for

false positives by application of a species-specific threshold: a species was only classed as present

at a given site if its sequence frequency exceeded a species-specific threshold. Thresholds for each

species were defined by analysing sequence data from PCR positive controls (N = 114) and

identifying the maximum sequence frequency for a given species across all PCR positive controls

(Supporting Information: Table S2). For example, the species-specific false positive sequence

threshold for great crested newt was 0.028% to omit all false detections in the PCR positive

controls. The resultant dataset was termed metabarcoding TA (Threshold Applied).

We tested whether mineral oil reduced contamination by analysing the distribution of

positive control sequences (*R. esox*) and human DNA in eDNA samples, and any DNA in NTCs,

across both sequencing runs using binomial generalised linear mixed effects models (GLMMs)

within the R package 'lme4' v1.1-12 (Bates *et al.* 2015). The response variable was presence-

absence of contamination and explanatory variables were PCR plate (random effect) and

sequencing run i.e. mineral oil sealed versus non-sealed (fixed effect). Human DNA may be

present in eDNA samples as a real environmental signal or contaminant prior to PCR and thus may

not be a true PCR contaminant. Consequently, contamination in eDNA samples was examined

using several model permutations, where contamination comprised both cichlid and human DNA,

cichlid DNA alone and human DNA alone. An information-theoretic approach using Akaike's

Information Criteria (AIC) to evaluate model fit was employed, where low AIC models are more

parsimonious than high AIC models (Akaike 1973). Significance of the fixed effect in the model

was tested by a Likelihood Ratio Test (LRT).

*Comparison of eDNA methods for crested newt detection*

We next tested the null hypothesis of no significant difference in sensitivity of qPCR and

metabarcoding. Overall agreement between eDNA metabarcoding and qPCR for crested newt

detection was measured using Cohen's kappa coefficient (Cohen 1960), following which

Pearson's Chi-squared Test for Independence was used to test equality of crested newt detection

between eDNA approaches.

The relationship between read count and qPCR score has not been examined previously,

and whether read production is indicative of DNA concentration remains unknown. qPCR score

is positively correlated with crested newt abundance and can be proxy for amount of DNA present

(Biggs *et al.* 2015). We hypothesised samples with higher qPCR score would have increased

crested newt read count. First, the average number of crested newt reads produced by eDNA

metabarcoding per qPCR score (1-12 out of 12) was calculated. A Spearman Rank Correlation was

then used to test for a relationship between average read count and qPCR score.

Following data exploration (see Supporting Information: Appendix 3), a negative binomial

GLMM was used to counter overdispersion and improve model fit. The GLMM examined read

count in relation to qPCR score, accounting for other variables that may affect metabarcoding

signal strength. Variation in crested newt read count was examined using the proportion of crested

newt reads within the total number of reads produced for each eDNA sample as the response

variable. Sequencing run and PCR plate were considered random effects and all other explanatory

variables as fixed effects (qPCR score, sample degradation, sample inhibition, post-PCR eDNA

concentration). Presence-absence of sample degradation and inhibition was determined by qPCR

in 2015 using methodology outlined by Biggs *et al.* (2015). Model fit was again evaluated using

AIC and significance of fixed effects in the model was tested with stepwise backward deletion of

terms from the model informed by LRTs. All values were bound in a new data frame and model

results plotted for evaluation using the R package 'ggplot2' v 2.1.0 (Wickham, 2009).

**Results**

TARGETED qPCR AND EGG SEARCHES

Targeted qPCR detected crested newt in 253 (49.80%) samples analysed by Fera (N = 508). Of

255 (50.20%) samples that were crested newt negative, one was inhibited and nine were degraded.

qPCR and egg searches produced consistent results for 297 (58.47%) ponds, with 51 (10.04%)

positive and 246 (48.43%) negative ponds by both methods. Of the 211 ponds where there was

disagreement between methods, 202 (39.76%) were qPCR positive but negative by egg searches,

and 7 (1.38%) were positive with egg searches but qPCR negative. Of 24 samples analysed by

ADAS, 12 (50.00%) were qPCR negative and 12 (50.00%) were qPCR positive for crested newt.

No egg search data were available for these ponds.

VERTEBRATE METABARCODING

The *in silico* and *in vitro* primer validation confirmed that crested newt, and other native UK

amphibians tested, can be reliably amplified and identified with the chosen assay (Supporting

Information: Appendix 4, Fig. S1). Furthermore, the *in silico* approach showed that the majority

of all UK vertebrates can be amplified (see Appendix 4 for details). Both sequencing runs had

comparable yield and sequencing quality score; summary statistics for each sequencing run and

read counts for taxonomic assignment levels are provided in Supporting Information: Appendix 4

(Tables S3, S4). A full summary of sequence read count data is also given in Supporting

Information: Appendix 5. eDNA metabarcoding identified a combined total of 60 species

(Supporting Information: Appendix 4, Fig. S2) across both sequencing libraries, with 375,954 and

508,879 sequences assigned to crested newt from each library. Analyses of overall pond species

compositions inferred by eDNA metabarcoding (Supporting Information: Appendix 4, Fig. S3,

Table S5) are reported separately (Harper *et al.* in preparation).

All samples (N=532) were sequenced and of 57 samples that did not produce visible PCR

bands, nine generated sequence reads. Notably, the 57 samples were not inhibited or degraded at

time of qPCR. Weak PCR bands were observed in some NTCs therefore all PCR controls were

sequenced (Supporting Information: Appendix 4, Fig. S4). Six NTCs contained crested newt DNA

but only one exceeded 100 crested newt reads (307/330 reads). Twelve other sources occurred in

NTCs (Supporting Information: Appendix 4, Table S6); seven occurred in more than one NTC and

eight had high maximum read counts (> 100 reads).

Contamination of NTCs (any DNA) and environmental samples (cichlid/human DNA) was

observed (Supporting Information: Appendix 4, Figs. S4, S5, S6). Read counts of NTC

contaminants were reduced between sequencing runs with the addition of mineral oil to PCR

reactions included on the second sequencing run (Supporting Information: Appendix 4, Figs. S4,

S5, S6) but this reduction was not statistically significant (GLMM: $\chi^2_1 = 2.083$ , $F_1 = 1.941$, $P >$

0.05). Mineral oil did not reduce human DNA signal in environmental samples between

sequencing runs either (GLMM: $\chi^2_1 = 3.608$ , $F_1 = 3.591$, $P > 0.05$); however, it did reduce human

DNA in combination with cichlid DNA (GLMM: $\chi^2_1 = 10.348$ , $F_1 = 21.143$, $P < 0.01$), and cichlid

DNA contamination alone (GLMM: $\chi^2_1 = 5.053$ , $F_1 = 6.978$, $P < 0.05$) of environmental samples.

eDNA METABARCODING VS qPCR FOR CRESTED NEWT DETECTION

Crested newt detection by metabarcoding NT (34.21%) was less sensitive than qPCR NT (49.81%)

but marginally higher than qPCR TA (32.71%) (N = 532 ponds, Fig. 1). Metabarcoding TA had

lower detection efficiency (28.01%) and failed to detect crested newt in 116 and 25 ponds where

the species was detected by qPCR NT and qPCR TA respectively. Nonetheless, both molecular

approaches attained higher crested newt detection than daytime egg searches (11.46%) in 506

ponds where all three approaches were implemented.

Overlap between survey methods for positive crested newt ponds (N = 277), and unique

detections by each method is summarised in Fig. 2. Negative crested newt ponds (N = 229) are

examined in combination with species positive ponds in Supporting Information: Appendix 4

(Table S7). Each survey method detected the species in ponds where other methods failed. Despite

lower crested newt detection efficiency, egg searches detected the species in six ponds where it

went undetected by qPCR and metabarcoding. Metabarcoding NT and metabarcoding TA revealed

crested newt in seven ponds which other methods did not, whilst qPCR NT and qPCR TA detected

crested newt in 33 ponds unique to other methods. All methods detected crested newt in 32 ponds,

and both metabarcoding and qPCR identified crested newt in 86 ponds. Disagreement between

molecular methods was more likely when samples were positive rather than negative by qPCR.

Without thresholds, 39.25% of qPCR positive ponds (N = 265) were negative by metabarcoding,

but 7.87% of qPCR negative ponds (N = 267) were positive by metabarcoding. With thresholds,

29.31% of qPCR positive ponds (N = 174) were negative by metabarcoding, whereas 7.26% of

qPCR negative ponds (N = 358) were positive by metabarcoding. Agreement between eDNA

approaches is summarised in Table 1. Agreement was strongest between eDNA approaches when

the qPCR detection threshold was applied, irrespective of whether the metabarcoding detection

threshold was applied. Metabarcoding (NT or TA) and qPCR TA did not significantly differ in

their detection of crested newt (Table 1).

An identical positive correlation was observed between qPCR score and the average

number of crested newt reads obtained for samples belonging to each qPCR score ($r_s = 0.648$, df

= 11, $P < 0.05$), regardless of threshold application to the metabarcoding data. Despite some

inconsistency across qPCR scores, samples with a higher qPCR score generally had more crested

newt reads, supportive of a relationship between metabarcoding and abundance of eDNA from

single species. Notably, metabarcoding produced crested newt reads for qPCR NT and qPCR TA

crested newt negative samples, but the crested newt metabarcoding signal of these (qPCR NT$_{negative}$

= 2639 reads max., qPCR TA$_{negative}$ = 3075 reads max.) was much lower than samples with higher

qPCR score (max. 65,325 reads; Supporting Information: Appendix 5). Further examination of the

relationship between qPCR score and metabarcoding TA revealed qPCR score and post-PCR

eDNA concentration of samples also influenced the proportion of crested newt reads i.e. relative

crested newt sequence read production (Table 2). A significant positive relationship was observed

between qPCR score and the proportion of crested newt reads within total reads per sample ($P <$

$0.001$) (Fig. 3a). Conversely, post-PCR eDNA concentration had a significant negative influence

on the proportion of crested newt reads ($P < 0.001$), where read proportion decreased as post-PCR

eDNA concentration increased (Fig. 3b).

COMPARISON OF METHOD COST AND INVESTIGATOR EFFORT

A full breakdown of expenditure per pond is given in Supporting Information (Table S8) and

summarised in Fig. 4. Cost of materials, and investigator effort and salary (assuming hourly rate

of £21.20) were calculated for eDNA samples but estimates do not include procedural controls,

qPCR standards, or consumables and reagents required for assay optimisation. Time required to

perform PCR for metabarcoding and qPCR was estimated assuming available machinery to run

four PCR plates in parallel and one qPCR plate.

**Discussion**

We have demonstrated eDNA metabarcoding is a highly sensitive tool for monitoring crested newt

alongside the wider biological community, corroborating other comparisons of eDNA

metabarcoding and qPCR for single-species monitoring (Lacoursière-Roussel *et al.* 2016;

Schneider *et al.* 2016). Despite marginal reduction in single-species detection, eDNA

metabarcoding revealed a wealth of biodiversity information and could enable more effective

freshwater monitoring networks and better understanding of community structure and ecosystem

function alongside crested newt monitoring (Biggs, von Fumetti & Kelly-Quinn 2016). However,

both eDNA approaches have advantages and drawbacks which must be considered for design and

implementation of biodiversity monitoring programmes.

SINGLE-SPECIES DETECTION BY qPCR AND METABARCODING

A direct comparison of sensitivity between qPCR and metabarcoding is not straightforward:

stochasticity in qPCR largely occurs during amplification (volume of template DNA and technical

replication), whereas stochastic variation during metabarcoding arises through PCR amplification

and sequencing (depth and replication) (Kelly *et al.* 2014; Thomsen *et al.* 2016; Deiner *et al.* 2017).

In our study, 12 independent qPCR replicates were performed for each sample but due to

limited resources, metabarcoding was based on three pooled PCR replicates which were sequenced

once only. Therefore, to enable a fair comparison between methods in terms of PCR effort, a

threshold of $\geq 4/12$ positive replicates (qPCR TA) was applied to the qPCR data. Indeed,

detection sensitivity was most similar between methods with the qPCR threshold and without the

metabarcoding threshold. Both eDNA metabarcoding and qPCR displayed reduced crested newt

detection when thresholds were applied; although, this may reflect reduced false positive

detections rather than decreased sensitivity. Lower sensitivity of the eDNA metabarcoding

approach used here may also stem from sample degradation during long-term storage. The samples

used were stored for more than 12 months at -80 ℃ before metabarcoding. However, long-term

storage and continual freeze-thawing of samples may allow aggregation of inhibitory substances

which impair PCR amplification and cause false negatives (Takahara, Minamoto & Doi 2015).

Despite lower sensitivity, strength of eDNA signal produced by metabarcoding was

correlated with that of qPCR, where both crested newt average read count and read proportion

broadly increased with qPCR score of eDNA samples. The correlation was inconsistent though,

where high average or proportional crested newt read count did not always correspond to high

qPCR score. Biggs *et al.* (2015) also found a variable positive association between qPCR and

visual counts of crested newt, where high qPCR score did not always correlate with high counts.

Our results indicate sequence reads may allow relative abundance to be estimated but this must be

statistically verified against established population census methods.

The comparison between qPCR and metabarcoding must also be examined in context of

the sequencing effort. Here, we sequenced a large number of samples (380 including positive and

negative controls) per run to provide a realistic cost scenario for routine monitoring. Yet,

metabarcoding detection sensitivity would likely improve with an increase in read depth per

sample (Kelly *et al.* 2014). In order to directly compare eDNA signal production by these

approaches, it may be necessary to perform sequencing replicates to verify true positives where

rare species are expected and generate an "eDNA metabarcoding score" system similar to qPCR

(Brandon-Mong *et al.* 2015; Port *et al.* 2016; Civade *et al.* 2016; Thomsen *et al.* 2016). PCR and

sequencing replication in metabarcoding may enhance species detection probability through

improved amplification of low abundance or highly degraded DNA (Ficetola *et al.* 2015; Port *et*

*al.* 2016) that is readily amplified by qPCR (Lacoursière-Roussel *et al.* 2016). Similarly,

sequencing of independent biological replicates, opposed to pseudoreplicates from a single water

sample (established crested newt eDNA sampling strategy), may improve detection and minimise

false negatives produced by eDNA metabarcoding (Andruszkiewicz *et al.* 2017; Bálint *et al.*

2017).

Metabarcoding assays are also susceptible to problems from taxon bias, DNA swamping

and bioinformatics related problems (Taberlet *et al.* 2012; Kelly *et al.* 2014; Shaw *et al.* 2016b).

Potential reduction in sensitivity of passive community sequencing versus targeted qPCR

approaches may relate to the performance of metabarcoding primers for target species. During

metabarcoding, DNA from rare species may be masked by highly abundant species (Schneider *et*

*al.* 2016), or under-represented due to disproportionate eDNA shedding rates across species and

preferential amplification of other species (Kelly *et al.* 2014). We found crested newt read

proportion was negatively associated with post-PCR concentration of eDNA samples. As a

positive relationship was not observed, this would suggest PCR amplification with our selected

marker and primers was not biased toward our focal species. However, we cannot conclude that

our metabarcoding assay was free of primer bias as post-PCR concentration of eDNA samples can

be influenced by PCR stochasticity.

FALSE NEGATIVES

The aim of this study was not to evaluate sensitivity of molecular methods against standard crested

newt survey methodologies. Egg searches were used to detect false negatives produced by qPCR

and metabarcoding and in doing so, revealed some interesting results. Biggs *et al.* (2015)

previously demonstrated that qPCR had higher detection rate than egg searches (as well as

torchlight, netting, and bottle trapping), but here we show this also holds true for metabarcoding.

Importantly, absence of eggs does not infer absence of adults, and this method is highly dependent

on weather conditions and water clarity (Rees *et al.* 2014a; Biggs *et al.* 2015). Despite considerably

higher detection rate of both eDNA approaches, eggs were recorded in a small number of ponds

that were eDNA negative. eDNA analysis can incorrectly infer absence or low abundance of

species if inhibition or interference from non-target DNA has occurred (Goldberg *et al.* 2016).

Alternatively, eDNA false negatives may have been a by-product of sampling strategy and effort

for crested newt. Larger water volumes and/or more biological replication instead of

pseudoreplication (6 x 15 mL sampled from 600 mL for crested newt) may improve detection

(Lopes *et al.* 2016; Andruszkiewicz *et al.* 2017; Bálint *et al.* 2017). All methods revealed crested

newt in ponds where other approaches failed, emphasising that these species monitoring tools are

complementary and should be used in combination to achieve maximum detection probability.

However, integrative strategies combining molecular and conventional tools are often not cost-

efficient for most applications.

FALSE POSITIVES

False positives may arise from field contamination and eDNA transport in the environment -

particularly by waterfowl (Shaw *et al.* 2016a). eDNA is retained by predators, discarded in faeces,

and transported by anthropogenic activity, combined with natural water currents and flow

(Hänfling *et al.* 2016). In the laboratory, PCR-accumulated and sequencing error, including primer

mismatch (Andersen *et al.* 2012) and 'tag jumps' (Schnell, Bohmann & Gilbert 2015), can induce

misassignment leading to false positives, cross-contamination between samples, or laboratory

contamination (Andruszkiewicz *et al.* 2017).

False positives can be identified using Site  Occupancy Modelling of metabarcoding data

(Ficetola *et al.* 2015) or risk of false positives minimised using a sequencing threshold i.e. the

number of sequence reads required for a sample to be species positive (Hänfling *et al.* 2016; Civade

*et al.* 2016; Evans *et al.* 2017a). However, such thresholds can reduce detection of rare species, a

primary goal of this study, and may fail where false and true positives occur at similar frequency

(Hänfling *et al.* 2016). Instead, we calculated species-specific sequence thresholds to more

accurately control for false positives in our dataset without compromising crested newt detection.

In our study, human DNA occurred at high frequency and abundance; this may have been

a true environmental signal from pond water, or real contaminant as encountered in other

metabarcoding research (Port *et al.* 2016; Valentini *et al.* 2016; Thomsen *et al.* 2016). Blocking

primers can prevent amplification of abundant non-target DNA like human (Valentini *et al.* 2016)

but may fail (Thomsen *et al.* 2016) or prevent amplification of target taxa (Port *et al.* 2016).

Alongside human, other aquatic and terrestrial vertebrate DNA occurred at high frequency in

NTCs (Supporting Information: Appendix 4), although these were not removed by addition of

mineral oil.

Our results also highlight the importance and impact of qPCR thresholds when inferring

species presence-absence. Similar to Smart *et al.* (2015), we found a stringent qPCR threshold

reduced detection sensitivity. As yet, there is no guidance to indicate how many samples or

replicates must be positive to class a site as species-positive (Goldberg *et al.* 2016; Smart *et al.*

2016) but this clearly needs to be addressed to improve standardisation and reproducibility of

eDNA research. Importantly, less stringent thresholds (and false positives inherent to these) are

somewhat precautionary and may enable better protection of crested newt by preventing

development. Therefore, whilst reduction or removal of false positives is desirable, detection

thresholds must not compromise protection of threatened species either. Until a suitable threshold

can be established, it may be more appropriate to re-analyse samples which yield one positive

qPCR replicate to prevent false positives (Rees *et al.* 2014b; Goldberg *et al.* 2016).

COST AND INVESTIGATOR EFFORT

Comparisons of qPCR and conventional survey for species monitoring have included cost

projections (Biggs *et al.* 2014; Davy *et al.* 2015; Smart *et al.* 2016; Evans *et al.* 2017b) but cost

has not been thoroughly assessed in qPCR and eDNA metabarcoding comparisons (Lacoursière-

Roussel *et al.* 2016; Schneider *et al.* 2016). We found eDNA metabarcoding was more costly than

qPCR but both approaches required similar investigator effort. qPCR scales to the number of

samples being processed (Schneider *et al.* 2016) whereas metabarcoding has fixed costs including

reagent kit for HTS platform (Bálint *et al.* 2017). eDNA metabarcoding becomes more cost-

efficient as more samples are processed (Bálint *et al.* 2017) but fewer replicates would reduce

qPCR cost (Davy *et al.* 2015; Smart *et al.* 2016). Travel was excluded from our cost estimate but

inclusion of this expense would further reduce cost efficiency of both approaches. Cost of eDNA

monitoring is influenced by sample size, methods, replication, laboratory, statistical power and

occupancy modelling (Davy *et al.* 2015; Evans *et al.* 2017b). Consequently, cost is proportional

to project requirements (Davy *et al.* 2015) and will vary depending on choice of qPCR or

metabarcoding workflow. Nevertheless, cost efficiency combined with the overarching aim of a

monitoring or conservation programme should always be considered.

**Conclusion**

eDNA metabarcoding holds promise for holistic biodiversity monitoring of freshwater ponds

opposed to targeted qPCR for flagship or indicator species such as crested newt. Metabarcoding

can reveal entire species assemblages from environmental samples without prior ecosystem

information, and provide broad-scale distribution data for multiple species simultaneously. This

diagnostic power over conventional monitoring has been repeatedly demonstrated for freshwater

vertebrates (Valentini *et al.* 2016; Hänfling *et al.* 2016; Shaw *et al.* 2016a; Civade *et al.* 2016;

Lacoursière-Roussel *et al.* 2016). Nonetheless, the method at present appears to be less sensitive

than qPCR for single-species monitoring, and species detection by molecular and conventional

methods was incongruent.

Comprehensive study of the influence of water volume, eDNA capture and extraction

method, and sample storage on single-species and community detection in lentic and lotic systems

is required. Currently, 90 mL water is sampled during crested newt eDNA survey, followed by

ethanol precipitation and Qiagen® DNeasy Blood and Tissue kit extraction (Biggs *et al.* 2015).

While this may be appropriate for highly-sensitive targeted qPCR, larger water volumes and

filtration may be required to capture eDNA from less abundant vertebrates and characterise

community diversity (Shaw *et al.* 2016b). Additionally, eDNA from different species can be

unevenly distributed throughout water bodies and may be concentrated in particular areas

(Hänfling *et al.* 2016; Evans *et al.* 2017a), thus sampling strategies must be carefully designed to

ensure eDNA samples are representative of biodiversity present. Indeed, how individuals within a

species are distributed in the habitat sampled can also influence detection (Biggs *et al.* 2015).

Minimising the risk of false positives and contamination remains a pressing issue in

metabarcoding, and standard contamination measures (Goldberg *et al.* 2016) may be insufficient

for analysis of vertebrate assemblages. An even stricter forensic laboratory set-up, such as that

employed for ancient DNA (aDNA) should be adopted to ensure robustness of the data. Positive

and negative controls should be included at each stage of metabarcoding workflows to monitor

contamination (Deiner *et al.* 2017). However, preventive measures inevitably increase research

cost and some degree of contamination is unavoidable in metabarcoding (Kelly *et al.* 2014;

Brandon-Mong *et al.* 2015; Port *et al.* 2016; Thomsen *et al.* 2016).

Currently, cost and investigator effort required for metabarcoding and qPCR are broadly

equivalent. Yet, where qPCR is an established technology and has reached its price ceiling, HTS

is relatively new technology and prices will continue to drop. This reduction in cost means higher

sample throughput and more technical replication will be possible. We therefore argue that

metabarcoding will become more cost-efficient in the long-term, providing more data at lower cost

and comparable sensitivity to qPCR. However, where samples cannot be processed in large

batches, qPCR may retain cost-efficiency.

We conclude that eDNA metabarcoding is not yet a replacement for targeted qPCR and

conventional survey but rather another tool in the ecologist toolbox. Ultimately, choice of

monitoring tool(s) is specific to aims of each conservation project. At present, qPCR retains

sensitivity for crested newt populations of all sizes, regardless of sample number processed. Under

a realistic conservation monitoring scenario, where funding is limited and samples must be

processed in large batches, metabarcoding may suffer from false negatives due to reduced

sequencing depth and replication. However, in many cases, the biodiversity information generated

by this approach, and its implications for community ecology and conservation, will eclipse lower

sensitivity. This passive screening approach would be most effective for initial survey of water

bodies to generate broad-scale multi-species distribution data. This holistic data can then inform

best use of funding and time for targeted species-specific survey.

**Acknowledgements**

This work was funded by University of Hull. We would like to thank Jennifer Hodgetts (Fera) for

assisting with sample collection and data dissemination, and Jianlong Li for primer design and

advice on laboratory protocols. Furthermore, Barbara Mabel, Elizabeth Kilbride (University of

Glasgow), Andrew Buxton and Richard Griffiths (DICE/University of Kent) provided tissue

samples for primer validation and Sanger sequencing to supplement the reference database.

**Author contributions**

B.H., L.R.H., L.L.H and N.B. conceived and designed the study. H.C.R., K.C.G., and N.B.

contributed samples for processing. H.C.R. performed extractions under licence from Natural

England and qPCR for eDNA samples from private contracts at The University of Nottingham.

N.B. performed qPCR for eDNA samples from Natural England's Great Crested Newt Evidence

Enhancement Program. L.R.H. performed remaining laboratory work and analysed the data. I.P.A.

and E.L. offered advice on and supervised sequencing. C.H. assisted with bioinformatics analysis.

P.B. and S.P. contributed datasets for analysis. L.R.H. wrote the manuscript, which all authors

revised.

**Data accessibility**

Raw sequence reads have been archived on the NCBI Sequence Read Archive. R scripts and

corresponding data are deposited in the dedicated GitHub repository: https://github.com/HullUni-

bioinformatics/Harper_et_al_2017.

**References**

Akaike, H. (1973) Maximum likelihood identification of Gaussian autoregressive moving
average models. *Biometrika*, **60**, 255-265.
Andersen, K., Bird, K.L., Rasmussen, M., Haile, J., Breuning-Madsen, H., Kjær, K.H., Orlando,
L., Gilbert, M.T.P. & Willerslev, E. (2012) Meta-barcoding of "dirt" DNA from soil reflects
vertebrate biodiversity. *Molecular Ecology*, **21**, 1966–1979.
Andruszkiewicz, E.A., Starks, H.A., Chavez, F.P., Sassoubre, L.M., Block, B.A. & Boehm, A.B.
(2017) Biomonitoring of marine vertebrates in Monterey Bay using eDNA metabarcoding.
*PLoS One*, **12**, e0176343.
Bálint, M., Nowak, C., Márton, O., Pauls, S., Wittwer, C., Aramayo, J.L., Schulze, A.,
Chambert, T., Cocchiararo, B. & Jansen, M. (2017) Twenty-five species of frogs in a liter of
water: eDNA survey for exploring tropical frog diversity. *bioRxiv*, 176065.
Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) lme4: Linear mixed-effects models
using Eigen and S4, 2014. *R package version*.
Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P. & Kauserud, H. (2010) ITS
as an environmental DNA barcode for fungi: an in silico approach reveals potential PCR
biases. *BMC Microbiology*, **10**, 189.
Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R.A., Foster, J.,
Wilkinson, J.W., Arnell, A., Brotherton, P., Williams, P. & Dunn, F. (2015) Using eDNA to
develop a national citizen science-based monitoring programme for the great crested newt
(*Triturus cristatus*). *Biological Conservation*, **183**, 19–28.
Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C. & Griffiths, R.A. (2014) Analytical and
methodological development for improved surveillance of the Great Crested Newt. *Defra
Project WC1067*.
Biggs, J., von Fumetti, S. & Kelly-Quinn, M. (2016) The importance of small waterbodies for
biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia*, 1–37.
Blackman, R.C., Constable, D., Hahn, C., Sheard, A.M., Durkota, J., Hänfling, B. & Handley,
L.L. (2017) Detection of a new non-native freshwater species by DNA metabarcoding of
environmental samples—first record of *Gammarus fossarum* in the UK. *Aquatic Invasions,*
**2**, 177-189.
Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*, **30**, 2114–2120.
Brandon-Mong, G.-J., Gan, H.-M., Sing, K.-W., Lee, P.-S., Lim, P.-E. & Wilson, J.-J. (2015)
DNA metabarcoding of insects and allies: an evaluation of primers and pipelines. *Bulletin of
Entomological Research*, **105**, 717–727.
Buxton, A.S., Groombridge, J.J., Zakaria, N.B. & Griffiths, R.A. (2017) Seasonal variation in
environmental DNA in relation to population size and environmental factors. *Scientific
Reports*, **7**, 46294.

Civade, R., Dejean, T., Valentini, A., Roset, N., Raymond, J.-C., Bonin, A., Taberlet, P. & Pont,
D. (2016) Spatial Representativeness of Environmental DNA Metabarcoding Signal for Fish
Biodiversity Assessment in a Natural Freshwater System. *PLoS One*, **11**, e0157366.
Cohen, J. (1960) A Coefficient of Agreement for Nominal Scales. *Educational and*
*Psychological Measurement*, **20**, 37–46.
Davy, C.M., Kidd, A.G. & Wilson, C.C. (2015) Development and Validation of Environmental
DNA (eDNA) Markers for Detection of Freshwater Turtles. *PLoS One*, **10**, e0130965.
Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer,
S., Bista, I., Lodge, D.M., de Vere, N., Pfrender, M.E. & Bernatchez, L. (2017)
Environmental DNA metabarcoding: transforming how we survey animal and plant
communities. *Molecular Ecology*, 1-24.
Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C. & Knight, R. (2011) UCHIME improves
sensitivity and speed of chimera detection. *Bioinformatics* , **27**, 2194–2200.
Evans, N.T., Li, Y., Renshaw, M.A., Olds, B.P., Deiner, K., Turner, C.R., Jerde, C.L., Lodge,
D.M., Lamberti, G.A. & Pfrender, M.E. (2017a) Fish community assessment with eDNA
metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of*
*Fisheries and Aquatic Sciences,* **74**, 1362-1374.
Evans, N.T., Shirey, P.D., Wieringa, J.G., Mahon, A.R. & Lamberti, G.A. (2017b) Comparative
Cost and Effort of Fish Distribution Detection via Environmental DNA Analysis and
Electrofishing. *Fisheries*, **42**, 90–99.
Ficetola, G.F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P. &
Pompanon, F. (2010) An *In silico* approach for the evaluation of DNA barcodes. *BMC*
*Genomics*, **11**, 434.
Ficetola, G.F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., Gielly, L.,
Lopes, C.M., Boyer, F., Pompanon, F., Rayé, G. & Taberlet, P. (2015) Replication levels,
false presences and the estimation of the presence/absence from eDNA metabarcoding data.
*Molecular Ecology Resources*, **15**, 543–556.
Goldberg, C.S., Turner, C.R., Deiner, K., Klymus, K.E., Thomsen, P.F., Murphy, M.A., Spear,
S.F., McKee, A., Oyler-McCance, S.J., Cornman, R.S., Laramie, M.B., Mahon, A.R., Lance,
R.F., Pilliod, D.S., Strickler, K.M., Waits, L.P., Fremier, A.K., Takahara, T., Herder, J.E. &
Taberlet, P. (2016) Critical considerations for the application of environmental DNA
methods to detect aquatic species. *Methods in Ecology and Evolution*, **7**, 1299–1307.
Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P., Blackman, R.C.,
Oliver, A. & Winfield, I.J. (2016) Environmental DNA metabarcoding of lake fish
communities reflects long-term data from established survey methods. *Molecular Ecology,*
**25**, 3101-3119.
Harrison, X.A. (2014) Using observation-level random effects to model overdispersion in count
data in ecology and evolution. *PeerJ*, **2**, e616.
Hosmer, D.W. & Lemeshow, S. (2000) Multiple Logistic Regression. *Applied Logistic*
*Regression*, pp. 31–46. John Wiley & Sons, Inc.

Kelly, R.P., Port, J.A., Yamahara, K.M. & Crowder, L.B. (2014) Using environmental DNA to
census marine fishes in a large mesocosm. *PLoS One*, **9**, e86175.
Lacoursière-Roussel, A., Dubois, Y., Normandeau, E. & Bernatchez, L. (2016) Improving
herpetological surveys in eastern North America using the environmental DNA method.
*Genome*, **59**, 991–1007.
Lawson Handley, L. (2015) How will the "molecular revolution" contribute to biological
recording? *Biological Journal of the Linnean Society*, **115**, 750–766.
Lopes, C.M., Sasso, T., Valentini, A., Dejean, T., Martins, M., Zamudio, K.R. & Haddad, C.F.B.
(2016) eDNA metabarcoding: a promising method for anuran surveys in highly diverse
tropical forests. *Molecular Ecology Resources,* **17**, 904-914.
Magoč, T. & Salzberg, S.L. (2011) FLASH: fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics* , **27**, 2957–2963.
Mazerolle, M.J. & Mazerolle, M. (2016) Package "AICcmodavg."
Natural England (2015) Great crested newts: surveys and mitigation for development projects.
Available at: https://www.gov.uk/guidance/great-crested-newts-surveys-and-mitigation-for-
development-projects [accessed: 7[th] November 2017].
Port, J.A., O'Donnell, J.L., Romero-Maraccini, O.C., Leary, P.R., Litvin, S.Y., Nickols, K.J.,
Yamahara, K.M. & Kelly, R.P. (2016) Assessing vertebrate biodiversity in a kelp forest
ecosystem using environmental DNA. *Molecular Ecology*, **25**, 527–541.
Rees, H.C., Bishop, K., Middleditch, D.J., Patmore, J.R.M., Maddison, B.C. & Gough, K.C.
(2014a) The application of eDNA for monitoring of the Great Crested Newt in the UK.
*Ecology and Evolution*, **4**, 4023–4032.
Rees, H.C., Maddison, B.C., Middleditch, D.J., Patmore, J.R.M. & and Gough, K.C. (2014b)
REVIEW: The detection of aquatic animal species using environmental DNA–a review of
eDNA as a survey tool in ecology. *Journal of Applied Ecology,* **51**, 1450-1459.
Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P. & Coissac, E. (2011) ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic*
*Acids Research*, **39**, e145.
Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. (2016) VSEARCH: a versatile open
source tool for metagenomics. *PeerJ*, **4**, e2584.
Schneider, J., Valentini, A., Dejean, T., Montarsi, F., Taberlet, P., Glaizot, O. & Fumagalli, L.
(2016) Detection of Invasive Mosquito Vectors Using Environmental DNA (eDNA) from
Water Samples. *PLoS One*, **11**, e0162493.
Schnell, I.B., Bohmann, K. & Gilbert, M.T.P. (2015) Tag jumps illuminated--reducing sequence-
to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources*, **15**,
1289–1303.
Shaw, J.L.A., Clarke, L.J., Wedderburn, S.D., Barnes, T.C., Weyrich, L.S. & Cooper, A. (2016a)
Comparison of environmental DNA metabarcoding and conventional fish survey methods in
a river system. *Biological Conservation*, **197**, 131–138.

Shaw, J.L.A., Weyrich, L. & Cooper, A. (2016b) Using environmental (e)DNA sequencing for
aquatic biodiversity surveys: a beginner's guide. *Marine and Freshwater Research*, 1-14.
Smart, A.S., Tingley, R., Weeks, A.R., van Rooyen, A.R. & McCarthy, M.A. (2015)
Environmental DNA sampling is more sensitive than a traditional survey technique for
detecting an aquatic invader. *Ecological Applications*, **25**, 1944–1952.
Smart, A.S., Weeks, A.R., van Rooyen, A.R., Moore, A., McCarthy, M.A. & Tingley, R. (2016)
Assessing the cost-efficiency of environmental DNA sampling. *Methods in Ecology and*
*Evolution*, **7**, 1291–1298.
Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C. & Willerslev, E. (2012) Towards next-
generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, **21**,
2045–2050.
Takahara, T., Minamoto, T. & Doi, H. (2015) Effects of sample processing on the detection rate
of environmental DNA from the Common Carp (*Cyprinus carpio*). *Biological Conservation*,
**183**, 64–69.
Thomsen, P.F., Iversen, L.L., Wiuf, C., Rasmussen, M., Gilbert, M.T.P., Orlando, L. &
Willerslev, E. (2012) Monitoring endangered freshwater biodiversity using environmental
DNA. *Molecular Ecology*, **21**, 2565–2573.
Thomsen, P.F., Møller, P.R., Sigsgaard, E.E., Knudsen, S.W., Jørgensen, O.A. & Willerslev, E.
(2016) Environmental DNA from Seawater Samples Correlate with Trawl Catches of
Subarctic, Deepwater Fishes. *PLoS One*, **11**, e0165252.
Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P.F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp,
G.H., Geniez, P., Pont, D., Argillier, C., Baudoin, J.-M., Peroux, T., Crivelli, A.J., Olivier,
A., Acqueberge, M., Le Brun, M., Møller, P.R., Willerslev, E. & Dejean, T. (2016) Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding.
*Molecular Ecology*, **25**, 929–942.
Ver Hoef, J.M. & Boveng, P.L. (2007) QUASI-POISSON VS. NEGATIVE BINOMIAL
REGRESSION: HOW SHOULD WE MODEL OVERDISPERSED COUNT DATA?
*Ecology,* **88**, 2766-2772.
Zhang, Z., Schwartz, S., Wagner, L. & Miller, W. (2000) A greedy algorithm for aligning DNA
sequences. *Journal of Computational Biology*, **7**, 203–214.
800

801  **Table 1.** Summary of analyses testing for agreement between eDNA approaches, with and without thresholds, for

802  crested newt detection. Cohen's kappa coefficient (k) represents strength of agreement between methods (1 = 100%).

803  Pearson's Chi-squared Test for Independence tested whether methods significantly differed for crested newt detection.

| Comparison | Probability of observed agreement | Probability of expected agreement | k | Overall agreement | $\chi^2$ | d.f. | $P$ |
|---|---|---|---|---|---|---|---|
| Metabarcoding NT qPCR NT | 0.77 | 0.50 | 0.53 | Moderate | 25.94 | 1 | < 0.001 |
| Metabarcoding TA qPCR NT | 0.74 | 0.50 | 0.48 | Moderate | 52.291 | 1 | < 0.001 |
| Metabarcoding NT qPCR TA | 0.84 | 0.56 | 0.63 | Good | 0.207 | 1 | > 0.05 |
| Metabarcoding TA qPCR TA | 0.86 | 0.58 | 0.66 | Good | 2.561 | 1 | > 0.05 |

806    **Table 2.** Summary of analyses testing for variation in proportion of crested newt sequence reads in a sample produced

807    by eDNA metabarcoding, attributable to qPCR score or post-PCR eDNA concentration. Test statistic is for LRT used.

| Model variables | *N* (ponds) | d.f. | AIC | Effect size | Standard error | $\chi^2$ | *F* | *P* |
|---|---|---|---|---|---|---|---|---|
| **qPCR score** | 532 | 1 | 1578.3 | 0.373 | 0.032 | 150.682 | 147.117 | *< 0.001* |
| **post-PCR eDNA concentration** | 532 | 1 | 1441.9 | -0.056 | 0.015 | 14.272 | 12.457 | *< 0.001* |

**Figure 1.** Comparison of survey methodology for crested newt detection in freshwater ponds across the UK. Bars

represent proportion of positive and negative crested newt ponds by each method with frequency displayed on bars.

**Figure 2.** Venn diagram which summarises the number of positive crested newt detections (N = 277) by each method

(egg search, qPCR NT, qPCR TA, metabarcoding NT, and metabarcoding TA), and overlap in crested newt detection

between methods, for 506 ponds where all methods were applied. Negative crested newt detections (N = 229) are

highlighted in red.

**Figure 3.** Relationship between fixed effects (qPCR score, post-PCR eDNA concentration) and response variable

(proportion of crested newt reads) in eDNA samples, as predicted by the negative binomial GLMM. The 95% CIs, as

calculated using the predicted proportions, and standard error for these predictions are given for each relationship.

The observed data (points) are also displayed against the predicted relationships (boxes, line). The proportion of

crested newt reads within eDNA samples increased as qPCR score increased **(a)**. Conversely, the proportion of crested

newt reads decreased as post-PCR eDNA concentration increased **(b)**.

**Figure 4.** Cost and investigator effort required for targeted qPCR of crested newt and eDNA metabarcoding of

830 vertebrate communities from pond water samples.