Knockdown of Liver-Derived C1 Inhibitor Results in Brain Pathology in mice

Running title: C1INH, the gatekeeper of the CNS

Dorit Farfara[1], Emily Feierman[1], Allison Richards[1], Alexey S. Revenko[2], Robert A. MacLeod[2], Erin H. Norris[1] and Sidney Strickland[1]

[1]Patricia and John Rosenwald Laboratory of Neurobiology and
Genetics, The Rockefeller University, New York, NY 10065, USA

[2]Department of Antisense Drug Discovery, IONIS Pharmaceuticals Inc, Carlsbad, CA, USA

Keywords: C1inhibitor, Kallikrein-kinin system, bradykinin, vascular permeability, neuro-inflammation, cognition

Abstract

Activation of the innate immune system, mediated by the complement and the contact systems, induces inflammation. The inhibition of these pathways by C1 inhibitor (C1INH) has been shown to decrease pro-inflammatory response and mediate vascular permeability in the periphery. However, the role of peripheral C1INH has not been associated with brain function. Using an antisense oligonucleotide (ASO) to deplete circulating liver-derived C1INH in wildtype (WT) mice, we induced the activation of the kallikrein-kinin system (KKS), which produced bradykinin in the plasma, resulting in hypotension. Interestingly, the complement system was quiescent. Depletion of liver-derived C1INH increased blood-brain barrier (BBB) permeability, increased expression of bradykinin 2 receptor, activated resident glial cells to secrete pro-inflammatory mediators such as Il-1beta, Il-6, TNF-alpha and iNOS, and induced cognitive decline. The results of this study emphasize the important role of circulating C1INH in mediating brain function through the activation of the KKS. Thus, manipulation of C1INH in neuro-vascular disorders might be therapeutically beneficial.

Significant statement

By using Antisenseoligonucleotide (ASO) techniques we knocked-down liver-derived C1INH for 12 weeks. This elimination led to the activation of the kallikrein-kinin system in the plasma to produce elevated levels of bradykinin leading to increases bradykinin 2 receptor expressions in the brain, blood-brain barrier permeability, activation of astrocytes and microglial cells towards a pro-inflammatory response, and cognitive decline. These results suggest that liver-derived C1INH must be monitored and properly mediated to maintain normal brain functions.

Introduction

The involvement of the peripheral immune system in brain pathology is now supported by mounting evidence. C1 inhibitor (C1INH) belongs to the super-family of serine protease inhibitors (serpins). C1INH inhibits the initiating proteases of the complement system (Ratnoff and Lepow, 1957, Ziccardi and Cooper, 1979) and the kallikrein system (Schapira et al., 1982; van der Graaf et al., 1983). The complement system and KKS share important features (Bossi et al., 2011), and both involve in proteolytic cascades resulting in a pro-inflammatory response (Ghebrehiwet et al., 2016).

Activation of the KKS is induced by plasma kallikrein which cleaves high molecular weight kininogen (HK) to liberate bradykinin, a 9-amino acid peptide, known to cause vasodilation, reduce blood pressure, and induce a pro-inflammatory response through activation of bradykinin receptors 1 and 2 (B1R and B2R) (Schmaier and McCrae, 2007).

The distribution of C1IHN is very similar in mouse and human tissues (Lener et al., 1998; Vinci et al., 2002). Hepatocytes are thought to be the main source for C1INH biosynthesis (Colten, 1972). In the CNS, C1INH is highly expressed in the cortex and the external capsule of the forebrain (Vinci et al., 2002). The synthesis and/or secretion of C1INH have been detected in astrocyte-derived tumor cell lines (Gasque et al., 1993), brain tissue, neuroblastoma cell lines, microglia, monocytes (Gasque et al., 1995; Gasque et al., 1996), and neurons (Walker et al., 1995).

In this study, we show the important role of balanced liver-derived C1INH in maintaining brain integrity and function. By using ASO-mediated knockdown, we eliminated liver-derived C1INH in mouse plasma, which resulted in kallikrein-kinin system (KKS) activation. KKS activation led to the production of bradykinin, which in turn led to hypotension and increased the expression of bradykinin 2 receptor (B2R) in the brain. Impaired blood-brain barrier (BBB) integrity, activated astrocytes and microglia cells to secrete pro-inflammatory mediators, and impaired cognition were the pathological result of this knockdown. These results highlights the important role of normal levels of liver-derived C1INH in maintaining proper brain function.

Results

Subcutaneous administration of C1INH ASO eliminates C1INH in the plasma and activates the KKS. In order to prevent C1INH production in the liver, we subcutaneously administered C1INH ASO or control ASO (CTRL ASO) to ten-week-old WT mice for three months. Plasma levels of C1INH were significantly downregulated in mice treated with C1INH ASO compared to CTRL ASO (Fig 1A and B and Supplementary figure 1A for C57/C3H). Similar results were obtained in non-reduced conditions (Supp Fig.1B-C). In addition, C1INH ASO treatment, targeted against liver C1INH did not eliminate the protein expression in splenocytes and macrophages measures by FACS (Supp. Fig 1D-F). These findings suggest that the reduction of C1INH in the circulation is due to the targeted treatment of C1INH in the liver which results in decreased expression of the secreted form of C1INH and that the ASO does not target other peripheral cells.

Since C1INH can also inhibit the KKS through plasma kallikrein, we analyzed the cleavage of high molecular weight kininogen (HK) after C1INH ASO administration. Using Western blot analysis, we found a 50% reduction in levels of intact HK (iHK; mHK+ΔD5) in plasma from C1INH ASO-treated mice compared to CTRL ASO-treated mice (Fig 1C and D and Supplementary figure 2A), indicative of increased HK cleavage as previously suggested (Zamolodchikov et al., 2015). Non-reduced HK levels were also significantly reduced in C1INH ASO mice (Supplementary figure 2B, C). Furthermore, levels of plasma kallikrein, which activates HK cleavage, were significantly increased in plasma from C1INH ASO-treated mice compared to controls (Fig 1E, F). However, when we evaluated the plasma after eight weeks of treatment, we found no significant differences in iHK and plasma kallikrein levels in C1INH targeted by ASO (Supplementary figure 2D). Therefore, we continued our evaluations using the chronic treatment time point (12 weeks of treatment).

We measured kallikrein activity in plasma of C1INH ASO-treated mice compared to CTRL ASO, as previously described (Zamolodchikov et al., 2015). Kallikrein activity levels were significantly increased by 48% (p=0.0244) compared to plasma from control treated mice (Fig 1G and H),

supporting the results of KKS activation seen in western blots.

The end product of HK cleavage is a 9-amino acid peptide called bradykinin. Bradykinin is unstable in the plasma and rapidly eliminated by degradation. Using a sensitive ELISA kit for detecting bradykinin, we found that levels of bradykinin were significantly increased in C1INH ASO-treated mice compared to CTRL ASO-treated mice (Fig 1I). These results suggest that chronic knockdown of liver-derived C1INH leads to activation of the KKS and to the generation of bradykinin.

Studies have shown that elevated levels of bradykinin correlate with low blood pressure (Guilarte et al., 2017; Sala-Cunill et al., 2015; Waeber et al., 1984). Since plasma from our C1INH ASO-treated mice showed a significant increase in bradykinin, we examined the blood pressure of these mice. We found a significant decrease in systolic, diastolic, and mean blood pressure in C1INH ASO-treated mice compared to CTRL ASO-treated mice (Fig 1J).

Fig.ure 1

Figure 1. Elimination of C1INH in plasma of C1INH ASO treated mice and activation of the KKS. (A) Representative Western blot of plasma from mice treated with CTRL ASO vs C1INH ASO, probed with an anti-C1INH (A), HK comprised of two bands (110 kDa and 85 kDa) (C) and plasma kallikrein (E) antibodies under reduced conditions. Quantification of C1INH (B) levels show a significant reduction in C1INH ASO treated mice compared to controls

(p<0.001, n=12/group). Quantification of Western blots showed a significant decrease in the total level of intact HK (110 kDa + 85 kDa; D; p=0.002, n=9-10/group) and a significant increase in the level of plasma kallikerin (F; p=0.003, n=9-10/group) between plasma from C1INH ASO treated mice compared to CTRL ASO treated mice. Samples were normalized to transferrin (TF). (G, H) Representative kallikrein activity plot and quantification of only males treated with CTRL ASO or C1INH ASO. Plasma from C1INH ASO treated mice show a significant increase in KKS activity (p=0.0244, n=5-7/group) compared to CTRL ASO. (I) ELISA analysis indicates there was significantly more bradykinin in the plasma of C1INH ASO treated mice compared to the control group (p=0.009, n=7-11/group). (J) Blood pressure measurement was determined by tail cuff plethysmography. Systolic, diastolic, and mean blood pressure readings indicate that C1INH ASO treated mice are hypotensive in comparison to the control group (p=0.032, p=0.014, p=0.036; n=9- 11/group).

C1INH was first characterized as the major inhibitor of the complement system initiation cascade, therefore we explored if the downregulation of peripheral C1INH induced complement system protein activation. Surprisingly, we found no changes in the levels of C1qA and C1r by Western blot (Supp Fig 3A-D and Supplementary figure 3H). Furthermore, levels of peripheral C3a were not changed when depleting peripheral C1INH, examined by Western blot and ELISA (Supp Fig 3E-G). These results suggest that knockdown of endogenous peripheral C1INH does not induce the activation of the complement system in the periphery.

ASO treatments target mRNA in the liver which may cause liver dysfunction resulting in pro-inflammatory responses (Lapinski, 2001; Mehta et al., 2015)ˑ(Liu, 2009), neurological deficits, brain edema, and brain inflammation (Jiang et al., 2009; O'Beirne et al., 2006; Sawhney et al., 2016). We ruled out liver dysfunction as a possible outcome of ASO treatment effects (Butterworth, 2013) by using alanine aminotransferase (ALT) assay and ELISA for pro-inflammatory cytokines in the plasma (Supp Fig 4A-D) which showed no significant changes between the treated groups. Furthermore, we measured the spleen as another indication of liver dysfunction and found no differences in weight between the groups (Supplementary figure 4E), suggesting no toxic liver dysfunction apparent due to the ASO treatment. We also ruled out stress, as a cause for KKS activation (Couto Alves et al., 2013; Kadoch et al., 2014), measured by corticosterone (CORT) levels (Supplementary figure 4F). These low levels of CORT indicate that chronic injections did not induce a stress response which might result in activation of the KKS.

Chronic knockdown of liver-derived C1INH increased bradykinin 2 receptor (B2R) expression in the brain and BBB dysfunction. The bradykinin system is known to be involved in BBB integrity (Bartus et al., 1996), and B2R antagonists can prevent BBB permeability induced by elevated levels of bradykinin (Marcos-Contreras et al., 2016). Bradykinin was suggested to be an important pro-inflammatory inducer in the nervous system (Wahl et al., 1996). Since we detected elevated levels of plasma bradykinin in C1INH ASO-treated mice, we measured the levels of B2R in the brain. Both gene and protein expression levels of B2R were significantly increased by two-fold in C1INH ASO mice compared to controls (Fig 2A-C), suggesting an activation of the bradykinin pathway in the brain.

Next, we examined BBB integrity by immunofluorescence staining of coronal brain sections for IgG. IgG is usually not expressed in healthy brain. There was increased staining of IgG in the parenchyma, outside laminin-positive vessels, specifically in borders of the lateral ventricle and the fimbria (Fig 2D). Using Evans blue, we evaluated the BBB leakage in the brains of C1INH ASO-treated mice compared to CTRL ASO-treated mice. We found an increase in blue staining in the freshly frozen brains of C1INH ASO mice (Fig 2E-F). Moreover, Evans blue dye was more readily detected by fluorescence microscopy in the brain tissue of C1INH ASO mice (Fig 2G), specifically in the margins of the ventricles and parenchyma. When we analyzed the brains for differences in water content, to test for brain edema, we did not find any changes between groups (Supp Fig 5). These results suggest a possible breakdown of the BBB specifically in the lateral ventricle region, with increased expression of B2R.

Figure 2. Increased B2R expression and BBB permeability is caused by depletion of peripheral C1INH. RTPCR of coronal brain samples showed elevated B2R gene expression in mouse brains from C1INH-ASO treated mice compared to controls (A; p=0.001, n=6-7/group). (B, C) Western blot of coronal brain homogenates was performed to compare B2R protein expression between groups. GAPDH was used for normalization. C1INH ASO group showed significantly higher levels of B2R in the brain than did the CTRL ASO group (p<0.001, n=5/group). (D) IgG immunofluorescence staining (green) at the fimbria of the hippocampus outside the vessels of C1INH ASO treated mice compared to that of CTRL ASO indicates a disruption of BBB integrity (arrows). Laminin staining (red) was used to outline blood vessels in the region (100μm scale). (E) Representative images of brain hemispheres from mice after Evans blue injection and perfusion. C1INH ASO mice administered Evans blue show a much greater amount of blue coloring in their brains than Evans blue-injected CTRL ASO mice and untreated mice. (F) Quantification of blue area from brain sectioned area in percentage using ImageJ (p=0.025, n=3) (G) Representative staining of Evans blue (red) in the brain parenchyma and the lateral ventricle. Blue signal is DAPI staining (100μm scale).

Chronic depletion of liver-derived C1INH induced activation of resident glial cell towards a pro-inflammatory response and cognitive decline. Emerging evidence suggests that peripheral inflammation can induce brain pathology leading to neuroinflammation and cognitive decline (Hu et al., 2010; Obasi et al., 2012; Simen et al., 2011). We therefore examined this phenomenon in our studies. We found increased expression of GFAP and CD11b, markers of astrocyte and microglial cells activation,

respectively, in the cortex of C1INH ASO-treated mice compared to CTRL ASO-treated mice (Fig 3). Expression levels of both genes and proteins were significantly increased in the cortex as shown by mRNA analysis (Fig 3A for GFAP; 3E for CD11b) and western blot analysis (Fig 3B&C for astrocytes; Fig 3F&G for microglia/macrophages). We have also confirmed the results using immunofluorescence staining for astrocytes and microglia (Fig 3D for astrocytes; Fig 3H for microglia/macrophages). These results suggest that the activation of resident immune cells in the brain is influenced by reduced levels of C1INH in the periphery.

Figure 3. Glial cell activation in the cortex is induced by depletion of peripheral C1INH. (A) Gene expression of GFAP is increased (p=0.024; n=4-10/group) in the cortex of mice treated with C1INH ASO compared to control treatment. (B) Representative Western blot of GFAP in the cortex of C1INH ASO treated mice vs. CTRL ASO treated mice. Samples were normalized to GAPDH. (C) GFAP protein expression is significantly increased (p=0.014, n=5) in the cortex of C1INH ASO treated mice compared to CTRL ASO treated mice. (D) Representative immunofluorescence staining of GFAP in the cortex of C1INH ASO vs. CTRL ASO treated mice (100μm scale). (E) Gene expression of CD11b is increased (p=0.048, n=4-10/group) in cortex of C1INH ASO vs. CTRL ASO treated mice. (F) Representative Western blot of CD11b in the cortex of C1INH ASO vs. CTRL ASO treated mice. Samples were normalized to GAPDH. (G) CD11b protein expression is increased (p=0.04, n=5) in the cortex of C1INH ASO treated mice compared to CTRL ASO treated mice. (H) Representative immunofluorescence staining of CD11b in the cortex of C1INH ASO vs CTRL ASO treated mice (100μm scale).

Activation of glial cells can trigger inflammatory pathways (Farfara et al., 2008). In order to understand the response of resident glial cell activation, we measured levels of pro-inflammatory cytokines in the brain. We first measured gene expression of pro-inflammatory cytokines such as IL-1β, IL-6 and TNF-α, since they are the major secreted cytokines from activated glial cells. Comparing brain samples from C1INH ASO- and CTRL ASO-treated mice, RT-PCR analysis confirmed the elevation in pro-inflammatory cytokine gene expression (Fig 4A-C) and ELISA confirmed an increase in their proteins (Fig 6D-F). Furthermore, since nitric oxide is known to be secreted from activated resident immune cells during pro-inflammatory response (Farfara et al., 2008), we used immunofluorescence and Western blotting to examine changes in inducible nitric oxide synthase (iNOS). We observed an increase in iNOS expression in brains of C1INH ASO mice compared to CTRL ASO-treated mice (Fig 4G-I).

At the end of the chronic C1INH ASO treatment, we performed behavioral tests to evaluate motor and cognitive function in mice. Using the fear conditioning paradigm, C1INH ASO-treated mice showed reduced percentage of total freezing time on the test day compared to the CTRL ASO group (Fig 4J), indicative of memory impairment. We have also examined cognitive function after 8 weeks of treatment, and found no significant changes (Supp Fig 6A), suggesting the chronic, rather than short term, depletion of liver-derived C1INH induced brain dysfunction. In order to evaluate motor abilities that might be affected by the ASO treatment, we quantified the average distance traveled on the first day of the fear conditioning test prior to foot shock. No motor dysfunction was observed between the groups (Supp Fig 6B). We also performed an open field test to compare the distance which mice traveled after treatment and found no differences (Supp Fig 6C). Since there was no difference observed in motor function between the C1INH ASO and CTRL ASO groups, the reduced freezing time during fear conditioning is most likely a result of cognitive impairment due to the chronic depletion of peripheral C1INH. Although cognitive impairment is often correlated with neurodegeneration, we did not observe evidence of neuronal degeneration

as measured by evaluation of apoptosis (Supplementary figure 6D). This result suggests that the cognitive decline observed in C1INH ASO-treated mice might be due to the loss of BBB integrity and the pro-inflammatory response in the brain, resulting from peripheral activation of the KKS. In conclusion, these results suggest that brain function is mediated by peripheral C1INH.

Figure 4. Increased pro-inflammatory response and cognitive decline in the CNS of mice treated with C1INH ASO. (A-C) Elevated gene expression of (A) IL-1β (p=0.015, n=5-7/group), (B) IL-6 (p=0.019, n=5- 7/group), and (C) TNF-α (p=0.017, n=5-7/group) in brains of C1INH ASO treated mice compared to CTRL ASO treated mice. (D-F) Protein expression levels of pro-inflammatory cytokines are increased in brains of C1INH ASO treated mice compared to controls: (D) Il-1β (p=0.038, n=6), (E) IL-6 (p=0.047, n=6), and (F) TNF-α (p=0.034, n=6). (G) Representative immunofluorescence staining of iNOS in the brains of C1INH ASO treated mice compared to CTRL ASO treated mice. (H) Representative Western blot of iNOS in brain homogenates of C1INH ASO treated mice compared to controls. (I) iNOS protein expression is increased (p<0.001, n=6) in C1INH ASO treated mice compared to controls. See my handwritten suggestions and notes on this Figure. (J) Fear conditioning test (day 2) demonstrates decreased freezing time in C1INH ASO treated mice compared to controls (74.80 ± 6.521 vs. 74.80 ± 6.521 N=8, p=0.022, n=8-10/group).

Discussion

Emerging evidence are raising regarding the effect the peripheral blood

system on brain function. Blood proteins and activated immune cells are known to play a role in many brain pathologies. However, the exact mechanism and the routes involved are still under investigation.

In this study, we show that chronic depletion of liver-derived C1INH lead to BBB permeability, brain inflammation, and cognitive dysfunction. The mechanistic link underlying these results is through the activation of the KKS to produce bradykinin in the periphery but spares the activation of the complement system.

By using an ASO technique that targets the production of C1INH in the liver, we reduced the protein expression in the periphery. We first tested whether C1INH ASO mice demonstrated changes in the initiation factors of the complement system and found no differences in the activation state of the complement system in the plasma or the brain. However, we observed increased activation of the KKS, through the cleavage of intact HK by plasma kallikrein, resulting in a production of bradykinin. This toxic protein is known to induce peripheral vascular permeability and hypotension mediated by bradykinin 2 receptor (Han et al., 2002; Sala-Cunill et al., 2015) however, no data was presented regarding brain integrity and function.

ASOs that target the liver may produce liver disorders which can correlate with an enlarged spleen, elevated pro-inflammatory cytokines(Reid, 1986), and induce neurological deficits and brain inflammation (Jiang et al., 2009; O'Beirne et al., 2006; Sawhney et al., 2016). We analyzed liver dysfunction and secretion of pro-inflammatory mediators in the plasma and found no liver stress measured by ALT or pro-inflammatory response such as IL-6, IL-1β, and TNF-α. Spleen weight was also unchanged between the treated groups, suggesting no toxic liver dysfunction apparent due to the ASO treatment. Altogether, these results indicate that the activation of the KKS in the periphery is not associated with the chronic treatment of the ASO but by the reduction of liver-derived C1INH alone. In the brain, we observed leakage of the BBB, pro-inflammatory response and cognitive decline which are not a result of liver dysfunction due to the chronic C1INH ASO treatment, but because of decreased C1INH levels in the plasma.

Our mouse model partially mimics the clinical pathology of Hereditary angioedema (HAE) type-1, a condition caused by mutations in the SERPING1 gene leading to depletion of C1INH production in humans. HAE type-1 and acquired angioedema (AAE) type-1 are diseases characterized by recurrent episodes of severe, localized inflammation and increased vascular permeability affecting soft tissues, including the gastrointestinal tract, upper airways, and the mucosa (Agostoni and Cicardi, 1992; Carugati et al., 2001). The acute attacks in HAE patients are thought to be mediated by the activation of the KKS and the complement system (Gao et al., 2013). Plasma from C1INH-deficient patients contains decreased levels of intact HK (Schapira et al., 1983). During inflammatory attacks, HK is cleaved (Berrettini et al., 1986) and bradykinin is released (Nussberger et al., 1998). Activation of the contact system was shown to mediate vascular permeability via B2R n in the C1INH-/- model (Han et al., 2002). Today, HAE patients are treated during attacks with either recombinant C1INH or kallikrein inhibitors, which halt the production of bradykinin (Cugno et al., 2009). Moreover, a prophylactic treatment with C1INH was shown to reduce the frequency of acute attacks (Ma et al., 2011). Previously, the term angioneurotic edema was used to characterize angioedema because of the psychological stress involved with this disorder (C.H Douglas, 1957) suggesting an involvement of stress hormones. Physiological stress is considered a trigger for acute attacks in the cases of HAE (Milder et al., 2007). In order to understand if the chronic treatment induced stress response and thus contribute to the activation of the KKS, we tested corticosteroid levels in the plasma of ASO-treated mice compared with that of restrained mice, which demonstrate high levels of plasma corticosteroid. ASO-treated mice did not show elevated levels of plasma corticosteroid, suggesting that the cause of KKS activation and bradykinin overproduction was due to the depletion of the C1INH itself, indicating an important role of controlled, balanced levels of C1INH in the plasma. Depletion of peripheral C1INH led to low blood pressure likely due to the high levels of bradykinin detected in the plasma, which causes blood vessels to dilate. Thus, we concluded KKS activation upon C1INH ASO treatment was not driven by inflammation in the periphery and was specific to C1INH depletion.

C1INH was first suggested to be involved in brain injury in 1983 (Spiteri et al., 1983). Twelve years later, C1INH was shown to be synthesized in the brain and was correlated with Alzheimer's disease (AD) pathology (Walker et al., 1995). Our group has also shown a decreased expression of C1INH in AD plasma compared to non-demented controls (Zamolodchikov et al., 2016). C1INH, as a modulator of inflammation, was increased in the brains of C1INH ASO treated mice as a result of the pro-inflammatory response and the permeable BBB affected by the down-regulated expression of the liver-derived C1INH expression.

The therapeutic roles of C1INH have been observed in several animal models of peripheral inflammatory disease including sepsis (Croner et al., 2004; Giebler et al., 1999; Guerrero et al., 1993; Jansen et al., 1998; Liu et al., 2007; Scherer et al., 1996), xenograft transplant rejection (Fiane et al., 1999; Hecker et al., 2002a; Hecker et al., 2002b), myocardium ischemia-reperfusion injury(Akita et al., 2003; Horstick, 2002; Horstick et al., 2001a; Nielsen et al., 2002), and hemorrhagic shock (Horstick et al., 2001b). Beyond clinical treatment with recombinant C1INH for HAE, clinical treatment with C1INH in humans revealed therapeutic effects in coronary and myocardial infarction (Bauernschmitt et al.,1998; de Zwaan et al., 2002), septic shock (Hack et al., 1993), vascular leak syndrome (Hack et al., 1993), and other acute inflammatory diseases (Niederau et al., 1995; Ogilvie et al., 1994; Tassani et al., 2001). Moreover, mice infected with *Streptococcus pneumonia* and treated with C1INH showed an improvement in clinical signs of bacterial clearance in the cerebrospinal fluid (CSF) and blood, decreased leukocyte infiltration to the CSF, meningitis, and reduced Il-6 levels (Zwijnenburg et al., 2007). Furthermore, administration of C1INH was shown to have neuroprotective roles in acute neurodegenerative diseases such as stroke and traumatic brain injury (Albert-Weissenberger et al., 2014; De Simoni et al., 2003; Heydenreich et al., 2012) by reducing the infarct volume and the neuronal damage in the temporal cortex, striatum, hippocampus, and thalamus (De Simoni et al., 2003).

Our data suggests that chronic depletion of C1INH can cause serious brain pathologies via the activation of the KKS alone, and treatment with

recombinant C1INH, KKS inhibitors, or B2R antagonists might be considered a necessary prophylactic treatment for down-regulated C1INH brain derived-disorders. Furthermore, manipulation of C1INH levels in the periphery might be used to intentionally open the BBB for drug administration. It will be also interesting to determine the role of peripheral C1INH protein in acute and chronic neurodegenerative diseases such as stroke and Alzheimer's disease and evaluate the therapeutic potential of recombinant C1INH, KKS inhibitors, or B2R antagonists in neuro-vascular disorders.

Materials and Methods

Animals. All animal experiments were conducted in accordance with the guidelines of the US NIH Guide for the Care and Use of Laboratory Animals and with approval from the Animal Care and Use Committee of The Rockefeller University. C57/Bl6J mice (males and females) were used for all experiments. A second WT model, C57/C3H, was also used. C1INH ASO (murine sequence)(Bhattacharjee et al., 2013) and control ASO (CTRL ASO, no homologies to the mouse genome) were provided by Ionis Pharmaceuticals.

ASO preparation and treatment. ASOs were dissolved in saline and injected subcutaneously at 150 mg/kg/week for two weeks and then 50 mg/kg/week for 10 weeks. Ten-week-old WT mice were treated with C1INH ASO or CTRL ASO for 12 weeks (n=6-14 mice/group).

Plasma processing. Plasma was collected the day before the treatment started, after two weeks of treatment with 150mg/kg/week (sub-mandibularly), and the day of sacrifice (12 weeks, cardiac puncture). Animals were anesthetized at the end of the treatment using regulated CO_2. Mice were then transcardially perfused with saline. Upon collection, blood was immediately processed in EDTA-containing tubes (BD Microtainer). Heparin was avoided as it has been shown to bind to C1INH (Rossi et al., 2010). Blood was centrifuged at 1300 rpm for 15 min at room temperature (RT). The upper phase of the supernatant was transferred to a second tube containing 0.5 M EDTA PH8 and centrifuged again. The

plasma was aliquoted and immediately frozen and stored at -80 ^{0}C until analysis. During the lifespan of the mice, blood was taken submandibularly. At the end of the experiment, blood was taken from the heart.

Splenocytes and peritoneal macrophages. Spleens and peritoneal macrophages were taken from CTRL ASO and C1INH ASO-treated mice, and single-cell suspensions were prepared (Farfara et al., 2011; Weiss et al., 2011). CD4+ cells and CD8+ cells were positively stained (BD Bioscience) along with C1INH-conjugated FITC (biorbyt) for FACS analysis.

Chronic restraint stress. Chronic restraint stress (CRS) was induced(Wood et al., 2008) with slight modifications. Mice were immobilized for 6 hours using a disposable netting. Blood was collected following the restraint period, and plasma was prepared. Levels of plasma corticosterone were determined using Corticosterone Enzyme Immunoassay Kit (K014-H1, Arbor Assays).

LPS treatment. Mice were injected intraperitoneally (i.p.) with 100 μl of 1 mg/ml lipopolysaccharide (LPS; L-2630 Sigma-Aldrich). Blood was collected and plasma prepared 12 hours after LPS injection. This source was used as a positive control during all ELISA examinations (see below).

Alanine aminotransferase (ALT) activity assay. Liver enzyme function was examined using the ALT Activity Assay (MAK052, Sigma-Aldrich). As a positive control group for this assay, we used lipopolysaccharide (LPS) to induce high levels of secreted ALT levels in the plasma (Baveja et al., 2002).

Blood pressure (Gonsette, 1978). Blood pressure was measured at the end of ASO treatment using tail-cuff plethysmography (Kent Scientific). An average of 3 readings was obtained for each animal during measurement.

Evans blue and brain edema (Lim et al., 1984). Twelve hours before the sacrifice, 2% Evans blue in saline was injected into mice intraperitoneally. After perfusion with saline, one hemisphere was taken to assess percent of H_2O volume using the wet/dry procedure(Hellal et al., 2004). Hemispheres were immediately weighed to obtain wet weight (WW) and heated to 100

^{0}C for 24h. Then samples were weighed again to obtain the dry weight (DW). Brain water content was calculated as %H_2O = (WW-DW) X 100/WW. The other brain hemisphere was collected and sectioned. Loss of BBB integrity was revealed by visualizing Evans blue under the fluorescence microscope.

Immunohistochemistry (Avrahami et al., 2013; Farfara et al., 2015). The primary antibodies used were: anti-GFAP (DAKO Z0334), anti-CD11b (Abcam ab-8878), IgG (Thermo scientific), anti-laminin (Fisher scientific RT-795-PO), and anti-iNOS (Abcam ab-129372). For secondary antibodies, we used IgG anti- mouse, rat, or rabbit, depending on the host of the first antibody (Thermo Scientific). Vectasheild-DAPI (Vector Labs) was used to seal the slides. Brain sections were visualized using Zeiss Axiovert200. Images were analyzed using ImageJ software.

Tunnel assay (Yao et al., 2014). Cell death by apoptosis was examined using the In Situ Cell Death Detection Kit, TMR red (Roche).

Western blot analysis (Chen et al., 2017)**.** Antibodies used for plasma samples: anti-HK light chain (R&D MAB22061); anti-FXII (HTI PAHFFXII-s); anti-plasma kallikrein (R&D AF2498); anti-transferrin (Abcam ab82411). Antibodies used for brain samples: anti-GFAP (DAKO Z0334); anti-CD11b (Abcam ab-133357); anti-B2R (LSBio LS-C405461); anti-GAPDH (Abcam ab9484 or Proteintech 60004-1-Ig). Protein levels were quantified using NIH Image J densitometry. Western blot results for plasma samples were normalized to transferrin, and brain protein extracts were normalized to GAPDH or Actin.

ELISA (Farfara et al., 2011). ELISA was used to determine level of C3a (Molecular Innovations) and bradykinin (ENZO) in plasma and brain homogenate. The samples were normalized to 100% from CTRL-ASO to pool results from three different tests. For detection of pro-inflammatory cytokines, we used Mouse DuoSet ELISA kits for IL-1b, IL-6, and TNF-α (R&D). For a positive control in these assays, we used plasma from an LPS-injected mouse (Qadri et al., 2016).

Gene expression (Farfara et al., 2011). RNA was extracted from fresh

cortex or frozen whole brain using RNeasy Lipid Tissue mini kit (74804, Qiagen). RNA was converted to cDNA using High Capacity cDNA Reverse Transcription Kit (4368814, Life technologies). Gene expression levels were amplified using Taqman enzyme (Applied Biosystems, 4370048) and primers (IL1-β Mm00434228_m1, IL-6 Mm00446190_m1, TNF-α Mm00443258_m1, GFAP Mm01253033_m1, CD11b Mm00434455_m1) and normalized to endogenous mouse GAPDH (Mm99999915_g1). $\Delta\Delta$CT was quantified and compared between samples.

Kallikrein activity assay (Zamolodchikov et al., 2015). Plasma was evaluated for KKS activity using the chromogenic substrate S-2303 (Diapharma) or Pefachrome PK8092 (Pentapharm).

Behavioral analysis (Farfara et al., 2015). All behavioral experiments were performed and analyzed with a researcher blinded to genotype and treatment. Fear conditioning was performed as previously described(Farfara et al., 2015) with slight changes. Two foot shocks were given (0.7 milliamp, 0.5 sec), one after the first three minutes and the second at the end of the five minutes of the first day. Open field test was done as previously described (Gould, 2009).

Statistical analysis. All statistical analyses as indicated in the figure legends were determined by two-tailed t-test. One-tailed t-test was performed when the results confirmed the hypothesis. All numerical values presented in graphs are mean ±SEM.

Acknowledgements:

The authors thank members of the Strickland laboratory for their help. This work was supported by The EGL foundation and NIH; Rudin Family Foundation; Mellam Family Foundation; Mr. John A. Herrmann Jr.; and Mary & James G. Wallach Foundation.

Authorship:

Contributions: D.F. designed the study, performed experiments, analyzed data, and wrote the manuscript; E.F. and A.R. performed

experiments and provided technical assistance; E.H.N and S.S participated in data analysis and manuscript preparation. Conflict-of-interest disclosure: Alexey S. Revenko and A. Robert MacLeod are employees and stockholders of Ionis Pharmaceuticals. The authors declare no additional competing financial interests.

References

Agostoni, A., and M. Cicardi. 1992. Hereditary and acquired C1-inhibitor deficiency: biological and clinical characteristics in 235 patients. *Medicine* 71:206-215.

Akita, N., H. Nakase, T. Kaido, Y. Kanemoto, and T. Sakaki. 2003. Protective effect of C1 esterase inhibitor on reperfusion injury in the rat middle cerebral artery occlusion model. *Neurosurgery* 52:395-400; discussion 400-391.

Albert-Weissenberger, C., S. Mencl, M.K. Schuhmann, I. Salur, E. Gob, F. Langhauser, S. Hopp,
N. Hennig, S.G. Meuth, M.W. Nolte, A.L. Siren, and C. Kleinschnitz. 2014. C1-Inhibitor protects from focal brain trauma in a cortical cryolesion mice model
by reducing thrombo-inflammation. *Frontiers in cellular neuroscience* 8:269.

Avrahami, L., D. Farfara, M. Shaham-Kol, R. Vassar, D. Frenkel, and H. Eldar-Finkelman. 2013.
Inhibition of glycogen synthase kinase-3 ameliorates beta-amyloid pathology
and restores lysosomal acidification and mammalian target of rapamycin
activity in the Alzheimer disease mouse model: in vivo and in vitro studies. *J Biol Chem* 288:1295- 1306.

Bartus, R.T., P. Elliott, N. Hayward, R. Dean, E.L. McEwen, and S.K. Fisher. 1996. Permeability of the blood brain barrier by the bradykinin agonist, RMP-7: evidence for a sensitive,
auto-regulated, receptor-mediated system. *Immunopharmacology* 33:270-278.

Bauernschmitt, R., H. Bohrer, and S. Hagl. 1998. Rescue therapy with C1-esterase inhibitor concentrate after emergency coronary surgery for failed PTCA. *Intensive care medicine* 24:635-638.

Baveja, R., S. Keller, Y. Yokoyama, N. Sonin, M.G. Clemens, and J.X. Zhang. 2002. LPS-induced imbalanced expression of hepatic vascular stress genes in cirrhosis: possible mechanism of increased susceptibility to endotoxemia. *Shock* 17:316-321.

Berrettini, M., B. Lammle, T. White, M.J. Heeb, H.P. Schwarz, B. Zuraw, J. Curd, and J.H. Griffin. 1986. Detection of in vitro and in vivo cleavage of high molecular weight kininogen in
human plasma by immunoblotting with monoclonal antibodies. *Blood* 68:455-462.

Bhattacharjee, G., A.S. Revenko, J.R. Crosby, C. May, D. Gao, C. Zhao, B.P. Monia, and A.R. MacLeod. 2013. Inhibition of vascular permeability by antisense-mediated inhibition of plasma kallikrein and coagulation factor 12. *Nucleic acid therapeutics* 23:175-187.

Bossi, F., E.I. Peerschke, B. Ghebrehiwet, and F. Tedesco. 2011. Cross-talk between the complement and the kinin system in vascular permeability. *Immunology letters* 140:7- 13.

Butterworth, R.F. 2013. The liver-brain axis in liver failure: neuroinflammation and encephalopathy. *Nature reviews. Gastroenterology & hepatology* 10:522-528.

C.H Douglas, J.R.B. 1957. Familial paroxismal dysfunction of the autonomic nervous

system (a periodic disease), often precipitated by emotional stress. *Pediatrics, 18 (1957), pp.*
782–793

Carugati, A., E. Pappalardo, L.C. Zingale, and M. Cicardi. 2001. C1-inhibitor deficiency and angioedema. *Mol Immunol* 38:161-173.

Chen, Z.L., A.S. Revenko, P. Singh, A.R. MacLeod, E.H. Norris, and S. Strickland. 2017. Depletion of coagulation factor XII ameliorates brain pathology and cognitive impairment in

Alzheimer's disease mice. *Blood*

Colten, H.R. 1972. Ontogeny of the human complement system: in vitro biosynthesis of individual complement components by fetal tissues. *J Clin Invest* 51:725-730.

Couto Alves, A., S. Bruhn, A. Ramasamy, H. Wang, J.W. Holloway, A.L. Hartikainen, M.R. Jarvelin, M. Benson, D.J. Balding, and L.J. Coin. 2013. Dysregulation of complement system and CD4+ T cell activation pathways implicated in allergic response. *PLoS One* 8:e74821.

Croner, R.S., T.G. Lehmann, C. Fallsehr, C. Herfarth, E. Klar, and M. Kirschfink. 2004. C1-inhibitor reduces hepatic leukocyte-endothelial interaction and the expression of VCAM-1 in

LPS-induced sepsis in the rat. *Microvascular research* 67:182-191.

Cugno, M., A. Zanichelli, F. Foieni, S. Caccia, and M. Cicardi. 2009. C1-inhibitor deficiency and angioedema: molecular mechanisms and clinical progress. *Trends Mol Med* 15:69-78.

De Simoni, M.G., C. Storini, M. Barba, L. Catapano, A.M. Arabia, E. Rossi, and L. Bergamaschini. 2003. Neuroprotection by complement (C1) inhibitor in mouse transient brain

ischemia. *J Cereb Blood Flow Metab* 23:232-239.

de Zwaan, C., A.H. Kleine, J.H. Diris, J.F. Glatz, H.J. Wellens, P.F. Strengers, M. Tissing, C.E. Hack,

M.P. van Dieijen-Visser, and W.T. Hermens. 2002. Continuous 48-h C1-inhibitor treatment, following reperfusion therapy, in patients with acute myocardial infarction. *European heart journal* 23:1670-1677.

Farfara, D., V. Lifshitz, and D. Frenkel. 2008. Neuroprotective and neurotoxic properties of glial cells in the pathogenesis of Alzheimer's disease. *Journal of cellular and molecular medicine* 12:762-780.

Farfara, D., D. Trudler, N. Segev-Amzaleg, R. Galron, R. Stein, and D. Frenkel. 2011. gamma- Secretase component presenilin is important for microglia beta-amyloid clearance.

Annals of neurology 69:170-180.

Farfara, D., H. Tuby, D. Trudler, E. Doron-Mandel, L. Maltz, R.J. Vassar, D. Frenkel, and U. Oron. 2015. Low-level laser therapy ameliorates disease progression in a mouse model of Alzheimer's disease. *Journal of molecular neuroscience : MN* 55:430-436.

Fiane, A.E., V. Videm, H.T. Johansen, O.J. Mellbye, E.W. Nielsen, and T.E. Mollnes. 1999. C1- inhibitor attenuates hyperacute rejection and inhibits complement, leukocyte and

platelet activation in an ex vivo pig-to-human perfusion model. *Immunopharmacology* 42:231-243.

Gao, Y.P., J. Dong, X. Zhang, Y. Liu, Q. Lu, J.N. Feng, X.R. Tan, and G. Yang. 2013. A novel peptide can mimic extracellular fibrinogen-binding protein to block the activation of complement system. *Cell biochemistry and biophysics* 66:753-757.

Gasque, P., M. Fontaine, and B.P. Morgan. 1995. Complement expression in human brain. Biosynthesis of terminal pathway components and regulators in human glial cells and cell lines. *J Immunol* 154:4726-4733.

Gasque, P., A. Ischenko, J. Legoedec, C. Mauger, M.T. Schouft, and M. Fontaine. 1993. Expression of the complement classical pathway by human glioma in

culture. A model for complement expression by nerve cells. *J Biol Chem* 268:25068-25074.

Gasque, P., A. Thomas, M. Fontaine, and B.P. Morgan. 1996. Complement activation on human neuroblastoma cell lines in vitro: route of activation and expression of functional complement regulatory proteins. *J Neuroimmunol* 66:29-40.

Ghebrehiwet, B., A.P. Kaplan, K. Joseph, and E.I. Peerschke. 2016. The complement and contact activation systems: partnership in pathogenesis beyond angioedema. *Immunological*

reviews 274:281-289.

Giebler, R., U. Schmidt, S. Koch, J. Peters, and R. Scherer. 1999. Combined antithrombin III and C1-esterase inhibitor treatment decreases intravascular fibrin deposition and

attenuates cardiorespiratory impairment in rabbits exposed to Escherichia coli endotoxin. *Crit Care Med* 27:597-604.

Gonsette, R.E. 1978. Complement system activation by contrast media in neuroradiology. *Neuroradiology* 16:253-255.

Gould, T.D. 2009. The Open Field Test. *Mood and Anxiety Related Phenotypes in Mice* Volume 42 pp 1-20.

Guerrero, R., F. Velasco, M. Rodriguez, A. Lopez, R. Rojas, M.A. Alvarez, R. Villalba, V. Rubio, A. Torres, and D. del Castillo. 1993. Endotoxin-induced pulmonary dysfunction is prevented by C1-esterase inhibitor. *J Clin Invest* 91:2754-2760.

Guilarte, M., A. Sala-Cunill, O. Luengo, M. Labrador-Horrillo, and V. Cardona. 2017. The Mast Cell, Contact, and Coagulation System Connection in Anaphylaxis. *Frontiers in immunology* 8:846.

Hack, C.E., A.C. Ogilvie, B. Eisele, A.J. Eerenberg, J. Wagstaff, and L.G. Thijs. 1993. C1-inhibitor substitution therapy in septic shock and in the vascular leak syndrome induced by high doses of interleukin-2. *Intensive care medicine* 19 Suppl 1:S19-28.

Han, E.D., R.C. MacFarlane, A.N. Mulligan, J. Scafidi, and A.E. Davis, 3rd. 2002. Increased vascular permeability in C1 inhibitor-deficient mice mediated by the bradykinin type 2 receptor. *J Clin Invest* 109:1057-1063.

Harpel, P.C., M.F. Lewin, and A.P. Kaplan. 1985. Distribution of plasma kallikrein between C-1 inactivator and alpha 2-macroglobulin in plasma utilizing a new assay for alpha 2- macroglobulin-kallikrein complexes. *J Biol Chem* 260:4257-4263.

Hecker, J.M., R. Lorenz, R. Appiah, B. Vangerow, M. Loss, R. Kunz, J. Schmidtko, M. Mengel, J. Klempnauer, S. Piepenbrock, G. Dickneite, H. Neidhardt, H. Ruckoldt, and M. Winkler. 2002a. C1-inhibitor for prophylaxis of xenograft rejection after pig to cynomolgus monkey kidney transplantation. *Transplantation* 73:688-694.

Hecker, J.M., M. Loss, J. Klempnauer, and M. Winkler. 2002b. C1-inhibitor for therapy and prophylaxis of acute vascular rejection of porcine kidneys in cynomolgus monkeys. *Transplantation proceedings* 34:2382.

Hellal, F., D. Bonnefont-Rousselot, N. Croci, B. Palmier, M. Plotkine, and C. Marchand- Verrecchia. 2004. Pattern of cerebral edema and hemorrhage in a mice model of diffuse brain injury. *Neuroscience letters* 357:21-24.

Heydenreich, N., M.W. Nolte, E. Gob, F. Langhauser, M. Hofmeister, P. Kraft, C. Albert- Weissenberger, M. Brede, C. Varallyay, K. Gobel, S.G. Meuth, B. Nieswandt, G. Dickneite, G. Stoll, and C. Kleinschnitz. 2012. C1-inhibitor protects from brain ischemia- reperfusion injury by combined antiinflammatory and antithrombotic mechanisms. *Stroke* 43:2457-2467.

Horstick, G. 2002. C1-esterase inhibitor in ischemia and reperfusion. *Immunobiology* 205:552- 562.

Horstick, G., O. Berg, A. Heimann, O. Gotze, M. Loos, G. Hafner, B. Bierbach, S. Petersen, S. Bhakdi, H. Darius, M. Horstick, J. Meyer, and O. Kempski. 2001a. Application of C1- esterase inhibitor during reperfusion of ischemic myocardium: dose-related beneficial versus detrimental effects. *Circulation* 104:3125-3131.

Horstick, G., T. Kempf, M. Lauterbach, S. Bhakdi, L. Kopacz, A. Heimann, M. Malzahn, M. Horstick, J. Meyer, and O. Kempski. 2001b. C1-esterase-inhibitor

treatment at early reperfusion of hemorrhagic shock reduces mesentry leukocyte adhesion and rolling.
Microcirculation 8:427-433.

Hu, Z., Y. Ou, K. Duan, and X. Jiang. 2010. Inflammation: a bridge between postoperative cognitive dysfunction and Alzheimer's disease. *Medical hypotheses* 74:722-724.

Jansen, P.M., B. Eisele, I.W. de Jong, A. Chang, U. Delvos, F.B. Taylor, Jr., and C.E. Hack. 1998. Effect of C1 inhibitor on inflammatory and physiologic response patterns in primates suffering from lethal septic shock. *J Immunol* 160:475-484.

Jiang, W., P. Desjardins, and R.F. Butterworth. 2009. Cerebral inflammation contributes to encephalopathy and brain edema in acute liver failure: protective effect of

minocycline. *Journal of neurochemistry* 109:485-493.

Johnson, A.M., C.A. Alper, F.S. Rosen, and J.M. Craig. 1971. C1 inhibitor: evidence for decreased hepatic synthesis in hereditary angioneurotic edema. *Science* 173:553-554.

Kadoch, C., J. Li, V.S. Wong, L. Chen, S. Cha, P. Munster, C.A. Lowell, M.A. Shuman, and J.L. Rubenstein. 2014. Complement activation and intraventricular rituximab distribution in recurrent central nervous system lymphoma. *Clinical cancer research : an official*

journal of the American Association for Cancer Research 20:1029-1041.

Lapinski, T.W. 2001. The levels of IL-1beta, IL-4 and IL-6 in the serum and the liver tissue of chronic HCV-infected patients. *Archivum immunologiae et therapiae experimentalis* 49:311-316.

Laurell, A.B., U. Martensson, and A.G. Sjoholm. 1976. C1 subcomponent complexes in normal and pathological sera studied by crossed immunoelectrophoresis. *Acta Pathol Microbiol Scand C* 84C:455-464.

Lener, M., G. Vinci, C. Duponchel, T. Meo, and M. Tosi. 1998. Molecular cloning, gene structure and expression profile of mouse C1 inhibitor. *Eur J Biochem* 254:117-122.

Lim, H.W., M.B. Poh-Fitzpatrick, and I. Gigli. 1984. Activation of the complement system in patients with porphyrias after irradiation in vivo. *J Clin Invest* 74:1961-1965.

Liu, D., F. Lu, G. Qin, S.M. Fernandes, J. Li, and A.E. Davis, 3rd. 2007. C1 inhibitor-mediated protection from sepsis. *J Immunol* 179:3966-3972.

Liu, Q. 2009. Role of cytokines in the pathophysiology of acute-on-chronic liver failure. *Blood purification* 28:331-341.

Ma, Y.J., A. Doni, M.O. Skjoedt, C. Honore, M. Arendrup, A. Mantovani, and P. Garred. 2011. Heterocomplexes of mannose-binding lectin and the pentraxins PTX3 or serum amyloid

P component trigger cross-activation of the complement system. *J Biol Chem* 286:3405- 3417.

Marcos-Contreras, O.A., S. Martinez de Lizarrondo, I. Bardou, C. Orset, M. Pruvost, A. Anfray, Y. Frigout, Y. Hommet, L. Lebouvier, J. Montaner, D. Vivien, and M. Gauberti. 2016.

Hyperfibrinolysis increases blood-brain barrier permeability by a

plasmin- and bradykinin-dependent mechanism. *Blood* 128:2423-2434.

Mehta, G., R.P. Mookerjee, V. Sharma, and R. Jalan. 2015. Systemic inflammation is associated with increased intrahepatic resistance and mortality in alcohol-related acute-on-

chronic liver failure. *Liver international : official journal of the International Association for the Study of the Liver* 35:724-734.

Milder, F.J., L. Gomes, A. Schouten, B.J. Janssen, E.G. Huizinga, R.A. Romijn, W. Hemrika, A. Roos, M.R. Daha, and P. Gros. 2007. Factor B structure provides insights into activation of the central protease of the complement system. *Nature structural & molecular biology* 14:224-228.

Morris, K.M., D.P. Aden, B.B. Knowles, and H.R. Colten. 1982. Complement biosynthesis by the human hepatoma-derived cell line HepG2. *J Clin Invest* 70:906-913.

Niederau, C., R. Brinsa, M. Niederau, R. Luthen, G. Strohmeyer, and L.D. Ferrell. 1995. Effects of C1-esterase inhibitor in three models of acute pancreatitis. *International journal of pancreatology : official journal of the International Association of*

Pancreatology
17:189-196.
Nielsen, E.W., T.E. Mollnes, J.M. Harlan, and R.K. Winn. 2002. C1-inhibitor reduces the ischaemia-reperfusion injury of skeletal muscles in mice after aortic cross-clamping.
Scandinavian journal of immunology 56:588-592.
Nussberger, J., M. Cugno, C. Amstutz, M. Cicardi, A. Pellacani, and A. Agostoni. 1998. Plasma bradykinin in angio-oedema. *Lancet* 351:1693-1697.

O'Beirne, J.P., M. Chouhan, and R.D. Hughes. 2006. The role of infection and inflammation in the pathogenesis of hepatic encephalopathy and cerebral edema in acute liver failure.
Nature clinical practice. Gastroenterology & hepatology 3:118-119.
Obasi, C.N., K.J. Cruickshanks, D.M. Nondahl, B.E. Klein, R. Klein, F.J. Nieto, A. Shankar, M.E. Fischer, M.Y. Tsai, and R. Chappell. 2012. Association of Biomarkers for Inflammation, Endothelial Dysfunction and Oxidative Stress with Cognitive Impairment. The
Epidemiology of Hearing Loss Study (EHLS). *Oxidants and antioxidants in medical science* 1:169-173.
Ogilvie, A.C., J.W. Baars, A.J. Eerenberg, C.E. Hack, H.M. Pinedo, L.G. Thijs, and J. Wagstaff. 1994. A pilot study to evaluate the effects of C1 esterase inhibitor on the toxicity of
high-dose interleukin 2. *British journal of cancer* 69:596-598.
Orsini, F., D. De Blasio, R. Zangari, E.R. Zanier, and M.G. De Simoni. 2014. Versatility of the complement system in neuroinflammation, neurodegeneration and brain homeostasis.
Frontiers in cellular neuroscience 8:380.
Pixley, R.A., M. Schapira, and R.W. Colman. 1985. The regulation of human factor XIIa by plasma proteinase inhibitors. *J Biol Chem* 260:1723-1729.
Qadri, F., F. Rimmele, L. Mallis, W. Hauser, A. Dendorfer, O. Johren, P. Dominiak, L.M. Leeb- Lundberg, and M. Bader. 2016. Acute hypothalamo-pituitary-adrenal axis response to
LPS-induced endotoxemia: expression pattern of kinin type B1 and B2 receptors. *Biological chemistry* 397:97-109.
Ratnoff, O.D., and I.H. Lepow. 1957. Some properties of an esterase derived from preparations of the first component of complement. *J Exp Med* 106:327-343.
Reid, K.B. 1986. Activation and control of the complement system. *Essays in biochemistry* 22:27-68.
Rossi, V., I. Bally, S. Ancelet, Y. Xu, V. Fremeaux-Bacchi, R.R. Vives, R. Sadir, N. Thielens, and G.J.
Arlaud. 2010. Functional characterization of the recombinant human C1 inhibitor serpin domain: insights into heparin binding. *J Immunol* 184:4982-4989.
Sala-Cunill, A., J. Bjorkqvist, R. Senter, M. Guilarte, V. Cardona, M. Labrador, K.F. Nickel, L. Butler, O. Luengo, P. Kumar, L. Labberton, A. Long, A. Di Gennaro, E. Kenne, A. Jamsa,
T. Krieger, H. Schluter, T. Fuchs, S. Flohr, U. Hassiepen, F. Cumin, K. McCrae, C. Maas, E. Stavrou, and T. Renne. 2015. Plasma contact system activation drives anaphylaxis in severe mast cell-mediated allergic reactions. *J Allergy Clin Immunol* 135:1031-1043 e1036.
Sawhney, R., P. Holland-Fischer, M. Rosselli, R.P. Mookerjee, B. Agarwal, and R. Jalan. 2016. Role of ammonia, inflammation, and cerebral oxygenation in brain dysfunction of acute-on-chronic liver failure patients. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 22:732-742.
Schapira, M., C.F. Scott, and R.W. Colman. 1982. Contribution of plasma protease inhibitors to the inactivation of kallikrein in plasma. *J Clin Invest* 69:462-468.
Schapira, M., L.D. Silver, C.F. Scott, A.H. Schmaier, L.J. Prograis, Jr., J.G. Curd, and R.W. Colman. 1983. Prekallikrein activation and high-molecular-weight kininogen consumption

in
hereditary angioedema. *N Engl J Med* 308:1050-1053.

Scherer, R.U., R.M. Giebler, U. Schmidt, D. Paar, and W.J. Kox. 1996. The influence of C1- esterase inhibitor substitution on coagulation and cardiorespiratory parameters in an endotoxin-induced rabbit model of hypercoagulability. *Seminars in thrombosis and hemostasis* 22:357-366.

Schmaier, A.H., and K.R. McCrae. 2007. The plasma kallikrein-kinin system: its evolution from contact activation. *Journal of thrombosis and haemostasis : JTH* 5:2323-2329.

Sim, R.B., G.J. Arlaud, and M.G. Colomb. 1979. C1 inhibitor-dependent dissociation of human complement component C1 bound to immune complexes. *Biochem J* 179:449-457.

Simen, A.A., K.A. Bordner, M.P. Martin, L.A. Moy, and L.C. Barry. 2011. Cognitive dysfunction with aging and the role of inflammation. *Therapeutic advances in chronic disease*
2:175-195.

Spiteri, D., L. Viola, G. Allegretti De Lista, F. Galloro, and G. Schisano. 1983. Changes of C1 esterase inhibitor, alpha2 globulin and sedimentation rate after severe brain injury.
Acta neurologica 5:420-425.

Tassani, P., R. Kunkel, J.A. Richter, H. Oechsler, H.P. Lorenz, S.L. Braun, G.P. Eising, F. Haas, S.U.
Paek, R. Bauernschmitt, M. Jochum, and R. Lange. 2001. Effect of C1-esterase-inhibitor on capillary leak and inflammatory response syndrome during arterial switch operations in neonates. *Journal of cardiothoracic and vascular anesthesia*
15:469-473.

van der Graaf, F., J.A. Koedam, J.H. Griffin, and B.N. Bouma. 1983. Interaction of human plasma kallikrein and its light chain with C1 inhibitor. *Biochemistry* 22:4860-4866.

Vinci, G., N.J. Lynch, C. Duponchel, T.M. Lebastard, G. Milon, C. Stover, W. Schwaeble, and M.
Tosi. 2002. In vivo biosynthesis of endogenous and of human C1 inhibitor in transgenic mice: tissue distribution and colocalization of their expression. *J Immunol* 169:5948- 5954.

Waeber, B., J. Nussberger, H.R. Brunner, A. de Agostini, and M. Schapira. 1984. [Role of bradykinin in hypotension induced by the active factor derived from factor XII].
Schweizerische medizinische Wochenschrift 114:1749-1751.

Wahl, M., E.T. Whalley, A. Unterberg, L. Schilling, A.A. Parsons, A. Baethmann, and A.R. Young. 1996. Vasomotor and permeability effects of bradykinin in the cerebral microcirculation. *Immunopharmacology* 33:257-263.

Walker, D.G., O. Yasuhara, P.A. Patston, E.G. McGeer, and P.L. McGeer. 1995. Complement C1 inhibitor is produced by brain tissue and is cleaved in Alzheimer disease. *Brain Res*
675:75-82.

Weiss, R., V. Lifshitz, and D. Frenkel. 2011. TGF-beta1 affects endothelial cell interaction with macrophages and T cells leading to the development of cerebrovascular amyloidosis.
Brain, behavior, and immunity 25:1017-1024.

Wood, G.E., E.H. Norris, E. Waters, J.T. Stoldt, and B.S. McEwen. 2008. Chronic immobilization stress alters aspects of emotionality and associative learning in the rat. *Behavioral neuroscience* 122:282-292.

Yao, Y., Z.L. Chen, E.H. Norris, and S. Strickland. 2014. Astrocytic laminin regulates pericyte differentiation and maintains blood brain barrier integrity. *Nature communications* 5:3413.

Zamolodchikov, D., Z.L. Chen, B.A. Conti, T. Renne, and S. Strickland. 2015. Activation of the factor XII-driven contact system in Alzheimer's disease patient and mouse model
plasma. *Proceedings of the National Academy of Sciences of the United States of America* 112:4068-4073.

Zamolodchikov, D., T. Renne, and S. Strickland. 2016. The Alzheimer's disease peptide beta- amyloid promotes thrombin generation through activation of coagulation factor XII. *Journal of thrombosis and haemostasis : JTH* 14:995-1007.

Ziccardi, R.J., and N.R. Cooper. 1979. Active disassembly of the first complement component, C- 1, by C-1 inactivator. *J Immunol* 123:788-792.

Zwijnenburg, P.J., T. van der Poll, S. Florquin, M.M. Polfliet, T.K. van den Berg, C.D. Dijkstra, J.J.

Roord, C.E. Hack, and A.M. van Furth. 2007. C1 inhibitor treatment improves host defense in pneumococcal meningitis in rats and mice. *The Journal of infectious diseases* 196:115-123.

Figure 4. Increased pro-inflammatory response and cognitive decline in the CNS of mice treated with C1INH ASO. (A-C) Elevated gene expression of (A) IL-1β (p=0.015, n=5-7/group), (B) IL-6 (p=0.019, n=5- 7/group), and (C) TNF-α (p=0.017, n=5-7/group) in brains of C1INH ASO treated mice compared to CTRL ASO treated mice. (D-F) Protein expression levels of pro-inflammatory cytokines are increased in brains of C1INH ASO treated mice compared to controls: (D) Il-1β (p=0.038, n=6), (E) IL-6 (p=0.047, n=6), and (F) TNF-α (p=0.034, n=6). (G) Representative immunofluorescence staining of iNOS in the brains of C1INH ASO treated mice compared to CTRL ASO treated mice. (H) Representative Western blot of iNOS in brain homogenates of C1INH ASO treated mice compared to controls. (I) iNOS protein expression is increased (p<0.001, n=6) in C1INH ASO treated mice compared to controls. See my handwritten suggestions and notes on this Figure. (J) Fear conditioning test (day 2) demonstrates decreased freezing time in C1INH ASO treated mice compared to controls (74.80 ± 6.521 vs. 74.80 ± 6.521 N=8, p=0.022, n=8-10/group).

Figure 4

A.

B2R mRNA
(% from control)

p=0.001

CTRL ASO   C1INH ASO

B.

CTRL ASO   C1INH-ASO

B2R   43 kDa

GAPDH   40 kDa

C.

B2R/GAPDH
(% from control)

p<0.001

CTRL ASO   C1INH ASO

D.

IgG   Laminin   Merge

CTRL
ASO

C1INH
ASO

E.

CTRL ASO   CTRL ASO   C1INH ASO

Evans Blue   -   +   +

F.

% Area

p=0.025

CTRL ASO   C1INH ASO

G.

CTRL ASO   C1INH ASO

Figure3

Figure 6