EpiGraph: an open-source platform to quantify epithelial organization.

Pablo Vicente-Munuera[1,†], Pedro Gómez-Gálvez[1,†], Robert J. Tetley[2], Cristina Forja[1], Antonio Tagua[1], Marta Letrán[1], Melda Tozluoglu[2], Yanlan Mao[2,3,*] and Luis M. Escudero[1,*].

1: Departamento de Biología Celular, Universidad de Sevilla and Instituto de Biomedicina de Sevilla (IBiS), Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla. 41013 Seville, Spain.

2: MRC Laboratory for Molecular Cell Biology, University College London, Gower Street, London WC1E 6BT, UK.

3: College of Information and Control, Nanjing University of Information Science and Technology, Nanjing, Jiangsu 210044, China

†: These authors contributed equally to this work.

*: Corresponding authors: Yanlan Mao (y.mao@ucl.ac.uk) and L.M. Escudero (lmescudero-ibis@us.es).

Lead contact: L.M. Escudero (lmescudero-ibis@us.es).

SUMMARY

During development, cells must coordinate their differentiation with their growth and organization to form complex multicellular structures such as tissues and organs. Healthy tissues must maintain these structures during homeostasis. Epithelia are packed ensembles of cells from which the different tissues of the organism will originate during embryogenesis. A large barrier to the analysis of the morphogenetic changes in epithelia is the lack of simple tools that enable the quantification of cell rearrangements. Here we present EpiGraph, an image analysis tool that quantifies epithelial organization. Our method combines computational geometry and graph theory to measure the degree of order of any packed tissue. EpiGraph goes beyond the traditional polygon distribution analysis, capturing other organizational traits that improve the characterization of epithelia. EpiGraph can objectively compare the rearrangements of epithelial cells during development and homeostasis to quantify how the global ensemble is affected. Importantly, it has been implemented in the open-access platform FIJI. This makes EpiGraph very user friendly, with no programming skills required.

KEYWORDS

Morphogenesis, Epithelial organization, Graph Theory.

HIGHLIGHTS

Epithelial images can be converted into graphs to quantify their topology.

EpiGraph is a new simple method to capture epithelial organization.

Epigraph can detect changes in tissue fluidity.

EpiGraph has been designed as a plugin of FIJI to be accessible to any biology lab.

INTRODUCTION

The development of any multicellular organism is based on coordinated changes that transform the embryo into the adult individual. During morphogenesis and growth, patterning, cell divisions and architectural changes must perfectly fit together for the correct development of the body plan. Any morphogenetic movement such as migration, extension or invagination of epithelial cells is coupled with dramatic changes in the organization of cells (Bertet et al., 2004; Blankenship et al., 2006; Escudero et al., 2007; Farhadifar et al., 2007; Girdler and Roper, 2014; Lecuit and Lenne, 2007; Pilot and Lecuit, 2005). After development, homeostatic tissues must maintain their complex organization of cells in order to function correctly. Changes in normal organization is often an indication of disease onset, for which early detection would be vital (Emmanuele et al., 2015; Park et al., 2015; Sáez et al., 2013).

The control of cell proliferation and its role in morphogenesis has been deeply studied and the main signalling pathways that control organ size and shape have been revealed. In contrast, how tissues modulate and maintain their organization during development and homeostasis is an important question that remains unsolved. This is mainly due to the lack of simple and general methods that can capture and quantify the arrangement of cells. It has been known for almost a hundred years that epithelial tissues exhibit a degree of order. The analysis of epithelial organization has been mainly based on the number of neighbours of the epithelial cells, considering the apical surface of these cells as convex polygons with the same number of sides as neighbours. In previous works, we have investigated several aspects of the organization of packed tissues using Voronoi tessellations to compare the polygon distributions of natural and mathematical tessellations (Sanchez-Gutierrez et al., 2016). We have described that the polygon distribution of natural tessellations is restricted to a series of frequencies of polygons that match the Voronoi diagrams that conform to the Centroidal Voronoi tessellation (CVT). This is what we call a "CVT path" and was used as a scale to compare the organization of different packed tissues. However, polygon

distribution is not sufficient to completely characterize tissue organization. Tissues with clearly different appearance can present very similar polygon distribution (Sanchez-Gutierrez et al., 2016).

As an alternative approach, we have proposed that Graph Theory could capture differences in the topology of tissues (Escudero et al., 2011; Sanchez-Gutierrez et al., 2013; Sánchez-Gutiérrez et al., 2017). This is based on the idea of converting the epithelium into a network of cell-to-cell contacts (Escudero et al., 2011). The resulting "epithelial graph" can be analysed by combining the tools of network theory and multivariable statistical analysis (Escudero et al., 2011; Kursawe et al., 2016; Sanchez-Gutierrez et al., 2013; Yamashita and Michiue, 2014). Finding features and patterns that can describe the graphs is key in many diverse fields, including biology (Benson et al., 2016; Costa et al., 2007; Hayes et al., 2013). A network can be split up into different subgraphs named graphlets. The graphlet composition of a network has been used to quantify differences between complex systems (Hayes et al., 2013; Ho et al., 2010; Kuchaiev et al., 2011; Pržulj et al., 2004). These measurements are based on the comparison of the number of these subgraphs in different networks and are able to provide an index of distance between them. This feature has the advantage of integrating the differences between diverse networks into a single value, simplifying the analyses and allowing multiple comparisons.

In summary, there is a clear need for a method to specifically quantify tissue organization and aid the interpretation of biophysical and mechanical aspects of morphogenesis and tissue homeostasis (Blankenship et al., 2006; Canela-Xandri et al., 2011; Farhadifar et al., 2007; Gibson et al., 2006, 2011; Gómez-Gálvez et al., 2018; Heisenberg and Bellaiche, 2013; Heller et al., 2016; Lecuit and Lenne, 2007; Mao et al., 2013; Sanchez-Gutierrez et al., 2016; Zallen and Zallen, 2004). The advances in imaging techniques, together with the appearance of powerful methods for automated image analysis (Heller et al., 2016; Khan et al., 2014; Kursawe et al., 2016) and new simulation resources (Bi et al., 2015, 2016; Blanchard et al., 2009; Etournay et al., 2016; Fletcher et al., 2014; Guirao et al., 2015; Mirams et al., 2013; Tanaka et al.,

2015) provide a large amount of source data that require analysis in terms of organization. Here we present the open source platform EpiGraph, a new image analysis method that uses segmented images from real epithelia or simulations, to easily quantify and compare the organization of packed tissues.

RESULTS

Graphlet measurements as an approach to capture organization of packed tissues.

In previous studies, a set of 29 graphlets was used to distinguish between different types of networks (Pržulj et al., 2004). This method calculated the Graphlet degree Distribution agreement Distance (GDD) between two networks (Pržulj, 2007). Therefore, the "GDD value", that in theory can range from 0 to 1, weighs the differences among the two distributions of graphlets; the higher the value, the more different the arrangements. (**Fig. S1** and **Methods**). Epithelial images can be considered as natural tessellations and converted into networks of cell-to-cell contacts (Escudero et al., 2011). We have used the "graphlet" approach to capture the topology of epithelial tissues. Tessellations give rise to "geographic networks" (Albert and Barabasi, 2002) that only make sense in a planar surface. For this reason, when we translated the set of graphlets to cellular patterns, some of them were redundant (G5 and G27) or not possible (G20, G22, G25). Therefore, in this study we have used a total of 26 graphlets corresponding to 29 different cellular motifs that account for the organization of groups of up to 5 cells (**Fig. 1A, Fig. S1**). Most of the analyses performed in this work were completed with only 17 motifs (Mo17, **Fig. 1A,** mauve). We found that, although all the motifs could be present in an actual tissue, Mo17 minimized the redundancy of the information provided by the graphlets. In addition, this set downplays the importance of rare cellular geometries that could excessively weight GDD calculations (for example, a high difference in GDD could appear when comparing an image with one or two square cells versus another image with no four-sided cells; this effect is minimized using Mo17). However, it would

be possible to use other combinations such as all the motifs (Mo29) or cellular motifs that account for the organization of groups of up to 4 cells (Mo10) (**Fig. 1A**).

Graphlet measurements capture differences beyond polygon distributions.

We tested the power of graphlet-based measurements in quantifying differences between sets of images with very similar polygon distributions (**Fig. 1B**). In third instar larvae of *Drosophila,* the photoreceptors are specified, giving rise to a particular repetitive arrangement of the presumptive eye cells (Eye, **Fig. 1C**). This arrangement is very different to the irregular distribution in a Voronoi tessellation where the initial seeds were placed in a random way (Sanchez-Gutierrez et al., 2016), (Diagram 1, **Fig. 1C**). We previously showed that it was not possible to discriminate between the polygon distributions of these two tessellations (Sanchez-Gutierrez et al., 2016). Using the graphlets approach, we obtained a GDD value of 0.086 when comparing these two sets of images (Mo17, **Table S1, Fig. S2**). In order to know if this difference was biologically relevant, we tried to set a baseline, by comparing other images with very similar polygon distribution that also presented an apparently similar arrangement. This was the case for Diagram 4 of the CVT vs. the *Drosophila* wing imaginal disc in larvae (dWL) and Diagram 5 of the CVT vs. the *Drosophila* wing imaginal disc in prepupae (dWP) (**Fig. 1B-C**). Both results were in the same range, with a GDD value of 0.042 for Diagram 4 vs. dWL and 0.049 for Diagram 5 vs. dWP (**Fig. 1C**). Similar results were obtained when comparing Diagram 4 vs. Diagram 5 and dWL vs. dWP (**Fig. 1-C**). These results suggested the existence of a baseline in the range of 0.04-0.05 values that correspond to similar cellular arrangements that cannot be well distinguished using the graphlets distribution. Therefore, we interpreted the value of 0.086 obtained in the Eye vs. Diagram 1 comparison as the reflection of actual differences between these two sets. In all the mentioned cases, the results obtained using Mo17 and Mo29 were equivalent (**Table S1**).

EpiGraph quantitatively compares the organization of multiple sets of images.

The GDD had the limitation of comparing only 2 samples each time. However, one of our objectives was to evaluate different types of images together. Therefore, we designed EpiGraph, a method that calculates the GDD of any epithelial tissue with another tessellation that serves as a reference. As an output, we obtained an index of distance for each image with respect to this reference. We used three different references: i) a tessellation formed by regular hexagons, representing the most ordered way to pave the space (**Fig. 2A**, GDDH). ii) the network motifs emerging from a random Voronoi tessellation (**Fig. 2B**, GDDRV). iii) a Voronoi Diagram 5 from the CVT path (**Fig. 2C**, GDDV5) that presents a polygon distribution similar to the one from multiple examples in nature (Gibson et al., 2006; Sanchez-Gutierrez et al., 2016).

We tested the method with epithelial images that have been previously compared with the CVT path in terms of polygon distribution: chicken neural tube (cNT), dWL, dWP, reduction of myosin II in the *Drosophila* prepupa wing disc epithelium (dMWP) and Eye (**Fig. 2D**)(Sanchez-Gutierrez et al., 2016). To have a scale and facilitate fast comparisons, we used the concept of the CVT path (Sanchez-Gutierrez et al., 2016). We calculated the values for GDDH, GDDRV and GDDV5 for all the Voronoi diagrams and used this property to visualize the results and plot the GDDH, GDDRV and GDDV5 with respect to the percentage of hexagons of the corresponding diagram. The CVT does not progress beyond the 70% of hexagons limiting the possibilities of analysis. Therefore, we extended the Voronoi scale spanning a wider range of polygon distributions. The algorithm that devises the CVT was modified to introduce "noise" in the positioning of the seed that produces the subsequent diagram. In this way, we obtained a "CVT noise" (CVTn) whose last diagrams reached 90% of hexagons (**Fig. 2E-G** and **Experimental Procedures**). We decided to compare GDDH, GDDRV and GDDV5 values with the percentage of hexagons, since it is indicative of the proportion of the other types of polygons along the CVTn (**Table S2**). Interestingly, the plot

obtained using CVT and CVTn diagrams was an optimum way to easily visualize these geometric scales as a continuous "CVT path" and a "CVTn path". Therefore, we used this framework to analyse the values of GDDH, GDDRV and GDDV5 for each diagram in the scale. As expected, the GDDH values were higher in the initial diagrams and progressively decreased with the increase in the percentage of hexagons of the Voronoi diagrams (**Fig. 2E, left panel**). The opposite happened in the case of the GDDRV values (**Fig. 2E, central panel**). In the plot of percentage of hexagons vs GDDV5, the CVTn path presented the shape of a walking stick (**Fig. 2E, right panel**). The GDDV5 values of Voronoi Diagrams 1, 2, 3, and 4 were decreasing progressively, with Diagram 5 the closest to the zero value. The values for the rest of the diagrams gradually increased, as in the case of the GDDRV.

We then plotted the values for the actual epithelia. We found that for cNT, dWL and dWP the GDDH, GDDRV and GDDV5 values were similar to the CVTn at the same percentage of hexagons of the polygon distribution (**Fig. 2E**). In agreement with our previous results using the GDD, the Eye images presented a higher GDDRV and GDDV5 values than the expected for a 30% of hexagons (**Fig. 2E**). The differences with respect to the CVTn were even more clear when plotting GDDH vs GDDRV and GDDVR vs GDDV5 (**Fig. 2F-G**). We obtained similar results when analysed the dMWP set of images. In this case, our previous work showed a small deviation of the dMWP polygon distribution with respect the CVT (Sanchez-Gutierrez et al., 2016). However, using Epigraph, we observed that GDDRV and GDDV5 captured the clear differences in organization between these images and the CVTn (**Fig. 2E-G, Fig. S3**). These results suggested that EpiGraph is able to distinguish between different tessellations with a similar polygon distribution. In this regard, we have developed a statistical output using an outlier detection approach whose quantitative results represent how similar the organization of a tissue is when compared with the CVTn scale (**Experimental Procedures**). The test confirmed that cNT, dWL, and dWP were close to the CVTn and similar to the Voronoi diagrams 1, 4, and 6 respectively. In contrast,

the Eye and dMWP samples were labelled as different (**Table S3**). In this way, EpiGraph provides a quantitative description to tissue morphogenesis.

Epigraph can capture different organization traits.

We further investigated the possible applications of EpiGraph and performed a series of experiments aimed at understanding what traits of tissue organization are being captured and quantified by the graphlet measurements. To this end, we have used images of different vertex model simulations that alter tissue organization by changing the biophysical properties of the cells (images taken from (Sanchez-Gutierrez et al., 2016) (**Experimental Procedures** and **Fig. 3A-F**).

First, we analysed samples with 10% of the cells having increased effective cell-cell adhesion (**Experimental Procedures, Fig. 3B**). This feature induced the formation of cells with a "squared shape" that often organized in motifs presenting four-way vertex configurations. These images were compared with simulations in which "elongated" cells appear (by simultaneously increasing cell-cell adhesion and reducing ideal area, **Experimental Procedures** and **Fig. 3C**). Epigraph analysis indicated that while control simulations gave similar values to the CVTn, the "squared" and "elongated" sets of images were different to the control and well separated from the CVTn. However, EpiGraph failed to find differences between the "squared" and "elongated" images (**Fig. 3G** and **Fig. S4**).

Second, we used a set of conditions to mimic the effect of a reduction of myosin II in the *Drosophila* prepupa wing disc epithelium (dMWP, **Fig. 2D**). In the control simulation (**Fig. 3D**), cells grow to double the original area and then divide into two cells. In case III and case IV simulations there was a random reduction of the tension parameter together with a requirement of a minimum tension threshold to be able to divide (**Fig. 3E-F**). If the cells do not reach this threshold they continue to grow without dividing the cell body. When this happens, the cells will be stuck in mitotic phase and will not start a second round of cell division (Sanchez-Gutierrez et al., 2016) (**Experimental Procedures**). The control simulation gave similar values to the CVTn, while

case III, case IV and dMWP images presented a clear deviation in the GDDRV vs GDDV5 graph (**Fig. 3G**). All these data-points distributed in the same zone of the graph (**Fig. 3G**). Interestingly, we found that both sets of simulations (squared and elongated vs Case III and Case IV) appeared in two complementary regions, suggesting that the regions in the graph can reflect the existence of different traits of organization in each condition.

EpiGraph: a method to capture epithelial organization implemented in FIJI.

Aiming to enhance the accessibility of the analysis of tissue organization to the biology community, we have implemented EpiGraph as a plugin for FIJI (Schindelin et al., 2012). EpiGraph consists of a pipeline of 5 very simple steps. First, the skeleton of an epithelial image is uploaded and the individual cells are identified. Second, the user selects the distance threshold to identify two cells as neighbours. Here it is possible to select different thresholds and to check the number of neighbours of every cell in each case. Third, a ROI is selected. There are several possibilities such as a default ROI from the image or the selection of individual cells. Fourth, the graphlet information for the selected cells is calculated. These data are used to obtain the GDDH, GDDRV and GDDV5. These values are incorporated into a table and serve as input data for a statistical analysis that indicates if a new image is inside or outside of the CVTn path and describes which Voronoi diagram presents the most similar organization to the sample (**Experimental procedures**). The fifth step includes the classification and labelling of different images in order to represent them in a new window. This final phase allows one to export the representation of the data in a three-dimensional graph. **Movie S1** shows an example of EpiGraph usage. A detailed description of EpiGraph can be found in the **Supplemental Experimental Procedures**. A full set of tutorials explaining how to install and use EpiGraph is available at EpiGraph's wiki (https://imagej.net/EpiGraph).

EpiGraph provides biological insights I: Development and morphogenesis

As a proof of principle of the functionality of EpiGraph, we used images obtained from the literature, not obtained by our group and not directly related with our work. We chose a tissue where the organization of cells clearly changes over time: the appearance of rosettes during germ-band extension of the *Drosophila* embryo (Blankenship et al., 2006). We selected eight representative time points from "Movie S1" of this article and quantified the graphlet content (**Fig. 4A**). After segmentation (**Experimental Procedures**), the images were uploaded to EpiGraph and plotted in the 3D graph (**Fig. 4B-C**). We found that the early frames from the original movie (001, 035) were located close to the CVT. Progressively, the appearance of rosettes (100, 135) correlated with the separation of these samples from the CVT. This distance was greatest in the case of the image corresponding with frame 170. In contrast, the last two frames did not increase the distance with the CVTn. We observed a reversion in the measurement of the organization, with image 235 located very close to the time point 135, presumably as multicellular rosettes had begun to resolve by this time (**Table S3**). Interestingly, this was captured using different combinations of the four measured parameters (percentage of hexagons, GDDH, GDDRV and GDDV5, **Fig. 4B-C**). The ability to compare different steps of a developmental process suggested that EpiGraph was a versatile method, able to capture and analyse the dynamic changes in organization of developing tissues.

EpiGraph provides biological insights II: Homeostasis and tissue fluidity transitions

In the case of *Drosophila* germ-band extension, we have shown that EpiGraph can detect clear differences in the organization of the cells during development. Epithelial tissues have the ability to behave as a fluid due to cellular rearrangements or to solidify as cellular rearrangements cease (Bi et al., 2015, 2016). The shape index is a characteristic of epithelial cells that has been shown, in vertex model simulations, to be able to capture the degree of rigidity, or fluidity, of a tissue (Bi et al., 2015). This study established the transition point between a soft (fluid) and a rigid (solid) tissue, described as a jamming transition, at the dimensionless shape index value of 3.81. We

calculated the shape index for the CVTn path, finding that from Voronoi diagrams 1 to 20, the tessellations were behaving as a fluid. Using this descriptor, all the real images were placed in the fluid part as well as the four altered vertex model simulations shown in **Fig. 3** (**Fig. S3**, **Fig. S4, Fig. S5** and **Table S4**).

We have investigated the dynamics of epithelial jamming in different conditions. First, to test the capabilities of EpiGraph in this regard, we analysed several snapshots from two simulations published by Bi and cols. as supplementary movies (Bi et al., 2016). These videos show the movements of cells in two conditions: rigid state (shape index less than 3.81) and soft state (shape index greater than 3.81) (**Fig. 5A**). As expected, the snapshots of the soft tissue analysed appeared in different positions, indicating that the simulated epithelia changed its organization during the experiment. On the other hand, the different frames from the rigid simulation were clustered (**Fig. 5B**), showing little cell rearrangements.

We next tested whether EpiGraph could detect changes in tissue fluidity in real epithelia, which may be more ambiguous and noisier than simulations. Real tissues also display fluid-to-solid jamming transitions which are important for large scale tissue shape changes as well as for refining and maintaining tissue shape (Curran et al., 2017; Mongera et al., 2018). In the *Drosophila* pupal notum, the level of tissue fluidity is controlled by the global level of myosin II activity (Curran et al., 2017). We wondered if the regulation of myosin II could similarly impact on the fluidity state of the wing disc epithelium and the cell rearrangements that have been described during the late stages of normal wing disc development, where the overall tissue shape does not dramatically change (Heller et al., 2016) (**Fig. 5C**). To this end, we compared the WT organization with the effect of increasing myosin II activity by knocking down Mbs (Myosin binding subunit of the myosin phosphatase, which dephosphorylates myosin regulatory light chain, **Fig. 5D**) by RNAi throughout the entire wing pouch. Based on work in the pupal notum (Curran et al., 2017), we would predict that *Mbs-RNAi* discs behave as solids. Interestingly, in the two cases, the shape index was greater than the

described shape index threshold of 3.81, suggesting that both tissues are in a fluid state.

We used EpiGraph to analyse the changes in organization of wing discs with perturbed myosin II activity along time and compared them with a WT condition. The snapshots for WT samples appeared clustered in the 3D graph, indicating that the epithelia were not changing their organization during the 30 minutes of analysis (**Fig. 5E**), despite previous work showing that cell intercalations do occur (Heller et al., 2016). This suggests that during this slow growing phase of wing disc development, any cell rearrangements that occur do not drive large-scale morphogenesis, but act to maintain a homeostatic tissue topology. The statistical analysis confirmed that all of the WT wing discs were close to CVTn diagrams 3 and 4 (**Table S3**). In the case of the three samples from the *Mbs-RNAi* genotype, the data points presented different organizations (from similar to diagram 3 to close to diagram 13, see **Table S3**). In some cases, the dispersion was not only between samples, but occurred between images from each movie (**Fig. 5E** and **Table S3**). EpiGraph therefore predicted that these *Mbs-RNAi* wing discs are behaving very differently from WT wing discs, likely by changing their degree of fluidity. Accordingly, quantification of intercalation rates demonstrated that cell rearrangements happen significantly more frequently in WT than in *Mbs-RNAi* wing discs (**Fig. 5F**, 0.1281±0.08 vs. 0.0076±0.01 intercalations/cell/hour, Kolmogorov-Smirnov test, *p*=0.0079). As predicted, this resulted in more cells 'jamming' at 4-way vertex configurations as they fail to complete intercalations (**Fig. 5G**, 0.0065±0.005 vs. 0.0116±0.006 fourfold vertices/cell, Kolmogorov-Smirnov test, *p*=0.029). Interestingly, EpiGraph was able to detect this solidification of the tissue in the *Mbs-RNAi* discs, even though the shape index predicted a fluid tissue. Taken together, these results indicate that the quantification of tissue organization using EpiGraph can be used to study the dynamics of an epithelium from fixed snapshots, without the need to laboriously track individual frames of a time-lapse video.

DISCUSSION

Textbook definitions of morphogenesis include the term "organization" as key to explaining this fundamental developmental process (Dai and Gilbert, 1991). The authors wondered, "How can matter organize itself so as to create a complex structure such as a limb or an eye?" Here we have provided a tool that can help the developmental biology community to solve that question.

The analysis of the polygon sides of epithelial cells has been shown to be insufficient to completely understand tissue organization. Some tessellations can present very different arrangements yet have the same frequencies of number of neighbours. A second problem is the lack of a simple value as an indicator of epithelial organization. This feature complicates the comparison between morphogenesis of normal development and that of genetically perturbed or diseased tissues. Our previous attempts to overcome this caveat were based on multi-statistical analyses of graph features (Sanchez-Gutierrez et al., 2013) and the creation of a Voronoi scale to statistically compare groups of images with the CVT reference (Sánchez-Gutiérrez et al., 2016). However, we are aware that these methods are difficult to incorporate into the average developmental biology lab.

We have developed EpiGraph, aiming to solve both problems. EpiGraph transforms the image into a graph of cell-to-cell contacts and extracts their graphlet content to later compare with other images. These complex algorithms are hidden behind the friendly user window of FIJI. This is the most popular open-source biological image analysis platform. In addition, the output data options of EpiGraph facilitate fast and clear representations and interpretations of the results. Therefore, our method can be easily used in any developmental biology lab.

One of the strengths of EpiGraph is the comparison of any tessellation with the hexagonal lattice, the "random" Voronoi tessellation and the Voronoi tessellation that presents the "conserved polygon distribution" (Gibson et al., 2006; Sánchez-Gutiérrez et al., 2016) (**Fig. 2A-C**). We have tested EpiGraph with different types of samples: as expected, the average of the natural

tessellations such as dWL, dWP and cNT matched the CVTn path position (**Fig. 2D-G**). We interpret that these three natural samples present similar polygon distributions and graphlet compositions to some Voronoi Diagrams from the CVTn. This was independent of the combination of GDD values plotted. On the other hand, the average of the Eye samples appeared far from the CVTn when GDDV5 or GDDRV values were plotted. These two GDD were capturing differences in organization between the Eye and any Voronoi Diagram (including Diagram 1, which presents a similar polygon distribution to the Eye). This result supports the utility of EpiGraph to quantify organizational traits that were not accessible until now. The same idea is reinforced by the results obtained with the mutant samples for myosin II (dMWP). In previous work, we showed that this set of samples slightly deviated the CVT scale in terms of polygon distribution (Sánchez-Gutiérrez et al., 2016). Here we show very clear differences in terms of the values of GDDV5 and GDDRV (**Fig. 2G**), suggesting a higher sensitivity of the new method when capturing differences in organization.

The output images from EpiGraph show the CVTn path as a clear reference for proliferative epithelia such the wing imaginal disc or the chicken neural tube and for vertex model control simulations. We have incorporated a statistical test into EpiGraph that indicates if a new tissue is within or outside of the CVTn path, and which is the Voronoi diagram with the closest organization. The different results comparing GDDH, GDDRV and GDDV5 values also suggested that GDDH had better resolution for images with a higher percentage of hexagons while GDDRV and GDDV5 were more sensitive to the differences between images with less than 40% of hexagons. For this reason, we have designed the visualization step of the program to easily change the three axes and check the different results using any combination of GDDs and the "percentage of hexagons".

Using different sets of simulations, we are able to distinguish two different types of organization: The cases where a subset of cells presents a particular arrangement inside a mostly ordered tissue (**Fig 3B, C, G**) and the cases where the global topology of the tissue is altered (**Fig. 3E, F, G**). These two

patterns create a "map" of arrangements that are out the CVTn, and they will

help to other researchers to study the degree of order in their samples.

As a proof of principle of the capabilities of our method, we have used a

previously published movie to analyse the changes of organization during

morphogenesis. We have chosen the emergence and resolution of rosettes

during germ-band extension, since it is a well-known event with a dynamic

and complex arrangement of cells (Blankenship et al., 2006). The selected

time lapse images included the formation and resolution of the rosettes (**Fig.**

4A). The GDD values of these images correlate with the developmental

process and the 3D visualization tool of EpiGraph captures the complete

morphogenetic event (**Fig. 4B-C**). The selected frames 001, 035 and 070,

appear near the CVTn diagrams since they share a similar composition of

motifs. This starts to change with the formation of rosettes (frames 100, 135

and 170) that progressively separate them from the CVTn. The rosettes

contain specific motifs characterising this developmental biology process and

they are captured by the graphlet analysis. Indeed, the two last frames (200

and 235) show that this tendency is being reversed, going back closer to the

CVTn. We think that this is reflecting the return to an epithelial organization

without multiple motifs required to describe a multicellular rosette; namely the

resolution of rosettes associated with tissue elongation. These analyses

support EpiGraph's ability to provide a quantitative link to morphogenesis.

The dynamics of the transition between a tissue behaving as a fluid or a

solid is an emerging problem in developmental biology and biomedicine

(Curran et al., 2017; Mongera et al., 2018; Park et al., 2015; Tetley and Mao,

2018). We have used the capabilities of EpiGraph to study how the fluidity

state can affect the organization of a tissue. The utility of EpiGraph in this

regard is supported by its ability to quantify dynamic changes in organization

due to cell rearrangements in a vertex model simulation of a soft tissue (**Fig.**

5A-B). However, cell rearrangements do not necessarily have to lead to

changes in tissue organization, as is often the case in more homeostatic

tissues. Although, it has been shown that the late third instar *Drosophila*

imaginal disc can exchange neighbours and rearrange during development

(Heller et al., 2016)**,** we were not able to see changes in organization combining live imaging of the WT discs and EpiGraph analysis (**Fig. 5E**). Therefore, we interpret that the multiple re-arrangements of the WT tissue conserve the organization of the tissue, at least in the time framework analysed (30 min). On the contrary, the hyperactivation of myosin II (*Mbs-RNAi*) produced a clear change in the organization of the tissue as detected by EpiGraph, suggesting a change in tissue fluidity. EpiGraph is therefore able to efficiently detect mutant phenotypes and provide predictions related to changes in tissue fluidity. Importantly, this can be done from a few snap shots in time, without the need for sophisticated time-lapse imaging and tracking. This may provide a simple detection tool for the early onset of disease, where changes in tissue fluidity can occur, and only limited tissue samples are available from patients.

EpiGraph limitations.

Although Epigraph accepts a wide range of images as inputs, we have specified some minimum requirements. It does not accept images bigger than 3000 pixels of width or 3000 pixels of height, since processing them could be computationally intensive. In addition, EpiGraph only accepts single images. Images from time series should be adapted to single frames before uploading them to EpiGraph.

Computers with little RAM memory (less than 16gb) will work but with a series of restrictions. To ensure usability, it is not recommended computing images with a high number of cells (more than 1000) due to a possible lack of memory. In the same way, we suggest skeletonizing the edges of the images and using a small radius (lower than 10 pixels) to calculate the cells neighbourhood. Choosing a high radius value could slow down the work queue, increasing the use of RAM memory.

If any of these requirements are not satisfied, the program alerts the user, allowing him/her to change the image provided. Importantly, the images and ROIs require a minimum number of cells in order to get coherent graphlets. Therefore, to get any result, EpiGraph must detect at least a 4' valid cells (a

cell that does not have a no valid cell within a distance of four cells connexions) in the case of Mo7 or Mo10 or a 5' valid cell (a cell that does not have a no valid cell within a distance of five cells connexions) in the case of Mo17 and Mo29. In any case, we strongly recommend having a greater number of 4' and 5' valid cells to get results that can be trusted in terms of capturing the organization of a tissue. Regarding the 3D visualization tool, it allows the user to see the position of the samples from different angles. However, the resolution of the exported file is only 72 pixels per inch (dpi). This could be too low for publications and therefore EpiGraph provides an excel table with all the information needed to represent it with other programs.

In summary, we have generated a very accessible, open source method to produce a quantitative description of developmental events. This quantitative aspect is reinforced by the statistical comparison with the CVT path that serves as a scale for tissue organization. We anticipate that our tool will improve the study of tissue dynamics and morphogenesis by permitting the comparative analysis of epithelial organization in genetically mutated or diseased tissues during time.

EXPERIMENTAL PROCEDURES

Source images used in the study.

Centroidal Voronoi Tessellation (CVT) diagrams and variations

For the generation of this set of paths we have used the software Matlab R2014b to iteratively apply Lloyd´s algorithm to a random Voronoi tessellation (Lloyd, 1982). This implies that the centroid of a cell in a Voronoi diagram is the seed for the same cell in the next iteration.

- Centroidal Voronoi Tessellation (CVT) diagrams

Centroidal Voronoi Tessellation diagrams were obtained as described previously by our group (Sanchez-Gutierrez et al., 2016). The original 20 original Voronoi diagrams were created placing 500 seeds randomly in an

image of 1024x1024 pixels. A total of 700 iterations were generated for each initial image.

- Centroidal Voronoi Tessellation noise (CVTn) diagrams

We have developed a variation of the CVT path, named the CVT noise (CVTn) path. We started from the same 20 initial random diagrams described above. The development process of the CVTn path was modified so that the new seeds were not strictly the centroid from the previous iteration. In even iterations, we selected a region of 5 pixels of radius from the centroid position, in which seeds could be placed randomly. In odd iterations, the system was stabilized, applying the original Lloyd algorithm. A total of 700 iterations were generated for each initial image.

<u>Natural packed tissues and vertex model simulations</u>

The details of the obtaining and processing of epithelial images were described in (Escudero et al., 2011). Control vertex model simulations include cell proliferation and are the basis for the other two cases. Case III corresponds to a vertex model simulation with heterogeneous reduction of line tension and an impairment of cell division when the tension value is under 30 percent of the initial value. Case IV is a similar simulation to Case III, with a threshold of 40 percent. The exact conditions for the vertex model simulations were described in (Sanchez-Gutierrez et al., 2016).

<u>Natural packed tissues and vertex model simulations</u>

The details of the obtaining and processing of the epithelial images were described in (Escudero et al., 2011). Control vertex model simulations include cell proliferation and are the basis for the other two cases. Case III corresponds to a vertex model simulation with heterogeneous reduction of line tension and an impairment of cell division when tension value is under 30 percent of the initial value. Case IV is a similar simulation to Case III with a threshold of 40 percent. Regarding simulations with no cell proliferation, as a baseline, the control had homogeneous parameters for contractility, line tension and ideal area. 'Elongated' simulations were as the control, but with

ten percent of cells having a reduced line tension and ideal area, while

'squared' ones had ten percent of cells with only line tension reduced. The

exact conditions for the vertex model simulations were described in

(Sanchez-Gutierrez et al., 2016).

<u>Perturbing myosin II activity in *Drosophila* wing discs and calculating</u>

<u>intercalation rates</u>

Drosophila were raised in standard conditions. Wing discs were dissected

from third instar larvae and cultured under filters as described by (Zartman et

al., 2013). Discs were cultured in Shields and Sang M3 media supplemented

with 2% FBS, 1% pen/strep, 3ng/ml ecdysone and 2ng/ml insulin. The

following alleles and transgenes were used; *shg*-GFP (Ecad-GFP, Huang et

al., 2009), *UAS-Mbs-RNAi* (KK library, VDRC), *rn*-GAL4 (RMCE-MiMIC

Trojan-GAL4 collection). The following experimental genotypes were used;

Ecad-GFP (WT) and Ecad-GFP/UAS-*Mbs-RNAi*; *rn*-GAL4/+ (*Mbs-RNAi*). For

EpiGraph analysis, discs were imaged on a Zeiss LSM 880 microscope with

Airyscan at 512x512 resolution with a 63x objective (NA 1.4) at 1.4x zoom for

a total of 30 minutes with 1 minute time intervals and a z-step of 0.5μm. Time-

lapse image sequences were segmented using Epitools (Heller et al., 2016).

To quantify intercalation rates, 5 WT and 5 *Mbs-RNAi* wing discs were

imaged using the same methods as above, except using 5x zoom and 3

minute intervals for a total of 2 hours. Intercalation rate data was exported

from the "EDGE_T1_TRANSITIONS" overlay in the "CellOverlay" plugin in

Epitools. To exclude mistakes generated when 4-way junctions were not

recognised, junctions less than 0.075μm in length were assigned a length of

0μm. A productive intercalation event was scored when a neighbour

exchange was stabilised for at least 2 time points (6 minutes). The total

number of tracked cells was also quantified, allowing the intercalation rate to

be expressed as the number of intercalations per cell per hour.

We also counted the number of fourfold vertices per cell in both WT and

Mbs-RNAi conditions. In particular, we quantified the number of vertices in

which four or more cells were touching each other, using Matlab R2014b. The

cells closest to the border of the image were excluded from the analysis. In this way, we obtained the percentage of fourfold vertices per valid cell for each image and calculated a Kolmogorov-Smirnov test to check if the distributions of both conditions were different.

<u>Multicellular Rosette Formation segmentation</u>

We have segmented a set of images taken from a movie that represents the evolution of the multicellular rosette formation. This is "Movie S1" from Blankenship and cols., 2006 article (url: <u>https://ars.els-cdn.com/content/image/1-s2.0-S153458070600400X-mmc2.mov</u>) (Blankenship et al., 2006). The segmentation procedure consisted of the following steps: 1) We used Trainable Weka Segmentation (Arganda-Carreras et al., 2017) a plugin developed for FIJI (Yoon et al., 1995). This plugin uses machine learning to correctly segment images. We trained the program with a representative image of the whole set, classifying two categories: the outlines and the body of the cells. After the training step, we applied the plugin classifier to all the chosen images from the film, getting a first set of segmented images. 2) Manual curation: This step was necessary to rigorously reflect the real cell outlines of the segmented images, and discard the zones with bad visibility or artefacts. The process was carried out by using Adobe Photoshop CS6. 3) Image processing: We developed a script in Matlab R2014b to clean the noise and the segmentation imperfections. We used several morphological operations for this purpose. Finally, we fixed the cells outlines with a width of 3 pixels to preserve the nature of the image.

Soft and Rigid tissue simulations

We have extracted a set of screenshots from two videos that simulated different dynamical behaviour of vertex model simulations. These videos are presented as Supplemental Material in (Bi et al., 2016). The first video represents a rigid behaviour in the simulation: <u>https://journals.aps.org/prx/supplemental/10.1103/PhysRevX.6.021011/solid tissue v0 0.2 p0 3.5 Dr 0.1.mp4</u>; the second one represents a soft behaviour:https://journals.aps.org/prx/supplemental/10.1103/PhysRevX.6.02

1011/fluid_tissue_v0_0.2_p0_3.8_Dr_0.1.mp4. In both videos, we selected a

total of 13 frames with steps of 3.333 seconds (from t = 0 to 40 seconds).

Graphlets and motifs selection.

The different images from the previous section were used to create a graph of cell-to-cell contacts ((Escudero et al., 2011) and **Supplemental Experimental Procedures**) that served as the source for the graphlet analysis (Pržulj, 2007; Przulj et al., 2004). First, we adapted the graphlet analysis performed by EpiGraph to the nature of our samples (tessellations). Three graphlets were discarded since they were not possible in the context of an epithelial tissue (**Fig. 1** and **Fig. S1**). Second, we used the computer program for graphlet identification and calculation ORCA (Orbit Counting Algorithm) (Hocevar and Demsar, 2014), to extract the different conformations of nodes assembling the graphlets, called orbits (Pržulj, 2007). We computed the Graphlet degree Distribution of the 73 given orbits from the 29 graphlets, and then we removed the non-used ones.

Shape index calculation

We have extracted the shape index, as an indicator of rigidity, from each natural and simulated image, based on (Bi et al., 2015). The global shape index in a tissue was measured as the median of the shape index of the individual valid cells. We quantified the cell area and perimeter using Matlab R2014b. We performed the following approach: We captured the vertex coordinates for each valid cell. Then, we calculated the Euclidean distance between each adjacent vertex, and adding all of them, we got the cell perimeter. From these vertices, a polygon was inferred and we calculated its contained area using the "polyarea" Matlab function.

Statistical analysis.

We have estimated the closest CVTn diagram of a given image in terms of the three GDDs measured in EpiGraph (GDDH, GDDRV and GDDV5). We computed the centre of the point cloud formed by the 20 randomizations of a particular CVTn diagram as the mean of those twenty images, obtaining a 3D point. Then, we calculated the Euclidean distance between all the CVTn

diagrams central points and the four calculated parameters of the input image, obtaining its closest point, which corresponds to its closest diagram. Furthermore, we checked if this image belonged to the closest diagram point cloud using an outlier detection approach. In particular, we tested if the inclusion of the image into a CVTn diagram point cloud would increase or decrease the standard deviation of the original group. We assigned the probability of being an inlier, which is defined as follows:

Where is the total number of coordinates, which in our case is 3 dues to the three-dimensional space; the parameter stands for every different coordinate (GDDH, GDDRV and GDDV5); represents the values of the 20 images from the closest CVTn diagram in a specific coordinate and is the value of the input image for the same coordinate. The values range from 0 (very far from point cloud) to $+\infty$ (inside point cloud). We have estimated that with a confidence of > 0.95 the input image is considered to be an inlier.

REFERENCES

Albert, R., and Barabasi, A.-L. (2002). Statistical mechanics of complex networks. Rev. Mod. Phys. *74*, 47–97.

Arganda-Carreras, I., Kaynig, V., Rueden, C., Eliceiri, K.W., Schindelin, J., Cardona, A., and Seung, H.S. (2017). Trainable Weka Segmentation: a machine learning tool for microscopy pixel classification. Bioinformatics.

Benson, A.R., Gleich, D.F., and Leskovec, J. (2016). Higher-order organization of complex networks. Science (80-.). *353*, 163–166.

Bertet, C., Sulak, L., and Lecuit, T. (2004). Myosin-dependent junction remodelling controls planar cell intercalation and axis elongation. Nature *429*, 667–671.

Bi, D., Lopez, J.H., Schwarz, J.M., and Manning, M.L. (2015). A density-independent rigidity transition in biological tissues. Nat. Phys. *11*, 1074–1079.

Bi, D., Yang, X., Marchetti, M.C., and Manning, M.L. (2016). Motility-driven glass and jamming transitions in biological tissues. Phys. Rev. X *6*.

Blanchard, G.B., Kabla, A.J., Schultz, N.L., Butler, L.C., Sanson, B., Gorfinkiel, N., Mahadevan, L., and Adams, R.J. (2009). Tissue tectonics: morphogenetic strain rates, cell shape change and intercalation. Nat

Methods *6*, 458–464.

Blankenship, J.T., Backovic, S.T., Sanny, J.S., Weitz, O., and Zallen, J.A. (2006). Multicellular rosette formation links planar cell polarity to tissue morphogenesis. Dev Cell *11*, 459–470.

Canela-Xandri, O., Sagués, F., Casademunt, J., and Buceta, J. (2011). Dynamics and mechanical stability of the developing dorsoventral organizer of the wing imaginal disc. PLoS Comput. Biol. *7*.

Costa, L.D.A.F., Rodrigues, F.A., Travieso, G., and Boas, P.R. V (2007). Characterization of complex networks: a survey of measurements. Adv. Phys. *56*, 167–242.

Curran, S., Strandkvist, C., Bathmann, J., de Gennes, M., Kabla, A., Salbreux, G., and Baum, B. (2017). Myosin II Controls Junction Fluctuations to Guide Epithelial Tissue Ordering. Dev. Cell *43*, 480–492.e6.

Dai, J.D., and Gilbert, L.I. (1991). Metamorphosis of the corpus allatum and degeneration of the prothoracic glands during the larval-pupal-adult transformation of Drosophila melanogaster: a cytophysiological analysis of the ring gland. Dev Biol *144*, 309–326.

Emmanuele, V., Kubota, A., Garcia-Diaz, B., Garone, C., Akman, H.O., Sánchez-Gutiérrez, D., Escudero, L.M., Kariya, S., Homma, S., Tanji, K., et al. (2015). Fhl1 W122S causes loss of protein function and late-onset mild myopathy. Hum. Mol. Genet. *24*.

Escudero, L.M., Bischoff, M., and Freeman, M. (2007). Myosin II regulates complex cellular arrangement and epithelial architecture in Drosophila. Dev Cell *13*, 717–729.

Escudero, L.M., Costa Lda, F., Kicheva, A., Briscoe, J., Freeman, M., and Babu, M.M. (2011). Epithelial organisation revealed by a network of cellular contacts. Nat Commun *2*, 526.

Etournay, R., Merkel, M., Popović, M., Brandl, H., Dye, N.A., Aigouy, B., Salbreux, G., Eaton, S., and Jülicher, F. (2016). TissueMiner: A multiscale analysis toolkit to quantify how cellular processes create tissue dynamics. Elife *5*, 773–786.

Farhadifar, R., Roper, J.C., Aigouy, B., Eaton, S., and Julicher, F. (2007). The influence of cell mechanics, cell-cell interactions, and proliferation on epithelial packing. Curr Biol *17*, 2095–2104.

Fletcher, A.G., Osterfield, M., Baker, R.E., and Shvartsman, S.Y. (2014). Vertex models of epithelial morphogenesis. Biophys. J. *106*, 2291–2304.

Gibson, M.C., Patel, A.B., Nagpal, R., and Perrimon, N. (2006). The emergence of geometric order in proliferating metazoan epithelia. Nature *442*, 1038–1041.

Gibson, W.T., Veldhuis, J.H., Rubinstein, B., Cartwright, H.N., Perrimon, N., Brodland, G.W., Nagpal, R., and Gibson, M.C. (2011). Control of the mitotic cleavage plane by local epithelial topology. Cell *144*, 427–438.

Girdler, G.C., and Roper, K. (2014). Controlling cell shape changes during salivary gland tube formation in Drosophila. Semin Cell Dev Biol *31*, 74–81.

Gómez-Gálvez, P., Vicente-Munuera, P., Tagua, A., Forja, C., Castro, A.M., Letrán, M., Valencia-Expósito, A., Grima, C., Bermúdez-Gallardo, M., Serrano-Pérez-Higueras, Ó., et al. (2018). Scutoids are a geometrical solution to three-dimensional packing of epithelia. Nat. Commun.

Guirao, B., Rigaud, S.U., Bosveld, F., Bailles, A., López-Gay, J., Ishihara, S., Sugimura, K., Graner, F., and Bellaïche, Y. (2015). Unified quantitative characterization of epithelial tissue development. Elife *4*, 773–786.

Hayes, W., Sun, K., and Pržulj, N. (2013). Graphlet-based measures are suitable for biological network comparison. Bioinformatics *29*, 483–491.

Heisenberg, C.P., and Bellaiche, Y. (2013). Forces in tissue morphogenesis and patterning. Cell *153*, 948–962.

Heller, D., Hoppe, A., Restrepo, S., Gatti, L., Tournier, A.L., Tapon, N., Basler, K., and Mao, Y. (2016). EpiTools: An Open-Source Image Analysis Toolkit for Quantifying Epithelial Growth Dynamics. Dev Cell *36*, 103–116.

Ho, H., Milenković, T., Memisević, V., Aruri, J., Przulj, N., and Ganesan, A.K. (2010). Protein interaction network topology uncovers melanogenesis regulatory network components within functional genomics datasets. BMC Syst. Biol. *4*, 84.

Hocevar, T., and Demsar, J. (2014). A combinatorial approach to graphlet counting. Bioinformatics *30*, 559–565.

Khan, Z., Wang, Y.-C., Wieschaus, E.F., and Kaschube, M. (2014). Quantitative 4D analyses of epithelial folding during Drosophila gastrulation. Development *141*, 2895–2900.

Kuchaiev, O., Stevanovi, A., and Hayes, W. (2011). GraphCrunch 2: Software tool for network modeling , alignment and clustering.

Kursawe, J., Bardenet, R., Zartman, J.J., Baker, R.E., and Fletcher, A.G. (2016). Robust cell tracking in epithelial tissues through identification of maximum common subgraphs. BioRxiv.

Lecuit, T., and Lenne, P.F. (2007). Cell surface mechanics and the control of cell shape, tissue patterns and morphogenesis. Nat Rev Mol Cell Biol *8*, 633–644.

Lloyd, S. (1982). Least square quantization in PCM. IEEE Trans. Inform. Theory *28*, 129–137.

Mao, Y., Tournier, A.L., Hoppe, A., Kester, L., Thompson, B.J., and Tapon, N. (2013). Differential proliferation rates generate patterns of mechanical tension that orient tissue growth. EMBO J *32*, 2790–2803.

Mirams, G.R., Arthurs, C.J., Bernabeu, M.O., Bordas, R., Cooper, J., Corrias, A., Davit, Y., Dunn, S.J., Fletcher, A.G., Harvey, D.G., et al. (2013). Chaste: an open source C++ library for computational physiology and biology. PLoS Comput Biol *9*, e1002970.

Mongera, A., Rowghanian, P., Gustafson, H.J., Shelton, E., Kealhofer, D.A., Carn, E.K., Serwane, F., Lucio, A.A., Giammona, J., and Campàs, O. (2018). A fluid-to-solid jamming transition underlies vertebrate body axis elongation. Nature.

Park, J.A., Kim, J.H., Bi, D., Mitchel, J.A., Qazvini, N.T., Tantisira, K., Park, C.Y., McGill, M., Kim, S.H., Gweon, B., et al. (2015). Unjamming and cell shape in the asthmatic airway epithelium. Nat. Mater. *14*, 1040–1048.

Pilot, F., and Lecuit, T. (2005). Compartmentalized morphogenesis in epithelia: from cell to tissue shape. Dev Dyn *232*, 685–694.

Przulj, N., Corneil, D.G., and Jurisica, I. (2004). Modeling interactome: scale-free or geometric? Bioinformatics *20*, 3508–3515.

Pržulj, N. (2007). Biological network comparison using graphlet degree distribution. Bioinformatics *23*, 177–183.

Pržulj, N., Corneil, D.G., and Jurisica, I. (2004). Modeling interactome : scale-free or geometric ? *20*, 3508–3515.

Sáez, A., Rivas, E., Montero-Sánchez, A., Paradas, C., Acha, B., Pascual, A., Serrano, C., and Escudero, L.M. (2013). Quantifiable diagnosis of muscular dystrophies and neurogenic atrophies through network analysis. BMC Med. *11*.

Sanchez-Gutierrez, D., Saez, A., Pascual, A., and Escudero, L.M. (2013). Topological progression in proliferating epithelia is driven by a unique variation in polygon distribution. PLoS One *8*, e79227.

Sanchez-Gutierrez, D., Tozluoglu, M., Barry, J.D., Pascual, A., Mao, Y., and Escudero, L.M. (2016). Fundamental physical cellular constraints drive self-organization of tissues. EMBO J *35*, 77–88.

Sánchez-Gutiérrez, D., Tozluoglu, M., Barry, J.D., Pascual, A., Mao, Y., Escudero, L.M., Aegerter-Wilmsen, T., Smith, A., Christen, A., Aegerter, C., et al. (2016). Fundamental physical cellular constraints drive self-organization of tissues. EMBO J. *35*, 77–88.

Sánchez-Gutiérrez, D., Sáez, A., Gómez-Gálvez, P., Paradas, C., Escudero, L.M., Honda, H., Chichilnisky, E.J., Hayashi, T., Carthew, R.W., Classen, A.K., et al. (2017). Rules of tissue packing involving different cell types: human muscle organization. Sci. Rep. *7*, 40444.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682.

Tanaka, S., Sichau, D., and Iber, D. (2015). LBIBCell: a cell-based simulation environment for morphogenetic problems. Bioinformatics *31*, 2340–2347.

Tetley, R.J., and Mao, Y. (2018). The same but different : cell intercalation as a driver of tissue deformation and fluidity.

Yamashita, S., and Michiue, T. (2014). Quantitative analysis of cell
arrangement indicates early differentiation of the neural region during
Xenopus gastrulation. J. Theor. Biol. *346*, 1–7.

Yoon, C.H., Lee, J., Jongeward, G.D., and Sternberg, P.W. (1995).
Similarity of sli-1, a regulator of vulval development in C. elegans, to the
mammalian proto-oncogene c-cbl. Science (80-.). *269*, 1102–1105.

Zallen, J.A., and Zallen, R. (2004). Cell-pattern disordering during
convergent extension in Drosophila. J. Phys. Condens. Matter. *16*, S5073–
S5080.

Zartman, J., Restrepo, S., and Basler, K. (2013). A high-throughput template
for optimizing *Drosophila* organ culture with response-surface methods.
Development *140*, 2848–2848.

13
14

Figure 1. Graphlets, cellular motifs and characterization of epithelial organization. A) A representation of the cellular motifs that correspond to graphlets of up to five nodes. There are 29 motifs corresponding to 26 different graphlets (**Fig. S1**). Note that one graphlet can represent two cellular motifs (G8, G23 and G26). Mauve motifs form the Mo17 set. Prussian Blue motifs stands for the set Mo26. In the first row are the motifs that account for the organization of groups of up to 4 cells (Mo10). Therefore, Mo7 set is formed by the mauve coloured graphlets at the first row. **B)** Polygon distribution comparison of images from: Voronoi diagram 1 (black bar); Eye (orange), *Drosophila* eye disc: 3 samples; Voronoi diagram 4 (grey); dWL (green), *Drosophila* larva wing disc: 15 samples; Voronoi diagram 5 (light grey), dWP (red), *Drosophila* prepupal wing imaginal disc epithelium: 16 samples. Data shown refer to the mean ± SEM. Diagram 1, 4 and 5: 20 replicates. **C)** GDD value calculation (Mo17) between natural images and Voronoi diagrams with similar polygon distribution. The data shown are the mean of the GDD between each pair of images.

Figure 2. Epithelial organization of biological tissues with respect the CVTn. A-C) Tessellations with the corresponding graph of cell-to-cell contacts for a perfect hexagonal arrangement (**A**) a Voronoi Diagram 1 (**B**) and a Voronoi Diagram 5 (**C**) from a CVTn. These tessellations represent the diagrams used as reference to calculate the GDDH, GDDRV and GDDV5 respectively. The light blue edges in these panels represent the cellular connectivity network. The colourful nodes mark the valid cells that were involved in the cellular motifs to measure graphlets presence. The dark blue and green nodes are the 4' valid cells (cells that did not have a no valid cell within a distance of four cells connexions), which were used to calculate the graphlets for Mo10 and Mo7. The green nodes are the 5' valid cells (cells that did not have a no valid cell within a distance of five cells connexions) that were used to quantify the graphlets for Mo29 and Mo17. Cells without nodes were no valid cells for graphlet calculation. **D)** Representative images from the natural tessellations. **E)** Plots showing the different combinations of the values for Mo17 of GDDH, GDDRV and GDDV5 with the percentage of hexagons. The diagrams of the CVTn path from the iteration 1 until the iteration 700 are represented as a greyscale beginning in black and reducing its darkness with the increase of the iterations (from 1 to 20, from 30 to 100 by stepwise of 10 and from 100 to 700 by stepwise of 100). **F-G)** Charts representing the comparisons GDDH against GDDRV, and GDDRV against GDDV5, respectively. The CVTn path in both scatter plots, is formed by the diagrams with numbers between 1 and 100, in a greyscale as in (**E**). The natural tessellations are: dMWP (violet), *Drosophila* mutant wing disc: 3 samples; cNT (light blue), chicken neural tube: 16 samples; Eye, dWL and dWP are the same replicates than **Fig.1** and preserve their colour codes. Circumferences are individual values, circles are the average value obtained from the individual samples from each category.

Figure 3. Comparison of different simulations and mutants with the CVTn. A-C) Representative images for non-proliferative simulations. Control with homogeneous parameters (**A**). The 'squared' simulations are similar to control, but a ten percent of cells (randomly chosen) have a reduced line tension (**B**). The 'elongated' simulations have a ten percent of cells (randomly chosen) with its line tension and ideal area reduced, and the another 90% of cells have the same parameters than control simulations (**C**). **D)** Cell arrangement resulting from the control simulation that includes cell proliferation. **E-F)** Diagrams resulting from a vertex model simulation with an increase of the ideal area value, with respect the control, in some cells. Case III and Case IV slightly differ in the line-tension parameter conditions (see **Experimental Procedures**). **G)** Plots showing the values of GDDRV vs GDDV5 and the percentage of hexagons vs GDDV5 (Mo17) for CVTn, dMWP, Eye, cNT, dWL, dWP; Proliferative Control (20 replicates, carnation pink), Case III (17 replicates, hot pink) and Case IV (15 replicates, purple); Non-proliferative control (20 replicates, blue bell), Squared (20 replicates, azure blue) and Elongated simulations (20 replicates, cornflower Blue). The diagrams of the CVTn path from the iteration 1 until the iteration 100 are represented as a greyscale beginning in black and reducing its darkness with the increase of the iterations; dMWP, Eye, cNT, dWL and dWP have the same replicates and colour codes than Fig.2; circumferences are individual values, circles are the average value obtained from the individual samples from each category.

Figure 4. EpiGraph can evaluate the changes in epithelial organization of a morphogenetic process. A) Time points selected from "Movie S1" from Blankenship and cols., 2006 article (url: https://ars.els-cdn.com/content/image/1-s2.0-S153458070600400X-mmc2.mov) (Blankenship et al., 2006). We do not show these panels in this version of the manuscript since we have not been able to obtain the editorial permission yet. B-C) Captures from the 3D visualization tool of EpiGraph showing the position of CVTn (1-700), Eye, cNT, dWL, dWP, and the eight frames showed in (A). GDDH, GDDRV and percentage of hexagons (Mo17) are the axes in (B). GDDH, GDDRV and GDDV5 (Mo17) are the axes in (C).

Figure 5. The rigidity/fluidity of a tissue can be assessed using EpiGraph. A) Initial and final frames of two simulations with different settings: a rigid and a soft tissue. **B)** Epigraph's 3D plot with GDDRV, GDDH and GDDV5 axes, showing the soft simulation tissue in green dots and the rigid simulation as orange dots. Each simulation is represented in 13 frames (see **Experimental Procedures**). **C-D)** Representative examples of segmented images from the third instar *Drosophila* imaginal disc in different conditions: Wild Type (**C**); a solid mutant, *Mbs-RNAi* in (**D**). **E)** Plot comparing the fluidity and organization of the tissues in (**C-E**). CVTn (until diagram 30) displayed in greyscale. Dots in scales of blue represent the WT condition: wing disc 1, aquamarine; wing disc 2, light blue; wing disc 3, dark blue. Represented with points in tones of orange-red, *Mbs-RNAi*: sample 1, salmon colour; sample 2, orange; sample 3, red. **F-G)** Boxplot of the intercalation rate (**F**), which is the number of T1 transitions per cell per hour, and the fourfold vertices found per cell (**G**) (note that no fivefold vertices, or beyond, was found on any sample). Boxes stand for the data inside the upper and lower quartiles, while the vertical dashed lines (whiskers) indicate the variability outside them. Mean (dashed line) and median (thick line) of each condition is represented inside each box. The actual values are also presented as circles (and the outlier values as circumferences) with its correspondent colour. In addition, statistical significance, by means of a Kolmogorov-Smirnov test, is shown in the top of both panels (**F**: '******' $p<0.01$, **G**: '*****' $p<0.05$). Each condition has 3 samples (different colour tone), and the numeric tag represents its frame. In WT have been taken 6 frames per sample in periods of 6 minutes. In the case of *Mbs-RNAi* were taken 3 frames per sample with time lapse of 15 minutes. All the conditions have been tracked for 30 minutes (see **Experimental procedures**).