Acute fluoxetine differently affects aggressive display in zebrafish phenotypes

Hellen Pereira Barbosa[1], Monica Gomes Lima[1,2], Caio Maximino[2,3,4*]

[1] Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff", Departamento de

Morfologia e Ciências Fisiológicas, Campus VIII/Marabá, Universidade do Estado do Pará,

Marabá/PA, Brazil

[2] Programa de Pós-Graduação em Biodiversidade e Biotecnologia – Rede BIONORTE

[3] Programa de Pós-Graduação em Neurociências e Comportamento, Universidade Federal do Pará,

Belém/PA, Brazil

[4] Laboratório de Neurociências e Comportamento, Instituto de Estudos em Saúde e Biológicas,

Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

* Corresponding author

Caio Maximino

Laboratório de Neurociências e Comportamento – Instituto de Estudos em Saúde e Biológicas,

Universidade Federal do Sul e Sudeste do Pará, Unidade III

Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

Abstract

Zebrafish has been introduced as a model organism in behavioral neuroscience and biological psychiatry, increasing the breadth of findings using fish to study the neurobiology of aggression. Phenotypic differences between leopard and longfin zebrafish were exploited in order to elucidate the role of phasic serotonin in aggressive displays on this species. The present study revealed differences in aggressive display between leopard and longfin zebrafish, and a discrepant effect of acute fluoxetine in both populations. In mirror-induced aggression, leopard animals showed higher display latencies than longfin, as well as lower display duration and frequency (Experiment 1). Moreover, 2.5 mg/kg fluoxetine decreased the duration and frequency of display in longfin, but not leopard; and 5 mg/kg fluoxetine increased display frequency in leopard, but not longfin (Experiment 2). It is suggested that zebrafish from the longfin phenotype show more aggressive motivation and readiness in the mirror-induced aggression test that leopard, and that acute fluoxetine increases aggression in leopard and decreased it in longfin zebrafish.

Keywords: Zebrafish; Aggressive display; Serotonin; Fluoxetine; Phenotypic differences

1. Introduction

The biological comprehension of factors underlying aggression is still limited (Miczek et al., 2007), even though a range of mental disorders present aggression as a symptom (Krakowski, Volavka, & Brizer, 1986). Despite the paucity of neurobiological data on aggression, a role for monoamines has been proposed (Miczek et al., 2007; Takahashi, Quadros, Almeida, & Miczek, 2011). In most animal models, acutely increasing the serotonergic transmission inhibits aggressive behavior (Takahashi et al., 2011); a metanalysis of preclinical studies demonstrated that, across species, pharmacologically increasing 5-HT levels inhibit aggression (Carrillo, Ricci, Coppersmith, & Melloni Jr., 2009).

While this observation appears to hold for most studies, some controversies and gaps appear

in the literature, especially in basal vertebrates such as fish. For example, in the metanalysis by

Carrillo et al. (2009), 5-HT decreased aggression in wrasses and trouts, but not in the Siamese

fighting fish. Zebrafish has been introduced as a model organism in behavioral neuroscience and

biological psychiatry (W. Norton & Bally-Cuif, 2010; Stewart et al., 2015), increasing the relevance

of findings using fish to study the neurobiology of aggression.

A role for 5-HT in zebrafish aggressive behavior has been suggested by neurochemical

studies. After eliciting an aggressive display towards a mirror (mirror-induced aggression, MIA), 5-

HT levels were increased in the telencephalon, while 5-HIAA was increased in the optic tectum of

zebrafish (Teles, Dahlbom, Winberg, & Oliveira, 2013). Male and female zebrafish respond to

agonistic encounters in a similar fashion; nonetheless, males present higher 5-HT turnover in the

forebrain in relation to females, suggesting that aggressive bouts could be more stressful to males

than females (Dahlbom, Backström, Lundstedt-Enkel, & Winberg, 2012). Filby et al.

(2010) demonstrated that dominant males show an overexpression of genes associated with the

serotonergic system in the hypothalamus, including *tph1b* and *htr1aa*, while females showed

overexpression of *tph2*, *htr1aa*, *slc6a4a*, and *mao* in the hypothalamus and *tph1a* and *tph2* in the

telencephalon.

While these results suggest that aggressive behavior can be linked to differences in the

serotonergic system – especially in the context of dominance hierarchies –, a causal relationship is

more tenuous. Filby et al. (2010) treated dominant male zebrafish with fluoxetine (3 or 4.5 µg/L),

without effects on aggressive behavior in a dyadic encounter; however, a similar concentration (3

µg/L) decreased aggressive displays in the MIA (W. H. J. Norton et al., 2011). Using a much higher

concentration (5 mg/L), Theodoridi et al. (2017) were able to inhibit attacks and chasing behavior in

dominant animals in dyads. The lack of consistency could be due to dosing, behavioral paradigms

(e.g., MIA vs. dyadic encounters), or other variables.

Recent studies also showed that 5-HT levels are lower in zebrafish with the leopard

phenotype than in animals with the longfin phenotype, an alteration that is accompanied by

increased monoamine oxidase activity (Maximino, Puty, Oliveira, & Herculano, 2013). These

neurochemical differences were accompanied by increased anxiety-like behavior that is rescued by

fluoxetine treatment (Maximino, Puty, Oliveira, et al., 2013). Interestingly, in longfin animals

fluoxetine *increases* anxiety, and 5-HT levels are negatively correlated with anxiety-like behavior; it

is possible that embryological differences in the serotonergic system produce opposite adult

phenotypes.

These phenotypic differences are exploited in the present work to clarify the role of phasic

serotonin on aggressive displays in zebrafish. We hypothesized that the hyposerotonergic phenotype

of leopard zebrafish would produce increased aggressive behavior, and that fluoxetine would rescue

this phenotype. The experimental evidence produced in the present work contradicted this

hypothesis, since longfin were shown to display more aggressive motivation and readiness in the

mirror-induced aggression test than leopard zebrafish, and since acute fluoxetine increased

aggression in leopard animals and decreased it in longfin zebrafish. This manuscript is a complete

report of all the studies performed to test the effect of skin phenotype and fluoxetine on aggressive

behavior. We report how we determined our sample size, all data exclusions (if any), all

manipulations, and all measures in the study.

2. Methods

2.1. Animals, housing, and baseline characteristics

Outbred populations were used due to their increased genetic variability, decreasing the effects of

random genetic drift which could lead to the development of uniquely heritable traits (Parra, Adrian

Jr, & Gerlai, 2009; Speedie & Gerlai, 2008). Thus, the animals used in the experiments are expected

to better represent the natural populations in the wild. Adult zebrafish from the wildtype strain

(longfin and leopard phenotypes) were used in this experiment. Animals were bought from a

commercial vendor, and arrived in the laboratory with an approximate age of 3 months (standard

length = 13.2 ± 1.4 mm), and were quarantined for two weeks; the experiment begun when animals

had an approximate age of 4 months (standard length = 23.0 ± 3.2 mm). Animals were kept in

mixed-sex tanks during acclimation, with an approximate ratio of 50 male:50 female. The breeder

was licensed for aquaculture under Ibama's (Instituto Brasileiro do Meio Ambiente e dos Recursos

Naturais Renováveis) Resolution 95/1993. Animals were group-housed in 40 L tanks, with a

maximum density of 25 fish per tank, for at least 2 weeks before experiments begun. Tanks were

filled with non-chlorinated water at room temperature (28 °C) and a pH of 7.0-8.0. Lighting was

provided by fluorescent lamps in a cycle of 14-10 hours (LD), according to standards of care for

zebrafish (Lawrence, 2007). Water quality parameters were as follows: pH 7.0-8.0; hardness 100-

150 mg/L $CaCO_3$; dissolved oxygen 7.5-8.0 mg/L; ammonia and nitrite < 0.001 ppm. All

manipulations minimized their potential suffering of animals, and followed Brazilian legislation

(Conselho Nacional de Controle de Experimentação Animal - CONCEA, 2017). Animals were used

for only one experiment and in a single behavioral test, to reduce interference from apparatus

exposure.

2.2. Mirror-induced aggression

The protocol for mirror-induced aggression was adapted from Norton et al. (2011). Animals were

individually transferred to a tank (15 x 10 x 30 cm) containing 1 L of holding tank water. The tank

was lit from above with white light (445 ± 56 lumens). Animals were allowed to acclimate to the

tank for 5 min; after that, a mirror was positioned on the outside of the tank (on the narrower side),

in an angle of 22.5º. All steps of the experiment were executed under constant Gaussian white noise

(55 ± 2.5 dB above the tank). Behavior was recorded with a digital video camera (Samsung ES68)

positioned in the wider side of the tank, and analyzed by observers blind to treatment using the

5   5

event-recording software X-Plo-Rat (https://github.com/lanec-unifesspa/x-plo-rat). The following

endpoints were analyzed: time in the square nearest to the mirror (s); frequency (N) and duration (s)

of aggressive display; total number of squares crossed (N). Aggressive display was defined as a

swimming posture with erect dorsal, caudal, pectoral, and anal fins (Gerlai, Lahav, Guo, &

Rosenthal, 2000).

2.3. Data availability

Datasets and scripts for all analyses are available from https://github.com/lanec-unifesspa/5HT-

aggression (doi: 10.5281/zenodo.1006701).

2.4. Experiment 1

Sample size calculation and groups. Sample sizes were calculated on results regarding the effects

of fluoxetine on total time in broadside display in *Betta splendens*, reported by Lynn et al. (2007);

as a result, the closest endpoint (time on display) was chosen as the primary endpoint, and

calculations for sample sizes are valid only for that endpoint. Calculations were based on Rosner's

(2016) method for comparing two means, and assumed $\alpha = 0.05$ and $\beta = 0.2$ on a two-tailed

analysis. Based on these calculations, 15 animals were used in each group in Experiment 1. Animals

were derived from the stock population described in section 2.1, and displayed either the longfin

(Group LOF) or leopard (Group LEO) phenotypes. Animals were randomly drawn from the tank

immediately before testing, and the order with which phenotypes were tested was randomized *via*

generation of random numbers using the randomization tool in http://www.randomization.com/.

Blinding was not possible, due to the obvious differences in skin phenotype.

Experimental design and statistical analysis. Animals were allocated to each group according to

phenotype. Immediately after being drawn from the tank, animals were individually transported to

the experiment room, and left undisturbed for 30 min. After this interval, animals were exposed to

the MIA test, described above. Differences between groups were analyzed using Approximative

Two-Sample Fisher-Pitman Permutation Tests 10,000 Monte-Carlo re-samplings, using the R

package 'coin' (Hothorn, Hornik, van de Wiel, & Zeileis, 2006). The data analyst was blinded to

phenotype by using coding to reflect treatments in the resulting datasets; after analysis, data was

unblinded. Data are presented using individual dot plots combined with boxplots.

2.5. Experiment 2

Sample size calculation and groups. In the absence of similar experiments in the literature, sample

sizes were calculated based on the assumption of fixed effect sizes for both the phenotype and the

dose factors, with a projected effect size of 0.4; calculations were made using the R package 'pwr2'

(Lu, Liu, & Koestler, 2017). Based on these calculations, 10 animals were used in each group in

Experiment 1. Animals were derived from the stock population described in section 2.1, and

displayed either the longfin (Group LOF) or leopard (Group LEO) phenotypes. Animals were

randomly drawn from the tank immediately before testing, and the order with which phenotypes

were tested was randomized *via* generation of random numbers using the randomization tool in

http://www.randomization.com/. Blinding for phenotype was not possible, due to the obvious

differences in skin phenotype. Animals from each phenotype were randomly allocated to treatment

(vehicle or either fluoxetine dose) *via* generation of random numbers using the randomization tool

in http://www.randomization.com/.

Drug treatments. Fluoxetine (FLX) was bought from EMS, dissolved in Cortland's salt solution

(Wolf, 1963), and injected injected intraperitoneally in cold-anesthetised animals (Kinkel, Eames,

Philipson, & Prince, 2010). FLX doses (2.5 and 5.0 mg/kg) were based on the demonstration of

effect on the light/dark test on longfin (Maximino, Puty, Benzecry, et al., 2013) and leopard

(Maximino, Puty, Oliveira, et al., 2013) zebrafish. Experimenters were blinded to treatment by

coding drug vials.

Experimental design and statistical analysis. Animals were allocated to each group according to

phenotype. Immediately after being drawn from the tank, animals were individually transported to

the experiment room, injected with vehicle or drug, and left undisturbed for 30 min. After this

interval, animals were exposed to the MIA test, described above. Differences between groups were

analyzed using two-way analyses of variance with robust estimators on Huber's M-estimators,

using the R package 'rcompanion' (Mangiafico, 2017). P-values were adjusted for the false

discovery rate. The data analyst was blinded to phenotype by using coding to reflect treatments in

the resulting datasets; after analysis, data was unblinded. Data are presented using individual dot

plots combined with boxplots.

3. Results

3.1 Experiment 1

LEO zebrafish showed longer latencies to display than LOF animals ($Z = 3.3925$, $p = 0.0005$;

Figure 1A), as well as shorter display durations ($Z = -2.5659$, $p = < 2.2e-16$; Figure 1B) and

frequency ($Z = -2.7073$, $p = 0.003$; Figure 1C), and time spent near the mirror ($Z = -3.2284$, $p = <$

2.2e-16; Figure 1D). No differences were found in total locomotion ($Z = -0.69887$, $p = 0.4965$;

Figure 1E).

3.2. Experiment 2

No main effects of phenotype ($p = 0.5736$) or FLX dose ($p = 0.5044$) were found for latency, but a

significant interaction was found ($p = 0.0412$); nonetheless, post-hoc tests did not detect any

8   8

differences between groups (Figure 2A). Main effects of phenotype (p = 0.0132) and FLX dose (p =

0.0062), as well as an interaction effect (p = 0.009), were found for display duration (Figure 2B).

Post-hoc tests suggested that FLX (2.5 mg/kg) decreased display duration on LOF, but not LEO (p

= 0.032 vs. 0 mg/kg). Similarly, main effects of phenotype (p = 0.0458) and FLX dose (p = 0.004),

as well as an interaction effect (p < 0.0001), were found for display frequency (Figure 2C); post-hoc

tests suggested that FLX (5 mg/kg) increased display frequency in LEO, but not LOF animals (p =

0.004 vs. 0 mg/kg). No main effects of phenotype (p = 0.2692) were found for time near mirror, but

a main effect of FLX dose (p = 0.0004) and an interaction effect (p < 0.0001); post-hoc tests

suggested a inverted-U-shaped curve for FLX-treated LOF (0 vs. 2.5 mg/kg: p = 0.0012; 0 vs. 5.0

mg/kg: p = 0.0167), while a monotonic increase for FLX-treated LEO (0 vs. 2.5 mg/kg: p = 0.0496;

0 vs. 5.0 mg/kg: p < 0.0001) (Figure 2D). No main or interaction effects were found for total

locomotion (Figure 2E).

4. Discussion

The present work demonstrated that zebrafish with the leopard skin phenotype show less aggressive

readiness and less aggression in relation to longfin animals. Moreover, a different pattern of

fluoxetine effects was observed, with fluoxetine decreasing aggressive display (but not readiness) in

longfin animals and increasing it in leopard animals. Evidence for dose-dependence was also

observed.

Serotonin (5-HT) has long been implicated in the neurobiological mechanisms of aggressive

behavior (Miczek et al., 2007; Summers & Winberg, 2006; Takahashi et al., 2011). In a metanalysis

of preclinical studies, Carrillo et al. (2009) demonstrated that, across species, pharmacologically

increasing 5-HT levels inhibit aggression. Interestingly, in their metanalysis a species-specific effect

was found in fish, with 5-HT decreasing aggression in wrasses and trouts, but not in the Siamese

fighting fish (Carrillo et al., 2009). The present study examined only aggressive displays, which was

elicited in the mirror-induced aggression test; as a result, dominance hierarchies were not induced.

Similar results were observed by Norton et al. (2011), which observed reduced aggressive displays

in zebrafish treated with fluoxetine. When zebrafish are allowed to form dominance hierarchies,

fluoxetine either produces no effect (Filby et al., 2010) or reduces aggression in dominant males,

but not in subordinates (Theodoridi et al., 2017).

Behavioral differences between leopard and longfin phenotypes were observed in zebrafish before

(Canzian, Fontana, Quadros, & Rosemberg, 2017; Maximino, Puty, Oliveira, et al., 2013; Quadros

et al., 2016; https://doi.org/10.1101/055657). Of special relevance is the observation that leopard

zebrafish present increased brain monoamine oxidase (MAO) activity that is associated with lower

serotonin levels and higher turnover of 5-HT in the brain (Maximino, Puty, Oliveira, et al., 2013).

This hyposerotonergic profile was also associated with increased anxiety-like behavior that was

rescued by fluoxetine treatment (Maximino, Puty, Oliveira, et al., 2013).

The results from the Carrillo et al. (2009) metanalysis suggested a inhibitory role for phasic

serotonin (i.e., 5-HT released by either the aggressive act itself, or by pharmacological

manipulations such as fluoxetine); a role for tonic 5-HT, in *Betta splendens*, was discarded, because

neither the 5-HT synthesis inhibitor para-chlorophenylalanine nor the 5-HT precursor L-tryptophan

changed display behavior (Clotfelter, O'Hare, McNitt, Carpenter, & Summers, 2007). If that was

also true for zebrafish, differences in aggressive display between leopard and longfin would not be

expected, given that these phenotypes differ in serotonergic tone (Maximino, Puty, Oliveira, et al.,

2013). While these differences were observed in the present work, they occur in the opposite

direction from what would be predicted from 5-HTergic tone alone (i.e., we should expect leopard

zebrafish to be more aggressive). It is possible that a developmental effect is responsible for these

discrepancies.

These results are also reminiscent of what is observed in different populations of *Astyanax*

mexicanus. In that species, different populations occupy different niches, and surface-dwelling

populations are much more aggressive than cave-dwelling populations (Rétaux & Elipot, 2013).

These differences are related to the density of serotonergic neurons in the hypothalamus, with

cavefish showing a higher number of 5-HT neurons in that region (Elipot, Hinaux, Callebert, &

Rétaux, 2013). Moreover, a mutation in the *mao* gene was found in cavefish that led to an

hyperserotonergic phenotype (Elipot et al., 2014). Treating surface fish with fluoxetine decreases

aggression, while in cavefish the drug slightly increases it (Elipot et al., 2013). In the present paper,

however, treatment with fluoxetine increased aggression in leopard animals (which show an

hyposerotonergic profile in relation to longfin animals; Maximino, Puty, Oliveira, et al., 2013) and

decreased it in longfin zebrafish. These differences might be due to the origin of serotonin, since, in

Astyanax mexicanus populations, raphe 5-HT levels are unchanged, while hypothalamic 5-HT is

increased (Elipot et al., 2013); further experiments are needed to untangle this hypothesis.

In conclusion, the present experiments revealed differences in aggressive behavior between leopard

and longfin zebrafish, and a discrepant effect of fluoxetine on both populations. These results are

relevant to understand the role of tonic and phasic serotonin neurotransmission on aggressive

behavior in preclinical models, and might contribute to a better appreciation of the complex roles of

this monoamine in controlling vertebrate aggression.

Acknowledgments

Data packages and statistical analysis scripts for this article can be found at

https://dx.doi.org/10.5281/zenodo.1006701.

References

Canzian, J., Fontana, B. D., Quadros, V. A., & Rosemberg, D. B. (2017). Conspecific alarm substance differently alters group behavior of zebrafish populations: Putative involvement of cholinergic and purinergic signaling in anxiety- and fear-like responses. *Behavioural Brain Research*, *320*, 255–263. https://doi.org/10.1016/j.bbr.2016.12.018

Carrillo, M., Ricci, L. A., Coppersmith, G. A., & Melloni Jr., R. H. (2009). The effect of increased serotonergic neurotransmission on aggression: A critical meta-analytical review of preclinical studies. *Psychopharmacology*, *205*, 349–368. https://doi.org/10.1007/s00213-009-1543-2

Clotfelter, E. D., O'Hare, E. P., McNitt, M. M., Carpenter, R. E., & Summers, C. H. (2007). Serotonin decreases aggression via 5-HT1A receptors in the fighting fish *Betta splendens*. *Pharmacology, Biochemistry & Behavior*, *87*, 222–231. https://doi.org/10.1016/j.pbb.2007.04.018

Conselho Nacional de Controle de Experimentação Animal - CONCEA. Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científica, Pub. L. No. Resolução Normativa 30/2016 (2017). Brasil.

Dahlbom, S. J., Backström, T., Lundstedt-Enkel, K., & Winberg, S. (2012). Aggression and monoamines: Effects of sex and social rank in zebrafish (Danio rerio). *Behavioural Brain Research*, *228*, 333–338. https://doi.org/10.1016/j.bbr.2011.12.011

Elipot, Y., Hinaux, H., Callebert, J., Launay, J.-M., Blin, M., & Rétaux, S. (2014). A mutation in the enzyme monoamine oxidase explains part of the *Astyanax* cavefish behavioural syndrome. *Nature Communications*, *5*, 3647. https://doi.org/10.1038/ncomms4647

Elipot, Y., Hinaux, H., Callebert, J., & Rétaux, S. (2013). Evolutionary shift from fighting to foraging in blind cavefish through changes in the serotonin network. *Current Biology*, *23*, 1–10. https://doi.org/10.1016/j.cub.2012.10.044

Filby, A. L., Paull, G. C., Hickmore, T. F., & Tyler, C. R. (2010). Unravelling the neurophysiological basis of aggression in a fish model. *BMC Genomics*, *11*, 498. https://doi.org/10.1186/1471-2164-11-498

Gerlai, R., Lahav, M., Guo, S., & Rosenthal, A. (2000). Drinks like a fish: Zebra fish (*Danio rerio*) as a behavior genetic model to study alcohol effects. *Pharmacology, Biochemistry and Behavior*, *67*, 773–782.

Hothorn, T., Hornik, K., van de Wiel, M. A., & Zeileis, A. (2006). A Lego system for conditional inference. *The American Statician*, *60*, 257–263. https://doi.org/10.1198/000313006X118430

Kinkel, M. D., Eames, S. C., Philipson, L. H., & Prince, V. E. (2010). Intraperitoneal injection into adult zebrafish. *Journal of Visualized Experiments*, *42*, 2126. https://doi.org/10.3791/2126

Krakowski, M., Volavka, J., & Brizer, D. (1986). Psychopathology and violence: A review of literature. *Comprehensive Psychiatry*, *27*, 131–148. https://doi.org/10.1016/0010-440X(86)90022-2

Lawrence, C. (2007). The husbandry of zebrafish (*Danio rerio*): A review. *Aquaculture*, *269*, 1–20. https://doi.org/10.1016/j.aquaculture.2007.04.077

Lu, P., Liu, J., & Koestler, D. (2017). pwr2: Power and Sample Size Analysis for One-way and Two-way ANOVA Models. Retrieved from https://cran.r-project.org/package=pwr2

Lynn, S. E., Egar, J. M., Walker, B. G., Sperry, T. S., & Ramenofsky, M. (2007). Fish on Prozac: A simple, noninvasive physiology laboratory investigating the mechanisms of aggressive behavior in Betta splendens. *Advances in Physiology Education*, *31*, 358–363. https://doi.org/10.1152/advan.00024.2007

Mangiafico, S. (2017). rcompanion: Functions to Support Extension Education Program Evaluation. Retrieved from https://cran.r-project.org/package=rcompanion

Maximino, C., Puty, B., Benzecry, R., Araujo, J., Lima, M. G., Batista, E. de J. O., … Herculano, A. M. (2013). Role of serotonin in zebrafish (*Danio rerio*) anxiety: Relationship with serotonin levels and effect of buspirone, WAY 100635, SB 224289, fluoxetine and para-chlorophenylalanine (pCPA) in two behavioral models. *Neuropharmacology*, *71*, 83–97. https://doi.org/10.1016/j.neuropharm.2013.03.006

Maximino, C., Puty, B., Oliveira, K. R. M., & Herculano, A. M. (2013). Behavioral and neurochemical changes in the zebrafish leopard strain. *Genes, Brain and Behavior*, *12*, 576–582. https://doi.org/10.1111/gbb.12047

Miczek, K. A., de Almeida, R. M. M., Kravitz, E. a, Rissman, E. F., de Boer, S. F., & Raine, A. (2007). Neurobiology of escalated aggression and violence. *Journal of Neuroscience*, *27*, 11803–11806. https://doi.org/10.1523/JNEUROSCI.3500-07.2007

Norton, W., & Bally-Cuif, L. (2010). Adult zebrafish as a model organism for behavioural genetics. *BMC Neuroscience*, *11*, 90. https://doi.org/10.1186/1471-2202-11-90

Norton, W. H. J., Stumpenhorst, K., Faus-Kessler, T., Folchert, A., Rohner, N., Harris, M. P., … Bally-Cuif, L. (2011). Modulation of *fgfr1a* signaling in zebrafish reveals a genetic basis for the aggression–boldness syndrome. *Journal of Neuroscience*, *31*, 13796–13807. https://doi.org/10.1523/JNEUROSCI.2892-11.2011

Parra, K. V, Adrian Jr, J. C., & Gerlai, R. (2009). The synthetic substance hypoxanthine 3-N-oxide elicits alarm reactions in zebrafish (*Danio rerio*). *Behavioural Brain Research*, *205*, 336–341. https://doi.org/10.1016/j.bbr.2009.06.037

Quadros, V. A., Silveira, A., Giuliani, G. S., Didonet, F., SIlveira, A. S., Nunes, M. E., … Rosemberg, D. B. (2016). Strain- and context-dependent behavioural responses of acute alarm substance exposure in zebrafish. *Behavioural Processes*, *122*, 1–11. https://doi.org/10.1016/j.beproc.2015.10.014

Rétaux, S., & Elipot, Y. (2013). Feed or fight. A behavioral shift in blind cavefish. *Communicative & Integrative Biology*, *6*, e23166. https://doi.org/10.4161/cib.23166

Rosner, B. (2016). *Fundamentals of biostatistics*. Cengage Learning.

Speedie, N., & Gerlai, R. (2008). Alarm substance induced behavioral responses in zebrafish (*Danio rerio*). *Behavioural Brain Research*, *188*, 168–177. https://doi.org/10.1016/j.bbr.2007.10.031

Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H., Gerlai, R., & Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular Psychiatry*, *20*, 2–17. https://doi.org/10.1038/mp.2014.128

Summers, C. H., & Winberg, S. (2006). Interactions between the neural regulation of stress and aggression. *Journal of Experimental Biology*, *209*, 4581–4589. https://doi.org/10.1242/jeb.02565

Takahashi, A., Quadros, I. M., Almeida, R. M. M. de, & Miczek, K. A. (2011). Brain serotonin receptors and transporters: Initiation vs . termination of escalated aggression. *Psychopharmacology*, *213*, 183–212. https://doi.org/10.1007/s00213-010-2000-y

Teles, M. C., Dahlbom, S. J., Winberg, S., & Oliveira, R. F. (2013). Social modulation of brain monoamine levels in zebrafish. *Behavioural Brain Research*. https://doi.org/10.1016/j.bbr.2013.07.012

Theodoridi, A., Tsalafouta, A., & Pavlidis, M. (2017). Acute exposure to fluoxetine alters aggressive behavior of zebrafish and expression of genes involved in serotonergic system regulation. *Frontiers in Neuroscience*, *11*, Article 223. https://doi.org/10.3389/fnins.2017.00223

Wolf, K. (1963). Physiological salines for freshwater teleosts. *Progress in Fish-Culture, 25*, 135–140.

Figure captions

Figure 1 – Phenotype differences in (A) latency to display, in s; (B) display duration, in s; (C),

display frequency; (D), time spent near the mirror, in s; and (E), total locomotion. Boxplots

represent median and interquartile range, with Tukey whiskers.

Figure 2 – Effects of fluoxetine on the aggressive display of longfin (LOF, blue) and leopard (LEO,

coral) zebrafish. (A) latency to display, in s; (B) display duration, in s; (C), display frequency; (D),

time spent near the mirror, in s; and (E), total locomotion. Boxplots represent median and

interquartile range, with Tukey whiskers. Dots joined by lines represent means.