Running head: FLUOXETINE AND ZEBRAFISH AGGRESSION

Acute fluoxetine differently affects aggressive display in zebrafish phenotypes

Hellen Pereira Barbosa[1], Monica Gomes Lima[1,2], Caio Maximino[2,3,4]

[1] Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff",

Departamento de Morfologia e Ciências Fisiológicas, Campus VIII/Marabá, Universidade do

Estado do Pará, Marabá/PA, Brazil

[2] Programa de Pós-Graduação em Biodiversidade e Biotecnologia – Rede BIONORTE

[3] Programa de Pós-Graduação em Neurociências e Comportamento, Universidade Federal do

Pará, Belém/PA, Brazil

[4] Laboratório de Neurociências e Comportamento, Instituto de Estudos em Saúde e

Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

Acute fluoxetine differently affects aggressive display in zebrafish phenotypes

Hellen Pereira Barbosa[1], Monica Gomes Lima[1,2], Caio Maximino[2,3,4*]

[1] Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff",

Departamento de Morfologia e Ciências Fisiológicas, Campus VIII/Marabá, Universidade do

Estado do Pará, Marabá/PA, Brazil

[2] Programa de Pós-Graduação em Biodiversidade e Biotecnologia – Rede BIONORTE

[3] Programa de Pós-Graduação em Neurociências e Comportamento, Universidade Federal do

Pará, Belém/PA, Brazil

[4] Laboratório de Neurociências e Comportamento, Instituto de Estudos em Saúde e

Biológicas, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

* Corresponding author

Caio Maximino

Laboratório de Neurociências e Comportamento – Instituto de Estudos em Saúde e

Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III

Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

Abstract

Zebrafish have been introduced as a model organism in behavioral neuroscience and

biological psychiatry, increasing the breadth of findings using fish to study the neurobiology

of aggression. Phenotypic differences between leopard and longfin zebrafish were exploited

in order to elucidate the role of phasic serotonin in aggressive displays on this species. The

present study revealed differences in aggressive display between leopard and longfin

zebrafish, and a discrepant effect of acute fluoxetine in both populations. In mirror-induced

aggression, leopard animals showed higher display latencies than longfin, as well as lower

display duration and frequency (Experiment 1). Moreover, 2.5 mg/kg fluoxetine decreased

the duration and frequency of display in longfin, but not leopard; and 5 mg/kg fluoxetine

increased display frequency in leopard, but not longfin (Experiment 2). It is suggested that

zebrafish from the longfin phenotype show more aggressive motivation and readiness in the

mirror-induced aggression test that leopard, and that acute fluoxetine increases aggression in

leopard and decreased it in longfin zebrafish.

Keywords: Zebrafish; Aggressive display; Serotonin; Fluoxetine; Phenotypic

differences

1. Introduction

The biological comprehension of factors underlying aggression is still limited

(Miczek et al., 2007), even though a range of mental disorders present aggression as a

symptom (Krakowski, Volavka, & Brizer, 1986). Despite the paucity of neurobiological data

on aggression, a role for monoamines has been proposed (Miczek et al., 2007; Takahashi,

Quadros, Almeida, & Miczek, 2011). In most animal models, acutely increasing the

serotonergic transmission inhibits aggressive behavior (Takahashi et al., 2011); a metanalysis

of preclinical studies demonstrated that, across species, pharmacologically increasing 5-HT

levels inhibits aggression (Carrillo, Ricci, Coppersmith, & Melloni Jr., 2009).

While this observation appears to hold for most studies, some controversies and gaps

appear in the literature, especially in basal vertebrates such as fish. For example, in the

metanalysis by Carrillo et al. (2009), 5-HT decreased aggression in wrasses and trouts, but

not in the Siamese fighting fish. Zebrafish have been introduced as a model organism in

behavioral neuroscience and biological psychiatry (Norton & Bally-Cuif, 2010; Stewart et al.,

2015), increasing the relevance of findings using fish to study the neurobiology of

aggression.

A role for 5-HT in zebrafish aggressive behavior has been suggested by

neurochemical studies. After eliciting an aggressive display towards a mirror (mirror-induced

aggression, MIA), 5-HT levels were increased in the telencephalon, while 5-HIAA was

increased in the optic tectum of zebrafish (Teles, Dahlbom, Winberg, & Oliveira, 2013). Male

and female zebrafish respond to agonistic encounters in a similar fashion; nonetheless, males

present higher 5-HT turnover in the forebrain in relation to females, suggesting that

aggressive bouts could be more stressful to males than females (Dahlbom, Backström,

Lundstedt-Enkel, & Winberg, 2012). Filby et al. (2010) demonstrated that dominant males

show an overexpression of genes associated with the serotonergic system in the

hypothalamus, including *tph1b* and *htr1aa*, while females showed overexpression of *tph2*,

htr1aa, *slc6a4a*, and *mao* in the hypothalamus and *tph1a* and *tph2* in the telencephalon.

While these results suggest that aggressive behavior can be linked to differences in the

serotonergic system – especially in the context of dominance hierarchies –, a causal

relationship is more tenuous. Filby et al. (2010) treated dominant male zebrafish with

fluoxetine (3 or 4.5 µg/L), without effects on aggressive behavior in a dyadic encounter;

however, a similar concentration (3 µg/L) decreased aggressive displays in the MIA (W. H. J.

Norton et al., 2011). Using a much higher concentration (5 mg/L), Theodoridi et al. (2017)

were able to inhibit attacks and chasing behavior in dominant animals in dyads. The lack of

consistency could be due to dosing, behavioral paradigms (e.g., MIA vs. dyadic encounters),

or other variables.

Recent studies also showed that 5-HT levels are lower in zebrafish with the leopard

phenotype than in animals with the longfin phenotype, an alteration that is accompanied by

increased monoamine oxidase activity (Maximino, Puty, Oliveira, & Herculano, 2013). These

neurochemical differences were accompanied by increased anxiety-like behavior that is

rescued by fluoxetine treatment (Maximino, Puty, Oliveira, et al., 2013). Interestingly, in

longfin animals fluoxetine *increases* anxiety, and 5-HT levels are negatively correlated with

anxiety-like behavior; it is possible that embryological differences in the serotonergic system

produce opposite adult phenotypes.

These phenotypic differences are exploited in the present work to clarify the role of

phasic serotonin on aggressive displays in zebrafish. We hypothesized that the

hyposerotonergic phenotype of leopard zebrafish would produce increased aggressive

behavior, and that fluoxetine would rescue this phenotype. The experimental evidence

produced in the present work contradicted this hypothesis, since longfin were shown to

display more aggressive motivation and readiness in the mirror-induced aggression test than

leopard zebrafish, and since acute fluoxetine increased aggression in leopard animals and

decreased it in longfin zebrafish. This manuscript is a complete report of all the studies

performed to test the effect of skin phenotype and fluoxetine on aggressive behavior. We

report how we determined our sample size, all data exclusions (if any), all manipulations, and

all measures in the study.

2. Methods

2.1. Animals, housing, and baseline characteristics

Outbred populations were used due to their increased genetic variability, decreasing

the effects of random genetic drift which could lead to the development of uniquely heritable

traits (Parra, Adrian Jr, & Gerlai, 2009; Speedie & Gerlai, 2008). Thus, the animals used in

the experiments are expected to better represent the natural populations in the wild. Adult

zebrafish from the wildtype strain (longfin and leopard phenotypes) were used in this

experiment. Animals were bought from a commercial vendor, and arrived in the laboratory

with an approximate age of 3 months (standard length = 13.2 ± 1.4 mm), and were

quarantined for two weeks; the experiment began when animals had an approximate age of 4

months (standard length = 23.0 ± 3.2 mm). Animals were kept in mixed-sex tanks during

acclimation, with an approximate ratio of 50 male:50 female. Both phenotypes were kept in

the same tank before experiments. The breeder was licensed for aquaculture under Ibama's

(Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) Resolution

95/1993. Animals were group-housed in 40 L tanks, with a maximum density of 25 fish per

tank, for at least 2 weeks before experiments begun. Tanks were filled with non-chlorinated

water at room temperature (28 °C) and a pH of 7.0-8.0. Lighting was provided by fluorescent

lamps in a cycle of 14-10 hours (LD), according to standards of care for zebrafish (Lawrence,

2007). Water quality parameters were as follows: pH 7.0-8.0; hardness 100-150 mg/L

CaCO3; dissolved oxygen 7.5-8.0 mg/L; ammonia and nitrite < 0.001 ppm. All manipulations

minimized their potential suffering of animals, and followed Brazilian legislation (Conselho

Nacional de Controle de Experimentação Animal - CONCEA, 2017). Animals were used for

only one experiment and in a single behavioral test, to reduce interference from apparatus

exposure.

2.2. Mirror-induced aggression

137   The protocol for mirror-induced aggression was adapted from Norton et al. (2011).

Animals were individually transferred to a tank (15 x 10 x 30 cm) containing 1 L of holding

tank water. The tank was lit from above with white light (445 ± 56 lumens). Tanks were not

aerated during testing, so as not to disturb the animals. However, system water, which shows

adequate D.O. levels (7.5-8.0 mg/L), was used thoughout the experiments. Animals were

allowed to acclimate to the tank for 5 min; after that, a mirror was positioned on the outside

of the tank (on the narrower side), in an angle of 22.5°. All steps of the experiment were

executed under constant Gaussian white noise (55 ± 2.5 dB above the tank). Behavior was

recorded with a digital video camera (Samsung ES68) positioned in the wider side of the

tank, and analyzed by observers blind to treatment using the event-recording software X-Plo-

Rat (https://github.com/lanec-unifesspa/x-plo-rat). The following endpoints were analyzed:

time in the square nearest to the mirror (s); frequency (N) and duration (s) of aggressive

display; total number of squares crossed (N). Aggressive display was defined as a swimming

posture with erect dorsal, caudal, pectoral, and anal fins (Gerlai, Lahav, Guo, & Rosenthal,

2000).

2.3. Data availability

Datasets and scripts for all analyses are available from https://github.com/lanec-

unifesspa/5HT-aggression (doi: 10.5281/zenodo.1006701).

2.4. Experiment 1

Sample size calculation and groups. Sample sizes were calculated on results

regarding the effects of fluoxetine on total time in broadside display in *Betta splendens*,

reported by Lynn et al. (2007); as a result, the closest endpoint (time on display) was chosen

as the primary endpoint, and calculations for sample sizes are valid only for that endpoint.

Calculations were based on Rosner's (2016) method for comparing two means, and assumed

$\alpha = 0.05$ and power 80% on a two-tailed analysis. Based on these calculations, 15 animals

were used in each group in Experiment 1. Animals were derived from the stock population

described in section 2.1, and displayed either the longfin (Group LOF) or leopard (Group

LEO) phenotypes. Animals were randomly drawn from the tank immediately before testing,

and the order with which phenotypes were tested was randomized *via* generation of random

numbers using the randomization tool in http://www.randomization.com/. Blinding was not

possible, due to the obvious differences in skin phenotype.

Experimental design and statistical analysis. Animals were allocated to each group

according to phenotype. Immediately after being drawn from the tank, animals were

individually transported to the experiment room, and left undisturbed for 30 min. After this

interval, animals were exposed to the MIA test, described above. Differences between groups

were analyzed using Approximative Two-Sample Fisher-Pitman Permutation Tests 10,000

Monte-Carlo re-samplings, using the R package 'coin' (Hothorn, Hornik, van de Wiel, &

Zeileis, 2006). The data analyst was blinded to phenotype by using coding to reflect

treatments in the resulting datasets; after analysis, data was unblinded. Data are presented

using individual dot plots combined with boxplots. Effect sizes are noted in the text as

Cohen's d.

2.5. Experiment 2

Sample size calculation and groups. In the absence of similar experiments in the

literature, sample sizes were calculated based on the assumption of fixed effect sizes for both

the phenotype and the dose factors, with a projected effect size of 0.4, and 80% power;

calculations were made using the R package 'pwr2' (Lu, Liu, & Koestler, 2017). Based on

these calculations, 10 animals were used in each group in Experiment 1. Animals were

derived from the stock population described in section 2.1, and displayed either the longfin

(Group LOF) or leopard (Group LEO) phenotypes. Animals were randomly drawn from the

tank immediately before testing, and the order with which phenotypes were tested was

randomized *via* generation of random numbers using the randomization tool in

http://www.randomization.com/. Blinding for phenotype was not possible, due to the obvious

differences in skin phenotype. Animals from each phenotype were randomly allocated to

treatment (vehicle or either fluoxetine dose) *via* generation of random numbers using the

randomization tool in http://www.randomization.com/.

Drug treatments. Fluoxetine (FLX) was bought from EMS, dissolved in Cortland's

salt solution (Wolf, 1963), and injected intraperitoneally in cold-anesthetised animals (Kinkel,

Eames, Philipson, & Prince, 2010). FLX doses (2.5 and 5.0 mg/kg) were based on the

demonstration of effect on the light/dark test on longfin (Maximino, Puty, Benzecry, et al.,

2013) and leopard (Maximino, Puty, Oliveira, et al., 2013) zebrafish. Experimenters were

blinded to treatment by coding drug vials.

Experimental design and statistical analysis. Animals were allocated to each group

according to phenotype. Immediately after being drawn from the tank, animals were

individually transported to the experiment room, injected with vehicle or drug, and left

undisturbed for 30 min. After this interval, animals were exposed to the MIA test, described

above. Differences between groups were analyzed using two-way analyses of variance with

robust estimators on Huber's M-estimators, using the R package 'rcompanion' (Mangiafico,

2017). P-values were adjusted for the false discovery rate. The data analyst was blinded to

phenotype by using coding to reflect treatments in the resulting datasets; after analysis, data

was unblinded. Data are presented using individual dot plots combined with boxplots. Effect

sizes are reported as partial ε^2 values, and were calculated using the R package 'lsr' (Navarro,

2015).

3. Results

3.1 Experiment 1

LEO zebrafish showed longer latencies to display than LOF animals (Z = 3.3925, p =

0.0005, *d* = 1.5779; Figure 1A), as well as shorter display durations (Z = -2.5659, p = < 2.2e-

16, *d* = -1.0605; Figure 1B) and frequency (Z = -2.7073, p = 0.003, *d* = -1.1372; Figure 1C),

and time spent near the mirror (Z = -3.2284, p = < 2.2e-16, *d* = -1.4593; Figure 1D). No

differences were found in total locomotion (Z = -0.69887, p = 0.4965, *d* = -0.2573; Figure

1E).

3.2. Experiment 2

No main effects of phenotype (p = 0.5736; partial ε^2 = 0.0326) or FLX dose (p =

0.5044; partial ε^2 = 0.0482) were found for latency, but a significant interaction was found (p

= 0.0412; partial ε^2 = 0.1622); nonetheless, post-hoc tests did not detect any differences

between groups (Figure 2A). Main effects of phenotype (p = 0.0132; partial ε^2 = 0.2429) and

FLX dose (p = 0.0062; partial ε^2 = 0.2297), as well as an interaction effect (p = 0.009; partial

ε^2 = 0.1986), were found for display duration (Figure 2B). Post-hoc tests suggested that FLX

(2.5 mg/kg) decreased display duration on LOF, but not LEO (p = 0.032 vs. 0 mg/kg).

Similarly, main effects of phenotype (p = 0.0458; partial ε^2 = 0.2744) and FLX dose (p =

0.004; partial ε^2 = 0.2476), as well as an interaction effect (p < 0.0001; partial ε^2 = 0.3598),

were found for display frequency (Figure 2C); post-hoc tests suggested that FLX (5 mg/kg)

increased display frequency in LEO, but not LOF animals (p = 0.004 vs. 0 mg/kg). No main

effects of phenotype (p = 0.2692; partial ε^2 = 0.3704) were found for time near mirror, but a

main effect of FLX dose (p = 0.0004; partial ε^2 = 0.0164) and an interaction effect (p <

0.0001; partial ε^2 = 0.5760); post-hoc tests suggested a inverted-U-shaped curve for FLX-

treated LOF (0 vs. 2.5 mg/kg: p = 0.0012; 0 vs. 5.0 mg/kg: p = 0.0167), while a monotonic

increase for FLX-treated LEO (0 vs. 2.5 mg/kg: p = 0.0496; 0 vs. 5.0 mg/kg: p < 0.0001)

(Figure 2D). No main or interaction effects were found for total locomotion (Figure 2E).

4. Discussion

The present work demonstrated that zebrafish with the leopard skin phenotype show

less aggressive readiness and less aggression in relation to longfin animals. Moreover, a

different pattern of fluoxetine effects was observed, with fluoxetine decreasing aggressive

display (but not readiness) in longfin animals and increasing it in leopard animals. Evidence

for dose-dependence was also observed.

Serotonin (5-HT) has long been implicated in the neurobiological mechanisms of

aggressive behavior (Miczek et al., 2007; Summers & Winberg, 2006; Takahashi et al., 2011).

In a metanalysis of preclinical studies, Carrillo et al. (2009) demonstrated that, across species,

pharmacologically increasing 5-HT levels inhibit aggression. Interestingly, in their

metanalysis a species-specific effect was found in fish, with 5-HT decreasing aggression in

wrasses and trouts, but not in the Siamese fighting fish (Carrillo et al., 2009). The present

study examined only aggressive displays, which was elicited in the mirror-induced aggression

test; as a result, dominance hierarchies were not induced. Similar results were observed by

Norton et al. (2011), which observed reduced aggressive displays in Tübingen zebrafish

treated with fluoxetine. When zebrafish are allowed to form dominance hierarchies,

fluoxetine either produces no effect (Filby et al., 2010, using WIK zebrafish) or reduces

aggression in dominant males, but not in subordinates (Theodoridi et al., 2017, undescribed

phenotype). Given that these studies used different phenotypes than those reported here,

conclusions are limited.

Behavioral differences between leopard and longfin phenotypes were observed in

zebrafish before (Canzian, Fontana, Quadros, & Rosemberg, 2017; Egan et al., 2008;

Maximino, Puty, Oliveira, et al., 2013; Quadros et al., 2016; https://doi.org/10.1101/055657).

Of special relevance is the observation that leopard zebrafish present increased brain

monoamine oxidase (MAO) activity that is associated with lower serotonin levels and higher

turnover of 5-HT in the brain (Maximino, Puty, Oliveira, et al., 2013; Quadros et al., 2018).

This hyposerotonergic profile was also associated with increased anxiety-like behavior that

was rescued by fluoxetine treatment (Maximino, Puty, Oliveira, et al., 2013). Moreover,

Quadros et al. (2018) also found that leopard to be less aggressive than shortfin zebrafish,

suggesting a consistent hypoaggressive phenotype across laboratories, conditions, and

background genetics.

These results suggest a serotonin-linked behavioral syndrome in zebrafish that varies

across populations. Indeed, the hyperanxious profile observed in leopard (Maximino et al.,

2013) is also rescued by fluoxetine treatment at the same dose range as that reported here. A

variety of studies in Siamese fighting fish (*Betta splendens*) suggest that fluoxetine reduces

aggressive behavior (Dzieweczynski & Hebert, 2012; Eisenreich & Szalda-Petree, 2015;

Kania, Gralak, & Wielgosz, 2012) and boldness (Dzieweczynski, Campbell, & Kane, 2016;

Dzieweczynski, Kane, Campbell, & Lavin, 2016), which could be interpreted as either

increased impulsivity or decreased anxiety. The presence of a aggression-boldness syndrome

has long been proposed in as a dimension in fish behavior (Conrad, Weinersmith, Brodin, &

Saltz, 2011), and the literature appears to point to serotonin as an important link in that.

Nonetheless, these results must be interpreted with caution, given that the *B. splendens*

experiments were made with chronic fluoxetine treatment (which, along with increased

serotonin levels, is thought to induce other long-term neuroadaptations; Castrén & Antila,

2017). Moreover, other experiments with zebrafish (Norton et al., 2010) failed to find an

effect of fluoxetine in aggression-boldness – although, again, the use of different strains and

phenotypes make it difficult to generalize.

The results from the Carrillo et al. (2009) metanalysis suggested a inhibitory role for

phasic serotonin (i.e., 5-HT released by either the aggressive act itself, or by pharmacological

manipulations such as fluoxetine); a role for tonic 5-HT, in *Betta splendens*, was discarded,

because neither the 5-HT synthesis inhibitor *para*-chlorophenylalanine nor the 5-HT

precursor L-tryptophan changed display behavior (Clotfelter, O'Hare, McNitt, Carpenter, &

Summers, 2007). If that was also true for zebrafish, differences in aggressive display between

leopard and longfin would not be expected, given that these phenotypes differ in serotonergic

tone (Maximino, Puty, Oliveira, et al., 2013). While these differences were observed in the

present work, they occur in the opposite direction from what would be predicted from 5-

HTergic tone alone (i.e., we should expect leopard zebrafish to be more aggressive if

aggression was linearly and negatively related to tone). It is more likely that the

normalization of 5-HT levels in leopard after fluoxetine treatment is responsible for increased

aggression, while the "extra" 5-HT levels after fluoxetine treatment in longfin reduce its basal

aggression levels; as a result, the relationship between aggression and 5-HT levels are to be

interpreted as following and inverted-U-shaped distribution (Figure 3). This is also reinforced

by the generally hormetic dose-response curves observed in longfin animals treated with

fluoxetine. Alternatively, it is possible that a developmental effect is responsible for these

discrepancies.

While it might be tempting to attribute these differences to genetic differences across

populations, the animals used were not derived from inbred strains. The altered pigmentation

observed in our leopard fish has previously been reported, in the Tupfel long-fin (TL) strain,

to be due to a mutation in *connexin41.8* (Watanabe et al., 2006); nonetheless, it is unknown

whether this mutation is present in our animals, or whether this genetic marker is one of

many loci that differ between leopard and longfin animals (Gerlai, 2018). These differences

make it difficult to make specific genetic inferences regarding the differences observed in the

present work. Nonetheless, behavioral differences between leopard and longfin zebrafish

were consistently observed across laboratories (and therefore across fish vendors) (Canzian,

Fontana, Quadros, & Rosemberg, 2017; Egan et al., 2008; Maximino et al., 2013; Quadros et

al., 2016, 2018), and the effect of phenotype on zMAO activity was observed independently

at least twice (Maximino et al., 2013; Quadros et al., 2018), suggesting an important link

between the serotonergic system and aggressive behaviors across zebrafish strains.

These results are also reminiscent of what is observed in different populations of

Astyanax mexicanus. In that species, different populations occupy different niches, and

surface-dwelling populations are much more aggressive than cave-dwelling populations

(Rétaux & Elipot, 2013). These differences are related to the density of serotonergic neurons

in the hypothalamus, with cavefish showing a higher number of 5-HT neurons in that region

(Elipot, Hinaux, Callebert, & Rétaux, 2013). Moreover, a mutation in the *mao* gene was

found in cavefish that led to an hyperserotonergic phenotype (Elipot et al., 2014). Treating

surface fish with fluoxetine decreases aggression, while in cavefish the drug slightly

increases it (Elipot et al., 2013). In the present paper, however, treatment with fluoxetine

increased aggression in leopard animals (which show an hyposerotonergic profile in relation

to longfin animals; Maximino, Puty, Oliveira, et al., 2013) and decreased it in longfin

zebrafish. These differences might be due to the origin of serotonin, since, in *Astyanax*

mexicanus populations, raphe 5-HT levels are unchanged, while hypothalamic 5-HT is

increased (Elipot et al., 2013); further experiments are needed to untangle this hypothesis.

Interestingly, a different dose-response profile was observed between phenotypes in

the present study, with the low dose (2.5 mg/kg) generally decreasing aggression in longfin

and the high dose (5.0 mg/kg) generally increasing it in leopard. While difficult to explain

presently, these results suggest either that an "optimal" serotonergic tone is needed to

maintain aggression levels, or that serotonin transporters are desensitized or downregulated in

the leopard population. While the first hypothesis is more likely, given the observation of an

hyposerotonergic profile in leopard zebrafish (Maximino et al., 2013; Quadros et al., 2018),

the current state of the literature and the current data are not enough to assess this.

In conclusion, the present experiments revealed differences in aggressive behavior

between leopard and longfin zebrafish, and a discrepant effect of fluoxetine on both

populations. These results are relevant to understand the role of tonic and phasic serotonin

neurotransmission on aggressive behavior in preclinical models, and might contribute to a

better appreciation of the complex roles of this monoamine in controlling vertebrate

aggression.

Acknowledgments

References

Canzian, J., Fontana, B. D., Quadros, V. A., & Rosemberg, D. B. (2017). Conspecific alarm

substance differently alters group behavior of zebrafish populations: Putative

involvement of cholinergic and purinergic signaling in anxiety- and fear-like responses.

Behavioural Brain Research, *320*, 255–263. https://doi.org/10.1016/j.bbr.2016.12.018

Carrillo, M., Ricci, L. A., Coppersmith, G. A., & Melloni Jr., R. H. (2009). The effect of

increased serotonergic neurotransmission on aggression: A critical meta-analytical

review of preclinical studies. *Psychopharmacology*, *205*, 349–368.

https://doi.org/10.1007/s00213-009-1543-2

Castrén, E., & Antila, H. (2017). Neuronal plasticity and neurotrophic factors in drug

responses. *Molecular Psychiatry, 22*, 1085–1095. https://doi.org/10.1038/mp.2017.61

Clotfelter, E. D., O'Hare, E. P., McNitt, M. M., Carpenter, R. E., & Summers, C. H. (2007).

Serotonin decreases aggression via 5-HT1A receptors in the fighting fish *Betta

splendens*. *Pharmacology, Biochemistry & Behavior*, *87*, 222–231.

https://doi.org/10.1016/j.pbb.2007.04.018

Conrad, J. L., Weinersmith, K. L., Brodin, T., & Saltz, J. B. (2011). Behavioural syndromes in

fishes: A review with implications for ecology and fisheries management. *Journal of*

Fish Biology, 78, 395–435. https://doi.org/10.1111/j.1095-8649.2010.02874.x

Conselho Nacional de Controle de Experimentação Animal - CONCEA. Diretriz brasileira

para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo

I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científica, Pub.

L. No. Resolução Normativa 30/2016 (2017). Brasil.

Dahlbom, S. J., Backström, T., Lundstedt-Enkel, K., & Winberg, S. (2012). Aggression and

monoamines: Effects of sex and social rank in zebrafish (Danio rerio). *Behavioural*

Brain Research, *228*, 333–338. https://doi.org/10.1016/j.bbr.2011.12.011

Dzieweczynski, T. L., Campbell, B. A., & Kane, J. L. (2016). Dose-dependent fluoxetine

effects on boldness in male Siamese fighting fish. *Journal of Experimental Biology, 219*,

797–804. https://doi.org/10.1242/jeb.132761

Dzieweczynski, T. L., & Hebert, O. L. (2012). Fluoxetine alters behavioral consistency of

aggression and courtship in male Siamese fighting fish, *Betta splendens*. *Physiology and*

Behavior, 107, 92–97. https://doi.org/10.1016/j.physbeh.2012.06.007

Dzieweczynski, T. L., Kane, J. L., Campbell, B. A., & Lavin, L. E. (2016). Fluoxetine

exposure impacts boldness in female Siamese fighting fish, *Betta splendens*.

Ecotoxicology, 25, 69–79. https://doi.org/10.1007/s10646-015-1568-8

Eisenreich, B. R., & Szalda-Petree, A. (2015). Behavioral effects of fluoxetine on aggression

and associative learning in Siamese fighting fish (*Betta splendens*). *Behavioural*

Processes, 121, 37–42. https://doi.org/10.1016/j.beproc.2015.10.008

Elipot, Y., Hinaux, H., Callebert, J., Launay, J.-M., Blin, M., & Rétaux, S. (2014). A mutation

in the enzyme monoamine oxidase explains part of the *Astyanax* cavefish behavioural

syndrome. *Nature Communications*, *5*, 3647. https://doi.org/10.1038/ncomms4647

Elipot, Y., Hinaux, H., Callebert, J., & Rétaux, S. (2013). Evolutionary shift from fighting to

foraging in blind cavefish through changes in the serotonin network. *Current Biology*,

23, 1–10. https://doi.org/10.1016/j.cub.2012.10.044

Filby, A. L., Paull, G. C., Hickmore, T. F., & Tyler, C. R. (2010). Unravelling the

neurophysiological basis of aggression in a fish model. *BMC Genomics*, *11*, 498.

https://doi.org/10.1186/1471-2164-11-498

Gerlai, R., Lahav, M., Guo, S., & Rosenthal, A. (2000). Drinks like a fish: Zebra fish (*Danio*

rerio) as a behavior genetic model to study alcohol effects. *Pharmacology, Biochemistry*

and Behavior, *67*, 773–782.

Gerlai, R. T. (2018). Reproducibility and replicability in zebrafish behavioral neuroscience

research. *Pharmacology, Biochemistry & Behavior, In press.*

https://doi.org/10.1016/j.pbb.2018.02.005

Hothorn, T., Hornik, K., van de Wiel, M. A., & Zeileis, A. (2006). A Lego system for

conditional inference. *The American Statician*, *60*, 257–263.

https://doi.org/10.1198/000313006X118430

Kania, B. F., Gralak, M. A., & Wielgosz, M. (2012). Four-week fluoxetine (SSRI) exposure

diminishes aggressive behaviour of male Siamese fighting fish (*Betta splendens*).

Journal of Behavioral and Brain Science, 2, 185–190.

https://doi.org/10.4236/jbbs.2012.22022

Kinkel, M. D., Eames, S. C., Philipson, L. H., & Prince, V. E. (2010). Intraperitoneal injection into adult zebrafish. *Journal of Visualized Experiments*, *42*, 2126. https://doi.org/10.3791/2126

Krakowski, M., Volavka, J., & Brizer, D. (1986). Psychopathology and violence: A review of literature. *Comprehensive Psychiatry*, *27*, 131–148. https://doi.org/10.1016/0010-440X(86)90022-2

Lawrence, C. (2007). The husbandry of zebrafish (*Danio rerio*): A review. *Aquaculture*, *269*, 1–20. https://doi.org/10.1016/j.aquaculture.2007.04.077

Lu, P., Liu, J., & Koestler, D. (2017). pwr2: Power and Sample Size Analysis for One-way and Two-way ANOVA Models. Retrieved from https://cran.r-project.org/package=pwr2

Lynn, S. E., Egar, J. M., Walker, B. G., Sperry, T. S., & Ramenofsky, M. (2007). Fish on Prozac: A simple, noninvasive physiology laboratory investigating the mechanisms of aggressive behavior in Betta splendens. *Advances in Physiology Education*, *31*, 358–363. https://doi.org/10.1152/advan.00024.2007

Mangiafico, S. (2017). rcompanion: Functions to Support Extension Education Program Evaluation. Retrieved from https://cran.r-project.org/package=rcompanion

Maximino, C., Puty, B., Benzecry, R., Araujo, J., Lima, M. G., Batista, E. de J. O., … Herculano, A. M. (2013). Role of serotonin in zebrafish (*Danio rerio*) anxiety: Relationship with serotonin levels and effect of buspirone, WAY 100635, SB 224289, fluoxetine and para-chlorophenylalanine (pCPA) in two behavioral models. *Neuropharmacology*, *71*, 83–97. https://doi.org/10.1016/j.neuropharm.2013.03.006

Maximino, C., Puty, B., Oliveira, K. R. M., & Herculano, A. M. (2013). Behavioral and

neurochemical changes in the zebrafish leopard strain. *Genes, Brain and Behavior*, *12*,

576–582. https://doi.org/10.1111/gbb.12047

Miczek, K. A., de Almeida, R. M. M., Kravitz, E. a, Rissman, E. F., de Boer, S. F., & Raine,

A. (2007). Neurobiology of escalated aggression and violence. *Journal of Neuroscience*,

27, 11803–11806. https://doi.org/10.1523/JNEUROSCI.3500-07.2007

Navarro, D. J. (2015). Learning Statistics with R: A Tutorial for Psychology Students and

Other Beginners, Version 0.5. [Lecture notes] School of Psychology, University of

Adelaide, Adelaide, Australia. ISBN: 978-1-326-18972-3. URL:

http://health.adelaide.edu.au/psychology/ccs/teaching/lsr/.

Norton, W., & Bally-Cuif, L. (2010). Adult zebrafish as a model organism for behavioural

genetics. *BMC Neuroscience*, *11*, 90. https://doi.org/10.1186/1471-2202-11-90

Norton, W. H. J., Stumpenhorst, K., Faus-Kessler, T., Folchert, A., Rohner, N., Harris, M. P.,

… Bally-Cuif, L. (2011). Modulation of *fgfr1a* signaling in zebrafish reveals a genetic

basis for the aggression–boldness syndrome. *Journal of Neuroscience*, *31*, 13796–

13807. https://doi.org/10.1523/JNEUROSCI.2892-11.2011

Parra, K. V, Adrian Jr, J. C., & Gerlai, R. (2009). The synthetic substance hypoxanthine 3-N-

oxide elicits alarm reactions in zebrafish (*Danio rerio*). *Behavioural Brain Research*,

205, 336–341. https://doi.org/10.1016/j.bbr.2009.06.037

Quadros, V. A., Silveira, A., Giuliani, G. S., Didonet, F., SIlveira, A. S., Nunes, M. E., …

Rosemberg, D. B. (2016). Strain- and context-dependent behavioural responses of acute

alarm substance exposure in zebrafish. *Behavioural Processes*, *122*, 1–11.

https://doi.org/10.1016/j.beproc.2015.10.014

Quadros, V. A., Costa, F. V., Canzian, J., Nogueira, C. W., & Rosemberg, D. B. (2018).

Modulatory role of conspecific alarm substance on aggression and brain monoamine

oxidase activity in two zebrafish populations. *Progress in Neuro-Psychopharmacology*

& Biological Psychiatry. https://doi.org/10.1016/j.pnpbp.2018.03.018.

Rétaux, S., & Elipot, Y. (2013). Feed or fight. A behavioral shift in blind cavefish.

Communicative & Integrative Biology, *6*, e23166. https://doi.org/10.4161/cib.23166

Rosner, B. (2016). *Fundamentals of biostatistics*. Cengage Learning.

Speedie, N., & Gerlai, R. (2008). Alarm substance induced behavioral responses in zebrafish

(*Danio rerio*). *Behavioural Brain Research*, *188*, 168–177.

https://doi.org/10.1016/j.bbr.2007.10.031

Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H., Gerlai, R.,

& Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular Psychiatry*, *20*,

2–17. https://doi.org/10.1038/mp.2014.128

Summers, C. H., & Winberg, S. (2006). Interactions between the neural regulation of stress

and aggression. *Journal of Experimental Biology*, *209*, 4581–4589.

https://doi.org/10.1242/jeb.02565

Takahashi, A., Quadros, I. M., Almeida, R. M. M. de, & Miczek, K. A. (2011). Brain

serotonin receptors and transporters: Initiation vs . termination of escalated aggression.

Psychopharmacology, *213*, 183–212. https://doi.org/10.1007/s00213-010-2000-y

Teles, M. C., Dahlbom, S. J., Winberg, S., & Oliveira, R. F. (2013). Social modulation of

brain monoamine levels in zebrafish. *Behavioural Brain Research*.

https://doi.org/10.1016/j.bbr.2013.07.012

Theodoridi, A., Tsalafouta, A., & Pavlidis, M. (2017). Acute exposure to fluoxetine alters

aggressive behavior of zebrafish and expression of genes involved in serotonergic

system regulation. *Frontiers in Neuroscience*, *11*, Article 223.

https://doi.org/10.3389/fnins.2017.00223

Watanabe, M., Iwashita, M., Ishii, M., Kurachi, Y., Kawakami, A., Kondo, S., & Okada, N.

(2006). Spot pattern of leopard Danio is caused by mutation in the zebrafish

connexin41.8 gene. *EMBO Reports, 7,* 893–897.

https://doi.org/10.1038/sj.embor.7400757

Wolf, K. (1963). Physiological salines for freshwater teleosts. *Progress in Fish-Culture*, *25*,

135–140.

Figure captions

Figure 1 – Phenotype differences in (A) latency to display, in s; (B) display duration, in s;

(C), display frequency; (D), time spent near the mirror, in s; and (E), total locomotion.

Boxplots represent median and interquartile range, with Tukey whiskers.

Figure 2 – Effects of fluoxetine on the aggressive display of longfin (LOF, dark gray) and

leopard (LEO, light gray) zebrafish. (A) latency to display, in s; (B) display duration, in s;

(C), display frequency; (D), time spent near the mirror, in s; and (E), total locomotion.

Boxplots represent median and interquartile range, with Tukey whiskers. Dots joined by lines

represent means.

Figure 3 – Hypothesized role of the serotonergic tone on the organization of aggressive

display in longfin and leopard zebrafish.

- More 5-HTergic tone
- Less 5-HTergic tone

- More aggressive displays
- More aggressive readiness

- Less aggressive displays
- Less aggressive readiness

Acute FLX

- Too much 5-HT
- Normal 5-HT

Display

Display