Microbial community structure and microbial networks correspond to nutrient gradients within coastal wetlands of the Great Lakes.

Dean J. Horton[1], Kevin R. Theis[2], Donald G. Uzarski[1], and Deric R. Learman[1]*

[1]Institute for Great Lakes Research, CMU Biological Station, and Department of Biology, Central Michigan University, Mt. Pleasant, MI, U.S.A.
[2]Department of Microbiology, Immunology, and Biochemistry, Wayne State University, Detroit, MI, U.S.A

*Correspondence:
Deric R. Learman
deric.learman@cmich.edu

Keywords: microbial, community structure, network analysis, Great Lakes, wetlands, nutrients, anthropogenic, Lake Erie

Word count: 7,202
Figure count: 7

Abstract

Microbial communities within the soil of Great Lakes coastal wetlands drive biogeochemical cycles and provide several other ecosystems services. However, there exists a lack of understanding of how microbial communities respond to nutrient gradients and human activity in these systems. This research sought to address the lack of understanding through exploration of relationships between nutrient gradients, microbial community diversity, and microbial networks among coastal wetlands throughout the Great Lakes. Significant differences in microbial community structure were found among coastal wetlands within the western basin of Lake Erie and all other wetlands studied (three regions within Saginaw Bay and one region in the Beaver Archipelago). Further, within Lake Erie wetlands, chemical and biological structure did not vary with increasing soil depth. Beyond this, alpha diversity levels were highest within Lake Erie coastal wetlands. These diversity differences coincided with higher nutrient levels within the Lake Erie region. Site-to-site variability also existed within the majority of the regions studied, suggesting site-scale heterogeneity may impact microbial community structure. Several subnetworks of microbial communities and individual community members were related to chemical gradients among wetland regions, revealing several candidate indicator communities and taxa which may be useful for Great Lakes coastal wetland management. This research provides an initial characterization of microbial communities among Great Lakes coastal wetlands and demonstrates that microbial communities could be negatively impacted by anthropogenic activities. Anthropogenic impacts to these coastal wetland communities could influence natural biogeochemical cycles which occur within coastal wetland soils, and by extension would directly influence the Great Lakes themselves.

1. Introduction

The Laurentian Great Lakes of North America are one of the largest freshwater systems on Earth, and are critical in supporting biogeochemical cycles, freshwater resources, biodiversity, and economic viability of the surrounding region. Notably, the Great Lakes region has been impacted by anthropogenic pressure, with cumulative stress having a particular impact on the western basin of Lake Erie (Danz *et al*., 2007; Uzarski *et al*., 2017). These negative impacts extend to ecological transition zones between upland and aquatic environments in the form of coastal wetlands which border the Great Lakes (Uzarski, 2009), as agricultural runoff, atmospheric deposition, and urbanization influence water chemistry, thereby reducing water quality and impairing wetlands (Trebitz *et al*., 2007; Morrice *et al*., 2008). This has stoked a surge in research assessing biodiversity and anthropogenic pressure on coastal wetlands of the Great Lakes since the Great Lakes Water Quality Agreement (GLWQA) was established in 1972 (Hackett *et al*., 2017). While much research on coastal wetlands has flourished in the wake of this international agreement, microbial communities within Great Lakes coastal wetlands remain almost entirely uncharacterized (Hackett *et al*., 2017). The few research studies on microbial communities in Great Lakes coastal wetlands have focused on the use of microbial enzymatic assays as a tool to explore decomposition rates and nutrient limitation (Jackson *et al*., 1995; Hill *et al*., 2006). Community diversity, structure, and taxonomic composition have been largely overlooked. As the microbial communities within Great Lakes coastal wetlands have yet to be fundamentally described, it is important to gather baseline data on what microbes exist within these systems, to elucidate how

these microbes could be interacting, and to determine to what extent microbial diversity may already be impacted by anthropogenic chemical disturbance.

Microbial communities contribute substantially to the ecological functioning of coastal wetlands, and these wetlands are vital in the retention of chemical pollutants (e.g., heavy metals), sediments, and excess nutrients (e.g., N and P) caused by anthropogenic activity from impacting the Great Lakes themselves (Wang & Mitsch, 1998; Sierszen *et al*., 2012). Coastal wetlands border much of the Great Lakes coastline, where they make up nearly 200,000 ha of habitat between the United States and Canada, despite reduction of habitat by approximately 50% since European colonization (Dahl, 1990; Hecnar, 2004; Sierszen *et al*., 2012). Further, the economy of the Great Lakes is contingent on the existence and proper functioning of coastal wetlands. In providing ecosystem services and promoting biodiversity, these wetlands have an estimated annual worth of $69 billion USD; the value of recreational fishing alone is valued at $7.4 billion USD per year (Krantzberg & de Boer, 2008; Campbell *et al*., 2015). As such, negative anthropogenic impacts on microbial communities could influence the economic viability of the Great Lakes region, biodiversity retention, and the functioning of critical elemental cycles which commonly occur within freshwater wetlands.

Most notably, carbon mineralization occurs within wetland soils via redox processes mediated by microbial communities, and these processes contribute to pollution mitigation and atmospheric greenhouse gas flux (Conrad, 1996; Reddy & DeLaune, 2008). Wetland soils often become chemically structured with increasing depth through sequential reduction of electron acceptors that decrease in metabolic favorability to microbes due to thermodynamic constraints (Conrad, 1996; Reddy & DeLaune, 2008; Kögel-Knabner *et al*., 2010). As microbial community metabolism changes in concert with soil chemical profiles, microbial community compositional shifts commonly reflect functional changes of the community (Lüdemann *et al*., 2000; Edlund *et al*., 2008; Lipson *et al*., 2015). However, while availability of electron acceptors may influence chemical and biological structure within wetland soils, concentration of carbon electron donors can influence the vertical stratification of redox processes (Achtnich *et al*., 1995; Alewell *et al*., 2008), and by extension, vertical microbial community structure (defined as relative proportions of microbial taxa within a community). As an example of how this may apply to natural environments, increased carbon and nutrient influx from anthropogenic activities (such as agricultural pressure) may impact microbial community structure within coastal wetlands. Impacts to microbial community composition may extend to shifts in chemical cycles and redox processes as consequence, as disturbance to microbial community structure can often lead to a shift in community function (Shade *et al*., 2012). However, while community structure may be indicative of environmental gradients within wetlands, taxonomic identification of microbes which respond to human pressures is necessary to appreciate which fraction of wetland microbial communities are most sensitive to environmental disturbances.

Networks of microbial taxa exist within microbial communities, and impacts to individual members could affect entire networks (Faust & Raes, 2012). Thus, it is important explore hypothetical microbial networks within natural environments, and their relationships to changing environmental conditions. Understanding how microbial networks respond to physicochemical shifts could aide in predicting how a future change in environmental conditions (perhaps caused by anthropogenic activity) may impact local microbial communities. Further, identifying microbial

taxonomic and diversity responses to environmental stressors caused by human activity is the first
step in developing biological indicators that can predict levels of anthropogenic stress on natural
environments, such as wetlands. Physicochemical and biological indicators have been
continuously developed to determine which biological taxa are most sensitive to anthropogenic
pressures within freshwater wetlands, and by extension, how these biological responses can inform
scientists and managers about the health of coastal wetlands along the Great Lakes (Uzarski *et al*.,
2017). These indices have been established for physical and chemical attributes (such as nutrient
levels, urbanization, land use, etc.), as well as several eukaryotic taxonomic groups (e.g.,
macrophytes, macroinvertebrates, fish, anurans, and birds) (Uzarski *et al*., 2017). However, as
different taxonomic indicators highlight unique pressures on wetland systems, indicators based on
different biological groups can often conflict in their assessment of wetland ecosystem health. As
such, it is necessary to examine a wide range of biological indicators to assess different aspects of
wetland ecosystem health. A biological index for bacteria and archaea has yet to be developed for
responses to human impacts within freshwater coastal wetlands (Uzarski et al., 2017). A first step
in establishing a microbial index is to uncover specific networks of microbial taxa (Sims *et al*.,
2013; Urakawa & Bernhard, 2017) and diversity patterns found to be related to environmental
gradients linked to anthropogenic activity (e.g., soil nutrient levels) among Great Lakes coastal
wetlands.

This study sought to provide an initial characterization of microbial communities within
soils of Great Lakes coastal wetlands bordering the western basin of Lake Erie, Saginaw Bay of
Lake Huron, and northern Lake Michigan. Wetland sites explored in this study have been
extensively researched over multiple years and vary widely in the degree to which they are
impacted by human activity (Uzarski *et al*., 2017). This study explored how environmental
gradients among these coastal wetlands were related to microbial community structure among
wetlands. Additionally, relationships among microbial communities and changing environmental
conditions with increasing soil depth were also explored within each wetland site. It was predicted
that microbial community structure would be related to environmental gradients among and within
coastal wetland regions of the Great Lakes, and elevated nutrient levels within wetlands would
decouple the relationship between microbial community structure and soil depth with respect to
coastal wetlands lower in nutrient levels, as has been suggested in previous studies (Achtnich *et*
al., 1995; Alewell *et al*., 2008). Through high-throughput sequencing of the 16S rRNA gene and
microbial network analyses, variations in key microbial taxa and subcommunities related to
environmental gradients established by wetlands were identified.

2. Material and Methods

In the summer of 2014, wetland soil cores were collected within Laurentian Great Lakes
coastal wetland ecosystems across several sites within several regions. Specifically, soil cores were
collected from ten sites across five regions, including two sites in the western basin of Lake Erie
(LE), three sites in eastern Saginaw Bay (ESBT), two sites in northern Saginaw Bay (NSB), two
sites in western Saginaw Bay (WSB) in Lake Huron, and one site in the Beaver Island archipelago
(BA) in Lake Michigan (Fig. 1). These sites were selected as they corresponded to environmental
gradients, as well as human impact gradients based upon SumRank scores (an index assessing land
use and water quality) as described in Uzarski *et al*. (2017) (Supplemental Fig. 1). Soil cores were
collected by hand-driving plastic core tubes (~ 5 cm diameter) vertically into the soil. Among

wetlands, samples were collected within the same vegetation zone across sites (either dominated
by cattails, genus *Typhus*, or bulrush, genus *Shoenoplectus*) as an attempt to control for collection
bias, as different vegetation zones can harbor microbial communities distinct from other vegetation
zones (Tang *et al*., 2011). Cores were sampled to a depth of at least 6 cm (except for one core
which was sampled to a depth of 4 cm) and were immediately flash frozen in a dry ice ethanol
bath. Samples were transported on dry ice to Central Michigan University wherein they were
stored at -80 ˚C.

Triplicate cores were taken at five wetland sites while duplicate cores were taken at five
other wetland sites. For sample extraction and sectioning, cores were extruded while still frozen
via a custom-built core extruder. The edge of the core was warmed with a heat gun to allow the
soil core to pass efficiently through the plastic container, however, the inner-core did not thaw
during extrusion. Ice was applied to the plastic core liner to prevent accelerated thawing. Beginning
from the top surface of soil, 2 cm sections were cut via an ethanol and flame-sterilized hacksaw
blade and the sectioned core samples were placed into Whirl-Pak bags and stored at -80 ˚C. The
extruder face plate was sterilized between cuts of the same core with ethanol. The extruder device
was fully cleaned and sterilized between cores with physical scrubbing and ethanol sterilization.

2.1 Microbial community analysis

Each soil sample was analyzed independently for microbial community analyses. DNA
was extracted from ~ 0.25 g of soil using a MoBio PowerSoil DNA Isolation Kit (Mo Bio,
Carlsbad, CA) following the standard manufacturer's protocol. Concentrations of extracted DNA
were assessed using a Qubit® 2.0 fluorometer (Life Technologies, Carlsbad, CA) to ensure
successful DNA extraction and quantification for sequence library preparation. DNA samples were
sent to Michigan State University (East Lansing, MI) for library preparation and sequence analysis
at the Research Technology Support Facility. The V4 region of the 16S rRNA gene was amplified
for downstream sequencing with the commonly used primers 16Sf-V4 (515f) and 16Sr-V4 (806r)
and a previously developed protocol (Caporaso *et al*., 2012; Kozich *et al*., 2013). Paired-end 250
bp sequencing was accomplished via a MiSeq high-throughput sequencer (Illumina, San Diego,
CA). Acquired DNA sequences were filtered for quality and analyzed using MOTHUR v 1.35.1
(Schloss *et al*., 2009) following the MiSeq SOP (available at https://www.mothur.org/) with
modifications. Scripts used for sequence processing can be found at the GitHub repository
associated with this study (https://github.com/horto2dj/GLCW/). Briefly, paired end sequences
were combined into single contigs. Sequences that contained homopolymers > 8 bases, and those
less than 251 or greater than 254 bp were removed. Sequences were aligned against the Silva (v
119) rRNA gene reference database (Quast *et al*., 2012). Sequences which did not align with the
V4 region were also subsequently removed from analysis. Chimeric DNA was searched for and
removed via UCHIME (Edgar *et al*., 2011). Sequences were classified via the Ribosomal Database
Project (training set v 9; Cole *et al*., 2013) with a confidence threshold of 80. Sequences classified
as chloroplast, mitochondria, eukaryotic, or unknown were removed. Remaining sequences were
clustered into Operational Taxonomic Units (OTUs) at 0.03 sequence dissimilarity using the
opticlust clustering algorithm. Sequence data associated with this research have been submitted to
the GenBank database under accession numbers SRR6261304 – SRR6261377 (Horton et al.,
2017).

2.2 Chemical analysis

Each soil layer (top, middle, and bottom) was analyzed separately for local chemistry at each site. Within each site, soil samples of the same depth (i.e., top, middle, and bottom soil samples) among duplicate/triplicate cores were combined and homogenized to obtain enough soil for chemical analyses. For chemical analysis, soil samples were sent to Michigan State University Soil & Plant Nutrient Lab (East Lansing, MI) to analyze for percent total N ("TN"), total P ("TP", ppm), total S ("TS", ppm), NO_3^- (ppm), NH_4^+ (ppm), percent organic matter ("OM"), percent organic carbon ("OC"), and C:N. In the field, a YSI multiprobe (YSI Inc., Yellow Springs, OH) was used to measure pH of the water residing directly above each collected soil core.

2.3 Statistical analyses

Statistical analyses were completed using R statistical software version 3.2.2 (R Core Team, 2015) unless otherwise stated. Code used for statistical analyses (and bioinformatic workflow) in this study can be found in the associated GitHub repository (https://github.com/horto2dj/GLCW/).

2.3.1 Physicochemical analysis

Differences in chemical profiles between samples within and among wetland regions were visualized using Principal Component Analysis (PCA). Prior to PCA, percentages were arcsin square root transformed and ratios were log transformed. Additionally, Pearson correlation analyses were performed to search for significant correlations between chemical variables. Collinearity in the dataset was addressed by combining highly correlated environmental variables ($r > 0.7$, $p \leq 0.001$). Only one of the correlated variables was included in PCA to remove exaggeration of correlated variables in PCA structure. Permutational Multivariate Analysis of Variance (perMANOVA; Anderson, 2001) was used to determine the influences of region and soil depth on physicochemical composition of samples, and 95% confidence intervals were established to compare differences among groups. Chemical depth profiles were also visualized for each wetland site to understand shifts in measured environmental variables with increasing soil depth.

2.3.2 Alpha diversity analysis

Alpha diversity analyses were performed to explore variation in OTU richness and evenness among wetland sites, regions, and soil depths, as well as to determine whether observed trends were driven by environmental variables. Prior to alpha diversity analyses, sequence abundance for each sample was subsampled to the lowest sequence abundance for any one sample ($n = 48,226$ sequences). Singletons were maintained within the sequence dataset for alpha diversity analyses, as alpha diversity indices can be reliant on the presence of singletons for proper estimation. Alpha diversity was calculated for each site using MOTHUR, including Chao1 richness and non-parametric Shannon diversity. Linear mixed-effect models and ANOVAs were used to test influences of wetland site, region, and soil depth on alpha diversity, controlling for wetland site as a random effect. Linear models and ANOVAs were used to test for variation in alpha diversity among wetland sites. If significant variation was found within an ANOVA result, post-hoc comparisons were implemented between sample groups using Tukey's Honest

249 Significant Differences (HSD) tests with Bonferroni adjustments (p-values obtained by number of
250 comparisons) for pairwise comparisons.
251
2.3.3 Beta diversity analysis
252
253
Beta diversity analyses were used to evaluate variation in microbial community structure
among wetland sites, regions, and soil depths, and to assess the extent to which observed variation
was explained by environmental conditions. Singletons and doubletons were removed from the
dataset for beta diversity analyses. All sequence data were maintained for beta diversity analyses
and transformed using the *DeSeq2* (Love *et al*., 2014) package, which normalized OTU
abundances among samples using a variance stabilizing transformation (VST) (McMurdie &
Holmes, 2014). The *phyloseq* (McMurdie & Holmes, 2013) and *Vegan* (Oksanen *et al*., 2007)
packages were used to compare beta diversity among samples. Dissimilarity in microbial
community structure among samples within and among sites was visualized using Non-metric
Multidimensional Scaling (NMDS) plots based on pairwise Bray-Curtis dissimilarity estimates.
The function *envfit* of the Vegan package was used to evaluate correlation between chemical
parameters and microbial community structure among samples according to NMDS. "Depth" was
also implemented as a dummy variable to test correlation between depth and microbial community
structure.
268
To test for significant differences in beta diversity among wetland sites, regions, and soil
depth, perMANOVA were implemented. Specifically, these tests evaluate significant variation
among within group and between group means (Clarke, 1993; Anderson, 2001; Anderson &
Walsh, 2013). If perMANOVA found significant differences among groups at the global level,
pairwise perMANOVA tests between groups were implemented with Bonferroni significance
adjustments to control for multiple pairwise comparisons. Anderson's permutation of dispersions
test (PERMDISP; Anderson, 2006; Anderson *et al*., 2006) was used to test for differences in
variance of community structure among sample groups (i.e. sites, regions, soil depths). Tukey's
Honest Significant Difference (HSD) tests were implemented with adjusted p-values for multiple
pairwise comparisons if significant differences in dispersion were found among groups.
279
To explore relationships between regional microbial community structure and
environmental variables, NMDS plots were generated for each individual region. Applying NMDS
to each region also allowed for the assessment of the correlational relationship between community
structure and soil depth (as a dummy variable) and other environmental variables (using the *envfit*
function) within individual regions. To test for differences in microbial community structure
between/among sites within a region, as well as among depths within a region, perMANOVA was
implemented individually for each region.
287
2.3.4 Network analyses
288
289
To explore relationships between microbial sub-communities and individual OTUs to
environmental variables, Weighted Correlation Network Analysis (WGCNA) was implemented
on OTU relative abundances using the *WGCNA* package (Langfelder & Horvath, 2008; Langfelder
& Horvath, 2012), executed as previously described (Guidi *et al*., 2016; Henson *et al*., 2016) with
modifications. OTUs which did not possess at least 2 sequences across 10% of samples were

removed from network analyses. These OTUs were removed to eliminate OTUs with potentially
spurious correlations to environmental variables or other OTUs, as well as to reduce computational
stress of analyses. Remaining OTU abundances across samples were normalized using variance
stabilizing transformation (VST) performed as described previously for beta diversity analyses.
To ensure scale-free topology of the network, the dissimilarity matrix generated through VST was
transformed to an adjacency matrix by raising this dissimilarity matrix to a soft threshold power.
A threshold power of $p = 4$ was chosen to meet scale-free topology assumptions based upon
criterion established by Zhang & Horvath (2005). Scale-free topology of network relationships
was further ensured through regression of the frequency distribution of node connectivity against
node connectivity; a network is scale-free if an approximate linear fit of this regression is evident
(see Zhang & Horvath, [2005] for more in-depth explanation). A topological overlap matrix
(TOM) was generated using the adjacency matrix, and subnetworks of highly connected and
correlated OTUs were delineated with the TOM and hierarchical clustering. Representative
eigenvalues of each subnetwork (i.e., the first principal component) were correlated (Pearson) with
values of measured environmental variables to identify the subnetworks most related to said
environmental variables. The subnetworks with the highest positive correlations to environmental
variables of interest (e.g., NO_3^-, C:N, etc.) were selected for further analyses of relationships among
subnetwork structure, individual OTUs, and environmental variables. Partial Least Square
regression (PLS) was used to test predictive ability of subnetworks in estimating variability of
environmental parameters, which allowed for delineation of potential indicator subnetworks and
OTUs. Pearson correlations were calculated between response variables and leave-one-out cross-
validation (LOOCV) predicted values. If PLS found that regression between actual and predicted
values was below minimum threshold of $R^2 = 0.3$, WGCNA analysis was halted for that network,
as the network was deemed to lack predictive ability of that environmental variable. Variable
Importance in Projection (VIP) (Chong & Jun, 2005) analysis was used to determine the influence
of individual OTUs in PLS. A high VIP value for an OTU indicates high importance in prediction
of the environmental response variable for that OTU. For network construction and visualization
purposes, the minimum correlation value required between two OTUs to constitute an "edge"
between them was delineated at different r values for each network related to an individual
environmental variable (ranging between 0.1 – 0.25), as co-correlations between OTUs within
some networks were stronger than others. The number of co-correlations an OTU has with other
OTUs within a network defines its "node centrality" (as described by Henson *et al.*, 2016).

3. Results

3.1 Chemical analyses

Significant correlations (r > 0.7, p ≤ 0.001) were found among NH_4^+, OM, OC, and TN.
Thus, downstream analyses combined these values into one parameter, "NUTR", represented by
OM values as this variable was the most strongly correlated with each of the other variables.
Environmental data were analyzed with a PCA and PC1 and PC2 explained 56.2% and 20.6% of
the variation among samples, respectively (Fig. b). perMANOVA found significant differences in
physicochemical profiles based on region ($R^2 = 0.570$, p ≤0.001) and depth ($R^2 = 0.058$, p ≤ 0.01).
Lake Erie coastal wetlands were chemically distinct from other wetland regions (ESBT and NSB;
adjusted p = 0.01) according to perMANOVA and pairwise perMANOVA based on Euclidean
distance. Ninety-five percent confidence intervals demonstrated no overlap between Lake Erie

coastal wetlands and other coastal wetlands (Fig. 2). This separation was related to increased
NUTR, NO_3^-, and S.
Increasing depth within cores showed a consistent shift in environmental variables, except
in those sites located in the western basin of Lake Erie (Supplemental Fig. 2). Specifically, OM,
OC, and TN consistently decreased with increasing depth within each region except Lake Erie.
Similarly, C:N increased with depth in each region except Lake Erie, wherein the C:N ratio
remained relatively low (~ 12) and stable with increasing soil depth. Within the Lake Erie wetland
region, pH was more acidic in the overlying water with respect to all other wetland regions
(Supplemental Table 1). However, pH was still relatively neutral within Lake Erie (average pH =
7.26 ± 0.24), whereas other wetland regions (regions within Saginaw Bay and Beaver Archipelago)
experienced slightly more basic pH, with average pH among these regions ranging between 7.72
– 8.39.

3.2 Alpha diversity

Sufficient depth of sampling was reinforced by rarefaction curve analysis (Supplemental
Fig. 3). Good's coverage values ranged between 89.3 – 93.5% for each region at the subsampled
value of 48,226 sequences. Chao1 richness estimates varied significantly among wetland regions
(F = 8.38, $p \leq 0.05$), as well as wetland sites (F = 16.78, $p \leq 0.001$). Pairwise comparisons found
that the LE region had significantly higher ($p \leq 0.01$) Chao1 estimates than NSB and WSB regions
(Fig. 3; Supplemental Table 2). Additionally, pairwise comparisons found a high degree of
significant variability ($p \leq 0.01$) in Chao1 estimates among wetland sites (Supplemental Table 2).
Further, Shannon diversity levels also significantly varied among wetland sites (F = 4.57, $p \leq$
0.001), with site LE_D having significantly higher ($p \leq 0.01$) Shannon diversity levels than sites
ESBT_A and WSB_B (Supplemental Table 2). Soil depth did not influence alpha diversity levels.

Shannon diversity and Chao1 were both positively correlated with measured environmental
variables (Table 1). Specifically, Chao1 estimates increased with NO_3^-, P, and S concentrations (p
≤ 0.01), and were weakly positively correlated ($p \leq 0.05$) with NUTR. Additionally, Shannon
diversity levels increased alongside NUTR and S ($p \leq 0.001$), and were weakly positively
correlated with NO_3^- ($p \leq 0.05$). There were no significant relationships between alpha diversity
and C:N, and alpha diversity was not negatively correlated with any of the measured environmental
variables.

3.3 Beta diversity
3.3.1 Beta diversity among regions

Multivariate analyses were implemented to explore relationships between microbial
communities and environmental gradients among wetland regions. NMDS demonstrated
separation of microbial communities based on wetland site, region, and soil depth (Fig. 4).
Substantiating this result, perMANOVA confirmed that differences in microbial community
structure were significantly related to wetland region ($R^2 = 0.220$, $p \leq 0.001$), site ($R^2 = 0.119$, p $\leq$
0.001), and soil depth ($R^2 = 0.070$; $p \leq 0.001$). Post-hoc pairwise perMANOVA found that
community structure within the LE region was significantly distinct ($p \leq 0.01$) from all other
wetland regions (Table 2). No significant differences in community structure were found between
any other wetland regions compared. Additionally, microbial community beta diversity was

distinct ($p \leq 0.003$) between the top soil depth and the middle and bottom soil depths. However, no significant differences in microbial community structure were found between the middle and bottom soil depths (Table 2). Variation in microbial community structure was significantly correlated ($p \leq 0.001$) to depth ($r = 0.41$), NO_3^- ($r = 0.20$), NUTR ($r = 0.60$), and S ($r = 0.41$), and also correlated ($p \leq 0.016$) with C:N ($r = 0.11$) and P ($r = 0.14$) (Supplemental Table 3).

Beta dispersion tests suggested significant variation in structural variance among regions ($p \leq 0.05$), however, Tukey's HSD test using adjusted p-values for multiple comparisons did not find any significance ($p > 0.05$) between pairwise comparisons of regional groups. There were no differences in community structural dispersion among soil depths.

3.3.2 Beta diversity within regions

Microbial community associations with environmental variables were also explored within regions to examine variation among wetland sites. Individual NMDS plots of each region identified relationships between microbial community structure and several environmental variables using vector-fitting regression, and strengths of these relationships were dependent upon the wetland region explored (Fig. 5; Supplemental Table 3). Depth was significantly related ($p \leq 0.05$) to microbial community structure in all wetland regions except NSB and LE. However, microbial community structure may have been more strongly related to depth in NSB ($r = 0.35$, $p = 0.071$) than LE ($r = 0.19$, $p = 0.40$). NUTR was significantly related ($p \leq 0.01$) to community structure within regions BA ($r = 0.82$), ESBT ($r = 0.51$), and LE ($r = 0.66$). C:N was related ($p \leq 0.01$) to community structure within regions of Saginaw Bay (i.e., ESBT [$r = 0.65$], NSB [$r = 0.58$], and WSB [$r = 0.58$]). Beta diversity was not significantly associated with concentrations of NO_3^- in any region.

To test for significant differences in microbial beta diversity within regions, perMANOVA was used to evaluate differences in microbial community structure among soil depths and sites within wetland regions (Supplemental Table 3). Depth did not significantly explain microbial community structure within the region LE ($p = 0.65$), however, it did explain differences in microbial community structure within the other wetland regions, specifically BA ($R^2 = 0.414$; $p = 0.006$), ESBT ($R^2 = 0.154$; $p = 0.001$), NSB ($R^2 = 0.161$; $p = 0.093$), and WSB ($R^2 = 0.259$; $p = 0.014$). Significant differences in microbial community structure were found among different wetland sites within regions ESBT ($R^2 = 0.192$; $p = 0.001$), LE ($R^2 = 0.236$; $p = 0.004$), and NSB ($R^2 = 0.140$; $p = 0.003$). As only one site was sampled within the BA region, testing for differences among wetland sites within the BA region could not be accomplished.

3.4 Network analyses

Weighted Correlation Network Analysis (WGCNA) was used to explore strong relationships between subcommunities and individual OTUs with environmental parameters within Great Lakes coastal wetlands. After removal of OTUs that did not have at least two representative sequences in at least 10% of samples, a total of 7,562 OTUs remained for WGCNA. In determining scale-free topology of the OTU network, a soft power threshold of 4 was reached, and an R^2 of 0.87 was established as linear fit from the regression of the frequency distribution of node connectivity against node connectivity (Supplemental Fig. 4). Of the 33 constructed

subnetworks, the same one (subnetwork "orange") was found to be most strongly correlated to
both NUTR (r = 0.94) and NO_3^- (r = 0.55) (Supplemental Fig. 5). A separate subnetwork ("pink")
was strongly correlated (r = 0.74) to C:N. All correlations of subnetworks to environmental
variables were significant (p ≤ 0.001). OTU VIP values ≤ 1 were removed due to the large amount
of OTUs within subnetworks correlated with C:N for visualization purposes.

For subnetwork relationships to NUTR (including OM, OC, NH_4^+, and TN), partial least
squares analysis (PLS) found that 69 OTUs were 93.8% predictive of variance in NUTR
(Supplemental Fig. 6). OTU co-correlation networks were constructed using an OTU co-
correlation threshold of 0.25, with strong correlations (r > 0.59) between all OTUs and NUTR
(Fig. 6). Of the top 15 OTUs contributing to PLS regression by VIP score, seven were related to
Betaproteobacteria, five were related to *Anaerolineaceae* (within *Chloroflexi*), and one
representative OTU was related to each of *Bellilinea* (*Chloroflexi*), *Desulfobacterales*
(*Deltaproteobacteria*), and *Rhizobiales* (*Alphaproteobacteria*).

For subnetwork relationships to C:N, PLS found that 144 OTUs were 59.0% predictive of
variance in C:N (Supplemental Fig. 7). Networks were constructed using an OTU co-correlation
threshold of 0.1, within positive or negative correlations (r > +/- 0.2) between OTUs (VIP > 1) and
C:N (Fig. 7). Of the top 15 OTUs by VIP score within the network, two OTUs related to
Bacteroidetes were negatively correlated with C:N. Other top OTUs were positively related to
C:N, including seven OTUs related to *Anaerolineaceae*, four OTUs which were unclassified
Bacteria, and one representative OTU related to each of *Bacillus* (*Firmicutes*) and *Chloroflexi*.

4. Discussion

4.1 Microbial diversity driven by chemistry within Great Lakes coastal wetlands

This study is the first to suggest that anthropogenic disturbance patterns correspond to
microbial community differences in Great Lakes coastal wetlands as consistent with other
taxonomic groups such as plants, birds, fish, and invertebrates (Howe *et al*., 2007; Tulbure *et al*.,
2007; Uzarski *et al*., 2009; Cooper *et al*., 2012; Uzarski *et al*., 2017). Microbial community
structure was significantly dissimilar between LE and all other wetland regions, and these
differences were related to physicochemical differences among coastal wetlands (Fig. 2, Fig. 4,
Table 2). As the wetlands within the Lake Erie region maintained the highest nutrient
concentrations within the soil, it is possible that anthropogenic stressors related to nutrient loading
(and potentially other pollutants) could be driving structural differences in microbial communities
among Great Lakes coastal wetlands. Further, network analysis found several taxa/sub
communities that were highly correlated to nutrient levels across wetlands explored in this study.
Previous research has found that nutrient levels (e.g., C, N, P, etc.), to varying degrees, can
influence microbial community composition and structure (Hartman *et al*., 2008; Peralta *et al*.,
2013; Ligi *et al*., 2014; Arroyo *et al*., 2015). Lake Erie coastal wetlands (and the watershed which
drains into them) have been historically impacted by anthropogenic pollution and agricultural
practices, particularly in comparison to other coastal wetlands within the Laurentian Great Lakes
region. This has been demonstrated by multiple ecological indices (e.g., Cvetkovic & Chow-
Fraser, 2011; Uzarski *et al*., 2017) and physicochemical uniqueness (increased levels of nutrients
and particulate matter) within the western basin of Lake Erie (Danz *et al*., 2007; Trebitz *et al*.,

2007; Cvetkovic & Chow-Fraser, 2011; Uzarski *et al*., 2017). Data presented in this study
corroborate this historical evidence of human impact and nutrient loading in the western basin of
Lake Erie (Fig. 2., Supplemental Fig. 1), which may be influencing the Lake Erie wetlands
explored in this study.

High nutrient influx could also be influencing the chemical and microbial vertical structure
within coastal wetland soils. Microbial community and chemical (e.g., C, N, P) vertical structure
was not evident within the first 6 cm of soil of coastal wetlands with elevated nutrient levels (e.g.
Lake Erie sites). The lack of vertical chemical gradients is unlikely to exclusively explain a
corresponding lack of vertical microbial community structure, as some wetland sites lower in
nutrient levels also did not experience vertical chemical gradients in this study (e.g. West Saginaw
Bay). One possibility is that a lack of vertical chemical structure in conjunction with high nutrient
levels in wetland soils could reduce vertical microbial community structure. It has been previously
demonstrated that concentrations of carbon electron donors may influence redox gradients within
wetland soils (Achtnich *et al*., 1995), and wetland microbial communities have been demonstrated
to correspond with soil redox gradients (Lüdemann *et al*., 2000; Edlund *et al*., 2008; Lipson *et al*.,
2015). However, connections between microbial community metabolic shifts with soil depth and
levels of dissolved organic carbon *in situ* remain unresolved in freshwater wetlands (Alewell *et*
al., 2008). Alternatively, another explanation for lack of vertical community structure could be
microsite heterogeneity throughout the soil matrix. Previous research in freshwater wetland soils
has suggested that microsite heterogeneity may explain coexistence of microbial functional guilds
(Alewell *et al*., 2008; Angle *et al*., 2017), which could substantially reduce vertical microbial
community structural gradients. However, it is necessary to better link microbial community
diversity, microbial activity, chemical structure, and microsite heterogeneity to establish
relationships between microbial communities and freshwater soil structure. As a caveat, it is
possible that chemical and microbial structuring still exists within wetlands with high nutrient
levels, yet is not evident within the first 6 cm of soil or at the spatial scale measured in this study.
Nevertheless, microbial communities within coastal wetlands with high nutrient levels did not
follow the same pattern of vertical structure evident in other comparable coastal wetlands, either
chemically or biologically, further suggesting that the microbial integrity of coastal wetland
systems may be susceptible to negative anthropogenic pressure.

While relationships between microbial diversity and nutrient levels among coastal
wetlands are strong, other unexplored variables unique to Lake Erie (such as geologic history)
could also be influencing uniqueness of chemical and microbial profiles in Lake Erie coastal
wetlands. The Lake Erie coastal wetland sites explored here were barrier (protected) wetlands,
while other wetland sites explored in this study are all classified as lacustrine (open water)
wetlands (www.greatlakeswetlands.org). As such, wave action from the Great Lakes impacted
wetlands within the western basin of Lake Erie to a lesser degree than other wetlands, thereby
reducing sediment export rates into the Great Lakes themselves. Hydrologic energy was found to
impact wetland primary productivity and respiration in Lake Huron coastal wetlands, suggesting
Great Lakes ecosystems may exert unique environmental forces on wetland microbial
communities (Cooper *et al.,* 2013). Low carbon export rates or elevated sedimentation rates may
exist in the western basin of Lake Erie as consequence of low wave action in these wetlands, which
may influence the chemical and biological structure (such as vertical microbial community
structure) within wetland soils of this region. Nevertheless, previous research at the same wetland

locations explored in this study have demonstrated that wetlands within the western basin of Lake
Erie are highly degraded with respect to other wetlands (Uzarski *et al.*, 2017), particularly with
respect to physicochemical conditions. Additionally, the same vegetation zone (dominated by
cattails or bulrush) was sampled among all wetlands explored in this study as an attempt to reduce
bias in distinct environmental conditions which may exist in other vegetation zones among wetland
sites. Burton et al. (2002) suggested that soil organic content was related to plant zonation in Great
Lakes coastal wetlands. Further research would be necessary to fully tease apart the effects of
anthropogenic stress and other natural contributions to differences in microbial communities
among coastal wetlands.
534
535
536 **4.2 Relationships between microbial subnetworks and environmental gradients**
537
Through network analyses, multiple subcommunities were delineated which were
significantly related to environmental gradients (such as nutrients C, N, and P) among coastal
wetlands sampled in this study. Specifically, a subnetwork of 69 microbial taxa was 93.8%
predictive of nutrient level variation among coastal wetland soils. Several microbial taxa within
this subcommunity were individually predictive of nutrient levels to a high degree, including
several OTUs related to *Anaerolineaceae*, one OTU related to genus *Anaerolinea*, and another
related to genus *Bellilinea*. From the genus *Anaerolinea*, two thermophilic chemoorganotrophs
(*Anaerolinea thermophila* and *Anaerolinea thermolimosa*) have been isolated (Sekiguchi *et al.*,
2003; Yamada *et al.*, 2006). Only one isolated member has been established within the genus
Bellilinea (*Bellilinea caldifistulae*); it has been described as a thermophilic, fermentative, obligate
anaerobe which thrives in co-culture with methanogens (Yamada *et al.*, 2007). It is unlikely that
the OTUs found in our study are the same species as the isolated *Anaerolinea* and *Bellilinea*
species, as coastal wetland soils are not high-temperature environments necessary for thermophilic
species. Additionally, no OTUs related to methanogenic archaea were found within this
subnetwork, suggesting that *Anaerolineacea* OTUs within coastal wetland soils may fluctuate
independently of any specific methanogenic OTUs. It is possible that the *Bellilinea* OTU found
within the subnetwork is related to nutrient level concentrations. This would support fermentative
metabolism as noted within *Bellilinea caldifistulae*. It is important to note that several other studies
have discovered OTUs related to *Anaerolineaceae* within wetland soils, with upwards of 90%
relative abundance among *Chloroflexi* OTUs within these systems (Ansola *et al.*, 2014; Deng *et*
al., 2014; Hu *et al.*, 2016). This suggests that there are probable mesophilic species yet to be
isolated within this ubiquitous family of bacteria, which may be of high importance within wetland
soils.
561
Betaproteobacteria were also found to significantly predict nutrient levels among coastal
wetlands. Hu *et al.* (2016) found that both *Betaproteobacteria* and *Anaerolineae* were positively
related to TN levels, which is consistent with the data presented here, and these two taxa were
suggested to contribute to higher levels of heterotrophic activity. Further, *Anaerolineaceae* OTUs
were consistently related to increasing C:N, suggesting that many taxa within this family have
preference for recalcitrant carbon sources. As C:N also tends to increase with soil depth, it is also
probable that the putatively obligate anaerobic *Anaerolineaceae* are coinciding with decreasing
oxygen levels and/or changing metabolism requirements with increasing soil depth.
570

Development of biological indices and establishment of indicator taxa have been suggested
as necessary for microbial communities within wetlands (Uzarski *et al.*, 2017), particularly through
the use of high-throughput sequencing technologies which now allow for deep assessment of
microbial community composition and structure within environmental samples (Sims *et al.*, 2013;
Urakawa & Bernhard, 2017). Specifically within Great Lakes coastal wetlands, it is integral to
develop ecosystem health indicators based upon multiple different groups of taxonomy, as separate
biological indices can present contrasting assessments of wetland health (Uzarski *et al.*, 2017). As
microbial indicators have yet to be established in Great Lakes coastal wetlands, this research
begins the first steps in exploring how microbial communities can be used as an additional and
potentially important ecosystem health indicator. In addition to their importance as biological
signals for environmental health, microbial indicator taxa may play prominent roles in
bioremediation of excess nutrients and pollutants found within anthropogenically impacted coastal
wetlands. Network analyses in this study have allowed for the generation of hypothetical
subcommunities of diverse microbial taxa related to nutrient levels among Great Lakes coastal
wetlands, and could assist in further understanding of which microbial taxa may be responding to
anthropogenic stress in these ecosystems.

5. Conclusions

This study marks the first characterization of microbial communities within Great Lakes
coastal wetlands. Coastal wetlands are integral in the proper functioning of biogeochemical cycles
and environmental sustainability of the Great Lakes. While it has long been known that
anthropogenic pressure can impact animal and plant communities within these coastal wetlands,
this is the first evidence that these pressures may also be influencing microbial communities and
may be influencing biogeochemical cycles by extension. Alpha and beta diversity were both
related to nutrient gradients among and within regions, suggesting that variability in microbial
community structure is highly coupled to geochemistry within wetland soils. We propose that
wetland microbial community structure can also potentially be used to assess a wetland for
monitoring purposes. As illustrated within this study, wetland microbial community structure and
depth are decoupled within the wetlands experiencing the highest nutrient levels, likely originating
from terrestrial inputs due to human activity. As such, multivariate statistics (as used in the
methods of this study) may prove useful in examining relationships between wetland soil depth
and microbial community structure alongside microbial network analyses, which could provide
biological indicators of nutrient loading stress on coastal wetland habitats. We propose that
wetland microbial community structure can also potentially be used to assess a wetland for
monitoring purposes.

Further, this study provides insight on microbial community subnetworks and individual
OTUs, which were predictive of chemical concentrations, and may be useful for future
management of Great Lakes coastal wetland systems. Within subnetworks existed multiple taxa
with strong individual relationships to environmental gradients among coastal wetlands throughout
the Great Lakes. Even further, several community members within these subnetworks were
taxonomically related (such as OTUs related to *Anaerolineaceae* within *Chloroflexi*), suggesting
that specific taxonomic groups of microbes may be useful to explore further as potential biological
indicator groups. This study highlights the strength of network analyses (such as WGCNA) in

delineating hypothetical networks of interacting microbes, and whether these networks are
predictive of physical or chemical gradients measured within an environment.
618

619 **Acknowledgements**
620
Special thanks to Alexandra Bozimowski and Thomas Langer for their assistance in
collecting the wetland samples and data acquisition, as well as Dr. Matthew Cooper for providing
valuable insight and conversation on wetland ecology and the custom-built core extruder built by
Mr. Gary Cooper. Special thanks also to Mike Henson for providing assistance on network
statistical analyses. This paper is Contribution Number XX of the Central Michigan University
Institute for Great Lakes Research.
627

628 **Author Contributions Statement**
629

630 DH processed samples, analyzed data, and wrote the manuscript. DU provided access to the
631 samples, aided sample selection, and statistical analysis. DL and KT provided guidance on
632 microbial analysis and KT, DU, and DL assisted in editing the manuscript.
633

634 **Conflict of Interest Statement**
635

636   The authors declare no conflicts of interest.
637

638 **Funding Statement**
639
Funding was provided, in part, by the CMU College of Science and Engineering.
Additional funding for this work was provided by the Great Lakes National Program Office under
the United States Environmental Protection Agency, grant number GL-00E00612-0 as part of the
US federal government's Great Lakes Restoration Initiative. Although the research described in
this work has been partly funded by the United States Environmental Protection Agency, it has not
been subjected to the agency's required peer and policy review, and therefore, does not necessarily
reflect the views of the agency and no official endorsement should be inferred.
647

648 **Data Availability Statement**
649

650 Sequence data generated for this study can be found in the GenBank repository at
651 https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA417157. Other data generated for this
652 study, along with R code for replication of statistical methodology, can be found in the GitHub
653 repository at https://github.com/horto2dj/GLCW/.
654

655 **References**
656

657 Achtnich, C., Bak, F., and Conrad, R. (1995). Competition for electron donors among nitrate
658 reducers, ferric iron reducers, sulfate reducers, and methanogens in anoxic paddy soil. Biol Fert
659 Soils. 19(1), 65-72.
660

Alewell, C., Paul, S., Lischeid, G., and Storck, F.R. (2008). Co-regulation of redox processes in freshwater wetlands as a function of organic matter availability?. Sci Total Environ 404(2), 335-342.

Anderson, M.J. (2001). A new method for non- parametric multivariate analysis of variance. Austral Ecol 26(1), 32-46.

Anderson, M.J. (2006). Distance- based tests for homogeneity of multivariate dispersions. Biometrics 62(1), 245-253.

Anderson, M.J., Ellingsen, K.E., and McArdle, B.H. (2006). Multivariate dispersion as a measure of beta diversity. Ecol Lett 9(6), 683-693.

Anderson, M.J., and Walsh, D.C. (2013). PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing?. Ecol Monogr 83(4), 557-574.

Angle, J.C., Morin, T.H., Solden, L.M., Narrowe, A.B., Smith, G.J., Borton, M.A., *et al*. (2017). Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions. Nat Commun 8(1), 1567.

Ansola, G., Arroyo, P., and de Miera, L.E.S. (2014). Characterisation of the soil bacterial community structure and composition of natural and constructed wetlands. Sci Total Environ 473, 63-71.

Arroyo, P., de Miera, L.E.S., and Ansola, G. (2015). Influence of environmental variables on the structure and composition of soil bacterial communities in natural and constructed wetlands. Sci Total Environ 506, 380-390.

Burton, T.M., Stricker, C.A., and Uzarski, D.G. (2002). Effects of plant community composition and exposure to wave action on invertebrate habitat use of Lake Huron coastal wetlands. Lakes & Reservoirs: Research and Management. 7: 255-269.

Campbell, M., Cooper, M.J., Friedman, K., and Anderson, W.P. (2015). The economy as a driver of change in the Great Lakes–St. Lawrence River basin. J Great Lakes Res 41, 69-83.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., *et al*. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J 6(8), 1621-1624.

Chong, I.G., and Jun, C.H. (2005). Performance of some variable selection methods when multicollinearity is present. Chemometr Intell Lab 78(1), 103-112.

Clarke, K.R. (1993). Non-parametric multivariate analyses of changes in community structure. Aust J Ecol 18, 117-143.

707   Cole J.R., Wang Q., Fish J.A., Chai B., Mcgarrell D.M., Sun Y., *et al*. (2013) Ribosomal Database
708   Project: data and tools for high throughput rRNA analysis. Nucleic Acids Res 42(D1),
709   doi:10.1093/nar/gkt1244
710
711   Conrad, R. (1996). Soil microorganisms as controllers of atmospheric trace gases (H_2, CO, CH_4,
712   OCS, N_2O, and NO). Microbiol Rev 60(4), 609-640.
713
714   Cooper, M.J., Gyekis, K.F., and Uzarski, D.G. (2012). Edge effects on abiotic conditions,
715   zooplankton, macroinvertebrates, and larval fishes in Great Lakes fringing marshes. J Great Lakes
716   Res 38(1), 142-151.
717
718   Cooper, M.J., Steinman, A.D., and Uzarski, D.G. (2013). Influence of geomorphic setting on the
719   metabolism of Lake Huron fringing wetlands. Limnol Oceanogr 58(2), 452-464.
720
721   Cvetkovic, M., and Chow-Fraser, P. (2011). Use of ecological indicators to assess the quality of
722   Great Lakes coastal wetlands. Ecological Indicators 11(6), 1609-1622.
723
724   Dahl, T.E. (1990). Wetlands losses in the United States 1780's to 1980's. U.S. Department of the
725   Interior, Fish and Wildlife Service, Washington. D.C. 13pp.
726
727   Danz, N.P., Niemi, G.J., Regal, R.R., Hollenhorst, T., Johnson, L.B., Hanowski, J.M., *et al*. (2007).
728   Integrated measures of anthropogenic stress in the US Great Lakes basin. Environ Manage 39(5),
729   631-647.
730
731   Deng, Y., Cui, X., Hernández, M., and Dumont, M.G. (2014). Microbial diversity in hummock
732   and hollow soils of three wetlands on the Qinghai-Tibetan Plateau revealed by 16S rRNA
733   pyrosequencing. PLoS One 9(7), e103115.
734
735   Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011). UCHIME improves
736   sensitivity and speed of chimera detection. Bioinformatics 27(16), 2194-2200.
737
738   Edlund, A., Hårdeman, F., Jansson, J.K., and Sjöling, S. (2008). Active bacterial community
739   structure along vertical redox gradients in Baltic Sea sediment. Environ Microbiol 10(8), 2051-
740   2063.
741
742   Faust, K., & Raes, J. (2012). Microbial interactions: from networks to models. Nat Rev Microbiol
743   10(8), 538.
744
745   Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., *et al*. (2016). Plankton
746   networks driving carbon export in the oligotrophic ocean. Nature 532(7600), 465.
747
748   Hackett, R.A., Babos, H.B., Collins, E.E., Horton, D., Schock, N., and Schoen, L.S. (2017).
749   Researcher disciplines and the assessment techniques used to evaluate Laurentian Great Lakes
750   coastal ecosystems. J Great Lakes Res 43(1), 9-16.
751

Hartman, W.H., Richardson, C.J., Vilgalys, R., and Bruland, G.L. (2008). Environmental and
anthropogenic controls over bacterial communities in wetland soils. P Natl Acad Sci 105(46),
17842-17847.

Hecnar, S.J. (2004). Great Lakes wetlands as amphibian habitats: a review. Aquat Ecosyst Health
7(2), 289-303.

Henson, M.W., Hanssen, J., Spooner, G., Flemming, P., Pukonen, M., Stahr, F., and Thrash, J.C.
(2016). Nutrient dynamics and stream order influence microbial community patterns along a 2914
km transect of the Mississippi River. bioRxiv, 091512.

Hill, B.H., Elonen, C.M., Jicha, T.M., Cotter, A.M., Trebitz, A.S., and Danz, N.P. (2006).
Sediment microbial enzyme activity as an indicator of nutrient limitation in Great Lakes coastal
wetlands. Freshwater Biol 51(9), 1670-1683.

Horton, D.J., Theis, K.R., Uzarski, D.G., Learman, D.R. Data from: Microbial community
structure corresponds to nutrient gradients and human impact within coastal wetlands of the Great
Lakes. GenBank. (2017). Accession: PRJNA417157

Howe, R.W., Regal, R.R., Hanowski, J., Niemi, G.J., Danz, N.P., and Smith, C.R. (2007). An
index of ecological condition based on bird assemblages in Great Lakes coastal wetlands. J Great
Lakes Res 33(sp3), 93-105.

Hu, Y., Wang, L., Fu, X., Yan, J., Wu, J., Tsang, Y., *et al*. (2016). Salinity and nutrient contents
of tidal water affects soil respiration and carbon sequestration of high and low tidal flats of
Jiuduansha wetlands in different ways. Sci Total Environ 565, 637-648.

Jackson, C.R., Foreman, C.M., and Sinsabaugh, R.L. (1995). Microbial enzyme activities as
indicators of organic matter processing rates in a Lake Erie coastal wetland. Freshwater biol 34(2),
329-342.

Kögel-Knabner, I., Amelung, W., Cao, Z., Fiedler, S., Frenzel, P., Jahn, R., *et al*. (2010).
Biogeochemistry of paddy soils. Geoderma 157(1), 1-14.

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D. (2013) Development of
a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data
on the MiSeq Illumina Sequencing Platform. Appl Environ Microb 79(17), 5112-5120.

Krantzberg, G., and De Boer, C. (2008). A valuation of ecological services in the Laurentian Great
Lakes Basin with an emphasis on Canada. Am Wat Works Assoc J 100(6), 100.

Lamers, L.P., Van Diggelen, J.M., den Camp, H.J.O., Visser, E.J., Lucassen, E.C., Vile, M.A., *et
al*. (2012). Microbial transformations of nitrogen, sulfur, and iron dictate vegetation composition
in wetlands: a review. Front Microbiol, 3:156.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics 9(1), 559.

Langfelder, P., and Horvath, S. (2012). Fast R functions for robust correlations and hierarchical clustering. J Stat Softw 46(11).

Ligi, T., Oopkaup, K., Truu, M., Preem, J.K., Nõlvak, H., Mitsch, W.J., *et al*. (2014). Characterization of bacterial communities in soil and sediment of a created riverine wetland complex using high-throughput 16S rRNA amplicon sequencing. Ecol Eng 72, 56-66.

Lipson, D.A., Raab, T.K., Parker, M., Kelley, S.T., Brislawn, C.J., and Jansson, J. (2015). Changes in microbial communities along redox gradients in polygonized Arctic wet tundra soils. Env Microbiol Rep 7(4), 649-657.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15(12), 550.

Lüdemann, H., Arth, I., and Liesack, W. (2000). Spatial changes in the bacterial community structure along a vertical oxygen gradient in flooded paddy soil cores. Appl Environ Microb 66(2), 754-762.

McMurdie, P.J., and Holmes, S. (2013) Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8(4).

McMurdie, P.J., and Holmes, S. (2014). Waste not, want not: why rarefying microbiome data is inadmissible. PLoS Comput Biol, 10(4), e1003531.

Morrice, J.A., Danz, N.P., Regal, R.R., Kelly, J.R., Niemi, G.J., Reavie, E.D., *et al*. (2008). Human influences on water quality in Great Lakes coastal wetlands. Environ Manage 41(3), 347-357.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J., and Suggests, M.A.S.S. (2007). The vegan package. Community ecology package, 10, 631-637.

Peralta, R.M., Ahn, C., and Gillevet, P.M. (2013). Characterization of soil bacterial community structure and physicochemical properties in created and natural wetlands. Sci Total Environ 443, 725-732.

Quast C., Pruesse E., Yilmaz P., Gerken J., Schweer T., Yarza P., *et al*. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41(D1), D590-D596.

R Core Team (2015) R: A language and environment for statistical computing. Vienna, Austria. https://www.R-project.org/.

Reddy, K.R., and DeLaune, R.D. (2008). Biogeochemistry of wetlands: science and applications. Boca Raton, FL: CRC Press.

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., *et al*. (2009).
Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for
Describing and Comparing Microbial Communities. Appl Environ Microbiol 75(23), 7537-7541.

Shade, A., Peter, H., Allison, S. D., Baho, D., Berga, M., Bürgmann, H., *et al*. (2012).
Fundamentals of microbial community resistance and resilience. Front Microbiol 3, 417.

Sekiguchi, Y., Yamada, T., Hanada, S., Ohashi, A., Harada, H., and Kamagata, Y. (2003).
Anaerolinea thermophila gen. nov., sp. nov. and Caldilinea aerophila gen. nov., sp. nov., novel
filamentous thermophiles that represent a previously uncultured lineage of the domain Bacteria at
the subphylum level. Int J Syst Evol Micr 53(6), 1843-1851.

Sierszen, M.E., Morrice, J.A., Trebitz, A.S., and Hoffman, J.C. (2012). A review of selected
ecosystem services provided by coastal wetlands of the Laurentian Great Lakes. Aquat Ecosyst
Health 15(1), 92-106.

Sims, A., Zhang, Y., Gajaraj, S., Brown, P.B., and Hu, Z. (2013). Toward the development of
microbial indicators for wetland assessment. Water Res 47(5), 1711-1725.

Tang, Y.S., Wang, L., Jia, J.W., Fu, X.H., Le, Y.Q., Chen, X.Z., and Sun, Y. (2011). Response of
soil microbial community in Jiuduansha wetland to different successional stages and its
implications for soil microbial respiration and carbon turnover. Soil Biol Biochem 43(3), 638-646.

Trebitz, A.S., Brazner, J.C., Cotter, A.M., Knuth, M.L., Morrice, J.A., Peterson, G.S., *et al*. (2007).
Water quality in Great Lakes coastal wetlands: basin-wide patterns and responses to an
anthropogenic disturbance gradient. J Great Lakes Res 33(sp3), 67-85.

Tulbure, M.G., Johnston, C.A., and Auger, D.L. (2007). Rapid invasion of a Great Lakes coastal
wetland by non-native Phragmites australis and Typha. J Great Lakes Res 33(sp3), 269-279.

Urakawa, H., and Bernhard, A.E. (2017). Wetland management using microbial indicators. Ecol
Eng 108(B), 456-476.

Uzarski, D.G. (2009). Wetlands of Large Lakes. Encyclopedia of Inland Waters. Oxford: Elsevier.
p. 599-606.

Uzarski, D.G., Brady, V.J., Cooper, M.J., Wilcox, D.A., Albert, D.A., Axler, R.P, *et al*. (2017).
Standardized measures of coastal wetland condition: Implementation at a Laurentian Great Lakes
basin-wide scale. Wetlands 37(1), 15-32.

Uzarski, D.G., Burton, T.M., Kolar, R.E., and Cooper, M.J. (2009). The ecological impacts of
fragmentation and vegetation removal in Lake Huron's coastal wetlands. Aquat Ecosyst Health
12(1), 45-62.

Wang, N., and Mitsch, W.J. (1998). Estimating phosphorus retention of existing and restored
coastal wetlands in a tributary watershed of the Laurentian Great Lakes in Michigan, USA. Wetl
Ecol Manag 6(1), 69-82.
891
892 Yamada, T., Sekiguchi, Y., Hanada, S., Imachi, H., Ohashi, A., Harada, H., and Kamagata, Y.
893 (2006). Anaerolinea thermolimosa sp. nov., Levilinea saccharolytica gen. nov., sp. nov. and
894 Leptolinea tardivitalis gen. nov., sp. nov., novel filamentous anaerobes, and description of the new
895 classes Anaerolineae classis nov. and Caldilineae classis nov. in the bacterial phylum Chloroflexi.
896 Int J Syst Evol Micr 56(6), 1331-1340.
897
898 Yamada, T., Imachi, H., Ohashi, A., Harada, H., Hanada, S., Kamagata, Y., and Sekiguchi, Y.
899 (2007). Bellilinea caldifistulae gen. nov., sp. nov. and Longilinea arvoryzae gen. nov., sp. nov.,
900 strictly anaerobic, filamentous bacteria of the phylum Chloroflexi isolated from methanogenic
901 propionate-degrading consortia. Int J Syst Evol Micr 57(10), 2299-2306.
902
903 Zhang, B., and Horvath, S. (2005). A general framework for weighted gene co-expression network
904 analysis. Stat Appl Genet Mol 4(1).

910 **Tables**

912 Table 1. Correlations between alpha diversity metrics and measured environmental variables.

913 Asterisks represent significance values where $p \leq 0.001$ (***), $p \leq 0.01$ (**), and $p \leq 0.05$ (*).

	Chao1	Shannon
P	0.31**	-0.09
S	0.42***	0.45***
NO_3^-	0.42***	0.24*
C:N	-0.03	-0.2
NUTR	0.24*	0.41***

916 Table 2. Pairwise perMANOVA results comparing pairwise differences between wetland regions

917 and differences between wetland soil depths. Values represent significant ($p \leq 0.01$) R^2 results,

918 and *n.s.* represents lack of significance ($p > 0.01$).

Region	BA	ESBT	LE	NSB	WSB
BA	-				
ESBT	*n.s.*	-			
LE	0.507	0.401	-		
NSB	*n.s.*	*n.s.*	0.524	-	
WSB	*n.s.*	*n.s.*	0.435	*n.s.*	-

Depth	Top	Middle	Bottom
Top	-		
Middle	**	-	
Bottom	**	*n.s.*	-

Figures

Figure 1. Geographic map displaying location of sites sampled within this study. Colors of points
correspond to region sampled.

Figure 2. Principal Component Analysis (PCA) illustrating separation of samples based upon soil
geochemistry. Shapes and colors correspond to different wetland depths and regions, respectively,
as listed in the legend. Percentages on axes represent explained variance of that principal
component. Vectors represent impact of specific environmental variables on sample distribution.
NUTR represents OM values, which correlated significantly ($p \leq .01$, $r > 0.56$) to NO_3^-, OC, OM,
S, and TN. Ellipses represent 95% confidence intervals of region groupings.

Figure 3. Boxplot diagram comparing Chao1 diversity among wetland regions. Boxes with the
same letter are not significantly different, while those with no common letters are significantly
different ($p \leq 0.01$). Lines within boxes represent the median, hinges represent +/- 25% quartiles,
whiskers represent up to 1.5x the interquartile range. Colors represent wetland region.

Figure 4. Nonmetric Multidimensional Scaling (NMDS) plot illustrating separation of samples
based upon differences in microbial community structure. Shapes and colors correspond to
different depths and wetland regions, respectively, as listed in the legend. Vectors represent
correlations of environmental variables to the distribution of the microbial communities
represented in the plot.

Figure 5. NMDS plots of each wetland region demonstrating separation of samples based upon
differences in microbial community structure, including **(A)** BA, **(B)** ESBT, **(C)** LE, **(D)** NSB, and
(E) WSB. Shapes and colors correspond to different depths and wetland sites, respectively, as
listed in the legends. Vectors represent correlations of environmental variables to the distribution
of microbial communities represented in the plots.

Figure 6. Network visualization and results of partial least squares analysis on the subnetwork
most correlated with NUTR. The y-axis represents correlation of OTU to OC values, whereas the
x-axis represents the node centrality. Points represent OTUs, and the color of points corresponds
to the phylum to which an OTU belongs. Point size corresponds to VIP score of that OTU. The
top 15 OTUs are labeled within the graph with corresponding lowest taxonomic identification
possible, and the level of that classification. D = Domain; P = Phylum, C = Class, O = Order, F =
Family, G = Genus.

Figure 7. Network visualization and results of partial least squares analysis on the subnetwork
most correlated with C:N. The y-axis represents correlation of OTU to C:N, whereas the x-axis
represents the node centrality. Points represent OTUs, and the color of points corresponds to the
phylum to which an OTU belongs. Point size corresponds to VIP score of that OTU. Only OTUs
with a VIP score > 1 were displayed for visualization purposes. The top 15 OTUs are labeled
within the graph with corresponding lowest taxonomic identification possible, and the level of that
classification. D = Domain; P = Phylum, C = Class, O = Order, F = Family, G = Genus.

Figure 7.JPEG