Title: Microbial community structure corresponds to nutrient gradients and human impact within

coastal wetlands of the Great Lakes.

Dean J. Horton[1], Kevin R. Theis[2], Donald G. Uzarski[1], and Deric R. Learman[1]*

[1]Institute for Great Lakes Research and Department of Biology, Central Michigan University,

Mt. Pleasant, MI, 48859, U.S.A.

[2]Department of Microbiology, Immunology, and Biochemistry, Wayne State University, Detroit,

MI 48201, U.S.A

*Corresponding author: Department of Biology, 3111 Biosciences Building, Central Michigan

University, Mt. Pleasant, MI, 48859. Email: deric.learman@cmich.edu; Telephone: 989-774-

4409

Originality-Significance Statement

This research is original in providing an initial, geographically wide-ranging

characterization of microbial communities of the Great Lakes coastal wetlands, an understudied

system of wetlands directly bordering the North American Laurentian Great Lakes. Relationships

between microbial communities within coastal wetland soils and geochemistry highlighted that

anthropogenic impacts may be significantly influencing microbial community structure, as well

as unique subnetworks of microbial taxa. Negative anthropogenic impacts to these coastal

wetland communities could influence natural biogeochemical cycles which occur within coastal

wetland soils, and by extension would directly influence the Great Lakes themselves.

Summary

Microbial communities within the soil of Great Lakes coastal wetlands drive

biogeochemical cycles and provide several other ecosystems services. However, there exists a

lack of understanding of how microbial communities respond to nutrient gradients and human

activity in these systems. This research sought to address this lack of understanding through

exploration of microbial community diversity and networks among coastal wetlands throughout

the Great Lakes. Significant differences in microbial community structure were illuminated

between Lake Erie and all other wetlands, and chemical and biological structure did not vary

within Lake Erie with increasing soil depth. Beyond this, alpha diversity levels were highest

within Lake Erie coastal wetlands. These diversity differences coincided with higher nutrient

levels within the Lake Erie region. Site-to-site variability existed within Lake Erie, East and

North Saginaw Bay regions, suggesting site-scale heterogeneity may impact microbial

community structure. Several subnetworks of microbial communities and individual OTUs were

related to chemical gradients among wetland regions, revealing several candidate indicator

communities and taxa which may be useful for Great Lakes coastal wetland management. This

research provides an initial characterization of microbial communities among Great Lakes

coastal wetlands, and demonstrates that microbial communities could be negatively impacted by

anthropogenic activities.

Introduction

The Laurentian Great Lakes of North America are one of the largest freshwater systems

on Earth, and are critical in supporting biogeochemical cycles, freshwater resources,

biodiversity, and economic viability of the surrounding region. Coastal wetlands bordering the

Great Lakes are integral in sustaining the proper functioning of the Great Lakes themselves,

making up nearly 200,000 ha of habitat between the United States and Canada despite reduction

of habitat by approximately 50% since European colonization (Dahl, 1990; Hecnar, 2004;

Sierszen *et al.*, 2012). In addition, Great Lakes coastal wetlands are vital in the retention of

chemical pollutants (e.g., heavy metals), sediments, and excess nutrients (e.g., N and P) caused

by anthropogenic activity, and provide critical habitat for countless biological species (Wang &

Mitsch, 1998; Sierszen *et al.*, 2012). The economy of the Great Lakes is contingent on the

existence and proper functioning of coastal wetlands. In providing ecosystem services and

promoting biodiversity, these wetlands have an estimated annual worth of $69 billion USD; the

value of recreational fishing alone is valued at $7.4 billion USD per year (Krantzberg & de Boer,

2008; Campbell *et al.*, 2015). Notably, the Great Lakes region has been impacted by negative

anthropogenic pressure, with cumulative stress having a particular impact on the western basin of

Lake Erie (Danz *et al.*, 2007; Uzarski *et al.*, 2017). These negative impacts extend to the coastal

wetlands, as pressure from agricultural runoff, atmospheric deposition, and urbanization

influence water chemistry, thereby reducing water quality and impairing wetlands (Trebitz *et al.*,

2007; Morrice *et al.*, 2008). This has stoked a surge in research assessing biodiversity and

anthropogenic pressure on coastal wetlands of the Great Lakes since the Great Lakes Water

Quality Agreement (GLWQA) was established in 1972 (Hackett *et al.*, 2017). While much

research on coastal wetlands has flourished in the wake of this international agreement, microbial

communities within Great Lakes coastal wetlands remain almost entirely uncharacterized

(Hackett *et al.*, 2017).

As ecological transition zones between upland and aquatic environments (Uzarski, 2009),

wetlands host a unique suite of chemical cycles while providing ecosystem services to bordering

environments. Most notably, carbon mineralization occurs within wetland soils via redox

processes mediated by microbial communities, and these processes contribute to pollution

mitigation and atmospheric greenhouse gas flux (Conrad, 1996; Reddy & DeLaune, 2008).

Wetland soil often becomes chemically structured with increasing depth through sequential

reduction of electron acceptors that decrease in metabolic favorability to microbes due to

thermodynamic constraints (Conrad, 1996; Reddy & DeLaune, 2008; Kögel-Knabner *et al.*,

2010). As microbial community metabolism will change in concert with soil chemical profiles,

microbial community structural shifts commonly result (Lüdemann *et al.*, 2000; Edlund *et al.*,

2008; Lipson *et al.*, 2015). However, concentration of carbon electron donors can influence the

vertical stratification of redox processes (Achtnich *et al.*, 1995; Alewell *et al.*, 2008), and by

extension, vertical microbial community structure (defined as relative proportions of microbial

taxa within a community). As an example of how this may apply to natural environments,

increased nutrient influx from anthropogenic activities (such as agricultural pressure) may

impact microbial community structure within coastal wetlands, which may extend to redox

processes. While sequential reduction chains and vertical microbial community structuring can

be predictable in lab experiments, complex heterogeneity within wetland soil can often influence

microbial community processes and structure to a large degree (Alewell *et al.*, 2008; Lamers *et*

al., 2012). As such, it is relevant to understand relationships between microbial community

structure and vertical soil depth *in situ* along a gradient of nutrient input, as microbial community

beta diversity could be indirectly linked to carbon mineralization.

As the microbial communities within Great Lakes coastal wetlands have yet to be

fundamentally described, it is of importance to gather baseline data on what microbes exist

within these systems, elucidate how these microbes could be interacting, and determine to what

extent microbial diversity may already be impacted by anthropogenic chemical disturbance.

Additionally, if specific groups of microbial taxa are found to be predictive of environmental

gradients, this could aid in understanding specific impacts of anthropogenic activity in

generating these gradients and selecting for specific microbial networks or taxa (Sims *et al.*,

2013; Urakawa & Bernhard, 2017). This study sought to provide an initial characterization of

microbial communities within soils of Great Lakes coastal wetlands bordering the western basin

of Lake Erie, Saginaw Bay of Lake Huron, and northern Lake Michigan. Additionally, this study

explored how environmental conditions of these coastal wetlands could act as potential drivers of

microbial community structure among and within wetlands. It was predicted that microbial

community structure would be related to environmental gradients among and within coastal

wetland regions of the Great Lakes. Further, it was anticipated that high nutrient levels within

wetlands would decouple the relationship between microbial community structure and soil depth,

as has been suggested in previous studies (Achtnich *et al.*, 1995; Alewell *et al.*, 2008). Through

high-throughput sequencing of the 16S rRNA gene and network analyses, variations in key

microbial taxa and subcommunities related to environmental gradients established by wetlands

were identified.

Results

Chemical analyses

Significant correlations ($r > 0.7$, $p \leq 0.001$) were found among NH_4^+, OM, OC, and TN.

Thus, downstream analysis combined these values into one parameter, "NUTR", represented by

OM values as this variable was the most strongly correlated with each of the other variables. PC1

and PC2 explained 56.2% and 20.6% of the variation among samples, respectively (Fig. 1).

MANOVA found significant differences in physicochemical profiles based on region ($F = 2.71$,

$p \leq 0.001$) and depth ($F = 6.85$, $p \leq 0.01$). Ninety-five percent confidence intervals suggested that

significant variation among regions was driven primarily by physicochemical distinctness of

Lake Erie coastal wetlands. This separation was related to increased NUTR, NO_3^-, and S.

Increasing depth within cores showed a consistent shift in environmental variables,

except those sites located in the western basin of Lake Erie (Supplemental Fig. 1). Specifically,

OM, OC, and TN consistently decreased with increasing depths within all wetlands except Lake

Erie sites. Similarly, C:N increased with depth in all wetlands except Lake Erie sites, where C:N

ratio remained relatively low (~ 12) and stable with increasing soil depth. Within the Lake Erie

wetland region, pH was more acidic in the overlying water with respect to all other wetland

regions (Supplemental Table 1). However, pH was still relatively neutral within Lake Erie

(average pH = 7.26, standard deviation = 0.24), where other wetland regions experienced slightly

more basic pH, with averages ranging between 7.72 – 8.39.

Alpha Diversity

Sufficient depth of sampling was also reinforced by rarefaction curve analysis

(Supplemental Fig. 2). Good's coverage reported averages ranging between 89.3 – 93.5% for

each region at the subsampled value of 48,226 sequences. Chao 1 richness estimates varied

significantly among wetland regions (F = 8.38, $p \leq 0.05$), as well as wetland sites (F = 16.78, $p \leq$

0.001). Pairwise comparisons found that the LE region had significantly higher ($p \leq 0.01$) Chao1

estimates than NSB and WSB regions (Fig. 2; Supplemental Table 2). Additionally, pairwise

comparisons found a high degree of significant variability ($p \leq 0.01$) in Chao1 estimates among

wetland sites (Supplemental Table 2). Further, Shannon diversity levels also significantly varied

among wetland sites (F = 4.57, $p \leq 0.001$), with site LE_D having significantly higher ($p \leq 0.01$)

Shannon diversity levels than sites ESBT_A and WSB_B (Supplemental Table 2). Soil depth

was not found to be significant in influencing alpha diversity levels.

Shannon diversity and Chao1 both significantly correlated with measured environmental

variables (Table 1). Specifically, Chao1 estimates increased with NO_3^-, P, and S concentrations

($p \leq 0.01$), and were weakly positively correlated ($p \leq 0.05$) with NUTR. Additionally, Shannon

diversity levels increased alongside NUTR and S ($p \leq 0.001$), and were weakly positively

correlated with NO_3^- ($p \leq 0.05$). There were no significant relationships between alpha diversity

and C:N, and there were no negative relationships between alpha diversity and any of the

measured environmental variables.

Beta Diversity

Beta diversity among regions

NMDS demonstrated separation of microbial communities based on wetland site, region,

and soil depth (Fig. 3). perMANOVA demonstrated that differences in microbial community

structure were highly significantly related to wetland site (pseudo-F = 2.57, p ≤ 0.001), region

(pseudo-F = 5.91, p ≤ 0.001), and soil depth (pseudo-F = 6.35; p ≤ 0.001). Post-hoc pairwise

perMANOVA found that community structure within the LE region was significantly distinct (p

≤ 0.01) from all other wetland regions (Table 2). No significant differences in community

structure were found between any other wetland regions compared. Additionally, microbial

community beta diversity was distinct (p ≤ 0.003) between the top soil depth and the middle and

bottom soil depths. However, no significant differences in microbial community structure were

found between the middle and bottom soil depths (Table 2). Variation in microbial community

structure was strongly significantly correlated (p ≤ 0.001) to depth (r = 0.41), NO_3^- (r = 0.20),

NUTR (r = 0.60), and S (r = 0.41), and weakly correlated (p ≤ 0.016) with C:N (r = 0.11) and P

(r = 0.14) (Supplemental Table 3).

Beta dispersion tests suggested weakly significant variation in structural variance among

regions (p ≤ 0.05), however, Tukey's HSD test using adjusted p-values for multiple comparisons

did not find any significance (p > 0.05) between pairwise comparisons of regional groups. There

were no differences in community structural dispersion among soil depths.

Beta diversity within regions

Individual NMDS plots of each region identified a high degree of variability of

significant correlations (p ≤ 0.05) between microbial community structure and several

environmental variables, as well as depth, dependent upon the wetland region explored (Fig. 4;

Supplemental Table 3). Depth was significantly correlated with microbial community structure

in all wetland regions except NSB and LE. However, microbial community structure was more

strongly related to depth in NSB (r = 0.35, p = 0.071) than LE (r = 0.19, p = 0.40). NUTR was

significantly correlated (p ≤ 0.01) with community structure within regions BA (r = 0.82), ESBT

(r = 0.51), and LE (r = 0.66). C:N was significantly correlated (p ≤ 0.01) with community

structure within regions within Saginaw Bay (i.e., ESBT [r = 0.65], NSB [r = 0.58], and WSB [r

= 0.58]). Beta diversity was not significantly associated with concentrations of NO_3^- in any

region.

To test for significant differences in microbial beta diversity within regions,

perMANOVA was used to evaluate differences in microbial community structure among soil

depths and sites within wetland regions (Supplemental Table 3). Depth did not significantly

explain microbial community structure within the region LE (pseudo-F = 0.87; p = 0.65),

however, it did explain differences in microbial community structure within the other wetland

regions. Specifically, BA (pseudo-F = 2.12; p = 0.006), ESBT (pseudo-F = 2.49; p = 0.001),

NSB (pseudo-F = 1.24; p = 0.093), and WSB (pseudo-F = 1.53; p = 0.014). Significant

differences in microbial community structure were found among different wetland sites within

regions ESBT (pseudo-F = 3.11; p = 0.001), LE (pseudo-F = 2.89; p = 0.004), and NSB (pseudo-

F = 2.16; p = 0.003). As only one site was sampled within the BA region, testing for differences

among wetland sites within the BA region could not be accomplished.

Network Analyses

Weighted Correlation Network Analysis (WGCNA) was used to explore strong

relationships between subcommunities and individual OTUs with environmental parameters

within Great Lakes coastal wetlands. After removal of OTUs which did not have at least two

representative sequences in at least 10% of samples, a total of 7,562 OTUs remained for

WGCNA. In determining scale-free topology of the OTU network, a soft power threshold of 4

was reached, and an R^2 of 0.87 was established as linear fit from the regression of the frequency

distribution of node connectivity against node connectivity (Supplemental Fig. 3). Of the 33

constructed subnetworks, the same one (subnetwork "orange") was found to be most strongly

correlated to both NUTR (r = 0.94) and NO_3^- (r = 0.55) (Supplemental Fig. 4). A separate

subnetwork ("pink") was strongly correlated (r = 0.74) to C:N. All correlations of subnetworks to

environmental variables were significant (p ≤ 0.001). OTU VIP values ≤ 1 were removed due to

the large amount of OTUs within subnetworks correlated with C:N for visualization purposes.

For subnetwork relationships to NUTR (including OM, OC, NH_4^+, and TN), partial least

square analysis (PLS) found that 69 OTUs were 93.8% predictive of variance in NUTR

(Supplemental Fig. 5). OTU co-correlation networks were constructed using an OTU co-

correlation threshold of 0.25, with strong correlations (r > 0.59) between all OTUs and NUTR

(Fig. 5). Of the top 15 OTUs contributing to PLS regression by VIP score, seven were related to

Betaproteobacteria, five were related to *Anaerolineaceae* (within *Chloroflexi*), and one

representative OTU was related to each of *Bellilinea* (*Chloroflexi*), *Desulfobacterales*

(*Deltaproteobacteria*), and *Rhizobiales* (*Alphaproteobacteria*).

For subnetwork relationships to C:N, PLS found that 144 OTUs were 59.0% predictive of

variance in C:N (Supplemental Fig. 6). Networks were constructed using an OTU co-correlation

threshold of 0.1, within positive or negative correlations (r > +/- 0.2) between OTUs (VIP > 1)

and C:N (Fig. 6). Of the top 15 OTUs by VIP score within the network, two OTUs related to

Bacteroidetes were negatively correlated with C:N. Other top OTUs were positively related to

C:N, including seven OTUs related to *Anaerolineaceae*, four OTUs which were unclassified

Bacteria, and one representative OTU related to each of *Bacillus* (*Firmicutes*) and *Chloroflexi*.

While a correlation strength of $r = 0.55$ was found between a subnetwork and NO_3^-

concentrations, PLS did not find a strong relationship between predicted values and actual values

of NO_3^- (Supplemental Fig. 7). As such, individual OTUs were correlated to NO_3^- (Fig. 7) and

were further explored through this method rather than WGCNA. Correlation strength to NO_3^-

ranged from +/-.37 to +/-.55 for positive and negative correlations, respectively. OTUs related to

Anaerolineaceae simultaneously were most positively and most negatively related to NO_3^-

concentration. OTUs related to *Acidobacteria*, *Chlorobi*, *Bacteroidetes*, *Proteobacteria*, OP11,

and *Verrucomicrobia* were predominantly positively related to NO_3^-. OTUs related to

Actinomycetales (*Actinobacteria*) and *Euryarchaeota* were negatively related to NO_3^-.

Discussion

Great Lakes coastal wetlands are biological diversity hotspots that mediate fluxes of

biogeochemical cycles, as well as buffer the Great Lakes themselves from terrestrial pollutants.

Despite the importance of coastal wetland microbial communities in mediating many of these

ecosystem services, a lack of fundamental information on these communities has hindered full

understanding of these critical environments (Hackett *et al.*, 2017). Previous research on

microbial communities in Great Lakes coastal wetlands has focused on the use of microbial

enzymatic assays as a tool to explore decomposition rates and nutrient limitation in wetlands

(Jackson *et al.*, 1995; Hill *et al.*, 2006). This study is the first to deeply characterize microbial

communities among multiple regions across the Laurentian Great Lakes basin, while

simultaneously delineating the importance of environmental conditions in structuring these

microbial communities. Additionally, this study highlights important keystone subcommunities

of potentially interacting OTUs, which may serve as indicators of ecosystem health.

246   *Microbial diversity driven by chemistry within Great Lakes coastal wetlands*

247   Taxonomic groups such as plants, birds, fish, and invertebrates within the Great Lakes

coastal wetlands have been impacted by human practices within the Great Lakes watershed

(Howe *et al.*, 2007; Tulbure *et al.*, 2007; Uzarski *et al.*, 2009; Cooper *et al.*, 2012; Uzarski *et al.*,

2017). This study is the first to establish that these patterns appear consistent with microbial

communities in these ecosystems as well. Microbial community structure was significantly

dissimilar between LE and all other wetland regions. It is therefore likely that anthropogenic

stressors related to nutrient loading (and potentially other pollutants) are driving these trends, as

nutrient levels were substantially elevated within LE sites. Likewise, Lake Erie costal wetland

sites had distinct microbial communities. Previous research has found that nutrient levels (e.g.,

C, N, P, etc.), to varying degrees, can influence microbial community composition and structure

(Hartman *et al.*, 2008; Peralta *et al.*, 2013; Ligi *et al.*, 2014; Arroyo *et al.*, 2015). Indeed, PCA

and MANOVA found that Lake Erie coastal wetlands were chemically distinct from all other

wetland regions, primarily driven by elevated nutrient (C, N, P), NO_3^-, and S concentrations

within the soil. Additionally, Lake Erie coastal wetlands (and the watershed which drains into

them) have been historically impacted by anthropogenic pollution and agricultural practices,

particularly in comparison to other coastal wetlands within the Laurentian Great Lakes region.

This has been demonstrated by multiple ecological indices (e.g., Cvetkovic & Chow-Fraser,

2011; Uzarski *et al.*, 2017) and physicochemical uniqueness (increased levels of nutrients and

particulate matter) within the western basin of Lake Erie (Danz *et al.*, 2007; Trebitz *et al.*, 2007;

Cvetkovic & Chow-Fraser, 2011; Uzarski *et al.*, 2017). Data presented in this study corroborate

this historical evidence of human impact and nutrient loading in the western basin of Lake Erie.

High nutrient influx could also be influencing chemical and microbial vertical structure

within coastal wetland soils. Vertical microbial community structure was not evident within the

first 6 cm of soil of Lake Erie coastal wetlands, unlike other coastal wetlands regions. Also

unique to Lake Erie, vertical chemical structure was not evident for nutrient levels (C, N, P) or

C:N. These data provide evidence that microbial community structural shifts corresponding to

wetland vertical profiles are related to concentrations of organic carbon and other nutrients (e.g.,

NO_3^-, NUTR, P, S). This is meaningful, as it has been previously postulated that concentrations

of carbon electron donors may influence redox gradients within wetland soils (Achtnich *et al.*,

1995). Redox gradients are uniquely tied to carbon cycling rates, and as microbial community

structure corresponds with redox gradients (Lüdemann *et al.*, 2000; Edlund *et al.*, 2008; Lipson

et al., 2015), distinct patterns of vertical community structure within the soil may be indicative of

biogeochemical processes being affected within Great Lakes coastal wetlands. Connections

between microbial community metabolic shifts with soil depth and levels of dissolved organic

carbon *in situ* were incompletely resolved (Alewell *et al.*, 2008). The results presented here

suggest that a connection between microbial community metabolism and organic carbon

concentration may exist within Great Lakes coastal wetlands, however, it is necessary to better

link microbial community diversity, microbial activity, and chemical cycles to establish this

connection. As a caveat, it is possible that chemical and microbial structuring still existed within

Lake Erie wetlands, yet they were not evident within the first 6 cm of soil or at the spatial scale

measured. Nevertheless, microbial communities within Lake Erie coastal wetlands did not follow

the same pattern of vertical structure, either chemically or biologically, evident in other regions,

suggesting that the biological integrity of coastal wetland systems is susceptible to negative

anthropogenic pressure. Furthermore, carbon and nutrient levels are stable with increasing soil

depth in Lake Erie coastal wetlands, which could be indicative of low carbon cycling rates or

elevated sedimentation rates.

Similar to beta diversity, patterns in alpha diversity of microbial communities may also

be related to soil chemical characteristics within the sampled freshwater coastal wetlands.

Microbial community alpha diversity was highest within Lake Erie soils, which experienced the

highest levels of nutrient concentrations of the sampled wetland regions. Significant correlations

between alpha diversity and physicochemical properties (Table 1) suggest that the higher

diversity in Lake Erie coastal wetlands could be driven by its distinct soil characteristics. It has

been established in some cases that productivity and alpha diversity increase in a linear

relationship on regional scales, but not at local scales (Chase & Leibold, 2002). This pattern of

diversity is commonly the result of relatively few species existing within productive habitats on

the local scale, while dissimilarity in beta diversity among productive habitat patches increases

alpha diversity at larger spatial resolution. It is possible that this relationship exists within our

data, with the Lake Erie region being the most OTU-diverse, coinciding with high nutrient

concentrations. Habitat patches within Lake Erie wetlands may be smaller in spatial scale than

was sampled within this study (where we might expect lower alpha diversity), and high

dissimilarity in OTU composition among these hypothetical habitat patches may be responsible

for higher alpha diversity in Lake Erie coastal wetlands. Substantiating this, taxa-area

relationships have been established for microbial communities which suggest that microbial

community turnover can occur at distances as short as millimeters to centimeters within soils and

sediments (Grundmann & Debouzie, 2000; Ettema & Wardle, 2002; Horner-Devine *et al.*, 2004;

Martiny *et al.*, 2011). Microbial diversity has also been found to have a strong relationship with

pH within forest and wetland soils (Fierer & Jackson, 2005; Hartman *et al.*, 2008; Lauber *et al.*,

2009). The water directly above the soil in the LE region had a pH that was relatively more

acidic (although circumneutral) with respect to other sampled wetland regions. This may suggest

that more OC was being oxidized as CO_2 reacts with water to form carbonic acid. This idea

would then suggest that LE sites had a higher deposition rate replacing OC as oxidation

occurred. While previous research has recognized links between pH and microbial community

diversity within wetland soils, these relationships have been highly variable, ranging from a lack

of relationship to a strong predictive relationship (Hartman *et al.*, 2008; Peralta *et al.*, 2013; Ligi

et al., 2014; Arroyo *et al.*, 2015). To date, research has been variable with respect to

environments studied and methods by which pH was measured; this may partly explain the wide

range of estimates of relationships between wetland soil pH and microbial community structure.

As such, more research is needed to appreciate the influence of pH on microbial communities

within wetland soils. Standardization of methodology for assessing environmental characteristics

such as pH may also be required.

328   *Relationships between microbial subnetworks and environmental gradients*

329   Development of biological indices and establishment of indicator taxa have been

suggested as necessary for microbial communities within wetlands, particularly through the use

of high-throughput sequencing technologies which now allow for deep assessment of microbial

community composition and structure within environmental samples (Sims *et al.*, 2013; Urakawa

& Bernhard, 2017). In addition to their importance as biological signals for environmental

health, indicator taxa may play prominent roles in bioremediation of excess nutrients and

pollutants found within anthropogenically impacted coastal wetlands. Through network analyses,

we have delineated multiple subcommunities which were significantly related to environmental

gradients (such as nutrients C, N, and P) measured by Uzarski *et al.* (2017) among coastal

wetlands sampled in this study. Specifically, a subnetwork of 69 microbial taxa was 93.8%

predictive of nutrient level variation among coastal wetland soils. Several microbial taxa within

this subcommunity were individually predictive of nutrient levels to a high degree, including

several OTUs related to *Anaerolineaceae*, one OTU related to genus *Anaerolinea*, and another

related to genus *Bellilinea*. From the genus *Anaerolinea*, two thermophilic chemoorganotrophs

(*Anaerolinea thermophila* and *Anaerolinea thermolimosa*) have been isolated (Sekiguchi *et al.*,

2003; Yamada *et al.*, 2006). Only one isolated member has been established within the genus

Bellilinea (*Bellilinea caldifistulae*); it has been described as a thermophilic, fermentative,

obligate anaerobe which thrives in co-culture with methanogens (Yamada *et al.*, 2007). It is

unlikely that the OTUs found in our study are the same species as the isolated *Anaerolinea* and

Bellilinea species, as coastal wetland soils are not high-temperature environments necessary for

thermophilic species. Additionally, no OTUs related to methanogenic archaea were found within

this subnetwork, suggesting that *Anaerolineacea* OTUs within coastal wetland soils may

fluctuate independently of any specific methanogenic OTUs. It is possible that the *Bellilinea*

OTU found within the subnetwork is related to nutrient level concentrations. This would support

fermentative metabolism as noted within *Bellilinea caldifistulae*. It is important to note that

several other studies have discovered OTUs related to *Anaerolineaceae* within wetland soils,

with upwards of 90% relative abundance among *Chloroflexi* OTUs within these systems (Ansola

et al., 2014; Deng *et al.*, 2014; Hu *et al.*, 2016). This suggests that there are probable mesophilic

species yet to be isolated within this ubiquitous family of bacteria, which may be of high

importance within wetland soils.

Betaproteobacteria were also found to significantly predict nutrient levels among coastal

wetlands. Hu *et al.* (2016) found that both *Betaproteobacteria* and *Anaerolineae* were positively

related to TN levels, which is consistent with the data presented here, and these two taxa were

suggested to contribute to higher levels of heterotrophic activity. Further, *Anaerolineaceae*

OTUs were consistently related to increasing C:N, suggesting that many taxa within this family

have preference for recalcitrant carbon sources. As C:N also tends to increase with soil depth, it

is also probable that the putatively obligate anaerobic *Anaerolineaceae* are coinciding with

decreasing oxygen levels and/or changing metabolism requirements with increasing soil depth.

While there was no subnetwork of microbial taxa predictive of NO_3^- in the studied coastal

wetland systems, there were several taxonomic groups that were either positively or negatively

related to NO_3^- independently. Taxa spanning ten phyla were found to either positively or

negatively correlate to NO_3^- gradients (Fig. 8). As nitrate reduction is a process which is

undertaken by a wide breadth of phylogenetic groups (Reddy & DeLaune, 2008), it is possible

that several taxa are capable of nitrate respiration within Great Lakes coastal wetlands. Notably,

some taxa which positively related to NO_3^- are known nitrate reducers in other systems, such as

members of *Deltaproteobacteria*, including *Myxococcales* and *Anaeromyxobacter* (Sanford *et*

al., 2002). Also noteworthy are the high number of OTUs within *Anaerolineaceae* which

positively correlated with NO_3^- . However, several *Anaerolineaceae* OTUs were negatively

related to NO_3^- gradients, suggesting that functional idiosyncrasy may exist among members of

this family.

Conclusions

This study marks the first characterization of microbial communities within Great Lakes

coastal wetlands. Coastal wetlands are integral in the proper functioning of biogeochemical

cycles and environmental sustainability of the Great Lakes. While it has long been known that

anthropogenic pressure can impact animal and plant communities within these coastal wetlands,

this is the first evidence that these pressures may also be influencing microbial communities, and

may be influencing biogeochemical cycles by extension. Alpha and beta diversity were both

related to nutrient gradients among and within regions, suggesting that variability in microbial

community structure is highly coupled to geochemistry within wetland soil. Further, this study

provides insight on microbial community subnetworks and individual OTUs, which were

predictive of chemical concentrations, and may be useful for future management of Great Lakes

coastal wetland systems.

Beyond taxonomic assessments of microbial communities with relation to wetland

monitoring, we propose that wetland microbial community structure can also potentially be used

to assess a wetland for management purposes. As illustrated within this study, wetland microbial

community structure and depth are decoupled within the wetlands experiencing the highest

nutrient levels, likely originating from terrestrial inputs due to human activity. As such,

multivariate statistics (as used in the methods of this study) may prove useful in examining

relationships between wetland soil depth and microbial community structure alongside

taxonomic analyses, which could provide indicators of nutrient loading stress on coastal wetland

habitats.

Experimental Procedures

In the summer of 2014, wetland soil cores were collected within Laurentian Great Lakes

coastal wetland ecosystems across several sites within several regions. Specifically, soil cores

were collected from ten sites across five regions, including two sites in the western basin of Lake

Erie (LE), three sites in eastern Saginaw Bay (ESBT), two sites in northern Saginaw Bay (NSB),

two sites in western Saginaw Bay (WSB) in Lake Huron, and one site in the Beaver Island

archipelago (BA) in Lake Michigan (Fig. 8). These sites were selected as they corresponded to

environmental gradients, as well as a human impact gradients based upon SumRank scores as

described in Uzarski *et al.* (2017) (Supplemental Fig. 8). Soil cores were collected by hand-

driving plastic core tubes (~ 5 cm diameter) vertically into the soil. Among wetlands, samples

were collected within the same vegetation zone (either dominated by cattails, genus *Typhus*, or

bulrush, genus *Shoenoplectus*) as an attempt to control for collection bias across sites. Cores

were sampled to a depth of at least 6 cm (except for one core which was sampled to a depth of 4

cm) and were immediately flash frozen in a dry ice ethanol bath. Samples were transported on

dry ice to Central Michigan University wherein they were stored at -80 ˚C.

Triplicate cores were taken at five wetland sites while duplicate cores were taken at five

other wetland sites. For sample extraction and sectioning, cores were extruded while still frozen

via a custom-built core extruder. The edge of the core was warmed with a heat gun to allow the

soil core to pass efficiently through the plastic container, however, the inner-core did not thaw

during extrusion. Ice was applied to the plastic core liner to prevent accelerated thawing.

Beginning from the top surface of soil, 2 cm sections were cut via an ethanol and flame-sterilized

hacksaw blade and the sectioned core samples were placed into whirl pak bags and stored at -80

˚C. The extruder face plate was sterilized between cuts of the same core with ethanol. The

extruder device was fully cleaned and sterilized between cores with physical scrubbing and

ethanol sterilization.

Microbial community analysis

Each soil sample was analyzed independently for microbial community analyses. DNA

was extracted from ~ 0.25 g of soil using a MoBio PowerSoil DNA Isolation Kit (Mo Bio,

Carlsbad, CA) following the standard manufacturer's protocol. Concentrations of extracted DNA

were assessed using a Qubit® 2.0 fluorometer (Life Technologies, Carlsbad, CA) to ensure

successful DNA extraction and quantification for sequence library preparation. DNA samples

were sent to Michigan State University (East Lansing, MI) for library preparation and sequence

analysis at the Research Technology Support Facility. The V4 region of the 16S rRNA gene was

amplified for downstream sequencing with commonly used primers 16Sf-V4 (515f) and 16Sr-V4

(806r) and a previously developed protocol (Caporaso *et al.*, 2012; Kozich *et al.*, 2013). Paired-

end 250 bp sequencing was accomplished via a MiSeq high-throughput sequencer (Illumina, San

Diego, CA). Acquired DNA sequences were filtered for quality and analyzed using MOTHUR v

1.35.1 (Schloss *et al.*, 2009) following the MiSeq SOP (available at https://www.mothur.org/)

with modifications. Scripts used for sequence processing can be found at the GitHub repository

associated with this study (https://github.com/horto2dj/GLCW/). Briefly, paired end sequences

were combined into single contigs. Sequences that contained homopolymers > 8 bases, and those

less than 251 or greater than 254 bp were removed. Sequences were aligned against the Silva (v.

119) rRNA gene reference database (Quast *et al.*, 2012). Sequences which did not align with the

V4 region were also subsequently removed from analysis. Chimeric DNA was searched for and

removed via UCHIME (Edgar *et al.*, 2011). Sequences were classified via the Ribosomal

Database Project (training set v. 9; Cole *et al.*, 2013) using the 'wang' method with a confidence

threshold of 80. Sequences classified as chloroplast, mitochondria, eukaryotic, or unknown were

removed. Remaining sequences were clustered into Operational Taxonomic Units (OTUs) at

0.03 sequence dissimilarity using the opticlust clustering algorithm. Sequence data associated

with this research have been submitted to the GenBank database under accession numbers

SRR6261304 – SRR6261377.

Chemical analysis

Each soil layer (top, middle, and bottom) was analyzed separately for local chemistry at

each site. Within each site, soil samples of the same depth (i.e., top, middle, and bottom soil

samples) among duplicate/triplicate cores were combined and homogenized to obtain enough

soil for chemical analyses. Soil samples were analyzed for percent total N ("TN"), total P ("TP",

ppm), total S ("TS", ppm), NO_3^- (ppm), NH_4^+ (ppm), percent organic matter ("OM"), percent

organic carbon ("OC"), and C:N at the Michigan State University Soil & Plant Nutrient Lab

(East Lansing, MI). A YSI multiprobe (YSI Inc., Yellow Springs, OH) was used to measure pH

of the water residing directly above each collected soil core.

Statistical analyses

Statistical analyses were completed using R statistical software version 3.2.2 (R Core

Team, 2015) unless otherwise stated. Code used for statistical analyses (and bioinformatic

workflow) in this study can be found in the associated GitHub repository

(https://github.com/horto2dj/GLCW/).

Physicochemical Analysis

Differences in chemical profiles between samples within and among wetland regions

were visualized using Principal Component Analysis (PCA). Prior to PCA, percentages were

arcsin square root transformed and ratios were log transformed. Additionally, Pearson correlation

analyses were performed to search for significant correlations between chemical variables.

Collinearity in the dataset was addressed by combining highly correlated environmental

variables ($r > 0.7$, $p \leq 0.001$). Only one of the correlated variables was included in PCA to

remove exaggeration of correlated variables in PCA structure. MANOVA was used to determine

the influences of region and soil depth on physicochemical composition of samples, and 95%

confidence intervals were established to compare differences among groups. Chemical depth

profiles were also visualized for each wetland site to understand shifts in measured

environmental variables with increasing soil depth.

Alpha-diversity Analysis

Alpha diversity analyses were performed to explore variation in OTU richness and

evenness among wetland sites, regions, and soil depths, as well as to determine whether observed

trends were driven by environmental variables. Prior to alpha diversity analyses, sequence

abundance for each sample was subsampled to the lowest sequence abundance for any one

sample ($n = 48,226$ sequences). Singletons were maintained within the sequence dataset for

alpha diversity analyses, as alpha diversity indices can be reliant on the presence of singletons

for proper estimation. Alpha diversity was calculated for each site using MOTHUR, including

Chao 1 richness, non-parametric Shannon diversity, and Good's coverage indices. Linear mixed-

effect models and ANOVAs were used to test influences of wetland site, region, and soil depth

on alpha diversity, controlling for wetland site as a random effect. Linear models and ANOVAs

were used to test for variation in alpha diversity among wetland sites. If significant variation was

found within an ANOVA result, post-hoc comparisons were implemented between sample

groups using Tukey's Honest Significant Differences (HSD) tests with Bonferroni adjustments

(p-values obtained by number of comparisons) for pairwise comparisons.

500   *Beta diversity Analysis*

501   Beta diversity analyses were used to evaluate variation in microbial community structure

among wetland sites, regions, and soil depths, and to assess the extent to which observed

variation was explained by environmental conditions. Singletons and doubletons were removed

from the dataset for beta diversity analyses. All sequence data were maintained for beta diversity

analyses and transformed using the *DeSeq2* (Love *et al.*, 2014) package, which normalized OTU

abundances among samples using a variance stabilizing transformation (VST) (McMurdie &

Holmes, 2014). The *phyloseq* (McMurdie & Holmes, 2013) and *Vegan* (Oksanen *et al.*, 2007)

packages were used to compare beta diversity among samples. Dissimilarity in microbial

community structure among samples within and among sites was visualized using Non-metric

Multidimensional Scaling (NMDS) plots based on pairwise Bray-Curtis dissimilarity estimates.

The function *envfit* of the Vegan package was used to evaluate correlation between chemical

parameters and microbial community structure among samples according to NMDS. "Depth"

was also implemented as a dummy variable to test correlation between depth and microbial

community structure.

515   To test for significant differences in beta diversity among wetland sites, regions, and soil

depth, Permutational Multivariate Analysis of Variance (perMANOVA) (Anderson, 2001) were

implemented. Specifically, these tests evaluate significant variation among within group and

between group means (Clarke, 1993; Anderson, 2001; Anderson & Walsh, 2013). If

perMANOVA found significant differences among groups at the global level, pairwise

perMANOVA tests between groups were implemented with Bonferroni significance adjustments

to control for multiple pairwise comparisons. Anderson's permutation of dispersions test

(PERMDISP; Anderson, 2006; Anderson *et al.*, 2006) was used to test for differences in variance

of community structure among sample groups (i.e. sites, regions, soil depths). Tukey's Honest

Significant Difference (HSD) tests were implemented with adjusted p-values for multiple

pairwise comparisons if significant differences in dispersion were found among groups.

To explore relationships between regional microbial community structure and

environmental variables, NMDS plots were generated for each individual region. Applying

NMDS to each region also allowed for the assessment of the correlational relationship between

community structure and soil depth (as a dummy variable) and other environmental variables

(using the *envfit* function) within individual regions. To test for differences in microbial

community structure between/among sites within a region, as well as among depths within a

region, perMANOVA was implemented individually for each region.

Network and OTU Correlation Analyses

To explore relationships between microbial sub-communities and OTUs to environmental

variables, Weighted Correlation Network Analysis (WGCNA) was implemented on OTU

relative abundances using the *WGCNA* package (Langfelder & Horvath, 2008; Langfelder &

Horvath, 2012), executed as previously described (Guidi *et al.*, 2016; Henson *et al.*, 2016) with

modifications. OTUs which did not possess at least 2 sequences across 10% of samples were

removed from network analyses. These OTUs were removed to eliminate OTUs with potentially

spurious correlations to environmental variables or other OTUs, as well as to reduce

computational stress of analyses. Remaining OTU abundances across samples were normalized

using variance stabilizing transformation (VST) performed as described previously for beta

diversity analyses. To ensure scale-free topology of the network, the dissimilarity matrix

generated through VST was transformed to an adjacency matrix by raising this dissimilarity

matrix to a soft threshold power. A threshold power of $p = 4$ was chosen to meet scale-free

topology assumptions based upon criterion established by Zhang & Horvath (2005). Scale-free

topology of network relationships was further ensured through regression of the frequency

distribution of node connectivity against node connectivity; a network is scale-free if an

approximate linear fit of this regression is evident (see Zhang & Horvath, [2005] for more in-

depth explanation). A topological overlap matrix (TOM) was generated using the adjacency

matrix, and subnetworks of highly connected and correlated OTUs were delineated with the

TOM and hierarchical clustering. Representative eigenvalues of each subnetwork (i.e., the first

principal component) were correlated (Pearson) with values of measured environmental variables

to identify the subnetworks most related to said environmental variables. The subnetworks with

the highest positive correlations to environmental variables of interest (e.g., NO_3^-, C:N, etc.)

were selected for further analyses of relationships among subnetwork structure, individual

OTUs, and environmental variables. Partial Least Square regression (PLS) was used to test

predictive ability of subnetworks in estimating variability of environmental parameters, which

allowed for delineation of potential indicator subnetworks and OTUs. Pearson correlations were

calculated between response variables and leave-one-out cross-validation (LOOCV) predicted

values. If PLS found that regression between actual and predicted values was below minimum

threshold of $R^2 = 0.3$, WGCNA analysis was halted for that network, as the network was deemed

to lack predictive ability of that environmental variable. Variable Importance in Projection (VIP)

(Chong & Jun, 2005) analysis was used to determine the influence of individual OTUs in PLS. A

high VIP value for an OTU indicates high importance in prediction of the environmental

response variable for that OTU. For network construction and visualization purposes, the

minimum correlation value required between two OTUs to constitute an "edge" between them

was delineated at different r values for each network related to an individual environmental

variable (ranging between 0.1 – 0.25), as co-correlations between OTUs within some networks

were stronger than others. The number of co-correlations an OTU has with other OTUs within a

network is defined as "node centrality" (as described by Henson *et al.*, 2016).

If an environmental variable of interest was not found to have a network strongly

associated with it, Spearman's correlations were implemented between individual OTUs and that

environmental variable. Significance was only considered for OTUs with strong relationships (r

> 0.3, p ≤ 0.001) to environmental variables to limit the impact of potentially spurious

correlations. Correlations to OTUs with unknown taxonomic identification were excluded from

correlational analysis.

Acknowledgments

Special thanks to Alexandra Bozimowski and Thomas Langer for their assistance in collecting

the wetland samples and data acquisition, as well as Dr. Matthew Cooper for providing valuable

insight and conversation on wetland ecology and the custom-built core extruder built by Mr.

Gary Cooper. Special thanks also to Mike Henson for providing assistance on network statistical

analyses. Funding was provided, in part, by the CMU College of Science and Engineering.

Additional funding for this work was provided by the Great Lakes National Program Office

under the United States Environmental Protection Agency, grant number GL-00E00612-0 as part

of the US federal government's Great Lakes Restoration Initiative. Although the research

described in this work has been partly funded by the United States Environmental Protection

Agency, it has not been subjected to the agency's required peer and policy review, and therefore,

does not necessarily reflect the views of the agency and no official endorsement should be

inferred. This paper is contribution number XX of the Central Michigan University Institute for

Great Lakes Research.

References

Achtnich, C., Bak, F., and Conrad, R. (1995). Competition for electron donors among nitrate

reducers, ferric iron reducers, sulfate reducers, and methanogens in anoxic paddy soil. Biol Fert

Soils. 19(1): 65-72.

Alewell, C., Paul, S., Lischeid, G., and Storck, F.R. (2008). Co-regulation of redox processes in

freshwater wetlands as a function of organic matter availability?. Sci Total Environ 404(2): 335-

342.

Anderson, M.J. (2001). A new method for non‐parametric multivariate analysis of variance.

Austral Ecol 26(1): 32-46.

Anderson, M.J. (2006). Distance‐based tests for homogeneity of multivariate dispersions.

Biometrics 62(1): 245-253.

Anderson, M.J., Ellingsen, K.E., and McArdle, B.H. (2006). Multivariate dispersion as a

measure of beta diversity. Ecol Lett 9(6): 683-693.

Anderson, M.J., and Walsh, D.C. (2013). PERMANOVA, ANOSIM, and the Mantel test in the

face of heterogeneous dispersions: what null hypothesis are you testing?. Ecol Monogr 83(4):

557-574.

Ansola, G., Arroyo, P., and de Miera, L.E.S. (2014). Characterisation of the soil bacterial

community structure and composition of natural and constructed wetlands. Sci Total Environ

473: 63-71.

Arroyo, P., de Miera, L.E.S., and Ansola, G. (2015). Influence of environmental variables on the

structure and composition of soil bacterial communities in natural and constructed wetlands. Sci

Total Environ 506: 380-390.

Campbell, M., Cooper, M.J., Friedman, K., and Anderson, W.P. (2015). The economy as a driver

of change in the Great Lakes–St. Lawrence River basin. J Great Lakes Res 41: 69-83.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., *et al.*

(2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq

platforms. ISME J 6(8): 1621-1624.

Chase, J.M., and Leibold, M.A. (2002). Spatial scale dictates the productivity–biodiversity

relationship. Nature 416(6879): 427-430.

Chong, I.G., and Jun, C.H. (2005). Performance of some variable selection methods when

multicollinearity is present. Chemometr Intell Lab 78(1): 103-112.

Clarke, K.R. (1993). Non-parametric multivariate analyses of changes in community structure.

Aust J Ecol 18: 117-143.

Cole J.R., Wang Q., Fish J.A., Chai B., Mcgarrell D.M., Sun Y., *et al.* (2013) Ribosomal

Database Project: data and tools for high throughput rRNA analysis. Nucleic Acids Res 42(D1):

doi:10.1093/nar/gkt1244

Conrad, R. (1996). Soil microorganisms as controllers of atmospheric trace gases (H2, CO, CH4,

OCS, N2O, and NO). Microbiol Rev 60(4): 609-640.

Cooper, M.J., Gyekis, K.F., and Uzarski, D.G. (2012). Edge effects on abiotic conditions,

zooplankton, macroinvertebrates, and larval fishes in Great Lakes fringing marshes. J Great

Lakes Res 38(1): 142-151.

Cvetkovic, M., and Chow-Fraser, P. (2011). Use of ecological indicators to assess the quality of

Great Lakes coastal wetlands. Ecological Indicators 11(6): 1609-1622.

Dahl, T.E. (1990). Wetlands losses in the United States 1780's to 1980's. U.S. Department of the

Interior, Fish and Wildlife Service, Washington. D.C. 13pp.

Danz, N.P., Niemi, G.J., Regal, R.R., Hollenhorst, T., Johnson, L.B., Hanowski, J.M., *et al*.

(2007). Integrated measures of anthropogenic stress in the US Great Lakes basin. Environ

Manage 39(5): 631-647.

Deng, Y., Cui, X., Hernández, M., and Dumont, M.G. (2014). Microbial diversity in hummock

and hollow soils of three wetlands on the Qinghai-Tibetan Plateau revealed by 16S rRNA

pyrosequencing. PLoS One 9(7): e103115.

Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011). UCHIME improves

sensitivity and speed of chimera detection. Bioinformatics 27(16): 2194-2200.

Edlund, A., Hårdeman, F., Jansson, J.K., and Sjöling, S. (2008). Active bacterial community

structure along vertical redox gradients in Baltic Sea sediment. Environ Microbiol 10(8): 2051-

2063.

Ettema, C.H., and Wardle, D.A. (2002). Spatial soil ecology. Trends in ecology and evolution

17(4): 177-183.

Fierer, N., and Jackson, R.B. (2006). The diversity and biogeography of soil bacterial

communities. P Natl Acad Sci USA 103(3): 626-631.

Grundmann, G.L., and Debouzie, D. (2000). Geostatistical analysis of the distribution of NH4+

and NO2−-oxidizing bacteria and serotypes at the millimeter scale along a soil transect. FEMS

Microbiol Ecol 34(1): 57-62.

Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., *et al*. (2016). Plankton

networks driving carbon export in the oligotrophic ocean. Nature 532(7600): 465.

Hackett, R.A., Babos, H.B., Collins, E.E., Horton, D., Schock, N., and Schoen, L.S. (2017).

Researcher disciplines and the assessment techniques used to evaluate Laurentian Great Lakes

coastal ecosystems. J Great Lakes Res 43(1): 9-16.

Hartman, W.H., Richardson, C.J., Vilgalys, R., and Bruland, G.L. (2008). Environmental and

anthropogenic controls over bacterial communities in wetland soils. P Natl Acad Sci 105(46):

17842-17847.

Hecnar, S.J. (2004). Great Lakes wetlands as amphibian habitats: a review. Aquat Ecosyst Health

7(2): 289-303.

Henson, M.W., Hanssen, J., Spooner, G., Flemming, P., Pukonen, M., Stahr, F., and Thrash, J.C.

(2016). Nutrient dynamics and stream order influence microbial community patterns along a

2914 km transect of the Mississippi River. bioRxiv, 091512.

Hill, B.H., Elonen, C.M., Jicha, T.M., Cotter, A.M., Trebitz, A.S., and Danz, N.P. (2006).

Sediment microbial enzyme activity as an indicator of nutrient limitation in Great Lakes coastal

wetlands. Freshwater Biol 51(9): 1670-1683.

Horner-Devine, M.C., Lage, M., Hughes, J.B., and Bohannan, B.J. (2004). A taxa-area

relationship for bacteria. Nature 432(7018): 750.

Howe, R.W., Regal, R.R., Hanowski, J., Niemi, G.J., Danz, N.P., and Smith, C.R. (2007). An

index of ecological condition based on bird assemblages in Great Lakes coastal wetlands. J Great

Lakes Res 33(sp3): 93-105.

Hu, Y., Wang, L., Fu, X., Yan, J., Wu, J., Tsang, Y., *et al*. (2016). Salinity and nutrient contents

of tidal water affects soil respiration and carbon sequestration of high and low tidal flats of

Jiuduansha wetlands in different ways. Sci Total Environ 565: 637-648.

Jackson, C.R., Foreman, C.M., and Sinsabaugh, R.L. (1995). Microbial enzyme activities as

indicators of organic matter processing rates in a Lake Erie coastal wetland. Freshwater biol

34(2): 329-342.

Kögel-Knabner, I., Amelung, W., Cao, Z., Fiedler, S., Frenzel, P., Jahn, R., *et al*. (2010).

Biogeochemistry of paddy soils. Geoderma 157(1): 1-14.

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D. (2013) Development

of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence

Data on the MiSeq Illumina Sequencing Platform. Appl Environ Microb 79(17): 5112-5120.

Krantzberg, G., and De Boer, C. (2008). A valuation of ecological services in the Laurentian

Great Lakes Basin with an emphasis on Canada. Am Wat Works Assoc J 100(6): 100.

Lamers, L.P., Van Diggelen, J.M., den Camp, H.J.O., Visser, E.J., Lucassen, E.C., Vile, M.A., *et*

al. (2012). Microbial transformations of nitrogen, sulfur, and iron dictate vegetation composition

in wetlands: a review. Front Microbiol, 3:156.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation

network analysis. BMC Bioinformatics 9(1): 559.

Langfelder, P., and Horvath, S. (2012). Fast R functions for robust correlations and hierarchical

clustering. J Stat Softw 46(11).

Lauber, C.L., Hamady, M., Knight, R., and Fierer, N. (2009). Pyrosequencing-based assessment

of soil pH as a predictor of soil bacterial community structure at the continental scale. Appl

Environ Microb 75(15): 5111-5120.

Ligi, T., Oopkaup, K., Truu, M., Preem, J.K., Nõlvak, H., Mitsch, W.J., *et al.* (2014).

Characterization of bacterial communities in soil and sediment of a created riverine wetland

complex using high-throughput 16S rRNA amplicon sequencing. Ecol Eng 72: 56-66.

Lipson, D.A., Raab, T.K., Parker, M., Kelley, S.T., Brislawn, C.J., and Jansson, J. (2015).

Changes in microbial communities along redox gradients in polygonized Arctic wet tundra soils.

Env Microbiol Rep 7(4): 649-657.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15(12): 550.

Lüdemann, H., Arth, I., and Liesack, W. (2000). Spatial changes in the bacterial community

structure along a vertical oxygen gradient in flooded paddy soil cores. Appl Environ Microb

66(2): 754-762.

Martiny, J.B., Eisen, J.A., Penn, K., Allison, S.D., and Horner-Devine, M.C. (2011). Drivers of

bacterial β-diversity depend on spatial scale. P Natl Acad Sci 108(19): 7850-7854.

McMurdie, P.J., and Holmes, S. (2013) Phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. PLoS One 8(4).

McMurdie, P.J., and Holmes, S. (2014). Waste not, want not: why rarefying microbiome data is

inadmissible. PLoS Comput Biol, 10(4): e1003531.

Morrice, J.A., Danz, N.P., Regal, R.R., Kelly, J.R., Niemi, G.J., Reavie, E.D., *et al.* (2008).

Human influences on water quality in Great Lakes coastal wetlands. Environ Manage 41(3): 347-

357.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J., and Suggests,

M.A.S.S. (2007). The vegan package. Community ecology package, 10, 631-637.

Peralta, R.M., Ahn, C., and Gillevet, P.M. (2013). Characterization of soil bacterial community

structure and physicochemical properties in created and natural wetlands. Sci Total Environ 443:

725-732.

Quast C., Pruesse E., Yilmaz P., Gerken J., Schweer T., Yarza P., *et al.* (2012). The SILVA

ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic

Acids Res 41(D1): D590-D596.

R Core Team (2015) R: A language and environment for statistical computing. Vienna, Austria.

https://www.R-project.org/.

Reddy, K.R., and DeLaune, R.D. (2008). Biogeochemistry of wetlands: science and applications.

Boca Raton, FL: CRC Press.

Sanford, R.A., Cole, J.R., and Tiedje, J.M. (2002). Characterization and description of

Anaeromyxobacter dehalogenans gen. nov., sp. nov., an aryl-halorespiring facultative anaerobic

myxobacterium. Appl Environ Microb 68(2): 893-900.

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., *et al.* (2009).

Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for

Describing and Comparing Microbial Communities. Appl Environ Microbiol 75(23): 7537-7541.

Sekiguchi, Y., Yamada, T., Hanada, S., Ohashi, A., Harada, H., and Kamagata, Y. (2003).

Anaerolinea thermophila gen. nov., sp. nov. and Caldilinea aerophila gen. nov., sp. nov., novel

filamentous thermophiles that represent a previously uncultured lineage of the domain Bacteria

at the subphylum level. Int J Syst Evol Micr 53(6): 1843-1851.

Sierszen, M.E., Morrice, J.A., Trebitz, A.S., and Hoffman, J.C. (2012). A review of selected

ecosystem services provided by coastal wetlands of the Laurentian Great Lakes. Aquat Ecosyst

Health 15(1): 92-106.

Sims, A., Zhang, Y., Gajaraj, S., Brown, P.B., and Hu, Z. (2013). Toward the development of

microbial indicators for wetland assessment. Water Res 47(5): 1711-1725.

Trebitz, A.S., Brazner, J.C., Cotter, A.M., Knuth, M.L., Morrice, J.A., Peterson, G.S., *et al.*

(2007). Water quality in Great Lakes coastal wetlands: basin-wide patterns and responses to an

anthropogenic disturbance gradient. J Great Lakes Res 33(sp3): 67-85.

Tulbure, M.G., Johnston, C.A., and Auger, D.L. (2007). Rapid invasion of a Great Lakes coastal

wetland by non-native Phragmites australis and Typha. J Great Lakes Res 33(sp3): 269-279.

Urakawa, H., and Bernhard, A.E. (2017). Wetland management using microbial indicators. Ecol

Eng 108(B): 456-476.

Uzarski, D.G. (2009). Wetlands of Large Lakes. Encyclopedia of Inland Waters. Oxford:

Elsevier. p. 599-606.

Uzarski, D.G., Brady, V.J., Cooper, M.J., Wilcox, D.A., Albert, D.A., Axler, R.P, *et al*. (2017).

Standardized measures of coastal wetland condition: Implementation at a Laurentian Great Lakes

basin-wide scale. Wetlands 37(1): 15-32.

Uzarski, D.G., Burton, T.M., Kolar, R.E., and Cooper, M.J. (2009). The ecological impacts of

fragmentation and vegetation removal in Lake Huron's coastal wetlands. Aquat Ecosyst Health

12(1): 45-62.

Wang, N., and Mitsch, W.J. (1998). Estimating phosphorus retention of existing and restored

coastal wetlands in a tributary watershed of the Laurentian Great Lakes in Michigan, USA. Wetl

Ecol Manag 6(1): 69-82.

Yamada, T., Sekiguchi, Y., Hanada, S., Imachi, H., Ohashi, A., Harada, H., and Kamagata, Y.

(2006). Anaerolinea thermolimosa sp. nov., Levilinea saccharolytica gen. nov., sp. nov. and

Leptolinea tardivitalis gen. nov., sp. nov., novel filamentous anaerobes, and description of the

new classes Anaerolineae classis nov. and Caldilineae classis nov. in the bacterial phylum

Chloroflexi. Int J Syst Evol Micr 56(6): 1331-1340.

Yamada, T., Imachi, H., Ohashi, A., Harada, H., Hanada, S., Kamagata, Y., and Sekiguchi, Y.

(2007). Bellilinea caldifistulae gen. nov., sp. nov. and Longilinea arvoryzae gen. nov., sp. nov.,

strictly anaerobic, filamentous bacteria of the phylum Chloroflexi isolated from methanogenic

propionate-degrading consortia. Int J Syst Evol Micr 57(10): 2299-2306.

Zhang, B., and Horvath, S. (2005). A general framework for weighted gene co-expression

network analysis. Stat Appl Genet Mol 4(1).

Tables

Table 1. Correlations between alpha diversity metrics and measured environmental variables.

Asterisks represent significance values where $p \leq 0.001$ (***), $p \leq 0.01$ (**), and $p \leq 0.05$ (*).

	Chao 1	Shannon
P	0.31**	-0.09
S	0.42***	0.45***
NO3	0.42***	0.24*
C:N	-0.03	-0.2
NUTR	0.24*	0.41***

Table 2. Pairwise perMANOVA results comparing pairwise differences between wetland regions

and differences between wetland soil depths. Asterisks represent significance values where $p \leq$

0.001 (***), $p \leq 0.01$ (**), $p \leq 0.05$ (*), and *n.s.* representing 'not significant' below p-value

thresholds.

Region	BA	ESBT	LE	NSB	WSB
BA	-				
ESBT	*n.s.*	-			
LE	**	**	-		
NSB	*n.s.*	*n.s.*	**	-	
WSB	*n.s.*	*n.s.*	**	*n.s.*	-

Depth	Top	Middle	Bottom
Top	-		
Middle	**	-	
Bottom	**	*n.s.*	-

Figures

Figure 1. Principal Component Analysis (PCA) illustrating separation of samples based upon soil

geochemistry. Shapes and colors correspond to different wetland depths and regions,

respectively, as listed in the legend. Percentages on axes represent explained variance of that

principal component. Vectors represent impact of specific environmental variables on sample

distribution. NUTR represents OM values, which correlated significantly (p ≤ .01, r > 0.56) to

NO_3^-, OC, OM, S, and TN. Ellipses represent 95% confidence intervals of region groupings.

Figure 2. Boxplot diagram comparing Chao1 diversity among wetland regions. Boxes with the

same letter are not significantly different, while those with no common letters are significantly

different (p ≤ 0.01). Lines within boxes represent the median, hinges represent +/- 25% quartiles,

whiskers represent up to 1.5x the interquartile range. Colors represent wetland region.

Figure 3. Nonmetric Multidimensional Scaling (NMDS) plot illustrating separation of samples

based upon differences in microbial community structure. Shapes and colors correspond to

different depths and wetland regions, respectively, as listed in the legend. Vectors represent

correlations of environmental variables to the distribution of the microbial communities

represented in the plot.

Figure 4. NMDS plots of each wetland region demonstrating separation of samples based upon

differences in microbial community structure, including a. BA, b. ESBT, c. LE, d. NSB, and e.

WSB. Shapes and colors correspond to different depths and wetland sites, respectively, as listed

in the legends. Vectors represent correlations of environmental variables to the distribution of

microbial communities represented in the plots.

Figure 5. Network visualization and results of partial least squares analysis on the subnetwork

most correlated with NUTR. The y-axis represents correlation of OTU to OC values, whereas the

x-axis represents the node centrality. Points represent OTUs, and the color of points corresponds

to the phylum to which an OTU belongs. Point size corresponds to VIP score of that OTU. The

top 15 OTUs are labeled within the graph with corresponding lowest taxonomic identification

possible, and the level of that classification. D = Domain; P = Phylum, C = Class, O = Order, F =

Family, G = Genus.

Figure 6. Network visualization and results of partial least squares analysis on the subnetwork

most correlated with C:N. The y-axis represents correlation of OTU to C:N, whereas the x-axis

represents the node centrality. Points represent OTUs, and the color of points corresponds to the

phylum to which an OTU belongs. Point size corresponds to VIP score of that OTU. Only OTUs

with a VIP score > 1 were displayed for visualization purposes. The top 15 OTUs are labeled

within the graph with corresponding lowest taxonomic identification possible, and the level of

that classification. D = Domain; P = Phylum, C = Class, O = Order, F = Family, G = Genus.

Figure 7. Significant (p ≤ 0.001) positive and negative correlations between microbial taxonomic

groups and NO_3^-. Abundance of OTUs corresponding to taxonomic groups is plotted on the y-

axis. Positive correlations are plotted above the "0" line, while negative correlations are plotted

below the "0" line. Colors of bars correspond to associated phylum of the taxonomic group on

the x-axis.

Figure 8. Geographic map displaying location of sites sampled within this study. Colors of points

correspond to region sampled.

Supplemental Material

Supplemental Figure 1. Depth profiles demonstrating trends in measured environmental

variables with increasing depth among wetland regions and sites. Colors represent wetland

regions, whereas point shapes represent distinct wetland sites.

Supplemental Figure 2. Rarefaction curve analysis demonstrating sequencing depth for alpha

diversity analyses. Different colored lines represent different samples. The vertical black line

represents the sequencing depth used to standardize all samples for alpha diversity analysis. The

dashed line represents the 1:1 slope.

Supplemental Figure 3. Plots demonstrating regression of the frequency distribution of node

connectivity against node connectivity.

Supplemental Figure 4. Individual correlations of established subnetworks with environmental

parameters.

Supplemental Figure 5. Partial least squares analysis results predicting OC (NUTR) values using

relative abundance values of OTUs within the module most connected to OC (NUTR).

Supplemental Figure 6. Partial least squares analysis results predicting C:N values using relative

abundance values of OTUs within the module most connected to C:N.

Supplemental Figure 7. Partial least squares analysis results predicting NO_3^- values using relative

abundance values of OTUs within the module most connected to NO_3^-.

Supplemental Figure 8. Principal Components Analysis demonstrating separation of coastal

wetland sampling locations by site water quality data. Points are color-coded by region.

Percentages on axes represent explained variance of that principal component. Vectors represent

impact of specific environmental variables on sample distribution. Choloro.a = chlorophyll A,

DO = dissolved oxygen, Temp = water temperature, Turb = water turbidity. SumRank represents

SumRank values calculated for each wetland site as outlined by Uzarski et al. (2017). DO also

represents redox potential values, which correlated significantly ($r > 0.7$, $p \leq .001$). Temp also

represents conductivity and total dissolved solids values, which correlated significantly ($r > 0.7$,

$p \leq .001$).