1 Don't throw out the sympatric species with the crater lake water:

2 fine-scale investigation of introgression provides weak support for

3 functional role of secondary gene flow in one of the clearest

4 examples of sympatric speciation

7 Emilie J. Richards[1*], Jelmer W. Poelstra[1,2], and Christopher H. Martin[1]

9 [1] Biology Department, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina,

10 United States of America

11 [2] Biology Department, Duke University, Durham, North Carolina, United States of America

12 *corresponding author: ejr@live.unc.edu

Running title: No functional role for secondary gene flow in sympatric cichlid radiation

Abstract: 232 words; Main text: 5,493 words; Figures: 7; Tables: 2

Abstract

Genomic data has revealed complex histories of colonization and repeated gene flow previously

unrecognized in some of the most celebrated examples of sympatric speciation and radiation.

However, much of the evidence for secondary gene flow into these radiations comes from

genome-wide tests, which tells us little about how gene flow potentially influenced sympatric

diversification. Here we investigated whole genomes of Barombi Mbo crater lake cichlids for

fine-scale patterns of introgression between species with neighboring riverine cichlid

populations. We did find evidence of secondary gene flow into the radiation scattered across <

0.24% of the genome; however, the functional and genetic diversity in these regions paint no

clear picture of how that variation could have contributed to the ecological and morphological

diversity found in the lake. Our results suggest that either variation in novel genetic pathways

introduced during secondary gene flow contributed to the radiation, or that secondary gene flow

was predominantly neutral with respect to the diversification processes. We also found evidence

for differential assortment of ancestral polymorphism found in riverine populations between

sympatric sister species, suggesting the presence of a hybrid swarm in the past. While the history

of gene flow and colonization appears to be more complicated than once thought, the lack of

compelling evidence for secondary gene flow influencing diversification suggests that we should

not yet rule out one of the most celebrated examples of sympatric speciation in nature.

Introduction

Sympatric speciation, the extreme endpoint on the speciation-with-gene-flow continuum, is traditionally defined as the evolution of reproductive isolation without the aid of geographic barriers (Coyne and Orr 2004). Sympatric speciation has fascinated evolutionary biologists since Darwin for its illustration of the power of complex interactions between natural and sexual selection to create new species. Despite intense searches, very few case studies have been able to meet the rigorous criteria for demonstrating sympatric speciation in nature (Coyne and Orr 2004; Bolnick and Fitzpatrick 2007). Even in some of the more convincing examples that do meet these criteria, genomic data have revealed more complex evolutionary histories of multiple colonizations and repeated gene flow than previously thought (Papadopulos et al. 2011; The Heliconius Genome Consortium et al. 2012; Geiger et al. 2013; Alcaide et al. 2014; Igea et al. 2015; Malinsky et al. 2015; Martin et al. 2015a; Kautt et al. 2016).

However, much of the support for complicated histories involving repeated gene flow events into radiations comes from genome-wide tests for gene flow (e.g. (Lamichhaney et al. 2015; Martin et al. 2015a; Meier et al. 2017)). One prediction of models of speciation with gene flow is that divergence between incipient species should be heterogeneous across the genome (Turner et al. 2005; Harr 2006; Feder et al. 2012; Nosil and Feder 2012a,b). Indeed, high heterogeneity in genomic differentiation has been found across the genomes of many recent or incipient sister species (e.g. Jones et al. 2012; Martin et al. 2013; Poelstra et al. 2014; Soria-Carrasco et al. 2014; Malinsky et al. 2015; McGirr and Martin 2016), although other processes besides differential gene flow across the genome can produce similar heterogeneous patterns (Noor and Bennett 2009; Nachman and Payseur 2012; Cutter and Payseur 2013; Cruickshank and Hahn 2014; Guerrero and Hahn 2017; Ravinet et al. 2017). Only a handful of genes may

directly contribute to the speciation process whereas the rest of the genome is porous to gene

flow while reproductive isolation is incomplete (Wu 2001; Wu and Ting 2004). Therefore, gene

flow detected at the genome-wide level from populations outside the sympatric radiation does

not by itself constitute evidence that secondary gene flow was involved in the divergence process

among incipient species and shaped the radiation.

The Cameroon crater lake cichlid radiations are some of the most compelling cases for

sympatric speciation in the wild (Coyne and Orr 2004). The most speciose of these radiations is

found in the isolated 2.3 km-wide volcanic crater lake Barombi Mbo (Trewavas et al. 1972;

Schliewen et al. 1994; Schliewen and Klee 2004). Barombi Mbo hosts a radiation of 11 endemic

cichlid species, many of which have clear morphological and ecological separation from other

sympatric species (Schliewen et al. 1994). Some endemics have evolved unique specializations,

such as the spongivore *Pungu maclareni* and deep-water hypoxia specialist *Konia dikume*

(Trewavas et al. 1972). Other endemics, such as *Stomatepia mariae* and *S. pindu,* appear to be

incipient or stalled species complexes with only slight morphological and ecological divergence

at the extremes of a unimodal distribution of phenotypes (Martin 2012). However, evidence of

differential introgression, weak support for Barombi Mbo monophyly, and differences in levels

of shared ancestry with outgroup riverine populations from genome-wide RAD-seq data suggest

additional secondary gene flow into the radiation after the initial colonization, casting doubt on

one of the best examples of sympatric speciation in the wild (Martin et al. 2015a).

Here we dissect those signals of repeated gene flow to investigate their role in the

radiation using whole-genome sequences. We performed exhaustive searches for all genetic

patterns consistent with secondary gene flow into the ancestral Barombi Mbo population or into

subclades after their initial divergence using machine learning to finely dissect phylogenetic

signal across the genome and genomic scans to test for differential introgression. We find

evidence of both shared introgression between sister species and across subclades in the radiation

as well as differential introgression among sister species across small regions of the genome.

However, functional and genetic diversity in these regions do not paint a clear picture of how

introgressed variants may have contributed to speciation in these groups. Our results suggest that

either 1) rare introgression of variants in novel genetic pathways contributed to the

morphological and ecological diversity of the radiation (speciation with an allopatric phase), 2)

secondary gene flow was predominantly or completely neutral and did not contribute to

diversification in Barombi Mbo (sympatric speciation with gene flow), or 3) multiple

colonizations of the lake before diversification brought in genetic variation that was then

differentially sorted among incipient species (sympatric speciation from a hybrid swarm).

Methods

Sampling and Genome Sequencing

We sequenced whole genomes of 1-3 individuals from 10 out of the 11 species within the

sympatric radiation of Oreochromini cichlids in Cameroon crater lake Barombi Mbo (excluding

Sarotherodon steinbachi which is morphologically and ecologically similar to the other three

Sarotherodon species), an endemic *Sarotherodon* species pair from Lake Ejagham, and outgroup

Sarotherodon individuals from all three river drainages flanking the lake: Cross, Meme, and

Mungo rivers (e.g. see map in (Schliewen et al. 1994)). Details on the collection, extraction,

alignment to the *Oreochromis niloticus* reference genome, and variant calling protocols

following the standard GATK pipeline are provided in the supplementary methods.

Characterization of introgression patterns across the genome

First, we exhaustively searched the genomes for patterns of non-monophyletic Barombi Mbo

relationships using the machine learning program SAGUARO (Zamani et al. 2013) to identify

regions of the genome that contained relationships consistent with expectations from multiple

colorizations and secondary gene flow into the radiation (i.e. paraphyletic/polyphyletic Barombi

Mbo radiations). This method infers relationships among individuals in the form of genetic

distance matrices and assigns segments across the genomes to different topologies without a

priori hypotheses about these relationships. We partitioned the genome into a total of 75 unique

topologies (well past the inflection point at 30 topologies where the percent of genome explained

by each additional topology plateaus; Fig S1) to exhaustively search for relationships where

subclades or individual Barombi Mbo species were more closely related to riverine populations

than other species in the crater lake, suggesting sympatric speciation after a hybrid swarm (i.e.

differential sorting of ancestral polymorphism) or secondary gene flow into this subclade

(introgression). Details on the SAGUARO analysis and filtering strategies for calculating

proportions are provided in the supplementary methods.

We also looked for evidence of differential introgression within subclades of the radiation

on both a genome-wide and local level using f_4 statistics (Reich et al. 2009; Patterson et al. 2012;

Pickrell and Pritchard 2012). The f_4 statistic tests if branches on a four-taxon tree lack residual

genotypic covariance (as expected in the presence of incomplete lineage sorting and no

introgression) by comparing allele frequencies among the three possible unrooted trees.

We focused on tests of introgression with the two outgroup clades from our sample that came

from two main clusters: riverine populations of *Sarotherodon galilaeus* in the Mungo and Meme

rivers (MM) and riverine populations of *S. galilaeus* from the more distant Cross River (CR).

Based on the tree ((P1, P2),(*S. galilaeus* MM, *S. galilaeus* CR)), f_4 statistics were calculated for

combinations of species among a) *Stomatepia*, b) the *Konia + Pungu* subclade, and c) *Myaka*

myaka with *S. linnelli* as a representative of its sister *Sarotherodon* group. This subset of

groupings was chosen to make these analyses more tractable by focusing on species with unique

trophic ecologies within the radiation. Genome-wide f_4 statistics were calculated using the

fourpop function in Treemix (Pickrell and Pritchard 2012). Standard error was estimated by

jackknifing in windows of 1,000 adjacent SNPs to account for linkage disequilibrium.

We characterized heterogeneity in introgression across the genome among these same

combinations and investigated whether differential introgression contributed variation potentially

important in the divergence between species by calculating f_4 statistics in 10-kb sliding windows.

We did this with a modified version of the ABBABABA.py and genomics.py scripts that use

population allele frequencies of biallelic SNPs

(https://github.com/simonhmartin/genomics_general;(Martin et al. 2015b); our modified version

is provided in the supplementary materials). Significance of f_4 values in sliding windows across

the genome were evaluated using the 1% tails of a null distribution generated from permutations

of the f_4 test. For more details on the sliding window calculations of f_4, see supplementary

methods.

For each of these regions, we looked for annotated genes using the well annotated NCBI

Oreochromis Annotation Release 102 and searched their gene ontology in the phenotype

database 'Phenoscape' (Mabee et al. 2012; Midford et al. 2013; Manda et al. 2015; Edmunds et

al. 2016) and AmiGO2 (Balsa-Canto et al. 2016) for pertinent functions to the specializations

and observed morphological differences among species, such as skeletal system or pigmentation.

Directionality of introgression

The sign of f_4 does not indicate the directionality of introgression because of the lack of an

explicit outgroup. For example, in the tree (P1,P2),(P3,P4)), a positive f_4 value indicates gene

flow either between P1 and P3 or P2 and P4. We narrowed down the directionality of

introgression detected in these regions using the f_d statistic, a modified version of the *D*-statistic

that looks at allele frequencies fitting two allelic patterns referred to as ABBA and BABA based

on the tree ((P1,P2),P3,O)), where O is an outgroup species in which no gene flow is thought to

occur with the other populations (Martin et al. 2015b). Using two individuals of *Coptodon kottae*

from another Cameroon crater lake as our distantly related outgroup population and the same

riverine and Barombi Mbo population combinations described above, f_d values were calculated in

10-kb windows across the genome using the same script and window settings as in the f_4 tests of

introgression. The variation in f_d values per region was higher than f_4 of similar window sizes,

perhaps due to the high variance between windows in number of sites that fit the desired

ABBA/BABA patterns given our outgroup, so we used only those f_d outliers (the top 2%) that

overlapped with significant f_4 outliers as potentially introgressed regions from which we could

narrow down the two populations in which gene flow occurred.

We also visualized the directionality of genome-wide introgression detected with the f_4

statistics using Treemix (v 1.13) (Pickrell and Pritchard 2012). Treemix estimates a maximum

likelihood phylogeny of the focal populations and then fits a user-specified number of migration

edges to the tree by comparing genetic covariances between populations. We ran Treemix with *S.*

galilaeus as root, and with 0 through 20 migration edges. To determine the most likely number

of migration events, we performed likelihood-ratio tests comparing each graph to that with one

fewer migration event, starting with 1 versus 0 events, and took as the most likely value the first

non-significant comparison.

Comparison of patterns of introgression to patterns of genetic divergence and diversity

Reduced levels of genetic polymorphism in a population may indicate a strong selective sweep.

We can look at introgressed regions found in only a single Barombi Mbo species for evidence

that they have been adaptive, suggesting that secondary gene flow brought in variation

potentially important for speciation. To examine genetic diversity in candidate introgressed

regions, we calculated between-population nucleotide divergence (D_{xy}) and within-population

nucleotide diversity (π) for pairwise species comparisons among the Barombi Mbo focal species

and the riverine outgroups. D_{xy} and π were calculated over the same 10-kb windows as the f_4 tests

using the python script popGenWindows.py

(https://github.com/simonhmartin/genomics_general; (Martin et al. 2015b); see supplementary

methods for more details on these calculations).

197
Results

Widespread polyphyletic relationships in Barombi Mbo are scattered across small regions of the

genome

After conservative filtering of segments to remove uninformative regions (see supplementary

methods and Table S1), the Barombi Mbo cichlid radiation was a monophyletic group across

53% of the genome and only 0.6% was assigned to topologies indicating a polyphyletic Barombi

Mbo. These polyphyletic relationships are consistent with many patterns, including secondary

gene flow, incomplete lineage sorting, divergent selection, and ancestral population structure.

The most prevalent topology spanned 38.2% of the genome and featured the expected species

phylogeny for this group, in which all Barombi Mbo individuals form a single clade with distant

relationships to outgroup riverine *Sarotherodon* populations in Cameroon (Fig. 1A). The second

most prevalent topology (spanning 11.8% of the genome) featured identical evolutionary

relationships, except for a much shorter branch leading to *S. galilaeus* Mungo and Meme River

populations (Fig. 1B). Branch lengths produced by SAGUARO have no direct interpretation as

an evolutionary distance (analogous to a neighbor-joining tree), but may be useful for

comparison to similar topologies with different branch lengths, e.g. regions with higher

divergence rates (Zamani et al. 2013).

In 0.6% of the genome indicating polyphyletic Barombi Mbo relationships, we found

evidence consistent with multiple colonizations of the lake. Since we were looking for patterns

consistent with secondary gene flow or a hybrid swarm for subclades of the radiation, we

focused on topologies where single species or entire subclades were more closely related to

outgroups than other Barombi Mbo species, which represented only 0.24% of the genome. Some

topologies featured an entire subclade (e.g. *Stomatepia*) as monophyletic, but more closely

related to the riverine populations than other Barombi Mbo species, consistent with a hybrid

swarm scenario before the diversification of the *Stomatepia* subclade. Other topologies featured

individual species more closely related to outgroup riverine populations than sister species,

consistent with secondary gene flow into that lineage after the initiation of divergence. For

example, in *Stomatepia* we found topologies that group multiple species with riverine

populations (Fig. 2A-B), but we also found topologies where individual *Stomatepia* species (*S.*

mariae and *S. pindu*; Fig. 2C-D) were more closely related to riverine outgroups than other

Stomatepia. In the *Konia* + *Pungu* subclade, we saw a similar pattern with topologies for the

hypoxia and sponge-eating specialists (*K. dikume* and *P. maclareni,* respectively; Fig. 3A-B) but

also a topology where the entire subclade was sister to the riverine outgroup populations (Fig.

3C). In the zooplanktivore *M. myaka*, we found topologies in which *M. myaka* was sister to the

riverine populations (Fig. 4A-B), but also topologies where *M. myaka*, along with all the

Barombi Mbo *Sarotherodon* species, were sister to the riverine outgroup populations (Fig. 4C-

D).

Genome-wide evidence for differential introgression into the radiation

Consistent with evidence of differential introgression from RAD-seq data (Martin et al. 2015a),

genome-wide f_4 tests provided evidence of genome-wide differential gene flow between some

Barombi Mbo sister species and the outgroup riverine species (Table 1). There was significant

evidence of genome-wide introgression in tests involving both *S. pindu* in the *Stomatepia* species

complex and the hypoxia specialist *K. dikume* in the *Konia + Pungu* subclade. Some species pair

combinations within these subclades did not show evidence of differential gene flow, suggesting

that there may still be sympatric speciation occurring for some species, if not entire subclades.

For example, there was no significant secondary gene flow detected genome-wide in the tests

involving sister species *S. mariae* and *S. mongo* or *M. myaka* and *S. linnelli* (Table 1).

We also found evidence for widespread gene flow connecting populations across

Barombi Mbo and neighboring riverine populations in highly interconnected population graphs;

the likelihood of each graph did not plateau until reaching 10 admixture events (Fig S5). On the

Treemix population graph with 10 admixture events, gene flow from the Mungo/Meme River

populations of *S. galilaeus* occurred directly into individual species *S. mongo* and *K. eisentrauti*

rather than the ancestral node of their respective subclades (Fig S6). The proportion of admixture

inferred for these two events (0.1% into *S. mongo* and 0.4% into *K. eisentrauti*) was similar to

the small proportions of the genome assigned to topologies consistent with secondary gene flow

in the SAGUARO analyses. These admixture events pointing to the tips of the graphs suggest

secondary gene flow events between nearby riverine populations and individual species within

the radiation. In all population graphs allowing up to 21 migration events, any admixture from

outgroup riverine populations appears to be coming from the Mungo and Meme rivers rather

than the Cross River, consistent with the closer geographic proximity of the former drainages.

Very few genomic regions contain signatures of differential introgression between sister species

Very few regions of the genome introgressed into single species from outgroup riverine

populations (Fig 5A-C). In *Stomatepia*, only one region introgressed from Mungo/Meme Rivers

into *S. pindu* and only three regions into *S. mariae*, respectively, suggesting secondary gene flow

after initial diversification of *Stomatepia* (Table 2). Similarly, secondary introgression occurred

into the *Konia + Pungu* subclade (Table 2). However, there was also evidence of shared

introgression signals among sister species across all three subclades, where two subclade sister

species shared introgressed regions from a riverine population. Only 0.000017- 0.0000354% of

the genome appears introgressed into a single species of a Barombi Mbo subclade. This number

is smaller than suggested in the SAGUARO analysis perhaps due to the conservative

significance cut-offs and window size choice for f_4 statistic and that relationships observed in the

polyphyletic topologies are consistent with other patterns besides introgression.

Evidence for sympatric sorting of ancestral polymorphism within a hybrid swarm

A few of these 10-40 kb regions with peak signals of introgression were also present in multiple

subclades, indicating differential assortment of introgressed variation shared among clades. For

example, two significant f_4 outliers on linkage group 20 out of the 35 found across the genome

appear within the *Stomatepia*, *Konia*, and *Pungu*, suggesting that some of this introgression may

have occurred in the ancestral stages of the radiation and differentially sorted among species.

We also found 11 regions across the sister species pairs in which one species was more

similar to one outgroup riverine population while its sister species was more similar to the other

riverine population (Table 2). This signal is consistent with a hybrid swarm scenario due to

multiple colonizations by riverine populations before diversification of some of the sister species

and the sorting of polymorphisms brought in by these populations among incipient Barombi Mbo

species. For example, two regions that appear to be differentially sorted between *S. galilaeus* and

S. mariae and *S. pindu* from *S. galilaeus* CR versus *S. mongo* from *S.galilaeus* MM. Similar

patterns were found scattered across the genomes for *K. eisentrauti* and *K. dikume* versus *P.*

maclareni and *M. myaka* versus *S. linelli*.

Weak support for functional importance of introgressed regions for species diversification

Although we did find evidence of differential introgression among sister species scattered across

a small proportion of the genome, the types of genes found in these regions painted no clear

picture of how introgressed variation may have contributed to speciation (Table 2). For example,

differential introgression in *Stomatepia* occurred in regions with genes involved in a large range

of biological processes, including intracellular signal transduction, immune system response, and

motor neuron axon development (Table 2), with no obvious links to the highly divergent

morphological, ecological, or patterning traits observed between these species (Martin 2012) nor

to those traits normally associated with adaptive radiation in cichlid fishes such as body shape,

pharyngeal jaw morphology, retinal pigments, or male coloration (Kocher 2004; Barluenga et al.

2006; Wagner et al. 2012; Brawand et al. 2014; Malinsky et al. 2015; Meier et al. 2017).

Similarly, in both the *Konia* + *Pungu* and *Myaka* + *Sarotherodon* subclades, introgressed regions

were near genes involved in a large range of biological processes not clearly associated with

adaptive ecological traits in these species, such as *K. dikume*'s hypoxia tolerance, *P. maclareni's*

spongivory, and *M. myaka's* zooplanktivory. For example, while there appears to be differential

introgression in *Konia* in a region containing *pafah1b3*, a gene involved in platelet activation

activity, and *K. dikume's* deep water specialization includes higher blood volume with higher

concentrations of hemoglobin (Green et al. 1973), it is not obvious how introgressed variation in

pafah1b3 would have played a role in the evolution of these traits from studies of its function in

model organisms, which includes spermatogenesis and sterility in mice (Prescott et al. 2000;

Koizumi et al. 2003; Yan et al. 2003).

Similarly, the amount of genetic diversity in introgressed regions does not suggest strong

divergent selection on introgressed genetic variation due to hard selective sweeps. In line with

the presence of peaks in f_4 values in these regions, between-population diversity (D_{xy}) was

typically high between one of the species and its sister species (Fig. 6). However, within-

population diversity across many of these regions was often greater or comparable to scaffold

and genome-wide averages (Table S2-4), suggesting these regions may not have experienced

hard selective sweeps that would support their role in adaptive divergence among species (Fig.

6). In summary, although we found evidence for differential secondary gene flow between sister

species in the radiation, we did not find strong functional support from gene ontology terms nor

signatures of selection that the introgressed alleles were important for sympatric species

diversification.

Discussion

Little evidence that secondary gene flow promoted the diversification of Barombi Mbo cichlids

Our fine-scale investigations of introgression across the genomes of a celebrated putative

example of sympatric speciation are consistent with two possible scenarios: 1) sympatric

speciation in the presence of continuous neutral secondary gene flow into the radiation, or 2)

speciation initiated by secondary gene flow. We found little support for the latter allopatric

scenario from both a learning machine and sliding-window approach. From the SAGUARO

analyses, our most conservative estimate of introgression into single species of the radiation

ranges from 0.013 -0.019% of the genome. Estimates are similarly small from the f_4 statistics,

ranging from 0.000017- 0.0000354% of the genome (Fig 7). Furthermore, even these significant

outliers may represent false positives. First, our method of selecting introgressed regions from

the 1% tails of a null distribution can always find outliers, even in the absence of introgression.

Second, it is also difficult to distinguish signatures of differential introgression from the biased

assortment of ancestral polymorphism into modern lineages, e.g. a hybrid swarm scenario that

would still result in sympatric divergence entirely within the crater lake. Finally, even if our

statistical outliers represent differentially introgressed regions, their importance to the speciation

process is equivocal. We found no evidence of selective sweeps in these regions that would

suggest they aided in divergence between species and they contain mainly housekeeping genes

that do not clearly suggest how introgressed variation would have contributed to the radiation.

This contrasts studies on other systems using similar approaches which found compelling

cases for adaptive introgression contributing to diversification (e.g. Abi-Rached et al. 2011; The

Heliconius Genome Consortium et al. 2012; Huerta-Sánchez et al. 2014; Lamichhaney et al.

2015; Stankowski and Streisfeld 2015; Arnold et al. 2016; Meier et al. 2017), including our own

previous work (Richards and Martin 2017). For example, several studies have found convincing

candidate genes/variants in introgressed regions to suggest that adaptive introgression played a

role in shaping ecological and morphological diversity. These include the detection of

introgressed alleles linked to wing-color patterning involved in mimicry and mate selection in

Heliconius butterflies (The Heliconius Genome Consortium et al. 2012), flower coloration

involved in pollinator preferences for *Mimulus* species (Stankowski and Streisfeld 2015), and

oral jaw size variation involved in scale-eating trophic specialization in *Cyprinodon* pupfishes

(Richards and Martin 2017).

Evidence for a hybrid swarm further complicates the role of gene flow in the speciation process

in Barombi Mbo cichlids

Beyond speciation scenarios involving secondary gene flow, our findings also suggest another

scenario for sympatric speciation in this system: sympatric speciation from a hybrid swarm

involving the differential sorting of ancestral polymorphism among incipient species. A hybrid

swarm is not easily detectable using the f_4 statistic because introgressed variation could be shared

among diverging sister species, leading to an f_4 value of zero (Reich et al. 2009; Patterson et al.

2012). However, many of the f_4 peaks appear to be shared across at least two of the sister species

in a subclade, shared between species of different subclades, or contain variation from both

riverine populations (Mungo/Meme and Cross Rivers) that has been differentially sorted among

sister species. All three of these patterns are consistent with an ancestral hybrid swarm before

divergence between sister species occurred. This pattern of differential sorting of variation from

a hybrid swarm from f_d analyses could also result from a lack of power in the statistic to

distinguish the directionality of the introgression detected in those regions when using biallelic

patterns and four populations (e.g. when two populations share similar allele patterns, the other

two populations can share the opposite allele pattern by default). However, we also found

evidence that entire subclades (e.g. *Stomatepia*) were more closely related to riverine populations

than other Barombi Mbo subclades from the SAGUARO analyses that are also consistent with a

hybrid swarm (e.g. Fig 2).

There are some caveats to our interpretations of secondary gene flow and its weak

functional role in the ecological and morphological diversity observed within the lake.

Recombination rate varies across genomes and determines the scale over which patterns of

admixture and differentiation vary (Smukowski and Noor 2011). In our fixed sliding window

size of 10-kb, we may have missed important patterns of introgression in regions of

recombination hotspots, where such patterns are expected to be very fine-scale. Shared variation

among species may reflect unsorted polymorphism from structured ancestral populations rather

than hybridization. Introgression events can also be hard to distinguish from ongoing balancing

selection of ancestral polymorphism that is sieved between species (Guerrero and Hahn 2017).

While we focused on searching for genetic signatures of hard selective sweeps in introgressed

regions, some of them with intermediate to high nucleotide diversity may have undergone soft

selective sweeps, when selection drives multiple adaptive haplotypes to fixation. Some of these

introgressed regions may have been adaptive and undergone soft selective sweeps, although the

relative contributions of hard sweeps versus soft sweeps during adaptation and speciation is still

the subject of much debate (Hermisson and Pennings 2005, 2017; Pritchard et al. 2010; Jensen

2014; Schrider et al. 2015).

Best remaining cases for sympatric speciation within Barombi Mbo cichlid radiation

While the radiation as a whole may not have entirely arisen from sympatric speciation, some

sister species within Barombi Mbo are better case studies of the process than others. Within the

three-species *Stomatepia* subclade, there is little evidence that secondary gene flow played an

important role in diversification. On a genome-wide level, we detected secondary gene flow in f_4

tests involving *S. pindu*. However, on a finer scale the one introgressed region unique to *S. pindu*

is unannotated and the three introgressed regions unique to *S. mariae* contain four housekeeping

genes involved in extracellular exosome activity and plasma membranes

(*jmjd8,prss1,cldn4,muc19*). Shared signals of introgression among the three species represent a

larger proportion of the genome than differentially introgressed regions, although both types of

introgressed material appear to be rare in the genome ($< 0.045\%$; Fig 7). The high ecological and

morphological overlap among *Stomatepia* species suggests that this species complex may be

stalled in the earliest stages of divergence. For example, *S. pindu* and *S. mariae* appear be at the

extremes of a unimodal distribution of phenotypes; this is one major prediction of sympatric

speciation models in the presence of only weak disruptive selection on ecological traits (e.g.

(Matessi et al. 2002; Burger et al. 2006)), as measured in this species pair (Martin 2012).

Even for the two monotypic specialist species *M. myaka* and *P. maclareni*, there is

minimal evidence for a role of secondary gene flow in the evolution of their trophic

specializations. On a genome-wide level, the tests for differential introgression with one of the

specialists and another species from the radiation were not significant. For the two regions which

appear to be differentially introgressed between riverine populations and the zooplantivore *M.*

myaka, one has no annotations, suggesting it may be neutral gene flow while the other is an

introgressed region in other subclades, a signal of differential sorting of variation after a hybrid

swarm. For spongivore specialist *Pungu maclareni*, we found only a single region differentially

introgressed with riverine populations. This region contains *ddn1*, a gene involved in serine-type

peptidase activity, proteins which are found ubiquitously in prokaryotes and eukaryotes.

Among all the ecologically divergent species pairs focused on in this study, *K. eisentrauti*

and *K. dikume* are the least convincing as a putative case for sympatric speciation between sister

species. Similar to *Stomatepia*, there is more evidence for shared introgression in regions of the

Konia genome than differentially introgressed regions (Fig 8). However, differential

introgression between the two *Konia* species occurs in regions with the best potential candidates

in this study for contributing to diversification (although these regions are still not as strong as

the 'smoking guns' observed in our past study of introgressed variation, e.g. Richards and Martin

2017). The gene *pafah1b3*, which is involved in platelet activation activity, is differentially

introgressed between *Konia* species and may function in *K. dikume's* deep water specialization

of higher blood volume with higher concentrations of hemoglobin (Green et al. 1973). However,

studies of the phenotypic effects of *pafah1b3* in model organisms do not hint at a role in

hemoglobin concentration. Instead, mutations in this gene have been suggested to play a role in

male spermatogenesis and fertility in mice (Koizumi et al. 2003; Yan et al. 2003). We also see

differential introgression in a region containing the gene *ehmt2*, which is involved in

neurogenesis and retinal cell differentiation. While it is not as directly clear how introgressed

variation in this gene would have contributed to divergence among the *Konia* sister species,

variation in visual traits such as color perception are important axes of diversification in other

cichlid radiations, particularly along a depth gradient (Terai et al. 2002, 2006; Seehausen et al.

2008). These two species also exist in microallopatry; *K. eisentrauti* is an abundant detritivore

along the shallow littoral region of the lake while *K. dikume* is a deep-water specialist on

Chaoborus midge larvae which is only collected in deep-water gill nets (Trewavas et al. 1972;

Schliewen 1994). Both species are mouthbrooders and likely breed in non-overlapping habitats.

Conclusion

The complex history of colonization in the crater evidenced in this and a previous genome-wide

study suggests allopatric phases of the speciation process in the radiation, which violates one of

the strict criteria for demonstrating sympatric speciation in the wild (Coyne and Orr 2004).

Nonetheless, from our fine-scale dissection of where in the genome these signals are coming

from, we cannot point to a functional role for secondary gene flow in the speciation process

across any of the subclades. This suggests that either variation in genes with undiscovered

functional effects underlies the divergent ecologies and morphologies seen in the lake or that any

secondary gene flow was neutral with regard to its role in the speciation process. We also find

evidence to support sympatric speciation after a hybrid swarm that resulted from multiple

colonizations of the lake, still consistent with a scenario of sympatric speciation through

differential sorting of ancestral polymorphism. Disentangling the effects of a putative hybrid

swarm from secondary contact events on the speciation process will require a better

understanding of the timing of gene flow events compared to the diversification times of

Barombi Mbo species. We found evidence for gene flow into the radiation both before and after

initial diversification of subclades within the lake. Even without this information, weak support

for a functional role of secondary gene flow in the radiation of Barombi Mbo cichlids suggests

that we should not rule out this example of a sympatric radiation just yet.

Acknowledgements

This study was funded by a National Geographic Society Young Explorer's Grant, a Lewis and

Clark Field Research grant from the American Philosophical Society, and the University of

North Carolina at Chapel Hill to CHM. We gratefully acknowledge the Cameroonian

government and the local chiefs of Barombi Mbo village and surrounding communities for

permission to conduct this research. We thank Cyrille Dening Touokong and Jackson Waite-

Himmelwright for field assistance and Nono Gonwuou for help obtaining permits.

Author contributions

EJR wrote the manuscript. EJR conducted all analyses except for Treemix; JP conducted

Treemix analyses. CHM collected samples and provided the genomic data. All authors

contributed to the development of ideas presented in the study and revised the manuscript.

Data Accessibility

All datasets used for this study will be deposited in Dryad and the NCBI Short Read Archive

(SRA).

Competing Interests

We declare no competing financial interests.

References

Abi-Rached, L., M. J. Jobin, S. Kulkarni, A. McWhinnie, K. Dalva, L. Gragert, F. Babrzadeh, B.

Gharizadeh, M. Luo, F. A. Plummer, J. Kimani, M. Carrington, D. Middleton, R.

Rajalingam, M. Beksac, S. G. E. Marsh, M. Maiers, L. A. Guethlein, S. Tavoularis, A.-M.

Little, R. E. Green, P. J. Norman, and P. Parham. 2011. The Shaping of Modern Human

Immune Systems by Multiregional Admixture with Archaic Humans. Science (80-.).

334:89 LP-94.

Alcaide, M., E. S. C. Scordato, T. D. Price, and D. E. Irwin. 2014. Genomic divergence in a ring

species complex. Nature 511:83–85. Nature Publishing Group.

Arnold, B. J., B. Lahner, J. M. DaCosta, C. M. Weisman, J. D. Hollister, D. E. Salt, K. Bomblies,

and L. Yant. 2016. Borrowed alleles and convergence in serpentine adaptation. Proc. Natl.

Acad. Sci. 113:8320–8325.

Balsa-Canto, E., D. Henriques, A. Gabor, and J. R. Banga. 2016. AMIGO2, a toolbox for

dynamic modeling, optimization and control in systems biology. Bioinformatics 32:1–2.

Barluenga, M., K. N. Stölting, W. Salzburger, M. Muschick, and A. Meyer. 2006. Sympatric

speciation in Nicaraguan crater lake cichlid fish. Nature 439:719–723.

Bolnick, D. I., and B. M. Fitzpatrick. 2007. Linked references are available on JSTOR for this

article : Sympatric Speciation : Models and Empirical Evidence. Annu. Rev. Ecol. Evol.

Syst. 38:459–487.

Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, I. Keller, S. Fan, O. Simakov, A. Y. Ng, Z.

W. Lim, E. Bezault, Turner-Maier, J. J. Johnson, R. Alcazar, H. J. Noh, P. Russell, B. Aken,

J. Alföldi, C. Amemiya, N. Azzouzi, J.-F. Baroiller, F. Barloy-Hubler, A. Berlin, R.

Bloomquist, K. L. Carleton, M. a. Conte, H. D'Cotta, O. Eshel, L. Gaffney, F. Galibert, H.

F. Gante, S. Gnerre, L. Greuter, R. Guyon, N. S. Haddad, W. Haerty, R. M. Harris, H. a.

Hofmann, T. Hourlier, G. Hulata, D. B. Jaffe, M. Lara, L. A.P., I. MacCallum, S. Mwaiko,

M. Nikaido, H. Nishihara, C. Ozouf-Costaz, D. J. Penman, D. Przybylski, M. Rakotomanga,

S. C. P. Renn, F. J. Ribeiro, M. Ron, W. Salzburger, L. Sanchez-Pulido, M. E. Santos, S.

Searle, T. Sharpe, R. Swofford, F. J. Tan, L. Williams, S. Young, S. Yin, N. Okada, T. D.

Kocher, E. a. Miska, E. S. Lander, B. Venkatesh, R. D. Fernald, A. Meyer, C. P. Ponting, J.

T. Streelman, K. Lindblad-Toh, O. Seehausen, and F. Di Palma. 2014. The genomic

substrate for adaptive radiation in African cichlid fish. Nature 513:375–381.

Burger, R., K. A. Schneider, and M. Willensdorfer. 2006. The Conditions for Speciation through

Intraspecific Competition. Evolution (N. Y). 60:2185–2206. [Society for the Study of

Evolution, Wiley].

Coyne, J. A., and H. A. Orr. 2004. Speciation. Sutherland, MA.

Cruickshank, T. E., and M. W. Hahn. 2014. Reanalysis suggests that genomic islands of

speciation are due to reduced diversity, not reduced gene flow. Mol. Ecol. 23:3133–3157.

Cutter, A. D., and B. A. Payseur. 2013. Genomic signatures of selection at linked sites: unifying

the disparity among species. Nat. Rev. Genet. 14:262–274. Nature Publishing Group.

Edmunds, R. C., B. Su, J. P. Balhoff, B. F. Eames, W. M. Dahdul, H. Lapp, J. G. Lundberg, T. J.

Vision, R. A. Dunham, P. M. Mabee, and M. Westerfield. 2016. Phenoscape: Identifying

candidate genes for evolutionary phenotypes. Mol. Biol. Evol. 33:13–24.

Feder, J. L., S. P. Egan, and P. Nosil. 2012. The genomics of speciation-with-gene-flow. Trends

Genet. 28:342–350.

Geiger, M. F., J. K. McCrary, and U. K. Schliewen. 2013. Crater Lake Apoyo Revisited -

Population Genetics of an Emerging Species Flock. PLoS One 8:1–17.

Green, J., S. A. Corbet, and E. Betney. 1973. Ecological studies on crater lakes in West

Cameroon The blood of endemic cichlids in Barombi Mbo in relation to stratification and

their feeding habits. J. Zool. 170:299–308. Blackwell Publishing Ltd.

Guerrero, R. F., and M. W. Hahn. 2017. Speciation as a sieve for ancestral polymorphism. Mol.

Ecol. 1–7.

Harr, B. 2006. Genomic islands of differentiation between house mouse subspecies Genomic

islands of differentiation between house mouse subspecies. 730–737.

Hermisson, J., and P. S. Pennings. 2005. Soft Sweeps. Genetics 169:2335 LP-2352.

Hermisson, J., and P. S. Pennings. 2017. Soft sweeps and beyond: understanding the patterns and

probabilities of selection footprints under rapid adaptation. Methods Ecol. Evol. 8:700–716.

Huerta-Sánchez, E., X. Jin, Asan, Z. Bianba, B. M. Peter, N. Vinckenbosch, Y. Liang, X. Yi, M.

He, M. Somel, P. Ni, B. Wang, X. Ou, Huasang, J. Luosang, Z. X. P. Cuo, K. Li, G. Gao,

Y. Yin, W. Wang, X. Zhang, X. Xu, H. Yang, Y. Li, J. Wang, J. Wang, and R. Nielsen.

2014. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA.

Nature 512:194–197.

Igea, J., D. Bogarín, A. S. T. Papadopulos, and V. Savolainen. 2015. A comparative analysis of

island floras challenges taxonomy-based biogeographical models of speciation. Evolution

(N. Y). 69:482–491.

Jensen, J. D. 2014. On the unfounded enthusiasm for soft selective sweeps. Nat. Commun.

5:5281.

Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, E. Mauceli, J. Johnson, R. Swofford, M.

Pirun, M. C. Zody, S. White, E. Birney, S. Searle, J. Schmutz, J. Grimwood, M. C. Dickson,

R. M. Myers, C. T. Miller, B. R. Summers, A. K. Knecht, S. D. Brady, H. Zhang, A. A.

Pollen, T. Howes, C. Amemiya, Broad Institute Genome Sequencing Platform & Whole

Genome Assembly Team, E. S. Lander, F. Di Palma, K. Lindblad-Toh, and D. M. Kingsley.

2012. The genomic basis of adaptive evolution in threespine sticklebacks. Nature 484:55–

61.

Kautt, A. F., G. Machado-Schiaffino, J. Torres-Dowdall, and A. Meyer. 2016. Incipient

sympatric speciation in Midas cichlid fish from the youngest and one of the smallest crater

lakes in Nicaragua due to differential use of the benthic and limnetic habitats? Ecol. Evol.

6:5342–5357.

Kocher, T. D. 2004. Adaptive evolution and explosive speciation: the cichlid fish model. Nat.

Rev. Genet. 5:288–298.

Koizumi, H., N. Yamaguchi, M. Hattori, T. O. Ishikawa, J. Aoki, M. M. Taketo, K. Inoue, and

H. Arai. 2003. Targeted disruption of intracellular type I platelet activating factor-

acetylhydrolase catalytic subunits causes severe impairment in spermatogenesis. J. Biol.

Chem. 278:12489–12494.

Lamichhaney, S., J. Berglund, M. S. Almén, K. Maqbool, M. Grabherr, A. Martinez-Barrio, M.

Promerová, C.-J. Rubin, C. Wang, N. Zamani, B. R. Grant, P. R. Grant, M. T. Webster, and

L. Andersson. 2015. Evolution of Darwin's finches and their beaks revealed by genome

sequencing. Nature 518:371–375.

Mabee, P., J. P. Balhoff, W. M. Dahdul, H. Lapp, P. E. Midford, T. J. Vision, and M.

Westerfield. 2012. 500,000 fish phenotypes: The new informatics landscape for

evolutionary and developmental biology of the vertebrate skeleton. J. Appl. Ichthyol.

28:300–305.

Malinsky, M., R. J. Challis, A. M. Tyers, S. Schiffels, Y. Terai, B. P. Ngatunga, E. A. Miska, R.

Durbin, M. J. Genner, and G. F. Turner. 2015. Genomic islands of speciation separate

cichlid ecomorphs in an East African crater lake. Science (80-.). 350:1493–1498.

Manda, P., J. P. Balhoff, H. Lapp, P. Mabee, and T. J. Vision. 2015. Using the phenoscape

knowledgebase to relate genetic perturbations to phenotypic evolution. Genesis 53:561–

571.

Martin, C. H. 2012. Weak Disruptive Selection and Incomplete Phenotypic Divergence in Two

Classic Examples of Sympatric Speciation: Cameroon Crater Lake Cichlids. Am. Nat.

180:E90–E109.

Martin, C. H., J. S. Cutler, J. P. Friel, C. Dening Touokong, G. Coop, and P. C. Wainwright.

2015a. Complex histories of repeated gene flow in Cameroon crater lake cichlids cast doubt

on one of the clearest examples of sympatric speciation. Evolution (N. Y). 69:1406–1422.

Martin, S. H., K. K. Dasmahapatra, N. J. Nadeau, C. Slazar, J. R. Walters, F. Simpson, M.

Blaxter, A. Manica, J. Mallet, and C. D. Jiggins. 2013. Genome-wide evidence for

speciation with gene flow in Heliconius butterflies. Genome Res. 23:1817–1828.

Martin, S. H., J. W. Davey, and C. D. Jiggins. 2015b. Evaluating the use of ABBA-BABA

statistics to locate introgressed loci. Mol. Biol. Evol. 32:244–257.

Matessi, C., A. Gimelfarb, and S. Gavrilets. 2002. Long-term Buildup of Reproductive Isolation

Promoted by Disruptive Selection: How Far Does it Go? Selection 2:41–64. Akadémiai

Kiadó.

McGirr, J. A., and C. H. Martin. 2016. Novel candidate genes underlying extreme trophic

specialization in Caribbean pupfishes. Mol. Biol. Evol. msw286.

Meier, J. I., D. A. Marques, S. Mwaiko, C. E. Wagner, L. Excoffier, and O. Seehausen. 2017.

Ancient hybridization fuels rapid cichlid fish adaptive radiations. Nat. Commun. 8:14363.

Nature Publishing Group.

Midford, P. E., T. Dececchi, J. P. Balhoff, W. M. Dahdul, N. Ibrahim, H. Lapp, J. G. Lundberg,

P. M. Mabee, P. C. Sereno, M. Westerfield, T. J. Vision, D. C. Blackburn, S. Federhen, P.

Mabee, J. Balhoff, W. Dahdul, H. Lapp, P. Midford, T. Vision, M. Westerfield, W. Dahdul,

J. Balhoff, J. Engeman, T. Grande, E. Hilton, C. Kothari, H. Lapp, J. Lundberg, P. Midford,

T. Vision, M. Westerfield, P. Mabee, F. Fang, M. Toledo-Piza, M. Uhen, A. Barnosky, B.

Bills, J. Blois, M. Carrano, M. Carrasco, G. Erickson, J. Eronen, M. Fortelius, R. Graham,

E. Grimm, M. O'Leary, A. Mast, W. Piel, P. Polly, L. Saila, J. Balhoff, P. Midford, H.

Lapp, M. Ghiselin, S. Schulz, H. Stenzhorn, M. Boeker, D. Thau, N. Franz, B. Smith, M.

Ashburner, C. Rosse, J. Bard, W. Bug, W. Ceusters, L. Goldberg, K. Eilbeck, A. Ireland, C.

Mungall, O. Consortium, N. Leontis, P. Rocca-Seraa, A. Ruttenberg, S. Sansone, R.

Scheuermann, N. Shah, P. Whetzel, S. Lewis, J. Beaulieu, R. Ree, J. Cavender-Bares, G.

Weiblen, and M. Donoghue. 2013. The vertebrate taxonomy ontology: a framework for

reasoning across model organism and species phenotypes. J. Biomed. Semantics 4:34.

Nachman, M. W., and B. A. Payseur. 2012. Recombination rate variation and speciation:

theoretical predictions and empirical results from rabbits and mice. Philos. Trans. R. Soc. B

Biol. Sci. 367:409–421.

Noor, M., and S. Bennett. 2009. Islands of speciation or mirages in the desert? Examingin the

role of restricted recombination in maintaining species. Heredity (Edinb). 103:439–444.

Nosil, P., and J. L. Feder. 2012a. Genomic divergence during speciation: causes and

consequences. Philos. Trans. R. Soc. B Biol. Sci. 367:332–342.

Nosil, P., and J. L. Feder. 2012b. Widespread yet heterogeneous genomic divergence. Mol. Ecol.

21:2829–2832.

Papadopulos, A. S. T., W. J. Baker, D. Crayn, R. K. Butlin, R. G. Kynast, I. Hutton, and V.

Savolainen. 2011. Speciation with gene flow on Lord Howe Island. Proc. Natl. Acad. Sci.

108:13188–13193.

Patterson, N., P. Moorjani, Y. Luo, S. Mallick, N. Rohland, Y. Zhan, T. Genschoreck, T.

Webster, and D. Reich. 2012. Ancient Admixture in Human History. Genetics 192:1069–

1093.

Pickrell, J. K., and J. K. Pritchard. 2012. Inference of Population Splits and Mixtures from

Genome-Wide Allele Frequency Data. PLoS Genet. 8.

Poelstra, J. W., N. Vijay, C. M. Bossu, H. Lantz, B. Ryll, I. Muller, V. Baglione, P. Unneberg,

M. Wikelski, M. G. Grabherr, and J. B. W. Wolf. 2014. The genomic landscape underlying

phenotypic integrity in the face of gene flow in crows. Science (80-.). 344:1410–1414.

Prescott, S. M., G. a Zimmerman, D. M. Stafforini, and T. M. Mcintyre. 2000. Platelet activating

factor. Annu. Rev. Biochem. 69:419–445.

Pritchard, J. K., J. K. Pickrell, and G. Coop. 2010. The Genetics of Human Adaptation: Hard

Sweeps, Soft Sweeps, and Polygenic Adaptation. Curr. Biol. 20:R208–R215.

Ravinet, M., R. Faria, R. K. Butlin, J. Galindo, N. Bierne, M. Rafajlović, M. A. F. Noor, B.

Mehlig, and A. M. Westram. 2017. Interpreting the genomic landscape of speciation: a road

map for finding barriers to gene flow. J. Evol. Biol. 30:1450–1477.

Reich, D., K. Thangaraj, N. Patterson, A. L. Price, and L. Singh. 2009. Reconstructing Indian

population history. Nature 461:489–94.

Richards, E. J., and C. H. Martin. 2017. Adaptive introgression from distant Caribbean islands

contributed to the diversification of a microendemic adaptive radiation of trophic specialist

pupfishes.

Schliewen, U. K., and B. Klee. 2004. Reticulate sympatric speciation in Cameroonian crater lake

cichlids. Front. Zool. 1:5.

Schliewen, U. K., D. Tautz, and S. Paabo. 1994. Sympatric speciation suggested by monophyly

of crater lake cichlids. Nature 368:629–631.

Schrider, D. R., F. K. Mendes, M. W. Hahn, and A. D. Kern. 2015. Soft Shoulders Ahead:

Spurious Signatures of Soft and Partial Selective Sweeps Result from Linked Hard Sweeps.

Genetics 200:267 LP-284.

Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. J. Mrosso, R. Miyagi, I. van der

Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai, and N. Okada. 2008. Speciation

through sensory drive in cichlid fish. Nature 455:620–626.

Smukowski, C. S., and M. A. F. Noor. 2011. Recombination rate variation in closely related

species. Heredity (Edinb). 107:496–508. The Genetics Society.

Soria-Carrasco, V., Z. Gompert, A. A. Comeault, T. E. Farkas, T. L. Parchman, J. S. Johnston, C.

A. Buerkle, J. L. Feder, J. Bast, T. Schwander, S. P. Egan, B. J. Crespi, and P. Nosil. 2014.

Stick Insect Genomes Reveal Natural Selection's Role in Parallel Speciation. Science (80-.

). 344:738 LP-742.

Stankowski, S., and M. A. Streisfeld. 2015. Introgressive hybridization facilitates adaptive

divergence in a recent radiation of monkeyflowers. Proc. R. Soc. London B 282:20151666.

Terai, Y., W. E. Mayer, J. Klein, H. Tichy, and N. Okada. 2002. The effect of selection on a long

wavelength-sensitive (LWS) opsin gene of Lake Victoria cichlid fishes. Proc. Natl. Acad.

Sci. 99:15501–15506.

Terai, Y., O. Seehausen, T. Sasaki, K. Takahashi, S. Mizoiri, T. Sugawara, T. Sato, M.

Watanabe, N. Konijnendijk, H. D. J. Mrosso, H. Tachida, H. Imai, Y. Shichida, and N.

Okada. 2006. Divergent selection on opsins drives incipient speciation in Lake Victoria

cichlids. PLoS Biol. 4:2244–2251.

The Heliconius Genome Consortium, K. K. Dasmahapatra, J. R. Walters, A. D. Briscoe, J. W.

Davey, A. Whibley, N. J. Nadeau, A. V. Zimin, D. S. T. Hughes, L. C. Ferguson, S. H.

Martin, C. Salazar, J. J. Lewis, S. Adler, S.-J. Ahn, D. a. Baker, S. W. Baxter, N. L.

Chamberlain, R. Chauhan, B. a. Counterman, T. Dalmay, L. E. Gilbert, K. Gordon, D. G.

Heckel, H. M. Hines, K. J. Hoff, P. W. H. Holland, E. Jacquin-Joly, F. M. Jiggins, R. T.

Jones, D. D. Kapan, P. Kersey, G. Lamas, D. Lawson, D. Mapleson, L. S. Maroja, A.

Martin, S. Moxon, W. J. Palmer, R. Papa, A. Papanicolaou, Y. Pauchet, D. a. Ray, N.

Rosser, S. L. Salzberg, M. a. Supple, A. Surridge, A. Tenger-Trolander, H. Vogel, P. a.

Wilkinson, D. Wilson, J. a. Yorke, F. Yuan, A. L. Balmuth, C. Eland, K. Gharbi, M.

Thomson, R. a. Gibbs, Y. Han, J. C. Jayaseelan, C. Kovar, T. Mathew, D. M. Muzny, F.

Ongeri, L.-L. Pu, J. Qu, R. L. Thornton, K. C. Worley, Y.-Q. Wu, M. Linares, M. L.

Blaxter, R. H. Ffrench-Constant, M. Joron, M. R. Kronforst, S. P. Mullen, R. D. Reed, S. E.

Scherer, S. Richards, J. Mallet, W. Owen McMillan, and C. D. Jiggins. 2012. Butterfly

genome reveals promiscuous exchange of mimicry adaptations among species. Nature

487:94–98.

Trewavas, E., J. Green, and S. A. Corbet. 1972. Ecological studies on crater lakes in West

Cameroon fishes of Barombi Mbo. J. Zool. 41–95.

Turner, T. L., M. W. Hahn, and S. V. Nuzhdin. 2005. Genomic islands of speciation in

Anopheles gambiae. PLoS Biol. 3:1572–1578.

Wagner, C. E., L. J. Harmon, and O. Seehausen. 2012. Ecological opportunity and sexual

selection together predict adaptive radiation. Nature 487:366–369. Nature Publishing

Group.

Wu, C.-I., and C.-T. Ting. 2004. Genes and speciation. Nat. Rev. Genet. 5:114–122.

Wu, C. I. 2001. The genic view of the process of speciation. J. Evol. Biol. 14:851–865.

Yan, W., A. H. Assadi, A. Wynshaw-Boris, G. Eichele, M. M. Matzuk, and G. D. Clark. 2003.

Previously uncharacterized roles of platelet-activating factor acetylhydrolase 1b complex in

mouse spermatogenesis. Proc. Natl. Acad. Sci. U. S. A. 100:7189–94.

Zamani, N., P. Russell, H. Lantz, M. P. Hoeppner, J. R. Meadows, N. Vijay, E. Mauceli, F. Di

Palma, K. Lindblad-Toh, P. Jern, and M. G. Grabherr. 2013. Unsupervised genome-wide

recognition of local relationship patterns. BMC Genomics 14:347.

715 **Table 1**. Genome-wide f_4 statistics supporting differential introgression within Barombi Mbo

716 radiation. Tests with significant evidence for differential introgression are highlighted in bold.

717 The f_4 statistic was calculated for pairwise combinations among sister species of Barombi Mbo

718 subclades and riverine populations of *S. galilaeus* from Mungo and Meme Rivers (MM) and

719 Cross River (CR).

Introgression with riverine outgroups: (A,B) <-> (*MM, CR*)	f_4 statistic	*Z*-score	*P*-value
S. mariae, S. mongo	$-2.04 \times 10^{-7} \pm -5.15 \times 10^{-7}$	-0.39	0.69
S. mariae, S. pindu	$\mathbf{-1.92 \times 10^{-6} \pm -4.48 \times 10^{-7}}$	**-4.29**	**1.8x10⁻⁵**
S. mongo, S. pindu	$\mathbf{-1.59 \times 10^{-6} \pm -4.98 \times 10^{-7}}$	**-3.19**	**0.0014**
K. dikume, K. eisentrauti	$\mathbf{-2.4 \times 10^{-6} \pm -6 \times 10^{-7}}$	**-4.01**	**6.3x10⁻⁵**
K. eisentrauti, P. maclareni	$-2.12 \times 10^{-7} \pm -6.15 \times 10^{-7}$	0.35	0.73
K. dikume, P. maclareni	$\mathbf{-2.56 \times 10^{-6} \pm -5.86 \times 10^{-7}}$	**-4.37**	**1.2x10⁻⁵**
M. myaka, S. linnelli	$-4.04 \times 10^{-7} \pm -7.11 \times 10^{-7}$	0.56	0.57

Table 2. Candidate introgressed regions in Barombi Mbo cichlid radiation. These regions feature

significant f_4 values between riverine populations of *S. galilaeus* (MM: Mungo and Meme River;

CR: Cross River) and the three subclades of the radiation focused on in this study. Directionality

was determined from f_d outlier values (above 98[th] percentile) that overlapped with significant f_4

peaks. Unannotated regions, regions with no GO terms, and regions that were not f_d outliers are

marked with (-). Regions introgressed into a single species within a subclade are highlighted in

bold.

Linkage Group	Gene regions	GO Terms	Direction (f_d)
Stomatepia			
LG2	*sirpb1*	intracellular signal transduction, extracellular exosome	-
LG3	**NA**	**-**	***S. galilaeus* MM <-> *S. pindu***
LG3	uncharacterized protein-coding gene	-	*S. galilaeus* MM <->S. *mongo, mariae*
LG4	***jmjd8*; *prss1***	**extracellular exosome; proteolysis,peptidase activity**	***S. galileaus* CR <-> *S. mariae***
LG7†	NA	-	*S. galileaus* CR <-> *S. pindu, mariae*; *S. galilaeus* MM <- > *S. mongo*
LG14	*cldn4*	plasma membrane; bicellular tight membrane	*S. galilaeus* MM <-> *S. mariae*
LG18	***muc19***	**hematopoietic progenitor cell differentiation; extracellular exosome**	***S. galilaeus* MM <-> *S. mariae***
LG20†	*tprxl*; *plod3*	pseudogene of *tprx1*;motor nueron axon guidance, nueral crest cell migration	*S. galileaus* CR <-> S. *pindu, mariae*; *S. galilaeus* MM<- > *S. mongo*
LG20	*samdh1*	dGTPASe, immune system process, brain hemorrhages	-
NT_168079.1†	*osbpl5*;*ptprj*	lipid transport ; artery morphogenesis, phosphatase activity	*S. galileaus* CR<-> S. *pindu,mariae*; *S. galilaeus* MM<- > *S. mongo*
Konia + Pungu			
LG2	*sirpb1*	intracellular signal transduction	-
LG4	*anapc2*	ubiquitin dependent catabolic process	*S. galilaeus* MM <-> *P. maclareni, K. dikume*
LG6†	*cd209e*	endocytosis,mannose binding	*S. galileaus* CR<-> *K. eisentrauti, P. maclareni*; *S. galilaeus* MM<-> *K. dikume*
LG15	***adgrf4***	**g-protien coupled receptor activity, signal transduction**	***S. galilaeus* MM<-> *K. dikume***

LG20†	*klhdc8b*	ubiquitin-protein transferase activity, cytoplasm	*S. galilaeus* CR<-> *P. maclareni*; *S. galilaeus* MM<-> *K. dikume, eisentrauti*
LG20	NA	-	*S. galilaeus* MM <-> *K. dikume, eisentrauti*
LG20	*tprxl; plod3;clec10a*	pseudogene of *tprx1*; motor nueron axon guidance, nueral crest cell migration ; immune system response	-
LG22	*ehmt2*	neurogenesis, camera-type eye photoreceptor cell differentiation	-
LG23	**ddn1**	**serine-type endopeptidase activity,proteolysis**	***S. galilaeus* MM<-> *P. maclareni***
NT_167508.1†	*fam159a*	integral component of membrane	*S. galileaus* CR<-> *K. dikume, eisentrauti*; *S. galilaeus* MM<-> *P. maclareni*
NT_167623.1	*mepce*	methyltransferase activity	*S. galilaeus* MM<-> *K. eisentrauti*
NT_167637.1	**NA**	**-**	***S. galilaeus* MM<-> *K. dikume***
NT_167671.1	**NA**	**-**	***S. galileaus* CR<-> *K. dikume***
NT_167702.1*	*pafah1b3; hmcn1*	**platelet-activating factor acetyltransferase activity,brain development;fin morphogenesis, calcium ion binding**	***S. galileaus* CR<-> *K. eisentrauti***
NT_168003.1	*p2ry14*	respirotory abnormal muscle physiology, g-protien coupled purinergic nucleotide receptor activity	-
Myaka + S. linnelli			
LG6†	*c3*	complement activation, inflammatory response	*S. galilaeus* MM<->*S. linnelli*; *S. galileaus* CR<-> *M.myaka*
LG8/LG24†	NA	-	*S. galileaus* CR<->*S. linnelli*; *S. galilaeus* MM<-> *M.myaka*
LG11†	*hfe2;txnip*	somite development, BMP signaling pathway; enzyme inhibitor activity	*S. galilaeus* MM<->*S. linnelli*; *S. galileaus* CR<-> *M.myaka*
LG11	*vmo1*	extracellular exosome	*S. galilaeus* MM<-> *M.myaka*
LG16/LG21	**parp4**	**ribosyltranferase activity**	***S. galileaus* CR<->*S. linnelli***
LG19†	NA	-	*S. galileaus* CR<->*S. linnelli*; *S. galilaeus* MM<-> *M.myaka*
LG20	**tprxl,plod3**	**psuedogene of tetra-peptide repeat homeobox ; motor nueron axon guidance, nueral crest cell migration**	***S. galilaeus* MM<-> *M.myaka***
LG20†	*klhdc8b; cd163l1*	ubiquitin-protein transferase activity, cytoplasm; receptor mediated endocytosis, scavenger receptor activity	*S. galileaus* CR<->*S. linnelli*; *S. galilaeus* MM<-> *M.myaka*
NT_167617.1†	*ssr1*	endoplasmic reticulum membrane; cotranslational protein targeting	*S. galileaus* CR<->*S. linnelli*; *S. galilaeus* MM<-> *M.myaka*
NT_168092.1	**NA**	**-**	***S. galilaeus* MM<-> *M.myaka***
NT_168092.1	*itgam*	integrin mediated signalling pathway, cell adhesion	-

737   *best candidate region for secondary gene flow contributing to diversification † regions with signatures of

738   differential sorting of polymorphism from putative hybrid swarm

Fig 1. The most common topologies feature a monophyletic Barombi Mbo radiation. Across 96.2% of the genome Barombi Mbo species (black) are more closely related to each other than riverine outgroup populations of *S. galilaeus* Mungo and Meme River (green) and *S. galileaus* Cross River (red), or the Lake Ejagham *Sarotherodon* radiation (blue).

Fig 2. Topologies featuring Barombi Mbo polyphyly with riverine populations involving

the *Stomatepia* three-species complex. Across small and independent proportions of the genome

A-B) the entire *Stomatepia* clade, C) only *S. pindu*, and D) only *S. mariae* are more closely

related to outgroups than other Barombi Mbo species.

Fig 3. Topologies featuring Barombi Mbo polyphyly with riverine populations involving

the *Konia + Pungu* subclade. Across small and independent proportions of the genome A) only

P. maclareni, B) only *K. dikume*, and C) the entire *Konia + Pungu* subclade are more closely

related to outgroups than other Barombi Mbo species.

Fig 4. Topologies featuring Barombi Mbo polyphyly with riverine populations involving

the *Myaka + Sarotherodon* subclade. Across small and independent proportions of the genome

A-B) only *M. myaka*, C) *M. myaka* and two Barombi Mbo *Sarotherdon* species (*S. linelli* and *S.*

lohbergi), and D) *M. myaka* and *S. caroli* are more closely related to outgroups than other

Barombi Mbo species.

Figure 5. Manhattan plots of f_4 values between riverine populations of *S. galilaeus* from

Mungo and Meme river and *S. galileaus* from Cross River and A) combinations of the three

species of *Stomatepia*, B) combinations of the three species in the *Konia+Pungu* subclade, and

C) *M. myaka* and a representative from its sister *Sarotherodon* clade: *S. linnelli*. Alternating

gray/black colors indicate different linkage groups. Dotted red lines mark the permutation-based

significance thresholds for each test (P = 0.02). Peaks highlighted in colors represent those

signals of introgression shared across different subclades. Manhattan plots for the scaffolds not

assigned to the 24 linkage groups are presented in Fig S2-4.

Figure 6. Candidate introgression region in the *Konia + Pungu* subclade of Barombi Mbo

region containing genes *pafah1b3* and *hmcn1*. Row 1 shows the peak signal of introgression

across scaffold NT_167702.1 detected from the f_4 statistic across the three test combinations

involving the three species in the *Konia + Pungu* subclade and riverine populations of *S.*

galilaeus from Mungo, Meme River, and Cross River in non-overlapping 10-kb windows. The

two genes in this peak are shown in red (*pafah1b3* on the left and *hmcn1* on the right). Row 2

shows between-population divergence (D_{xy}) among the three combinations of sister species in the

subclade calculated in non-overlapping 10-kb windows. Row 3 shows within-population

diversity (π) in the same non-overlapping 10-kb windows.

Figure 7. The contributions of differential and shared introgressed variation to the genomes of sister species in the genera *Stomatepia* and *Konia*. The left and right columns represent the best candidate sister species in the radiation for sympatric speciation and speciation

with secondary gene flow respectively. Species with no evidence for differential introgression

are highlighted with an asterick (*). The true sister species in the *Stomatepia* are unknown, so all

three potential relationships among the complex are shown. Outgroup riverine *S. galilaeus*

lineages from Mungo/Meme Rivers (MM) and Cross River (CR) are shown with gray branches.

The percentage of introgression consistent with secondary gene flow into single species and a

hybrid swarm from sliding window f_4 statistic tests is show by above arrows. Proportion of

genome from two riverine sources (red: Mungo/Meme River; blue: Cross River) estimated by

SAGUARO is shown below arrows. Introgression patterns not found in the SAGUARO analysis

are represented with dashes (--). Blue arrows with no information that are aligned with red

arrows represent patterns found consistent with hybrid swarm. Note that the separation of arrows

along a branch is for the purpose of clarity and doesn't represent known differences in the timing

of introgression.

810