1

# Genome-wide transcriptional responses of iron-starved

# *Chlamydia trachomatis* reveal prioritization of metabolic

# precursor synthesis over protein translation

Amanda J. Brinkworth[1], Mark R. Wildung[1,2], Rey A. Carabeo[1#]

1. School of Molecular Biosciences, College of Veterinary Medicine, Washington State University.

2. Laboratory of Biotechnology and Bioanalysis, College of Veterinary Medicine, Washington State University.

#. Corresponding Author

Running title: Global response of *C. trachomatis* to iron-starvation

Abstract word count: 250

Importance word count: 147

Text word count (Introduction, Results, Discussion): 5139

Text word count (Introduction, Results, Discussion, Materials and Methods, Acknowledgements): 6058

ABSTRACT (exactly 250 words)

Iron is essential for growth and development of *Chlamydia*. Its long-term starvation in

cultured mammalian cells leads to production of aberrant non-infectious chlamydial

forms, also known as persistence.  Immediate transcriptional responses to iron limitation

have not been characterized, leaving a knowledge gap of how *Chlamydia* regulates its

response to changes in iron availability. We used the fast-chelating agent 2,2'-Bipyridyl

(BPDL) to homogeneously starve *Chlamydia trachomatis* serovar L2 of iron, starting at

6 or 12h post-infection. Immediate transcriptional responses were monitored after only 3

or 6h of BPDL-treatment, well before formation of aberrant *Chlamydia.* The first

genome-wide transcriptional response of *C. trachomatis* to iron-starvation was

subsequently determined utilizing RNA-sequencing. Only 7% and 8% of the genome

was differentially expressed in response to iron-starvation at early and mid-stages of

development, respectively. Biological pathway analysis revealed an overarching theme.

Synthesis of macromolecular precursors (deoxynucleotides, amino acids, charged

tRNAs, and acetyl-coA) was up-regulated, while energy-expensive processes (ABC

transport and translation) were down-regulated.  A large fraction of differentially down-

regulated genes are involved in translation, including ribosome assembly, initiation and

termination factors, which could be analogous to the translation down-regulation

triggered by stress in other prokaryotes during stringent responses. Additionally,

transcriptional up-regulation of DNA repair, oxidative stress, and tryptophan salvage

genes reveals a possible coordination of responses to multiple antimicrobial and

immunological insults. These responses of replicative-phase *Chlamydia* to iron-

starvation indicate a prioritization of survival over replication, enabling the pathogen to

"stock the pantry" with ingredients needed for rapid growth once optimal iron levels are

restored.

IMPORTANCE (max 150 words, currently at 147)

By utilizing an experimental approach that monitors the immediate global response of

*Chlamydia trachomatis* to iron-starvation, clues to long-standing questions in *Chlamydia*

biology are revealed, including how *Chlamydia* adapts to this stress. We determined

that this pathogen initiates a transcriptional program that prioritizes replenishment of

nutrient stores over replication, possibly in preparation for rapid growth once optimal

iron levels are restored. Transcription of genes for biosynthesis of metabolic precursors

was generally up-regulated, while those involved in multiple steps of translation were

down-regulated. We also observed an increase in transcription of genes involved in

DNA repair and neutralizing oxidative stress, indicating that *Chlamydia* employs an "all-

or-nothing" strategy. Its small genome limits its ability to tailor a specific response to a

particular stress. Therefore, the "all-or-nothing" strategy may be the most efficient way

of surviving within the host, where the pathogen likely encounters multiple simultaneous

immunological and nutritional insults.

INTRODUCTION

The sexually transmitted bacterium *Chlamydia trachomatis* infects the mucosal

epithelium of the endocervix, urethra, and anogenital tract. These infections usually

resolve spontaneously, and most are asymptomatic and thus underreported. Over 1.5

million cases of *C. trachomatis* genital infections were reported in the US in 2015 alone

(1). As many as 17% of females infected with *C. trachomatis* develop long-term

infections in the genital tract, which can result in serious complications such as pelvic

inflammatory disease (PID), fallopian-tube scarring, and ectopic pregnancy, all of which

are major risk factors for tubal factor infertility (TFI) (2). Rectal infections with LGV

serovars of *C. trachomatis* can be invasive, and if untreated can lead to complications

such as proctocolitis, inguinal adenopathy, reactive arthropathy and colorectal ulcers

(3). In some patients, infection persists even after antibiotic treatment (4, 5). The ability

of *C. trachomatis* to survive long-term in some individuals despite host immunity and

antibiotic treatment is not well understood, and may be associated with *Chlamydia*'s

ability to become persistent (6). While aberrant chlamydial forms have been identified in

cervical samples, the clinical relevance of this phenomenon is not well understood (6,

7).

Chlamydiae are obligate intracellular Gram-negative bacteria that undergo a

biphasic developmental cycle that includes both non-replicative and replicative forms

(8). Infection begins when the small, metabolically quiescent chlamydial elementary

body (EB) binds to mucosal epithelial cells and translocates virulence factors that

induce its endocytic uptake. Within 2 hours of entry, the EB will differentiate into its

replicative form, the reticulate body (RB). Continued secretion of effectors leads to

modification of the endocytic vesicle such that it avoids fusion with the lysosome and

enables capture of nutrient-rich vesicles. This unique intracellular niche, called the

inclusion, continues to expand as RBs replicate. In response to unknown signals around

24h post-infection, RBs will then differentiate into infectious EBs, followed by EB release

36-72 hours post-infection (8). Under exposure to certain stress conditions in cell culture

(e.g. penicillin, interferon-gamma (IFN-g), iron-depletion or tryptophan-depletion) RBs

will not differentiate into EBs, but instead enter into a state of persistence, characterized

by aberrant, enlarged morphology (9–13). Persistent *Chlamydia* are resistant to both

antibiotics and host immunity mechanisms, and can recover from this state upon

removal of stress or addition of missing nutrients (14–17).

Chlamydiae have undergone reductive evolution as they have adapted to

intracellular growth in mammalian cells, discarding metabolic genes responsible for

synthesizing factors that could be acquired from the host (18). The core genome of *C.*

*trachomatis* serovar L2 encodes only 889 open-reading frames, making *Chlamydia*

dependent on its host for lipids, nucleotides, amino acids, and metal cofactors (18).

Exposure of *Chlamydia*-infected cells to immune mediators, such as IFN-g, reduces the

availability of these factors and results in reduced RB division and differentiation (12,

14). IFN-g induces intracellular depletion of tryptophan by increasing levels of

indolamine 2,3-dioxygenase, which is responsible for catabolizing tryptophan into

kynurenines, which cannot be utilized in tryptophan metabolism (19).

Induction of inflammatory cytokines such as IFN-g and IL-6 in response to

chlamydial infection likely causes sequestration of free iron by the mononuclear

phagocytic system, which includes both cellular and systemic regulatory pathways (20–

27). Readers are referred to two comprehensive reviews of the coordinated regulation

of iron homeostasis by systemic and cellular mechanisms (26, 28). In the context of

*Chlamydia* infection of the genital epithelium, iron availability in infected cells is likely

limited by down-regulation of transferrin receptor and upregulation of the iron-storage

factor, ferritin (29). Iron-levels in the female genital tract can also fluctuate throughout

the menstrual cycle, in part due to hormone-induced expression of lactoferrin (30, 31).

Iron is essential for growth and development of *Chlamydia*, and its acquisition and

accumulation must be carefully regulated. In mammals, the readily usable ferrous iron

($Fe^{2+}$) is tied up in molecular complexes, limiting their interaction with hydrogen peroxide

to form damaging hydroxyl radicals through the Fenton reaction (32). Eukaryotic stores

of ferric iron ($Fe^{3+}$) are strictly regulated to restrict access by pathogenic bacteria

(33). Extracellular bacteria such as *Pseudomonas* and *Yersinia* utilize multiple

redundant iron-binding molecules called siderophores that compete with mammalian

transferrin for ferrous iron (34, 35). Intracellular bacteria, such as *Mycobacterium*,

*Francisella* and *Chlamydia,* can obtain iron by subverting host vesicles that contain

holo-transferrin bound to transferrin receptor (36–39). Using a combination of endocytic

markers and chemical inhibitors, our laboratory discovered that *Chlamydia* specifically

recruits transferrin-containing vesicles from the slow-recycling endocytic pathway (38).

Once delivered into the inclusion, iron is likely imported into bacteria through an ABC

transporter system, encoded by *ytgABCD*, which is the only known iron acquisition

system in *Chlamydia* species (40–42). The C-terminus of the YtgC permease, referred

to as YtgR, is homologous to the *Cornyebacterium* repressor DtxR, and has been

recently characterized as an iron-dependent repressor of the *ytgABCD* iron-acquisition

operon (42). A recent review highlights the differences between iron-acquisition

strategies of *Chlamydia* with other intracellular bacteria (27).

Conversion to the aberrant phenotype in response to iron-starvation reduces the

infectious potential of *Chlamydia*, since only a portion of RBs recover from stress and

complete development into infectious EBs once iron is added back into the media (43).

Previous studies have characterized aberrant *C. trachomatis* after long-term treatment

with the iron-chelator deferoxamine, and have detected increased expression of the

iron-binding protein YtgA, indicating its role in iron-uptake (41, 44, 45). However, these

studies added deferoxamine at the time of infection, and did not monitor transcriptional

or proteomic patterns until ≥24 hours post-infection, making it difficult to determine if this

up-regulation is part of the initial response to iron starvation. Immediate genome-wide

transcriptional responses to iron limitation have not yet been characterized in detail,

leaving a gap in the current knowledge of how *Chlamydia* regulates its response to

changes in iron availability. We utilize the chelator 2,2-Bipyridyl (BPDL), which can

quickly and efficiently chelate free iron from both bacterial and mammalian cells, to

induce an immediate transcriptional response to iron-starvation by *C. trachomatis* (43,

46–48).

This study provides the first global profile of the *C. trachomatis* transcriptional

response to iron starvation. Our short-term, effective treatment regimen in combination

with deep RNA-sequencing reveals the immediate response of *Chlamydia* to iron-

starvation in the logarithmic phase of growth when the bacteria are in the RB form. Here

we utilize this dataset to map the specific biological pathways altered in response to

iron-starvation. Taken together our results provide important clues to how *Chlamydia*

survives iron limitation. Accumulation of metabolite precursors is prioritized over

macromolecular biosynthesis. In addition, the transcriptional induction of genes involved

in adaptation to other stress factors, e.g. oxidative stress and amino acid starvation,

points to the inability of *Chlamydia* to tailor its transcriptional response to a specific

stress. Lastly, the global transcriptomic profile of iron-starved *Chlamydia* provided

valuable insights into how the biphasic developmental cycle might irreversibly switch to

persistence.

RESULTS

**Treatment optimization to detect the immediate chlamydial response to iron-**

**starvation**

The bivalent chelator 2,2-Bipyridyl (BPDL) has been shown to deplete both ferrous

and ferric iron from *Chlamydia*-infected cells during long-term treatment, and it induces

the development of aberrant forms more consistently and homogenously than the

previously used ferric iron chelator, deferoxamine (43). Here, we determined the optimal

duration of BPDL treatment to induce iron-responsive transcription without inducing

morphological abnormalities in *Chlamydia*. We chose to begin starvation during mid-

cycle development (12h p.i.) instead of at the beginning of infection for two reasons: 1)

to test the response of actively replicating *Chlamydia* that are able to maximally respond

to stress, and 2) to ensure that both treated and mock-treated *Chlamydia* remain in the

same stage of development (RB). We monitored chlamydial morphology, growth, and

transcriptional responses after 3, 6, and 12h of BPDL treatment (Figure 1A). Indirect

immunofluorescent confocal microscopy revealed similar morphology between mock-

treated and BPDL-treated forms for up to 12h of BPDL-treatment (Figure 1B).

Interestingly, observation of BPDL-treated cultures by light microscopy revealed an

obvious decrease in Brownian movement within inclusions after 6 or more hours of

treatment (data not shown). This observation is consistent with our findings that

chlamydial growth is reduced compared to mock-treated after only 6h of BPDL-

treatment, as determined by quantitative PCR of chlamydial genomes (Figure 1C).

We monitored the transcriptional response of the known iron-responsive genes *ytgA*

and *ahpC* by reverse transcriptase quantitative polymerase chain reaction (RT-qPCR)

to validate the iron starvation protocol (40, 41, 43, 49). Elevated transcription of both

iron-starvation markers was detected after only 6h BPDL-treatment, compared to mock-

treated (1.5-fold and 1.7-fold). Maximum differences in transcription of both markers

were detected after 12h of BPDL treatment (2.8-fold and 3.7-fold; Figure 1D). In the

same experiment, we also monitored the transcriptional profile of the early gene, *euo*,

which decreases during late stages of normal development.  Multiple persistence

models have demonstrated dysregulated of *euo* transcription, with high levels of *euo*

mRNA detected late in development under persistence-inducing conditions (15, 43, 45).

After 12h BPDL treatment, we observed that *euo* transcript levels remained elevated

relative to the mock-treated control, indicating dysregulated transcription or a possible

delay in development (Figure 1E, top). This delay in development during longer BPDL-

treatment is supported by the lack of recoverable inclusion-forming units (IFUs)

detected after 12 or 24h BPDL-treatment compared to mock-treated controls (Figure

1F), indicating a possible lack of RB-to-EB differentiation. Because 6h BPDL-treatment

is sufficient to induce iron-responsive transcription without inducing the morphology and

transcriptional pattern associated with persistence, we chose it as the optimal duration

of iron-starvation for our genome-wide transcriptional studies. We also included 3h

BPDL-treatment to detect the earliest possible response to iron-starvation prior to

BPDL-induced changes in growth.

**Global transcriptional response of *C. trachomatis* to iron-starvation during mid-**

**cycle development**

The primary global response of *C. trachomatis* to mid-cycle iron starvation was

determined by RNA-sequencing (RNA-seq). We utilized an Ion Proton for sequencing,

which allowed for easy and rapid scaling of timepoints based on observed yield of

mapped reads. This approach is relevant to the study of *Chlamydia* transcription

because chlamydial mRNA represents a small proportion of the total RNA at the time

points analyzed, even after significant enrichment steps.  For mid-cycle iron starvation

studies, we aimed for greater than 10x coverage of 100% of the *C. trachomatis*

genome, with a minimum of 3 biological replicates per sample. The sequencing reads

were trimmed to exclude adaptor sequences and polyclonal reads, followed by

exclusion of reads less than 30 nucleotides in length. The remaining reads were aligned

to the *C. trachomatis* genome and plasmid, with 2 to 23% of trimmed reads mapping.

Average read lengths varied from 92 to 134 nucleotides, requiring an average of

108,837 mapped reads to reach our coverage goal.  A summary of the read and

mapping statistics for all of our samples can be found in Supplementary Table 1.

Alignments, comparisons, and normalization of aligned reads were done with CLC

Genomics Workbench version 9.0 according to default settings. All mid-cycle conditions

(12h untreated, 12+3h BPDL, 12+3h mock, 12h+6h BPDL, 12+6h mock) were

compared using the CLC Genomics experiment tool, normalized by quantile scaling and

analyzed for differential gene expression using EdgeR statistical analysis with false-

discovery rate (FDR) calculation. Because we included ribosomal rRNA, eukaryotic

mRNA, and small RNA (<100 nt) depletion steps when preparing chlamydial mRNA for

RNA-seq, we also excluded tRNAs, rRNAs, and genes with < 10 mean reads in a

sample group prior to normalization and analysis.

The genome-wide profile of mock-treated and BPDL-treated gene expression during

mid-cycle development (12h to 18h post-infection) is displayed as a heatmap of

normalized expression values (Figure 2A). The raw and normalized data for these

individual replicates can be found in Supplemental Table 4. Mock-treated (left) and

BPDL-treated (right) profiles are remarkably similar across all genes that significantly

change during normal mid-cycle development of *Chlamydia* (based on comparisons 12h

vs 15h, 15h vs 18h, or 12h vs 18h, p-value ≤ 0.01). The annotated expression heatmap

and EdgeR comparisons for normal growth can be found in Supplemental Figure 1 and

Supplemental Table 2, respectively. The entire RNA-seq dataset of normal development

can be found in Supplemental Table 3. The similarity between global expression profiles

indicates that *Chlamydia's* normal development is not dysregulated after only 3h or 6h

of BPDL-treatment. However, EdgeR analysis of BPDL-treated cultures compared to

mock-treated samples (at equivalent time points post-infection) revealed 8% (76/889) of

the genome was differentially expressed after 3h BPDL–treatment and 1% (12/889) was

differentially expressed after 6h BPDL-treatment. Genes that were differentially

expressed with a maximum p-value of 0.01 are displayed in a heatmap of fold-change

differences between BPDL-treated and mock-treated samples (Figure 2B). Examples of

decreased transcription after 3h and 6h BPDL-treatment include the ribosomal subunit

genes *rpsO* and *rpsT*, and the type III secretion genes *copB* and *scc2*, respectively.

Transcription of the tryptophan salvage pathway operon, *trpBA,* and the ribonucleotide

reductase operon, *nrdAB,* was significantly increased after both 3 and 6h of treatment.

Iron-responsive genes that were differentially expressed with a p-value $\leq$ 0.01 after 3 or

6h of BPDL-treatment are listed in Table 1 and Table 2, respectively. The fully

annotated heatmap can be found in Supplemental Figure 2, and the full set of RNA-

sequencing results for mid-cycle iron starvation can be found in Supplemental Table 4.

**Functional categorization of differentially expressed genes during mid-cycle**

**response to iron-starvation.**

Annotations and functional categories of differentially expressed genes during mid-

cycle iron-starvation were retrieved from UniProt and are listed in Table 1 and Table 2.

Genes differentially expressed, with a minimum p-value of 0.01, after only 3h of BPDL-

treatment are grouped by functional category of induced and reduced transcripts (Table

1, Figure 3) (50). Of the 39 genes significantly induced after only 3h iron-starvation,

representing 4% of the genome, five categories were equally represented with 3 genes

each: energy metabolism (*glgA, lpdA, glmS*), amino acid biosynthesis (*trpA, trpB, aroL*),

DNA replication and repair (*nrdA, recA, dnaQ*), type III secretion (*mcsC, ctl0085,*

*ctl0043*), and translation (*pheT, cysS, thrS*). Of the 37 genes that were significantly

reduced in response to 3h BPDL-treatment, representing 4% of the genome, the

majority of these (39%) are associated with translation (*prfA, rplW, rsfS, smpB, ctl0132,*

*rpsK, rplT, rplN, ctl0680, rpmI, rmpJ, infa2, rpsT, rpsO*), and 11% are associated with

nutrient transport (*ctl0061, ctl0485, gltT,* and *ytgD*). Transcript levels of only 5 genes

were significantly increased after 6h BPDL-treatment (*trpB, trpA, nrdA, ctl0071,* and

*nrdB*) while transcript levels of 7 genes were decreased (*ctl0185, ctl0619, tsp,* and the

entire *scc2-ctl0840-cobB-copD* operon) (Table 2).

To independently confirm the mid-cycle response detected by RNA-sequencing, we

utilized RT-qPCR. Increased transcription in response to iron-starvation was confirmed

for all of the transcripts tested by RT-qPCR, with the exception of *recA* (Figure 4A).

None of the tested down-regulated genes were significantly reduced compared to

controls by RT-qPCR, likely due to the fact that the genes tested were very low in

abundance at the timepoints tested (Figure 4B).

**Functional categorization of differentially expressed genes during early-cycle**

**response to iron-starvation.**

*Chlamydia* infections of the genital tract are asynchronous. Thus *Chlamydia* could

be exposed to host-induced stress at any point in the developmental cycle. For this

reason, we extended our analysis of the immediate response to iron starvation to an

earlier point in the developmental cycle. *Chlamydia*-infected cells were treated with

BPDL starting at 6h post-infection, which is after the initial EB-to-RB differentiation and

at the beginning of the logarithmic growth phase. RNA and genomic DNA were

collected at 9h post-infection for both treated and mock-treated samples. RNA-seq and

alignments were performed as described above. A summary of mapped reads and

coverage can be found in Supplemental Table 1.

Genes differentially expressed during the early cycle response (6+3h BPDL vs 6+3h

mock-treated), with a maximum p-value of 0.01, are grouped by functional categories of

induced and reduced transcripts (Figure 5). The full set of differentially expressed genes

and their annotations can be found in Table 3. Similar to results of the mid-cycle

response, transcription of 4% of the genome was induced, including genes involved in

DNA replication and repair (*nrdA, nrdB, mutS, dnaQ,* and *recA*), amino acid

biosynthesis (*trpB, trpA*, *aspC_1*, and *glyA*), and translation (*ctl0111*, *trpS*, *thrS*, and

*aspS*) during the early-cycle response to iron starvation. Uniquely, genes involved in

redox homeostasis (*pdi*, *ahpC* and *sodM*) were also up-regulated in response to iron-

starvation starting at 6h post-infection but not during the mid-cycle response. Of the 23

genes with reduced transcription during the early-cycle response to iron-starvation (3%

of the genome), 17% are associated with translation (*rplW, prfA, rplC, ftsY*), and 13%

with DNA replication and repair (*pGP8D*, *amn*, and *dnaX_1*).

Up-regulation of *trpA* transcription during early-cycle iron-starvation was confirmed

by RT-qPCR, while only modest increases were observed for the other up-regulated

genes tested (Figure 6A). Down-regulation of *ctl0430*, *ctl0063,* and *incD* during iron-

starvation could not be confirmed by RT-qPCR (Figure 6B). We reasoned that early-

cycle responses were not detected by RT-qPCR for most of our tested genes due to the

limit of detection of the technique. The raw values detected for most of our early-cycle

transcripts fell at or below the lowest concentrations of our standard curves. Between 6

and 9h post-infection, chlamydial mRNA represents a very small proportion of the total

RNA. This limitation was overcome for RNA-seq experiments by depleting rRNAs and

eukaryotic RNA prior to synthesizing cDNA. However, cDNA used in RT-qPCR was

prepared from total RNA because mRNA enrichment would have made it impossible for

us to normalize our RT-qPCR data to chlamydial genomes. The overwhelming

proportion of eukaryotic RNA present in the undiluted cDNA used as template may have

impeded accurate detection of transcripts.

**Network and biological pathway analysis**

To further analyze the relevance of these gene expression changes to chlamydial

survival, we utilized the bioinformatics tool STRING-db v.10.5 to generate networks of

functionally associated genes (51). Representation of differentially expressed gene sets

(p-value ≤ 0.05) from short-term iron starvation (3h) reveals gene networks with

intersecting pathway clusters (manually added grey circles). We chose to use the less

stringent p-value to allow entire pathways to emerge, which would not have been quite

as apparent with a more stringent cutoff. Consistent with our predicted functional

categories, network analysis of both early (Figure 7A) and mid-cycle (Figure 7B)

responses to iron-starvation revealed clusters that include amino acid biosynthesis,

DNA replication and repair, and translation. Functional clustering of the mid-cycle

response also revealed the entire cluster of genes necessary to convert pyruvate to

acetyl-CoA, as well as gene clusters involved in tRNA modification and charging.

The locus identifiers of genes in each identified cluster were submitted to

KEGGMapper v2.8 to determine possible roles in specific biological pathways (52). For

example, genes from the early-cycle DNA replication and repair cluster (Figure 7A)

were mapped to multiple pathways including purine metabolism (5), pyrimidine

metabolism (5), mismatch repair (5), replication (4), homologous recombination (3),

double-stranded breaks repair (2), and base excision repair (1).

*Nucleotide metabolism*

We modified the KEGGMapper output for pyrimidine metabolism to indicate the

direction of change in mid-cycle gene expression during iron starvation (Figure 7C).

Under all iron-starvation conditions, the ribonucleotide reductase genes *nrdA* was up-

regulated. Ribonucleotide diphosphates (NDPs) bound to NrdA are converted by NrdB

to deoxynucleotide diphosphates (dNDPs). These dNDPs are not likely further

converted to dNTPs, as indicated by the down-regulation of nucleotide diphosphate

kinase, *ndk*. Available dUMP would likely be derived from the UDP pool, instead of from

the dUTP pool, since transcription of the dUTP pyrophosphatase gene, *dut*, is down-

regulated during iron-starvation. Taken together, these transcriptional changes would

result in a net increase in dNDPs, enabling rapid DNA replication when iron levels and

*ndk* expression return to normal (Figure 7C).

*Amino acid biosynthesis*

Functional clustering also indicates that *Chlamydia* prioritizes maintenance of amino

acid pools during iron starvation. Multiple amino acid synthesis, inter-conversion, and

uptake mechanisms were up-regulated in response to short-term iron-starvation.

Transcriptional up-regulation of the branched chain amino acid transporter, *brnQ,* the

aspartate aminotransferase, *aspC*, and the serine hydroxymethyltransferase, *glyA* may

increase the diversity of the amino acid pool such that *Chlamydia* can quickly adapt to

fluctuations in amino acids. Surprisingly, the tryptophan salvage pathway genes, *trpB*

and *trpA,* were consistently up-regulated during short-term iron-starvation. The

tryptophan synthase subunit TrpB catalyzes the beta-replacement of indole with serine

to form tryptophan (Trp), while TrpA facilitates the interaction of TrpB with indole (53).

Their role in recovery from IFN-g and Trp-starvation stresses is well documented, but

differential regulation in response to iron-starvation is novel (54, 55). While the

biological relevance of *trpBA* induction during iron-starvation is unclear, we reason that

*Chlamydia* could in fact prepare for further immune insult (e.g. IFN-g induction of

indoleamine 2,3-dioxygenase expression) by increasing intracellular Trp levels. Taken

together, iron-starvation may increase levels of serine, aspartate, glutamate, branched-

chain amino acids, and tryptophan, many of which are essential for normal development

(56–60). Amino acid biosynthetic genes were significantly overrepresented (4.38-fold, p-

value=0.0464) in the set of differentially expressed mid-cycle genes as determined by

the Panther Overexpression Test (61).

*Translation*

The largest cluster generated from STRING-db included translation factors of the

mid-cycle response (Figure 7B).  Based on protein annotations in Uniprot and Biocyc

databases, it is evident that *C. trachomatis* responds to iron-starvation by shutting down

factors involved in every step of translation: ribosome assembly, initiation, elongation,

termination, ribosome recycling, and peptide targeting (46, 54; Table 4). While

preventing the assembly and function of translational machinery, *Chlamydia* also

responds to iron starvation by increasing factors important for synthesis and

modification of tRNAs, in addition to increasing transcription of *rnC,* the product of which

cleaves rRNA transcripts into ribosomal subunit precursors (Table 4). Translation genes

were significantly overrepresented (3.26-fold, p-value=0.0243) in the set of mid-cycle

differentially expressed genes as determined by the Panther Overexpression Test (61).

*Acetyl-coA synthesis*

Transcription of the entire set of genes necessary for conversion of pyruvate to

acetyl-CoA was induced during the mid-cycle response to BPDL-treatment (Figure 7D).

This includes the lipoylation enzymes *lipA* and *lpdA,* and the entire pyruvate

dehydrogenase complex, *pdhABC*. In addition, transcription of the TCA cycle gene

*mdhC* and glycolysis gene *eno* was induced, likely driving formation of pyruvate from

different carbon sources.  Acetyl-CoA can be converted to malonyl-CoA for fatty-acid

biosynthesis or utilized in the formation of N-acetylglucosamine-1-phosphate for

peptidoglycan synthesis, both of which are required for rapid growth of *Chlamydia* (63,

64). Since transcription of the peptidoglycan-modifying enzymes, *glmS* and *murB,* were

also increased during iron starvation*,* acetyl-coA is likely used to form new

peptidoglycan. Expression of fatty acid synthesis genes was unchanged during iron-

starvation.

Pathway analysis of both early and mid-cycle responses to iron-starvation (Figure

7A and 7B) reveals that down-regulation of translation and up-regulation of amino acid

synthesis and nucleotide synthesis may be important for surviving this stress. Similarly,

a core set of 13 genes (*trpB, trpA, nrdA, recA, dnaQ, CTL0704, CTL0102, thrS, prfA,*

*rplW, CTL0061, CTL0548, pGP8-D*) are differentially expressed after 3h BPDL

treatment, during both early and mid-cycle responses, while *trpB, trpA*, and *nrdA* are up-

regulated in all BPDL treatments (Fig S3). This overlap in differential gene expression is

displayed as a Venn diagram in Supplemental Figure 3.

DISCUSSION

We monitored the immediate global transcriptional response of *Chlamydia*

*trachomatis* serovar L2 to short-term iron-starvation during early and mid-cycle (RB

phase) development. In contrast to previous studies of iron starvation in *Chlamydia*, our

short-term treatment with BPDL did not cause the hallmark changes in morphology and

*euo* transcription associated with persistence. This approach enabled us to detect a

response specific to iron starvation as *Chlamydia* tries to adapt to stress, rather than the

transcriptome of the aberrant bacterium. By deep RNA-sequencing we were able to

identify novel primary transcriptional responses, representing 7-8% of the genome, after

only 3h of iron-starvation with BPDL. It is possible that a portion of the detected BPDL-

responsive regulon is actually due to chelation of metals other than iron. $Cu^{2+}$ is

chelated at similar affinities to $Fe^{2+}$ and $Fe^{3+}$, while $Zn^{2+}$ is chelated with 2-3 logs lower

affinity than iron ions (65). We expect that $Zn^{2+}$ is not efficiently depleted during the

short-term BPDL treatments used in this study, but cannot exclude the possibility that

we have detected transcriptional changes in response to altered availability of other

metals. It is also possible that a more immediate response could be detected with even

shorter BPDL treatments, though we expect a longer duration is required to chelate both

free iron and iron bound to protein complexes in intracellular *Chlamydia*. Since only 12

genes were differentially expressed after 6h BPDL-treatment, a longer duration of

treatment may be necessary to detect the full secondary response, which may not be

obvious until the effects of the primary transcriptional response are realized at the

protein level.  This is supported by the fact that 6h BPDL-treatment maintains induction

of the primary response operons, *trpBA* and *nrdAB,* while reducing or delaying

expression of some late cycle genes (*scc2-ctl0840-copB-copD, tsp*). Decreased or

delayed late gene expression has also been observed during long-term iron-starvation

(43, 49, 66, 67).

In agreement with proteomic observations of deferoxamine-treated *C. trachomatis*

after 24h and *C. pnuemoniae* after 48h post-infection, we observed up-regulation of

*ctl0874*/*cadd*, *ahpC, eno, and htrA* during short-term BPDL-treatment (45, 68). In

contrast to previous iron-starvation studies, we did not detect a significant increase in

*ytgA* expression in our RNA-seq results.  We expected the *ytgABCD* iron acquisition

operon to be immediately induced in response to iron-starvation, since its repression by

YtgR is dependent on available iron (42). Expression of the *ytgABCD* operon peaks

during mid-cycle development, indicating that the iron-dependent repressor, YtgR, may

be inactive or at low levels during early and mid-cycle (15, 42). It is possible that we do

not observe significant differences in the expression of the operon during iron-starvation

because it is already maximally expressed in the mock-treated controls (Table S3).

Global detection of YtgR repression by CHIP-sequencing or targeted analysis of specific

promoters will be necessary to delineate the contribution of YtgR activity to the detected

iron-responsive regulon. Recent work to define targets of known transcription factors in

*Waddlia chondophila* discovered binding-sites of YtgC by ChIP-sequencing (69).

Interestingly, the most frequent target, *hrcA*, was also increased during iron-starvation in

our study, indicating it may also be a target of YtgC in *C. trachomatis.*

Transcriptional responses to iron-starvation in most bacteria typically include up-

regulation of iron-acquisition systems and virulence factors (70–72). While the *ytgABCD*

operon was not up-regulated during short-term iron-starvation, other unidentified iron-

uptake and iron-dependent repression mechanisms may exist, and thus could be

represented in our set of iron-starvation induced genes. The virulence factor genes

*cadd* (*ctl0874*) and *macpF* (*ctl0408*) were induced in both the early and mid-cycle

response to iron-starvation. CADD overexpression induces apoptosis when expressed

in cultured human epithelial cells, but has also been demonstrated to play a role in

folate biosynthesis (73, 74). MACPF contains a domain that may enable perforin

activity, but so far has only been show to undergo cleavage upon infection and become

inserted into bacterial membranes (75). Several type III secretion structural components

(*mcsC, sctJ, sctR, fliF, cdsN/ctl0043, cdsD/ctl0033*) and effectors (*ctl0884, ctl0476,*

*ctl0184, ctl0081*) were also transcriptionally up-regulated during iron-starvation, which

could potentially alter interactions between the host and chlamydial inclusion.

Similar to the up-regulation of the ribonucleotide reductase operon *nrdHIEF* during

iron-starvation in *E. coli* and *Yersinia pestis*, the ribonucleotide reductase genes, *nrdA*

and *nrdB,* are consistently up-regulated during under short-term iron starvation (71, 76).

This up-regulation indicates that deoxynucleotides may be important for *Chlamydia* to

survive this stress. However, since NrdB requires iron for its function, deoxynucleotide

levels may not increase until iron becomes available. Instead, high levels of inactive

NrdA-B complexes may actually impede replication and development by inducing

stalling at replication forks, providing a possible explanation for the decreased

replication observed during iron-starvation (77, 78).

*Chlamydia's* immediate transcriptional response to iron-starvation is remarkably

similar to stringent responses in other bacteria, which enable rapid adaptation to various

stresses by diverting resources from macromolecular biosynthesis, e.g. translation, and

growth to immediate survival, often resulting in a quiescent state (79, 80). This rapid

transcriptional response is achieved through synthesis of the chemical alarmone,

(p)ppGpp, which interacts with RNA polymerase and DksA to globally modify

transcriptional activity (81, 82). During amino acid starvation in bacteria, uncharged

tRNAs in the A-site of ribosomes are sensed by RelA, which responds by synthesizing

(p)ppGpp from ATP and GDP or GTP (83, 84). (p)ppGpp can also be synthesized and

hydrolyzed by SpoT during other stress conditions. However, since *Chlamydia* lacks the

RelA and SpoT homologues necessary for (p)ppGpp synthesis, they likely evolved

alternative mechanisms to reduce growth and increase survival responses during stress

(17, 85, 86).   Iron-starvation has been shown to induce a stringent response in *Bacillus*

*subtilis* that up-regulates transcription of amino acid biosynthesis genes (87).

Multiple amino acid synthesis, inter-conversion, and uptake mechanisms were up-

regulated in response to short-term iron starvation.  Surprisingly, the primary response

includes an increase in transcripts involved in tryptophan salvage, *trpB* and *trpA*, but not

the tryptophan-dependent repressor *trpR*. TrpR-dependent regulation of the

polycistronic transcript, *trpRBA*, has been extensively studied during tryptophan

starvation and IFN-g treatment, but rarely, if ever, in the context of iron-starvation (54,

88, 89). Notably, *trpB*, but not *trpR* levels were also increased under estradiol-induced

persistence, suggesting a *trpR*-independent mechanism of inducing tryptophan salvage

transcription may exist (90).

Pathway analysis clearly indicates that transcripts involved in all steps of translation

from initiation to ribosome recycling are down-regulated during iron-starvation. This

reduction in translation factors may lead to an eventual shutdown or modification of

translation activity that could increase survival during stress. By shutting down energy

expensive protein synthesis, ATP and GTP pools can be rerouted to immediate survival

responses (tRNA-charging, transcription). Similarly, iron-starvation reduces the

transcription of several ABC transporter genes, which require ATP for their function.

Uncoupled RNA and protein levels in *Chlamydia* have also been observed during IFN-g

stress (17). The apparent decrease in translation during IFN-gamma exposure could be

exacerbated by decreases in the levels of components of the translation machinery in

response to simultaneous iron-starvation. However, decreased expression of translation

factors during the primary response to iron-starvation may not be apparent until pre-

existing ribosomal-protein complexes are degraded or destabilized. This may explain

why ≥24h of iron-starvation is required to induce the development of aberrant RBs (43).

Down-regulation of translation factors during iron starvation will have to be examined at

the protein level to determine its contribution to adaptation to iron starvation and

development of persistence.

In contrast to down-regulation of translation, iron starvation increases transcription of

amino-acyl synthesis genes (*cysS, pheT, glyQ, aspS, thrS*), which are responsible for

charging tRNAs with amino acids.  The apparent disconnect between increased amino-

acyl-tRNA pools and decreased translation indicates possible survival mechanisms.

Charged tRNAs might be utilized in an immediate survival response to iron-starvation,

prior to the turnover of ribosomal subunits. Alternatively, *Chlamydia* may be

accumulating charged tRNAs for recovery and resumption of development when normal

levels of iron and translation factors are restored.

A major theme that emerged from our gene expression analysis is that *Chlamydia*

likely perceives iron starvation as a signal to prepare for further nutrient deprivation and

immune insult. Transcriptional up-regulation of tryptophan salvage pathway (trpB, trpA),

oxidative stress (*ahpC*, *pdi,* and *sodM*) and DNA repair (*mutS, mutL, ssb, ung, recA*)

genes indicate a protective response against antimicrobial insults of the inflammatory

immune response (e.g. IDO activation, reactive oxygen species). As an obligate

intracellular pathogen, *Chlamydia* has undergone reductive evolution with constant

selective pressure from the host immune system and its multiple anti-chlamydial

effectors. Due to its small genome (~ 1 Mbp), *Chlamydia* may not have the capability to

induce a specific transcriptional response to each particular stressor, and the

simultaneous deployment of stress responses may have been the most parsimonious

route of adaptation to immune insult. In this case, we would expect that iron-starved

*Chlamydia* would be better protected from damage by antimicrobial insults than mock-

treated *Chlamydia*. Immediate transcriptional responses to other stress conditions will

need to be monitored to determine if this coordination of antimicrobial responses is

unique to iron starvation.

This study provides the first evidence of a global iron-dependent regulon for *C.*

*trachomatis*. By using a systems-approach to delineate *Chlamydia*'s transcriptional

response to iron starvation, we have been able to detect biological pathways and place

them in the context of chlamydial development. These findings are novel, and add to

previous studies of iron-dependent transcriptional and proteomic profiling in aberrant

RBs, revealing transcriptional adaptive strategies prior to the development of a

persistent state. Additionally, our results include a high-resolution profile of mid-cycle

development of *C. trachomatis* serovar L2, including relevant time points for monitoring

shifts in early, mid, and late-cycle gene expression. We expect this dataset will prove

useful for future studies that seek to determine *Chlamydia*'s immediate transcriptional

response to other chemical and/or nutrient stresses. Our findings include previously

unrecognized shifts in energy utilization and down-regulation of translation that

resemble a stringent-like survival response. *Chlamydia* may utilize a two-stage

approach of increasing transcription of survival genes in the short term to delay

development and survive during iron-starvation, followed by an eventual shutdown of

translation at later times of sustained stress. The latter might account for the observed

irreversibility of the persistent state during long-term starvation for iron or tryptophan.

MATERIALS AND METHODS

**Cell Culture and infection**

HeLa monolayers were infected with *C. trachomatis* strain L2 434/Bu in 6-well plates at

a multiplicity of infection (MOI) of 2 for RNA and genomic DNA (gDNA) collection

experiments, and on coverslips in 24-well plates for morphology studies. Cells were

grown in DMEM supplemented with 10% FBS, 2 mM glutamine, and 10 micrograms/mL

gentamycin in 5% $CO_2$ at 37° C. HeLa cells used in this study were started from P1

stocks from ATCC, and were regularly checked for contamination by DAPI-staining and

the Universal Mycoplasma detection kit (ATCC).

**RNA-sequencing**

RNA was collected and pooled from 2 or 4 T75 flasks of *C. trachomatis*-infected HeLa

monolayers that had been treated with 100 µM 2,2-Bipyridyl (BPDL) starting at 6 or 12h

hours post-infection (6h + 3h BPDL, 12h + 3h BPDL, 12h + 6h BPDL) and in mock-

treated samples at equivalent timepoints post-infection (9h, 12h, 15h, 18h). RNA was

purified using the RiboPure Bacteria (Ambion) kit as per manufacturers instructions.

Total RNA was further enriched for transcripts over 100 nucleotides in length with the

MegaClear kit (Ambion). Mammalian transcripts and rRNAs were removed with the

MicrobEnrich kit (Ambion) and bacterial rRNAs were removed using the MicrobExpress

kit (Ambion), repeating 2-3 times. The integrity and quantity of total and depleted RNA

was monitored with an AATI Fragment Analyzer. cDNA libraries were prepared with the

Ion Total RNA-seq Kit V2, and sequencing beads prepared using an ion Chef, and

sequencing performed on an Ion Proton chip with HiQ chemistry. Primary sequence

analysis, trimming, and binning of reads was performed using Torrent Suite Software

version 5.0.5. Remaining reads were mapped to the combined core genome of *C.*

*trachomatis* strain L2 434/Bu (Genbank Accession: AM884176) and the plasmid of *C.*

*trachomatis* L2b CS784/08 (NZ_CP009926) using CLC Genomics Workbench 9,

requiring reads be at least 30 nucleotides in length, with default alignment parameters.

The EdgeR algorithm in CLC Genomics was used to determine differential gene

expression during development and iron-starvation, assuming a false-discovery rate of

10% and p-values ≤ 0.05. tRNAs and ribosomal RNAs were filtered from the reads to

account for differences in depletion efficiency, and only genes with at least 5 mapped

reads were included in the analysis. Differentially expressed genes were confirmed for

selected transcripts by RT-qPCR.

**qPCR and RT-qPCR**

*C. trachomatis*-infected HeLa monolayers were treated with 100 micromolar BPDL

starting at 6 or 12h hours post-infection (6h + 3h BPDL, 12h + 3h BPDL, 12h + 6h

BPDL) and mock-treated samples at equivalent timepoints post-infection (6h, 9h, 12h,

15h, 18h). RNA and gDNA were collected with the RiboPure Bacteria and the DNeasy

Blood and Tissue (Qiagen) kits, respectively. cDNA was generated with Superscript IV

reverse transcriptase (Life Technologies) from 200-500 nanograms RNA as per

manufacturer's instruction, except the use of random nonamers instead of hexamers.

Transcripts were amplified with PowerUp SYBR Green system from undiluted cDNA for

early (6-9h) or diluted 1:10 in 10 mM Tris for midcycle (12-18h) samples and detected

with Applied Biosystems 7300 RT-qPCR system.

**Chlamydial morphology**

Chlamydiae were monitored for changes in morphology in response to mock-treatment

or treatment with 100 micromolar BPDL starting at 12h post-infection for 3, 6, or 12h.

Infected cultures were fixed on coverslips and stained with pooled human serum

(Sigma, H4522) at 1:750, followed by goat-anti-human antibody conjugated to Alexa

Fluor 488 (ThermoFisher) at 1:1000.  DNA was stained with DAPI at 5 micrograms/mL.

Images were taken on a Leica SP8 confocal microscope with a 63X oil-objective and 4X

zoom.

**IFU assay**

Chlamydiae were monitored for changes in infectivity in response to mock-treatment or

treatment with 100 micromolar BPDL starting at 12h post-infection for 12 or 24h, at an

MOI of 1. Infected cultures were scraped into 300 microliters SPG and stored at -80 C

for later testing.  Thawed lysates were serially diluted into complete DMEM, centrifuged

onto HeLa monolayers in 24-well plates, washed with HBSS, and allowed to infect for

24h.  Infected cultures were fixed and stained with pooled human serum at 1:750,

followed by goat-anti-human antibody conjugated to Alexa Fluor 488 (ThermoFisher) at

1:1000. Inclusions were counted by fluorescent microscopy and inclusion-forming units

(IFU) were calculated as previously described.

**Visual analysis of differentially expressed genes**

Functional categories were assigned for all genes differentially regulated with a p-value

≤ 0.01 by referring to the GO terms listed on UniProt. Pie charts were generated using

the "pie" function in Rstudio.  Heatmaps were generated in Rstudio using the package

"pheatmaps", with parameters set to average clustering and Euclidean distance. The

Panther overexpression test in Panther v12.0 was done on differentially regulated gene

sets (total) with p-values ≤ 0.01, using the default parameters and Bonferri correction.

Pathway analysis was performed on differentially regulated genesets with p-values ≤

0.05 and a minimum of 10 mapped reads, with STRING-db v.10.5 set to confidence ≥

0.7. StringDB maps were slightly modified to make space to increase font size, indicate

the direction of change by color-coding and add pathway labels, without altering

network relationships. Clustered genes detected with StringDB were further analyzed

using KeggMapper v.2.8, and pathway maps were generated based on KeggMapper

output using Affinity Designer v1.4.1.

**Data availability**

Raw and processed sequencing files were submitted to the NCBI Gene Expression

Omnibus (GEO) as a Superseries, and the mid-cycle and early-cycle projects can be

found using accession number GSE106763.

ACKNOWLEGEMENTS

We would like to acknowledge Dr. Scot Ouellette and Nicholas Pokorzynski for critical

reading of the manuscript. This work was funded by start-up funds for R.A.C. from the

School of Molecular Biosciences, College of Veterinary Medicine at Washington State

University, NIH grant 2R01AI065545-06A1 awarded to R.A.C., and NIH training grant

T32AI7025-36 for A.J.B.

REFERENCES

1.     CDC. 2016. Sexually Transmitted Disease Surveillance 2015. Atlanta: U.S.

Department of Health and Human Services.

2.     Price MJ, Ades AE, Welton NJ, Simms I, Macleod J, Horner PJ. 2016. Proportion

of Pelvic Inflammatory Disease Cases Caused by Chlamydia trachomatis:

Consistent Picture From Different Methods. J Infect Dis 214:617–24.

3.     Stoner BP, Cohen SE. 2015. Lymphogranuloma Venereum 2015: Clinical

Presentation, Diagnosis, and Treatment: Table 1. Clin Infect Dis 61:S865–S873.

4.     Kjaer HO, Dimcevski G, Hoff G, Olesen F, Ostergaard L. 2000. Recurrence of

urogenital Chlamydia trachomatis infection evaluated by mailed samples obtained

at home: 24 weeks' prospective follow up study. Sex Transm Infect 76:169–72.

5.  Batteiger BE, Tu W, Ofner S, Van Der Pol B, Stothard DR, Orr DP, Katz BP,

Fortenberry JD. 2010. Repeated Chlamydia trachomatis genital infections in

adolescent women. J Infect Dis 201:42–51.

6.  Wyrick PB. 2010. Chlamydia trachomatis Persistence In Vitro: An Overview. J

Infect Dis 201:88–95.

7.  Lewis ME, Belland RJ, AbdelRahman YM, Beatty WL, Aiyar AA, Zea AH, Greene

SJ, Marrero L, Buckner LR, Tate DJ, McGowin CL, Kozlowski PA, O'Brien M, Lillis

RA, Martin DH, Quayle AJ. 2014. Morphologic and molecular evaluation of

Chlamydia trachomatis growth in human endocervix reveals distinct growth

patterns. Front Cell Infect Microbiol 4:71.

8.  AbdelRahman YM, Belland RJ. 2005. The chlamydial developmental cycle. FEMS

Microbiol Rev 29:949–959.

9.  Matsumoto A, Manire GP. 1970. Electron microscopic observations on the effects

of penicillin on the morphology of Chlamydia psittaci. J Bacteriol 101:278–85.

10. Kazar J, Gillmore JD, Gordon FB. 1971. Effect of Interferon and Interferon

Inducers on Infections with a Nonviral Intracellular Microorganism, Chlamydia

trachomatis. Infect Immun 3:825–32.

11. Raulston JE. 1997. Response of Chlamydia trachomatis serovar E to iron

restriction in vitro and evidence for iron-regulated chlamydial proteins. Infect

Immun 65:4539–47.

12. Byrne GI, Lehmann LK, Landry GJ. 1986. Induction of tryptophan catabolism is

the mechanism for gamma-interferon-mediated inhibition of intracellular

Chlamydia psittaci replication in T24 cells. Infect Immun 53:347–51.

13.    Beatty WL, Byrne GI, Morrison RP. 1993. Morphologic and antigenic

704            characterization of interferon gamma-mediated persistent Chlamydia trachomatis

705            infection in vitro. Proc Natl Acad Sci U S A 90:3998–4002.

706    14.    Beatty WL, Morrison RP, Byrne GI. 1995. Reactivation of persistent Chlamydia

707            trachomatis infection in cell culture. Infect Immun 63:199–205.

708    15.    Belland RJ, Nelson DE, Virok D, Crane DD, Hogan D, Sturdevant D, Beatty WL,

709            Caldwell HD. 2003. Transcriptome analysis of chlamydial growth during IFN- -

710            mediated persistence and reactivation. Proc Natl Acad Sci 100:15971–15976.

711    16.    Reveneau N, Crane DD, Fischer E, Caldwell HD. 2005. Bactericidal activity of

712            first-choice antibiotics against gamma interferon-induced persistent infection of

713            human epithelial cells by Chlamydia trachomatis. Antimicrob Agents Chemother

714            49:1787–93.

715    17.    Ouellette SP, Hatch TP, AbdelRahman YM, Rose LA, Belland RJ, Byrne GI.

716            2006. Global transcriptional upregulation in the absence of increased translation

717            in Chlamydia during IFNgamma-mediated host cell tryptophan starvation. Mol

718            Microbiol 62:1387–401.

719    18.    Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W,

720            Olinger L, Tatusov RL, Zhao Q, Koonin E V., Davis RW. 1998. Genome

721            Sequence of an Obligate Intracellular Pathogen of Humans: Chlamydia

722            trachomatis. Science (80- ) 282.

723    19.    Chen W. 2011. IDO: more than an enzyme. Nat Immunol 12:809–811.

724    20.    Byrd TF, Horwitz MA. 1989. Interferon gamma-activated human monocytes

725            downregulate transferrin receptors and inhibit the intracellular multiplication of

Legionella pneumophila by limiting the availability of iron. J Clin Invest 83:1457–

1465.

21.  Nemeth E, Valore E V, Territo M, Schiller G, Lichtenstein A, Ganz T. 2003.

Hepcidin, a putative mediator of anemia of inflammation, is a type II acute-phase

protein. Blood 101:2461–2463.

22.  Ludwiczek S, Aigner E, Theurl I, Weiss G, Marx JJ, Brock J. 2003. Cytokine-

mediated regulation of iron transport in human monocytic cells. Blood 101:4148–

54.

23.  Mpiga P, Ravaoarinoro M. 2006. Chlamydia trachomatis persistence: An update.

Microbiol Res.

24.  Theurl I, Theurl M, Seifert M, Mair S, Nairz M, Rumpold H, Zoller H, Bellmann-

Weiler R, Niederegger H, Talasz H, Weiss G. 2008. Autocrine formation of

hepcidin induces iron retention in human monocytes. Blood 111:2392–9.

25.  Paradkar PN, De Domenico I, Durchfort N, Zohn I, Kaplan J, Ward DM. 2008. Iron

depletion limits intracellular bacterial growth in macrophages. Blood 112:866–74.

26.  Nairz M, Haschka D, Demetz E, Weiss G. 2014. Iron at the interface of immunity

and infection. Front Pharmacol 5:152.

27.  Pokorzynski ND, Thompson CC, Carabeo RA. 2017. Ironing Out the

Unconventional Mechanisms of Iron Acquisition and Gene Regulation in

Chlamydia. Front Cell Infect Microbiol 7:394.

28.  Hentze MW, Muckenthaler MU, Galy B, Camaschella C. 2010. Two to tango:

regulation of Mammalian iron metabolism. Cell 142:24–38.

29.  Vardhan H, Bhengraj AR, Jha R, Singh Mittal A. 2009. Chlamydia trachomatis

749  alters iron-regulatory protein-1 binding capacity and modulates cellular iron

750  homeostasis in HeLa-229 cells. J Biomed Biotechnol 2009:342032.

751 30. Kim I, Yetley EA, Calvo MS. 1993. Variations in iron-status measures during the

752  menstrual cycle. Am J Clin Nutr 58:705–9.

753 31. Kelver ME, Kaul A, Nowicki B, Findley WE, Hutchens TW, Nagamani M. 1996.

754  Estrogen regulation of lactoferrin expression in human endometrium. Am J

755  Reprod Immunol 36:243–7.

756 32. Aisen P, Enns C, Wessling-Resnick M. 2001. Chemistry and biology of eukaryotic

757  iron metabolism. Int J Biochem Cell Biol 33:940–959.

758 33. Skaar EP. 2010. The battle for iron between bacterial pathogens and their

759  vertebrate hosts. PLoS Pathog 6:e1000949.

760 34. Caza M, Kronstad JW. 2013. Shared and distinct mechanisms of iron acquisition

761  by bacterial and fungal pathogens of humans. Front Cell Infect Microbiol 3:80.

762 35. Hood MI, Skaar EP. 2012. Nutritional immunity: transition metals at the pathogen–

763  host interface. Nat Rev Microbiol 10:525–537.

764 36. Pan X, Tamilselvam B, Hansen EJ, Daefler S. 2010. Modulation of iron

765  homeostasis in macrophages by bacterial intracellular pathogens. BMC Microbiol

766  10:64.

767 37. Scidmore MA, Fischer ER, Hackstadt T. 2003. Restricted fusion of Chlamydia

768  trachomatis vesicles with endocytic compartments during the initial stages of

769  infection. Infect Immun 71:973–84.

770 38. Ouellette SP, Carabeo RA. 2010. A functional slow recycling pathway of

771  transferrin is required for growth of Chlamydia. Front Microbiol 1:1–12.

772    39.    Boradia VM, Malhotra H, Thakkar JS, Tillu VA, Vuppala B, Patil P, Sheokand N,

773           Sharma P, Chauhan AS, Raje M, Raje CI. 2014. Mycobacterium tuberculosis

774           acquires iron by cell-surface sequestration and internalization of human holo-

775           transferrin. Nat Commun 5:4730.

776    40.    Raulston JE, Miller JD, Davis CH, Schell M, Baldwin A, Ferguson K, Lane H.

777           2007. Identification of an iron-responsive protein that is antigenic in patients with

778           Chlamydia trachomatis genital infections. FEMS Immunol Med Microbiol 51:569–

779           76.

780    41.    Miller JD, Sal MS, Schell M, Whittimore JD, Raulston JE. 2009. Chlamydia

781           trachomatis YtgA is an iron-binding periplasmic protein induced by iron restriction.

782           Microbiology 155:2884–94.

783    42.    Thompson CC, Nicod SS, Malcolm DS, Grieshaber SS, Carabeo RA. 2012.

784           Cleavage of a putative metal permease in Chlamydia trachomatis yields an iron-

785           dependent transcriptional repressor. Proc Natl Acad Sci U S A 109:10546–51.

786    43.    Thompson CC, Carabeo RA. 2011. An optimal method of iron starvation of the

787           obligate intracellular pathogen, Chlamydia trachomatis. Front Microbiol 2:1–10.

788    44.    LaRue RW, Dill BD, Giles DK, Whittimore JD, Raulston JE. 2007. Chlamydial

789           Hsp60-2 is iron responsive in Chlamydia trachomatis serovar E-infected human

790           endometrial epithelial cells in vitro. Infect Immun 75:2374–80.

791    45.    Dill BD, Dessus-Babus S, Raulston JE. 2009. Identification of iron-responsive

792           proteins expressed by Chlamydia trachomatis reticulate bodies during intracellular

793           growth. Microbiology 155:210–9.

46.    Breuer W, Epsztejn S, Cabantchik ZI. 1995. Iron acquired from transferrin by

K562 cells is delivered into a cytoplasmic pool of chelatable iron(II). J Biol Chem

270:24209–15.

47.  Wilderman PJ, Sowa NA, FitzGerald DJ, FitzGerald PC, Gottesman S, Ochsner

UA, Vasil ML. 2004. Identification of tandem duplicate regulatory small RNAs in

Pseudomonas aeruginosa involved in iron homeostasis. Proc Natl Acad Sci

101:9792–9797.

48.  Klitgaard K, Friis C, Angen Ø, Boye M. 2010. Comparative profiling of the

transcriptional response to iron restriction in six serotypes of Actinobacillus

pleuropneumoniae with different virulence potential. BMC Genomics 11:698.

49.  Timms P. 2009. Differential transcriptional responses between the interferon-

gamma-induction and iron-limitation models of persistence for Chlamydia

pneumoniae. J Microbiol Immunol Infect 42:27–37.

50.  Consortium U. 2017. UniProt: the universal protein knowledgebase. Nucleic Acids

Res 45:D158–D169.

51.  Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J,

Simonovic M, Roth A, Santos A, Tsafou KP, Kuhn M, Bork P, Jensen LJ, von

Mering C. 2015. STRING v10: protein-protein interaction networks, integrated

over the tree of life. Nucleic Acids Res 43:D447–D452.

52.  Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration

and interpretation of large-scale molecular data sets. Nucleic Acids Res 40:D109-

14.

53.  Fehlner-Gardiner C, Roshick C, Carlson JH, Hughes S, Belland RJ, Caldwell HD,

McClarty G. 2002. Molecular basis defining human Chlamydia trachomatis tissue

tropism. A possible role for tryptophan synthase. J Biol Chem 277:26893–903.

54.  Wood H, Fehlner-Gardner C, Berry J, Fischer E, Graham B, Hackstadt T, Roshick

C, McClarty G. 2003. Regulation of tryptophan synthase gene expression in

Chlamydia trachomatis. Mol Microbiol 49:1347–1359.

55.  Caldwell HD, Wood H, Crane D, Bailey R, Jones RB, Mabey D, Maclean I,

Mohammed Z, Peeling R, Roshick C, Schachter J, Solomon AW, Stamm WE,

Suchland RJ, Taylor L, West SK, Quinn TC, Belland RJ, McClarty G. 2003.

Polymorphisms in Chlamydia trachomatis tryptophan synthase genes differentiate

between genital and ocular isolates. J Clin Invest 111:1757–69.

56.  Karayiannis P, Hobson D. 1981. Amino acid requirements of a Chlamydia

trachomatis genital strain in McCoy cell cultures. J Clin Microbiol 13:427–32.

57.  Al-Younes HM, Gussmann J, Braun PR, Brinkmann V, Meyer TF. 2006. Naturally

occurring amino acids differentially influence the development of Chlamydia

trachomatis and Chlamydia (Chlamydophila) pneumoniae. J Med Microbiol

55:879–886.

58.  Braun PR, Al-Younes H, Gussmann J, Klein J, Schneider E, Meyer TF. 2008.

Competitive Inhibition of Amino Acid Uptake Suppresses Chlamydial Growth:

Involvement of the Chlamydial Amino Acid Transporter BrnQ. J Bacteriol

190:1822–1830.

59.  Kane CD, Vena RM, Ouellette SP, Byrne GI. 1999. Intracellular Tryptophan Pool

Sizes May Account for Differences in Gamma Interferon-Mediated Inhibition and

Persistence of Chlamydial Growth in Polarized and Nonpolarized Cells. Infect

Immun 67:1666–1671.

60. Leonhardt RM, Lee SJ, Kavathas PB, Cresswell P. 2007. Severe tryptophan

starvation blocks onset of conventional persistence and reduces reactivation of

Chlamydia trachomatis. Infect Immun 75:5105–5117.

61. Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, Thomas PD. 2017.

PANTHER version 11: expanded annotation data from Gene Ontology and

Reactome pathways, and data analysis tool enhancements. Nucleic Acids Res

45:D183–D189.

62. Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM, Kothari A,

Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P,

Weaver DS, Karp PD. 2016. The MetaCyc database of metabolic pathways and

enzymes and the BioCyc collection of pathway/genome databases. Nucleic Acids

Res 44:D471–D480.

63. Liechti GW, Kuru E, Hall E, Kalinda A, Brun Y V., VanNieuwenhze M, Maurelli AT.

2013. A new metabolic cell-wall labelling method reveals peptidoglycan in

Chlamydia trachomatis. Nature 506:507–510.

64. Soupene E, Wang D, Kuypers FA. 2015. Remodeling of host phosphatidylcholine

by Chlamydia acyltransferase is regulated by acyl-CoA binding protein ACBD6

associated with lipid droplets. Microbiologyopen 4:235.

65. Smith R., Martell AE. 2002. NIST critically selected stability consants of metal

complexes database, NIST Standard Reference Database 46, Version 8.0.

66. Al-Younes HM, Rudel T, Brinkmann V, Szczepek AJ, Meyer TF. 2001. Low iron

availability modulates the course of Chlamydia pneumoniae infection. Cell

Microbiol 3:427–437.

67.    Mäurer AP, Mehlitz A, Mollenkopf HJ, Meyer TF. 2007. Gene expression profiles

of Chlamydophila pneumoniae during the developmental cycle and iron depletion-

mediated persistence. PLoS Pathog 3:0752–0769.

68.    Mukhopadhyay S, Miller RD, Sullivan ED, Theodoropoulos C, Mathews SA,

Timms P, Summersgill JT. 2006. Protein expression profiles of Chlamydia

pneumoniae in models of persistence versus those of heat shock stress

response. Infect Immun 74:3853–3863.

69.    de Barsy M, Frandi A, Panis G, Théraulaz L, Pillonel T, Greub G, Viollier PH.

2016. Regulatory (pan-)genome of an obligate intracellular pathogen in the PVC

superphylum. ISME J 10:2129–2144.

70.    Brickman TJ, Cummings CA, Liew S-Y, Relman DA, Armstrong SK. 2011.

Transcriptional profiling of the iron starvation response in Bordetella pertussis

provides new insights into siderophore utilization and virulence gene expression.

J Bacteriol 193:4798–812.

71.    Zhou D, Qin L, Han Y, Qiu J, Chen Z, Li B, Song Y, Wang J, Guo Z, Zhai J, Du Z,

Wang X, Yang R. 2006. Global analysis of iron assimilation and fur regulation in

Yersinia pestis. FEMS Microbiol Lett 258:9–17.

72.    Kunkle CA, Schmitt MP. 2003. Analysis of the Corynebacterium diphtheriae DtxR

regulon: identification of a putative siderophore synthesis and transport system

that is similar to the Yersinia high-pathogenicity island-encoded yersiniabactin

synthesis and uptake system. J Bacteriol 185:6826–40.

73.    Stenner-Liewen F, Liewen H, Zapata JM, Pawlowski K, Godzik A, Reed JC. 2002.

CADD, a Chlamydia protein that interacts with death receptors. J Biol Chem

277:9633–6.

74.   Adams NE, Thiaville JJ, Proestos J, Juárez-Vázquez AL, McCoy AJ, Barona-

Gómez F, Iwata-Reuyl D, de Crécy-Lagard V, Maurelli AT. 2014. Promiscuous

and adaptable enzymes fill "holes" in the tetrahydrofolate pathway in Chlamydia

species. MBio 5:e01378-14.

75.   Taylor LD, Nelson DE. 2014. Chlamydial MACPF Protein CT153, p. 255–269. *In*

Sub-cellular biochemistry.

76.   McHugh JP, Rodríguez-Quinoñes F, Abdul-Tehrani H, Svistunenko DA, Poole

RK, Cooper CE, Andrews SC. 2003. Global iron-dependent gene regulation in

Escherichia coli. A new mechanism for iron homeostasis. J Biol Chem

278:29478–86.

77.   Rosenkranz HS, Gutter B, Becker Y. 1973. Studies on the developmental cycle of

Chlamydia trachomatis: selective inhibition by hydroxyurea. J Bacteriol 115:682–

90.

78.   Odsbu I, Morigen S, Skarstad K, Beckwith J, Beckwith J. 2009. A Reduction in

Ribonucleotide Reductase Activity Slows Down the Chromosome Replication

Fork but Does Not Change Its Localization. PLoS One 4:e7617.

79.   Jain V, Kumar M, Chatterji D. 2006. ppGpp: stringent response and survival. J

Microbiol 44:1–10.

80.   Potrykus K, Murphy H, Philippe N, Cashel M. 2011. ppGpp is the major source of

growth rate control in E. coli. Environ Microbiol 13:563–575.

81.   Perederina A, Svetlov V, Vassylyeva MN, Tahirov TH, Yokoyama S, Artsimovitch

I, Vassylyev DG. 2004. Regulation through the Secondary Channel—Structural

Framework for ppGpp-DksA Synergism during Transcription. Cell 118:297–309.

82.  Cashel M. 1969. The control of ribonucleic acid synthesis in Escherichia coli. IV.

Relevance of unusual phosphorylated compounds from amino acid-starved

stringent strains. J Biol Chem 244:3133–41.

83.  Haseltine WA, Block R. 1973. Synthesis of guanosine tetra- and pentaphosphate

requires the presence of a codon-specific, uncharged transfer ribonucleic acid in

the acceptor site of ribosomes. Proc Natl Acad Sci U S A 70:1564–8.

84.  Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Hightower GA, Smith JT,

Conway T. 2008. The global, ppGpp-mediated stringent response to amino acid

starvation in Escherichia coli. Mol Microbiol 68:1128–1148.

85.  Mittenhuber G. 2001. Comparative genomics and evolution of genes encoding

bacterial (p)ppGpp synthetases/hydrolases (the Rel, RelA and SpoT proteins). J

Mol Microbiol Biotechnol 3:585–600.

86.  Ouellette SP, Rueden KJ, Rucks EA. 2016. Tryptophan Codon-Dependent

Transcription in Chlamydia pneumoniae during Gamma Interferon-Mediated

Tryptophan Limitation. Infect Immun 84:2703–2713.

87.  Miethke M, Westers H, Blom E-J, Kuipers OP, Marahiel MA. 2006. Iron starvation

triggers the stringent response and induces amino acid biosynthesis for

bacillibactin production in Bacillus subtilis. J Bacteriol 188:8655–7.

88.  Akers JC, Tan M. 2006. Molecular mechanism of tryptophan-dependent

transcriptional regulation in Chlamydia trachomatis. J Bacteriol 188:4236–4243.

89.  Carlson JH, Wood H, Roshick C, Caldwell HD, McClarty G. 2006. In vivo and in

vitro studies of Chlamydia trachomatis TrpR:DNA interactions. Mol Microbiol

59:1678–91.

90.    Amirshahi A, Wan C, Beagley K, Latter J, Symonds I, Timms P. 2011. Modulation

935            of the Chlamydia trachomatis in vitro transcriptome response by the sex

936            hormones estradiol and progesterone. BMC Microbiol 11:150.

964    TABLES

| Table 1. Differentially expressed genes after 3h BPD treatment during mid-cycle development.* | | | | | | |
|---|---|---|---|---|---|---|
| Feature ID | Locus tag | Fold change | P-value | Annotation | Functional category | UniProtKB ID |
| CTL0013 | CTL0013 | 3.45 | 3.10E-04 | hypothetical, YGGT family | Hypothetical | A0A0H3MB25 |
| trpB | CTL0423 | 3.18 | 3.23E-07 | tryptophan synthase subunit B | Amino acid biosynthesis | A0A0H3MD30 |
| glgA | CTL0167 | 2.95 | 1.24E-04 | glycogen synthase | Energy Metabolism | B0B925 |
| murB | CTL0203 | 2.52 | 6.41E-03 | UDP-N-acetylenolpyruvoylglucosamine reductase | Other | B0B960 |
| CTL0525 | CTL0525 | 2.38 | 4.53E-05 | TPR-containing domain | Hypothetical | A0A0H3MKX6 |
| recA | CTL0018 | 2.21 | 1.98E-03 | recombinase A | DNA replication and repair | B0B8M5 |
| CTL0339 | CTL0339 | 2.2 | 2.29E-03 | phosphatidylcholine-hydrolyzing phospholipase D | Other | A0A0H3MCY1 |
| aroL | CTL0621 | 2.13 | 6.56E-03 | shikimate kinase 2 | Amino acid biosynthesis | A0A0H3MDG7 |
| hemE | CTL0116 | 2.09 | 1.68E-03 | uroporphyrinogen decarboxylase | Cofactor biosynthesis | B0B8X3 |
| CTL0408 | CTL0408 | 2.08 | 1.36E-05 | MIR, MAC/perforin domain-containing protein | Other | A0A0H3MGT6 |
| nrdA | CTL0199 | 2 | 1.48E-05 | Ribonucleoside-diphosphate reductase | DNA replication and repair | A0A0H3MCP2 |
| mqnD | CTL0514 | 2 | 2.45E-03 | 1,4-dihydroxy-6-naphtoate synthase | Cofactor biosynthesis | A0A0H3MC13 |
| CTL0704 | CTL0704 | 1.99 | 0.01 | hypothetical | Hypothetical | A0A0H3MCH5 |
| trpA | CTL0424 | 1.94 | 4.15E-03 | tryptophan synthase subunit A | Amino acid biosynthesis | A0A0H3MKP4 |
| CTL0255 | CTL0255 | 1.93 | 4.37E-03 | hypothetical | Hypothetical | A0A0H3MGJ2 |
| pepF | CTL0367 | 1.9 | 6.17E-04 | endopeptidase F | Protein processing and folding | A0A0H3MKK2 |
| rnc | CTL0549 | 1.89 | 0.01 | ribonuclease III | Transcriptional regulation | B0B7L3 |
| CTL0823 | CTL0823 | 1.88 | 7.21E-04 | hypothetical | Hypothetical | A0A0H3MLF6 |
| CTL0301 | CTL0301 | 1.81 | 1.32E-04 | probable cytosol aminopeptidase pepA | Protein processing and folding | B0B9F3 |
| CTL0884 | CTL0884 | 1.81 | 7.04E-03 | hypothetical | Hypothetical | A0A0H3MCL0 |
| CTL0885 | CTL0885 | 1.77 | 3.48E-03 | hypothetical effector | Type III Secretion | A0A0H3MHM9 |
| lpdA | CTL0820 | 1.76 | 3.80E-03 | dihydrolipoyl dehydrogenase | Energy Metabolism | A0A0H3MHJ2 |
| CTL0096 | CTL0096 | 1.72 | 1.98E-04 | putative cation transporting ATPase | Nutrient transport | A0A0H3MCG9 |
| gp6D | CTL0846 | 1.72 | 2.77E-03 | virulence plasmid pGP6-D related protein | Hypothetical | A0A0H3MLH1 |
| dnaQ | CTL0513 | 1.69 | 0.01 | DNA polymerase III subunit epsilon | DNA replication and repair | A0A0H3MKW6 |
| CTL0512 | CTL0512 | 1.68 | 4.52E-03 | mcsc, secretion chaperone | Type III Secretion | A0A0H3MDA7 |
| CTL0847 | CTL0847 | 1.63 | 6.69E-03 | hypothetical | Hypothetical | A0A0H3MCR9 |
| oppA3 | CTL0450 | 1.61 | 9.20E-03 | oligopeptide transporter | Nutrient transport | A0A0H3MKR3 |
| CTL0102 | CTL0102 | 1.6 | 7.09E-03 | putative exported protein | Hypothetical | A0A0H3MG91 |
| lipA | CTL0821 | 1.59 | 8.90E-03 | lipioic acid synthase | Other | B0B8D2 |
| pheT | CTL0736 | 1.59 | 0.01 | phenylalanine−tRNA ligase beta subunit | Translation | A0A0H3MHF2 |
| cysS | CTL0151 | 1.56 | 2.47E-03 | cysteinyl-tRNA synthetase | Translation | A0A0H3MGC2 |
| CTL0055 | CTL0055 | 1.51 | 7.04E-03 | hypothetical | Hypothetical | A0A0H3MAN0 |
| pal | CTL0863 | 1.51 | 7.29E-03 | peptidogycan-associated lipoprotein | Other | A0A0H3MDU9 |
| thrS | CTL0844 | 1.48 | 6.31E-03 | threonine-tRNA ligase | Translation | B0B8F5 |
| CTL0476 | CTL0476 | 1.48 | 0.01 | candidate inclusion membrane protein | Hypothetical | A0A0H3MKT3 |
| CTL0043 | CTL0043 | 1.46 | 0.01 | type III secretion structural protein | Type III Secretion | A0A0H3MG59 |
| glmS | CTL0188 | 1.46 | 0.01 | Glutamine--fructose-6-phosphate aminotransferase | Energy Metabolism | A0A0H3MCN0 |
| CTL0684 | CTL0684 | 1.45 | 8.29E-03 | hypothetical | Hypothetical | A0A0H3MCF9 |
| nusA | CTL0352 | -1.48 | 4.90E-03 | transcription termination factor | Transcriptional regulation | A0A0H3MGQ1 |
| prfA | CTL0278 | -1.56 | 9.49E-03 | Peptide chain release factor RF1 | Translation | B0B9D0 |
| rplW | CTL0788 | -1.58 | 5.43E-04 | ribosomal subunit | Translation | B0B8A0 |
| CTL0326 | CTL0326 | -1.62 | 6.61E-03 | ytgD, ABC transport protein, membrane permease | Nutrient transport | A0A0H3MGN6 |
| trpF | CTL0581 | -1.67 | 2.26E-03 | N-(5'-phosphoribosyl)anthranilate isomerase | Cofactor biosynthesis | B0B7P4 |
| folX | CTL0878 | -1.68 | 7.23E-03 | Dihydroneopterin triphosphate 2'-epimerase | Cofactor biosynthesis | A0A0H3MCK6 |
| pgsA_2 | CTL0757 | -1.71 | 1.74E-03 | CDP-diacylglycerol-glycerol-3-phosphate 3-phosphatidyl-transferase | Other | A0A0H3MHG9 |
| sodM | CTL0546 | -1.74 | 1.46E-03 | superoxide dismutase | Redox homeostasis | A0A0H3MKY6 |
| CTL0138 | CTL0138 | -1.74 | 4.48E-03 | ribosomal silencing factor RsfS | Translation | A0A0H3MAQ8 |
| CTL0061 | CTL0061 | -1.75 | 2.26E-04 | inorganic phosphate transporter | Nutrient transport | A0A0H3MG71 |
| CTL0486 | CTL0486 | -1.77 | 3.48E-03 | putative membrane transport protein | Nutrient transport | A0A0H3MBY7 |
| gltT | CTL0658 | -1.85 | 3.07E-03 | sodium:dicarboxylate symport protein | Nutrient transport | A0A0H3MCC1 |
| smpB | CTL0332 | -1.88 | 0.01 | SsrA-binding protein | Translation | A0A0H3MBC6 |
| aroA | CTL0620 | -1.91 | 1.85E-04 | 3-phosphoshikimate 1-carboxyvinyltransferase | Energy Metabolism | B0B7T5 |
| CTL0132 | CTL0132 | -1.92 | 3.03E-03 | UPF0109-containing putative RNA-binding protein | Translation | A0A0H3MAQ7 |
| CTL0548 | CTL0548 | -1.98 | 1.14E-04 | dcrA, putative non-heme Fe(II) 2-oxoglutarate | Hypothetical | A0A0H3MBY2 |
| secG | CTL0606 | -2.01 | 0.01 | Protein-export membrane protein SecG | Protein processing and folding | A0A0H3MH61 |
| rpsK | CTL0770 | -2.07 | 1.59E-06 | ribosomal subunit | Translation | B0B882 |
| rplT | CTL0207 | -2.07 | 1.62E-06 | ribosomal subunit | Translation | B0B964 |
| rplN | CTL0780 | -2.08 | 3.98E-03 | ribosomal subunit | Translation | B0B892 |
| CTL0720 | CTL0720 | -2.12 | 5.09E-03 | SWIB-domain containing protein | Hypothetical | A0A0H3MC95 |
| gcsH | CTL0534 | -2.14 | 9.08E-04 | glycine cleavage system H protein | Amino acid biosynthesis | B0B7J8 |
| CTL0680 | CTL0680 | -2.14 | 1.59E-03 | putative rRNA processing peptide | Translation | A0A0H3MHB1 |
| rpmI | CTL0206 | -2.14 | 1.75E-03 | ribosomal subunit | Translation | A0A0H3MCP7 |
| fer | CTL0315 | -2.26 | 6.48E-05 | ferredoxin | Redox homeostasis | A0A0H3MCW5 |
| CTL0552 | CTL0552 | -2.27 | 1.09E-03 | TPR-containing domain | Hypothetical | pseudogene |
| CTL0222 | CTL0222 | -2.27 | 6.58E-03 | hypothetical | Hypothetical | A0A0H3MK96 |
| infA2 | CTL0575 | -2.29 | 6.70E-06 | Translation initiation factor IF-1 | Translation | A0A0H3MDD9 |
| rpsT | CTL0881 | -2.38 | 1.13E-04 | ribosomal subunit | Translation | B0B8J2 |
| CTL0335 | CTL0335 | -2.43 | 0.01 | Putative integral membrane protein | Hypothetical | A0A0H3MCX6 |
| ltuA | CTL0631 | -2.68 | 3.38E-03 | Late transcription unit A protein | Hypothetical | A0A0H3MH76 |
| rpsO | CTL0215 | -2.72 | 4.14E-04 | ribosomal subunit | Translation | B0B972 |
| ndk | CTL0762 | -2.76 | 7.97E-04 | nucleoside diphosphate kinase | DNA replication and repair | B0B874 |
| pGP8-D | L2bCS784 | -2.99 | 0.01 | Virulence plasmid integrase pGP8-D | DNA replication and repair | B0BCM4 |
| dut | CTL0544 | -3.02 | 6.06E-06 | deoxyuridine 5'-triphosphate nucleotidohydrolase | DNA replication and repair | B0B7K8 |
| rpmJ | CTL0154 | -3.03 | 3.18E-04 | ribosomal subunit | Translation | B0B912 |
| CTL0021 | CTL0021 | -3.16 | | hypothetical | Hypothetical | A0A0H3MG42 |

This data was exported from CLC Genomics Workbench 9.5.3. rRNAs, tRNAs, and Features (genes) with less than 10 reads in all samples were eliminated from the dataset prior to normalization and EDGE analysis, and only includes genes that were differentially expressed with a significance p-value ≤ 0.01. *FDR-corrected p-values can be found in Supplemental Table 4.

| Table 2. Differentially expressed genes after 6h BPD treatment during mid-cycle development.* | | | | | | |
|---|---|---|---|---|---|---|
| Feature ID | Locus Tag | P-value | Fold change | Annotation | Functional category | UniProtKB ID |
| *trpB* | CTL0423 | 2.61E-07 | 3.5 | tryptophan synthase subunit B | Amino acid biosynthesis | A0A0H3MD30 |
| *trpA* | CTL0424 | 9.26E-06 | 3.21 | tryptophan synthase subunit A | Amino acid biosynthesis | A0A0H3MKP4 |
| *nrdA* | CTL0199 | 3.50E-06 | 2.39 | ribonucleoside-diphosphate reductase | DNA replication and repair | A0A0H3MCP2 |
| *CTL0071* | CTL0071 | 6.73E-03 | 1.73 | hypothetical | Hypothetical | A0A0H3MCG5 |
| *nrdB* | CTL0200 | 9.34E-03 | 1.63 | ribonucleoside-diphosphate reductase | DNA replication and repair | A0A0H3MK81 |
| *CTL0619* | CTL0619 | 1.68E-03 | -1.83 | hypothetical integral membrane protein | Hypothetical | A0A0H3MH71 |
| *copD* | CTL0842 | 3.17E-03 | -2.13 | type III secretion system protein | Type III Secretion | A0A0H3MHK7 |
| *scc2* | CTL0839 | 1.37E-03 | -2.18 | type III secretion system chaperone | Type III Secretion | A0A0H3MLG7 |
| *tsp* | CTL0700 | 1.28E-03 | -2.34 | tail-specific protease | Protein processing and folding | A0A0H3MDM0 |
| *CTL0185* | CTL0185 | 6.85E-03 | -2.76 | hypothetical membrane protein | Hypothetical | A0A0H3MAV2 |
| *copB* | CTL0841 | 6.79E-05 | -2.81 | type III secretion system membrane protein | Type III Secretion | A0A0H3MCF0 |
| *CTL0840* | CTL0840 | 2.15E-03 | -2.97 | hypothetical | Hypothetical | A0A0H3MCQ4 |

This data was exported from CLC Genomics Workbench 9.5.3. rRNAs, tRNAs, and Features (genes) with less than 10 reads in all samples were eliminated from the dataset prior to normalization and EDGE analysis. This data only includes genes that were differentially expressed with a significance p-value ≤ 0.01. *FDR-corrected p-values can be found in Supplemental Table 4.

966
967

**Table 3. Differentially expressed genes after 3h BPD treatment during early-cycle development.***

| Feature ID | Locus Tag | Fold change | P-value | Annotation | Functional category | UniProtKB ID |
|---|---|---|---|---|---|---|
| CTL0149 | CTL0149 | 6.71 | 3.71E-03 | protein disulfide isomerase | Redox homeostasis | A0A0H3MAT1 |
| CTL0184 | CTL0184 | 3.4 | 4.44E-05 | hypothetical inclusion membrane protein | Hypothetical | A0A0H3MBD4 |
| trpA | CTL0424 | 3.12 | 3.27E-04 | tryptophan synthase subunit A | Amino acid biosynthesis | A0A0H3MKP4 |
| CTL0388 | CTL0388 | 2.93 | 3.02E-03 | hypothetical methyltransferase | Hypothetical | A0A0H3MKL6 |
| CTL0111 | CTL0111 | 2.89 | 0.000877 | rRNA methyltransferase TrmA | Translation | A0A0H3MAQ4 |
| hemN_1 | CTL0115 | 2.47 | 1.00E-02 | coproporphyrinogen-III oxidase | Cofactor biosynthesis | A0A0H3MJY5 |
| mip | CTL0803 | 2.37 | 3.97E-04 | peptidyl-prolyl cis-trans isomerase | Protein processing and folding | A0A0H3MDR1 |
| trpB | CTL0423 | 2.28 | 2.55E-03 | tryptophan synthase subunit B | Amino acid biosynthesis | A0A0H3MD30 |
| lpxB | CTL0668 | 2.28 | 0.00923 | lipid-A-disaccharide synthase | Other | A0A0H3MDJ6 |
| nrdB | CTL0200 | 2.12 | 1.57E-06 | ribonucleoside-diphosphate reductase subunit B | DNA replication and repair | A0A0H3MK81 |
| nrdA | CTL0199 | 2.09 | 4.10E-12 | ribonucleoside-diphosphate reductase subunit A | DNA replication and repair | A0A0H3MCP2 |
| CTL0874 | CTL0874 | 2.04 | 2.07E-06 | CADD,PABA synthase | Cofactor biosynthesis | A0A0H3MHM3 |
| CTL0360 | CTL0360 | 2.04 | 8.59E-03 | hypothetical | Hypothetical | A0A0H3MKJ7 |
| mutS | CTL0160 | 1.95 | 6.41E-06 | DNA mismatch repair protein | DNA replication and repair | B0B918 |
| dnaQ | CTL0513 | 1.88 | 2.83E-03 | DNA polymerase III subunit epsilon | DNA replication and repair | A0A0H3MKW6 |
| CTL0164 | CTL0164 | 1.86 | 1.20E-03 | hypothetical exported protein | Hypothetical | A0A0H3MBC3 |
| CTL0791 | CTL0791 | 1.82 | 1.78E-07 | hypothetical membrane protein | Hypothetical | A0A0H3MCL8 |
| trpS | CTL0848 | 1.81 | 1.46E-03 | tryptophan-tRNA ligase | Translation | A0A0H3MCF4 |
| aspC_1 | CTL0005 | 1.77 | 1.92E-05 | aminotransferase | Amino acid biosynthesis | A0A0H3MG09 |
| eno | CTL0850 | 1.75 | 2.51E-03 | enolase | Energy Metabolism | B0B8G1 |
| CTL0408 | CTL0408 | 1.73 | 3.50E-04 | MIR, MAC/perforin domain-containing protein | Other | A0A0H3MGT6 |
| recA | CTL0018 | 1.72 | 1.49E-03 | recombinase A | DNA replication and repair | B0B8M5 |
| sodM | CTL0546 | 1.65 | 7.17E-03 | superoxide dismutase | Redox homeostasis | A0A0H3MKY6 |
| brnQ | CTL0817 | 1.62 | 3.60E-04 | branched chain amino acid transporter | Nutrient transport | A0A0H3MLF2 |
| greA | CTL0004 | 1.59 | 2.63E-04 | transcription elongation factor | Transcriptional regulation | A0A0H3MAD9 |
| CTL0102 | CTL0102 | 1.58 | 4.63E-03 | hypothetical exported protein | Hypothetical | A0A0H3MG91 |
| ahpC | CTL0866 | 1.52 | 2.54E-03 | thio-specific antioxidant peroxidase | Redox homeostasis | A0A0H3MCJ5 |
| thrS | CTL0844 | 1.52 | 0.00501 | threonine-tRNA ligase | Translation | B0B8F5 |
| aspS | CTL0804 | 1.52 | 6.89E-03 | aspartate--tRNA ligase | Translation | B0B8B6 |
| rpoD | CTL0879 | 1.51 | 1.00E-02 | RNA polymerase sigma factor RpoD | Transcriptional regulation | A0A0H3MHM6 |
| glyA | CTL0691 | 1.47 | 0.00242 | serine hydroxymethyltransferase | Amino acid biosynthesis | B0B804 |
| sctJ | CTL0822 | 1.43 | 7.93E-03 | type III secretion protein | Type III Secretion | A0A0H3MDS0 |
| rpoC | CTL0566 | -1.28 | 7.87E-03 | DNA-directed RNA polymerase subunit beta' | Transcriptional regulation | B0B7N0 |
| pGP8-D | L2b_RS04755 | -1.36 | 4.91E-03 | virulence plasmid integrase pGP8-D | DNA replication and repair | B0BCM4 |
| rplW | CTL0788 | -1.41 | 0.0035 | ribosomal subunit | Translation | B0B8A0 |
| prfA | CTL0278 | -1.44 | 5.80E-03 | peptide chain release factor RF1 | Translation | B0B9D0 |
| rplC | CTL0790 | -1.52 | 4.30E-04 | ribosomal subunit | Translation | B0B8A2 |
| CTL0061 | CTL0061 | -1.52 | 2.95E-03 | inorganic phosphate transporter PHO4 | Nutrient transport | A0A0H3MG71 |
| CTL0659 | CTL0659 | -1.57 | 9.75E-04 | tetraacyldisaccharide 4'-kinase lpxK | Other | A0A0H3MC42 |
| CTL0473 | CTL0473 | -1.57 | 1.22E-03 | hypothetical exported protein | Hypothetical | A0A0H3MBQ9 |
| incD | CTL0370 | -1.59 | 3.33E-03 | inclusion membrane protein D | Other | B0B9M3 |
| plsX | CTL0182 | -1.59 | 1.00E-02 | phosphate acyltransferase | Other | B0B939 |
| CTL0613 | CTL0613 | -1.6 | 1.34E-03 | hypothetical inner membrane protein | Hypothetical | A0A0H3MC14 |
| pmpA | CTL0669 | -1.63 | 0.00534 | probable outer membrane protein PmpA | Other | A0A0H3ML49 |
| CTL0548 | CTL0548 | -1.66 | 1.01E-03 | hypothtetical non-heme Fe(II) 2-oxoglutarate | Hypothetical | A0A0H3MBY2 |
| CTL0541 | CTL0541 | -1.67 | 1.00E-02 | hypothetical membrane protein | Hypothetical | A0A0H3MC35 |
| sucB_2 | CTL0311 | -1.7 | 7.85E-03 | dihydrolipoyllysine-residue succinyltransferase | Energy Metabolism | A0A0H3ML42 |
| amn | CTL0120 | -1.71 | 4.71E-03 | AMP nucleosidase | DNA replication and repair | A0A0H3MGA3 |
| mrsA | CTL0547 | -1.77 | 1.00E-02 | phosphoglucomutase | Other | A0A0H3MC40 |
| ftsY | CTL0192 | -1.82 | 4.26E-04 | signal recognition particle receptor | Translation | A0A0H3MGE5 |
| CTL0609 | CTL0609 | -1.85 | 7.29E-06 | hypothetical exported protein | Hypothetical | A0A0H3MDF7 |
| dnaX_1 | CTL0439 | -1.92 | 5.15E-03 | DNA polymerase III subunit gamma/tau | DNA replication and repair | A0A0H3MBM8 |
| CTL0314 | CTL0314 | -2.04 | 2.67E-04 | hypothetical membrane protein | Hypothetical | A0A0H3MGM7 |
| CTL0430 | CTL0430 | -3.85 | 5.46E-05 | hypothetical integral membrane protein | Hypothetical | A0A0H3MBV0 |
| CTL0063 | CTL0063 | -3.89 | 2.61E-03 | hyptothetical | Hypothetical | A0A0H3MCG4 |

This data was exported from CLC Genomics Workbench 9.5.3. rRNAs, tRNAs, and Features (genes) with less than 10 reads in all samples were eliminated from the dataset prior to normalization and EDGE analysis. This data only includes genes that were differentially expressed with a significance p-value ≤ 0.01. *FDR-corrected p-values can be found in Supplemental Table 5.

968
969

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Table 3: Translation factors differentially expressed during mid-cycle iron-starvation. | | | | | | | |
| Gene | Expression change | Locus Tag | Protein ID | Annotation | Interacting component | Result of interaction | Expression change 6+3h BPD |
| **tRNA processing** | | | | | | | |
| rnpA_1 | decrease | CTL0153 | RnP1 | Ribonuclease P protein component | tRNA | Cleaves 5' end of pre-tRNA | |
| rnpA_2 | decrease | CTL0153A | RnP2 | Ribonuclease P protein component | tRNA | Cleaves 5' end of pre-tRNA | |
| **tRNA Biogenesis** | | | | | | | |
| cysS | increase | CTL0151 | CysS | Cysteine-tRNA ligase | tRNA(Cys) | Charges tRNA with Cysteine | |
| pheT | increase | CTL0736 | PheT | Phenylalanine-tRNA ligase beta | tRNA(Phe) | Charges tRNA with Phenylalanine | |
| glyQ | increase | CTL0165 | glyQ | Glycine-tRNA ligase alpha subunit | tRNA(Gly) | Charges tRNA with Glycine | |
| aspS | increase | CTL0804 | AspS | Aspartate-tRNA ligase | tRNA(Asp) | Charges tRNA with Aspartate | increase |
| thrS | increase | CTL0844 | ThrS | Threonine-tRNA ligase | tRNA(Thr) | Charges tRNA with Threonine | increase |
| truA | increase | CTL0723 | TruA | tRNA pseudouridine synthase A | tRNA-anticodon loop | Converts uridines at 38,39,40 to pseudouridine | |
| miaA | increase | CTL0135 | MiaA | tRNA dimethylallyltransferase | tRNA-anticodon loop | Converts adenine(37) to N6-(dimethylallyl)adenosine | |
| **rRNA processing** | | | | | | | |
| rnc | increase | CTL0549 | Rnc | Rnase III | 30s transcript | Cleaves 30s precursor transcript to 16S and 23S | increase |
| **Subunit assembly** | | | | | | | |
| rpsO | decrease | CTL0215 | S15 | ribosomal protein | 16s rRNA | Assembly of 30s subunit | |
| rpsT | decrease | CTL0881 | S20 | ribosomal protein | 16s rRNA | Assembly of 30s subunit | |
| rpsK | decrease | CTL0770 | S11 | ribosomal protein | 30s subunit | Forms Shine-Delgarno cleft | |
| rplT | decrease | CTL0207 | L20 | ribosomal protein | 23s rRNA | Assembly of 50s subunit | |
| rplN | decrease | CTL0780 | L14 | ribosomal protein | 23s rRNA | Fomrs bridge between 30s and 50s | |
| rplW | decrease | CTL0788 | L23 | ribosomal protein | 23s rRNA | Forms docking site for trigger factor | decrease |
| **Initiation** | | | | | | | |
| infA2 | decrease | CTL0575 | IF-1 | initiation factor | 30S-RpsA | Recruited to 30S by RpsA | |
| | decrease | | IF-1 | initiation factor | IF-3 | IF-1 and IF-3 recruit IF-2 to 30s, IF-2 recruits mRNA and tRNA | |
| rplN | decrease | CTL0780 | L14 | ribosomal protein | 23s rRNA | Forms bridge between 30s and 50s | decrease |
| rsfS | decrease | CTL0138 | RsfS | ribosomal silencing factor | RplN | Inhibits 70s assembly | |
| rplL | decrease | CTL0568 | L7/L12 | ribosomal protein | GTPases | Binds GTPases required for IF-3 recruitment | |
| **Elongation** | | | | | | | |
| rplL | decrease | CTL0568 | L7/L12 | ribosomal protein | GTPases | Binds GTPases required for EF-Tu and EF-G recruitment | |
| **Termination** | | | | | | | |
| prfA | decrease | CTL0278 | RF-1 | ribosome release factor | | Increases termination at UAA and UAG stop codons | decrease |
| rplL | decrease | CTL0568 | L7/L12 | ribosomal protein | GTPases | GTPase activity required for RF-3 recruitment | |
| **Recycling** | | | | | | | |
| rrf | decrease | CTL0046 | RrF | ribosome recycling factor | | Causes disassembly of stalled ribosomes | |
| CTL0634 | increase | CTL0634 | HflX | GTPase HflX | 50s subunit | Binds to E-site of 70s, disassembles ribosome | |
| **Nascent Peptide Folding/Targeting** | | | | | | | |
| ftsY | decrease | CTL0192 | FtsY | Signal recognition particle receptor | SRP-RNC | Targets nascent membrane proteins to Sec translocase | |
| secG | decrease | CTL0606 | SecG | Protein-export membrane protein | SecY | Forms SecYEG translocation channel | |
| smpB | decrease | CTL0332 | SmpB | SsrA-binding protein | tmRNA | Guides tmRNA into tRNA A site, rescuing stalled ribosomes and tagging nascent peptides for degradation | |
| **Unknown Function** | | | | | | | |
| rpmE | decrease | CTL0277 | L31 | ribosomal protein | 23s rRNA | Unknown | |
| rplQ | decrease | CTL0768 | L15 | ribosomal protein | 23s rRNA | Unknown | |
| rplY | decrease | CTL0168 | L25 | ribosomal protein | 5s rRNA | Binds to 5S in central protuberance | |

These genes were shown to be differentially regulated by EDGER analysis in CLC Genomics Workbench with p-value ≤ 0.05 during mid-cycle iron-starvation (12+3BPD vs 15h). Annotations and functions were retrieved from UniProt and BioCyc databases.

971    FIGURES
972
**Figure 1. Optimization of 2,2-Bipyrdyl (BPDL) treatment to induce iron-**

**responsiveness in the absence of persistence.** Timeline of BPDL-treatment A)

starting at 12h p.i., BPDL was supplemented to culture media for 3, 6, or 12h. Mock-

treated and BPDL-treated samples were tested for changes to B) morphology by

confocal microscopy, C) growth by qPCR, D) iron-responsive transcription (*ytgA, ahpC*)

D), and transcription of the developmental marker, *euo*, by RT-qPCR. Significant

changes with a p-value $\leq$ 0.05 in a one-tailed Student's t-test are indicated with an

asterisk, and are based on 3 biological replicates for the growth curve and 4 biological

replicates for RT-qPCR.

**Figure 2. Global and differential gene expression of the mid-cycle response to**

**iron-starvation.** The global response of *C. trachomatis* to iron starvation was detected

by RNA-sequencing and reads were aligned to the genome and plasmid. A) The

untreated expression profile is displayed for all genes that change significantly (p-value

$\leq$ 0.01) during mid-cycle development as a heatmap of log10 transformed normalized

expression means (left). Expression across the same genes are displayed for BPDL-

treated samples (right). The highest and lowest expression values are displayed in

green and red, respectively. B) Genes that are significantly changed in response to iron-

starvation, with a p-value $\leq$ 0.01, are displayed as a heatmap of fold-changes of BPDL-

treated compared to mock-treated equivalent samples. The most highly up-regulated

and down-regulated transcripts are displayed in green and red, respectively.

**Figure 3. Functional categorization of mid-cycle response to iron-starvation.**

Transcripts that were significantly up-regulated (left) or down-regulated (right) after 3h

BPDL treatment, starting at 12h p.i., are organized in pie charts by their functional

categories. Adjacent to each pie slice is the number of genes in that category, and in

parenthesis is the percentage of differentially expressed genes in the category.

N=number of differentially expressed genes, and the percentage of the total genome

that is represented.

**Figure 4. Confirmation of the mid-cycle response to iron starvation by RT-qPCR.**

Differentially expressed transcripts detected by RNA-sequencing were confirmed by RT-

qPCR. A) Up-regulated or B) down-regulated transcription is indicated as fold-changes

in transcripts after 3h BPD-treatment (solid gray bar) or 6h BPDL-treatment (striped bar)

compared to mock-treated samples at equivalent timepoints post-infection. An asterisk

indicates that the fold-change is significantly changed, with a p-value $\leq$ 0.05. Statistical

analysis was done with a one-tailed Student's t-test, based on at least 3 biological

replicates.

**Figure 5. Functional categorization of the early-cycle response to iron-starvation.**

Transcripts that were significantly up-regulated (left) or down-regulated (right) after 3h of

BPDL treatment, starting at 6h p.i., are organized in pie charts by their functional

categories. Adjacent to each pie slice is the number of genes in that category, and in

parenthesis is the percentage of differentially expressed genes in the category that

make up the pie. N=number of differentially expressed genes, and the percentage of the

total genome represented.

**Figure 6. Confirmation of the early-cycle response to iron starvation by RT-qPCR.**

Transcripts that were significantly changed by RNA-sequencing, in response to iron-

starvation starting at 6h p.i., were confirmed by RT-qPCR. A) Up-regulated or B) down-

regulated transcription is indicated as fold-changes in transcripts after 3h BPDL-

treatment in comparison to mock-treated at equivalent timepoints post-infection (solid

gray bar). An asterisk indicates that the fold-change is significantly changed, with a p-

value $\leq$ 0.05. Statistical analysis was done with a one-tailed Student's t-test, based on

two biological replicates.

**Figure 7. Pathway analysis of iron-starvation responses.** Association networks for

differentially expressed genes, with a p-value $\leq$ 0.05 and a minimum of 10 mapped

reads were generated for the A) early-cycle response (6+3h BPDL) and B) mid-cycle

response (12+3h BPDL) using STRING-db v.10.5.  Line thickness connecting nodes

(genes) correlates with confidence of gene association, with a minimum confidence

cutoff of 0.7. Mid-cycle (12+3h BPDL or 12+6h BPDL) clustered genes were mapped to

C) nucleotide metabolism and D) acetyl-CoA synthesis pathways using KEGGMapper

v.2.8. Up-regulated and down-regulated genes in C) and D) are shown with green and

red backgrounds, respectively. Unchanged genes have a white background.

SUPPLEMENTARY DATA

**Figure S1. Annotated heatmap of BPDL-treated and mock-treated gene**

**expression in *C. trachomatis* that corresponds to Figure 2A.**

**Figure S2. Annotated heatmap of mid-cycle iron-starvation corresponding to the**

**subset in Figure 2B.**

**Figure S3. Venn diagram of differential gene expression for all BPDL-treatments.**

Overlap in genes that were differentially up-regulated or down-regulated across multiple

treatments are displayed as a Venn diagram (p-value ≤ 0.01).

**Table S1. Summary of RNA-sequencing and mapping in this study.**

**Table S2. Normalized means of mock-treated and BPDL-treated transcription**

**during mid-cycle development of *C. trachomatis.*** The mean and $\log_{10}$ mean

expression values of genes that showed a significant change in gene expression during

normal mid-cycle development, p-value ≤ 0.01 are displayed for the following EdgeR

comparisons: 12h vs 18h, 12h vs 15h, 15h vs 18h) These values were used to create

the heatmaps in Figure 2A. Genes that had at least one value that was greater than the

4.5 threshold have an asterisk, and the values are displayed in bold.

**Table S3. Complete expression profile of *C. trachomatis* during normal**

**development.** The RNA-sequencing reads and EdgeR analysis of normal development

of *C. trachomatis* was exported from CLC Genomics Workbench 9.5.3.  Samples were

normalized across the entire dataset by quantile scaling. rRNAs, tRNAs, and Features

(genes) with less than 10 reads in all samples were eliminated from the dataset prior to

normalization and EDGE analysis. Unique Reads are raw values. Samples were

merged from multiple RNA-sequencing chips to obtain a minimum of 8X coverage.

**Table S4. Complete expression profile of *C. trachomatis* during mid-cycle iron**

**starvation.**

The RNA-sequencing reads and EdgeR analysis of iron-starved *C. trachomatis* was

exported from CLC Genomics Workbench 9.5.3.  Samples were normalized across the

entire dataset by quantile scaling. rRNAs, tRNAs, and Features (genes) with less than

10 reads in all samples were eliminated from the dataset prior to normalization and

EDGE analysis. Unique Reads are raw values. Samples were merged from multiple

RNA-sequencing chips to obtain a minimum of 8X coverage.

**Table S5. Complete expression profile of *C. trachomatis* during early-cycle iron**

**starvation.**

The RNA-sequencing reads and analysis of iron-starved *C. trachomatis* was exported

from CLC Genomics Workbench 9.5.3.  Samples were normalized across the entire

dataset by quantile scaling. rRNAs, tRNAs, and Features (genes) with less than 10

reads in all samples were eliminated from the dataset prior to normalization and EDGE

analysis. Unique Reads are raw values. Samples were merged from multiple RNA-

sequencing chips to obtain a minimum of 8X coverage.

**Table S6. Primers used in this study.**

1072

1073

**Induced**
**6+3h BPD vs 6+3h mock**

5 ( 16%)
2 ( 6%)
1 ( 3%)
4 ( 12%)
6 ( 19%)
1 ( 3%)
3 ( 9%)
2 ( 6%)
1 ( 3%)
2 ( 6%)
4 ( 12%)
1 ( 3%)

**N = 32 (4%)**
**p-value ≤ 0.01**

**Reduced**
**6+3h BPD vs 6+3h mock**

8 ( 35%)
1 ( 4%)
3 ( 13%)
1 ( 4%)
4 ( 17%)
5 ( 22%)
1 ( 4%)

**N = 23 (3%)**
**p-value ≤ 0.01**

Amino acid biosynthesis
DNA replication and Repair
Cofactor biosynthesis
Energy Metabolism
Hypothetical
Other

Protein processing and folding
Transcriptional regulation
Translation
Type III Secretion
Nutrient transport
Redox Homeostasis