1   An agent-based model of cattle grazing toxic Geyer's larkspur
2
3   Kevin E. Jablonski[1*], Randall B. Boone[2], Paul J. Meiman[1]
4
5   [1] Department of Forest and Rangeland Stewardship, Colorado State University, Fort Collins,
6   Colorado, United States of America.
7   [2] Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, Colorado, United
8   States of America
9
10   * Corresponding author
11   Email: kevin.jablonski@colostate.edu

Abstract

By killing cattle and otherwise complicating management, the many species of larkspur (*Delphinium* spp.) present a serious, intractable, and complex challenge to livestock grazing management in the western United States. Among the many obstacles to improving our understanding of cattle-larkspur dynamics has been the difficulty of testing different grazing management strategies in the field, as the risk of dead animals is too great. Agent-based models (ABMs) provide an effective method of testing alternate management strategies without risk to livestock. ABMs are especially useful for modeling complex systems such as livestock grazing management, and allow for realistic bottom-up encoding of cattle behavior. Here, we introduce a spatially-explicit, behavior-based ABM of cattle grazing in a pasture with a dangerous amount of Geyer's larkspur (*D. geyeri*). This model tests the role of herd cohesion and stocking density in larkspur intake, finds that both are key drivers of larkspur-induced toxicosis, and indicates that alteration of these factors within realistic bounds can mitigate risk. Crucially, the model points to herd cohesion, which has received little attention in the discipline, as playing an important role in lethal acute toxicosis. As the first ABM to model grazing behavior at realistic scales, this study also demonstrates the tremendous potential of ABMs to illuminate grazing management dynamics, including fundamental aspects of livestock behavior amidst ecological heterogeneity.

Introduction

The many species of larkspur (*Delphinium* spp. L.) present a serious, intractable, and complex challenge to livestock grazing management in the western United States [1–3]. Larkspur plants contain numerous norditerpinoid alkaloids, which are potent neuromuscular paralytics that, for reasons that are not entirely understood, are particularly effective at killing cattle, with yearly herd losses estimated at 2-5% for those grazing in larkspur habitat [3,4]. To avoid such losses, producers

will often abandon or delay grazing in pastures with larkspur, which creates a substantial opportunity

cost and an impediment to achieving management objectives [1,4].

Among the many challenges to improving our understanding of cattle-larkspur dynamics has

been the difficulty of testing different grazing management strategies in the field. Not only is risking

dead cattle impractical and unethical, but the complexity of livestock grazing management, especially

when considered across the wide range of habitats and management regimes in which larkspur is

found, suggests that results from individual field experiments would be unlikely to be broadly useful

anyway [5,6]. What is needed instead is a method of realistically testing grazing management

strategies without risk to livestock and with the flexibility to test multiple scenarios. Agent-based

models (ABMs) provide such a method.

ABMs are computational simulation tools that focus on the behavior of individual "agents"

as they interact with one another and the environment [7]. They differ from other types of

simulation models in being bottom-up (versus top-down) with group-level behaviors emerging from

(usually) realistic individual behaviors rather than deterministic formulae [8]. ABMs are thus

particularly useful in modeling complex systems, where the results of the interactions among system

elements are not easily predicted or understood [9,10]. Indeed, it has been suggested that bottom-

up-simulation may be the best way to increase our understanding of complex systems, which is one

of the most important challenges confronting modern science [9,11,12].

As noted by Dumont and Hill [11], ABMs are "particularly suited to simulate the behavior of

groups of herbivores foraging within a heterogeneous environment". The authors encourage the use

of ABMs in situations where experimentation is impractical, and those where comparison of

different management strategies is needed. Despite this encouragement, and despite the growing

enthusiasm for ABMs in other disciplines, they have been little used in livestock grazing

management research, despite the existence of relevant studies to parameterize such a model [e.g.,

13–18]. This is at least partly due to confusion about the purpose and role of models in improving

our understanding of complex systems.

Models can never be complete simulacra, and do not need to be in order to be useful.

Instead, "models are neither true nor false but lie on a continuum of usefulness for which credibility

can be built up only gradually" [19]. This credibility is built not just by model output but also, more

importantly, through thoughtful model development. This ensures that the necessary simplification

that occurs in modeling focuses in on rather than obscures the system processes of interest [20]. As

noted by Augusiak et al. [19], in well-designed models the important question is the extent to which

the model achieves its purpose in the light of existing evidence, rather than a binary yes or no

regarding its validity.

Previous research into the relationship between grazing management and larkspur toxicosis

has largely focused on timing of grazing, with some attention paid to mineral supplementation, pre-

grazing with sheep, and, increasingly, genetic susceptibility [3,4,21–24]. Some papers have suggested

that cattle behavior, influenced by management, can play a role in mitigating larkspur deaths [25,26],

but these ideas have received little empirical study. Only anecdotally has it been observed that,

regardless of timing of grazing, it may be possible to eliminate losses to larkspur by increasing

stocking density, due to a dilution effect (same amount of alkaloids, more cattle) or perhaps changes

in herd behavior [27].

In this paper, we introduce a spatially-explicit, behavior-based ABM of cattle grazing in a

pasture with a dangerous amount of Geyer's larkspur (*Delphinium geyeri* Green), in which MSAL-type

alkaloids are the dominant toxin [28,29]. This model provides significant management-relevant

insight for producers dealing with larkspur and demonstrates the great potential of ABMs to credibly

model livestock grazing management dynamics, including fundamental aspects of livestock behavior

amidst ecological heterogeneity.

Methods

85   The model description follows the updated Overview, Design Concepts, and Details (ODD)

86   protocol, an accepted method for standardizing published descriptions of ABMs [30].

Purpose

We developed this model to test the effect of co-varying instantaneous stocking density [31]

and herd cohesion (also known as troop length) [32] on cases of lethal acute alkaloid toxicosis

caused by *D. geyeri*. Cases of lethal acute toxicosis are a product of intensity of exposure to alkaloids

(via consumption) with passing time as a mitigating factor (via metabolism). Conceptually, this

model functions as a mechanistic effect model (MEM) aimed at understanding the processes

whereby toxic alkaloids kill grazing cattle. MEMs have been recognized for their potential to "close

the gap between laboratory tests on individuals and ecological systems in real landscapes" [20]. We

developed and executed the model in NetLogo 6.01, using the BehaviorSpace tool to implement

simulations [33].

Basic principles

Behavior-based encoding of cattle activities was the guiding principle of model design. As

noted by Mclane et al. [7], "the behavior-based approach leads to a more complex web of decisions,

and the responses of the animal to stimuli are often more multifaceted". We add that the behavior-

based approach is also more likely to allow for instructive emergent properties. In practice, the

behavior-based approach means that at every step of the coding process we sought literature on

actual cattle behavior and then encoded that behavior as realistically as possible. When literature was

lacking we used our knowledge of cattle behavior from our years as livestock managers and

researchers. The behavior-based approach also found expression in model evaluation, when one

mode of evaluation was whether the cows in the model "act like cows". This was achieved through a

lengthy process of visual debugging and other implementation verification [19,34].

A second core design principle was parsimony. Because this is the first ABM that we know

of to incorporate cattle at the individual scale of interaction with the environment (1 m^2) and

extended to a realistic pasture size, we were initially tempted to include every cattle behavior we

could. However, our focus on parsimony to the question at hand meant that we instead included

only those behaviors relevant to the consumption of larkspur. A final guiding principle was that

when a judgement call was needed, we erred on the side of making the effects of alkaloid toxicosis

more prominent. If the model was to show an effect of grazing management on reducing larkspur-

induced toxicosis, we wanted to be sure that we had taken every precaution against preconditioning

it to do so.

Overall, we followed as closely as possible the process of "evaludation" laid out by Augusiak

et al. [19], which is aimed at moving beyond insufficient and often counterproductive ideas about

model validation to a more thorough process of generating credible models. Specifically, we

incorporated data evaluation, conceptual model evaluation, implementation verification, output

verification, and other analysis of model output.

Entities and state variables

The model has two kinds of entities: pixels representing 1 m^2 patches of land and agents

representing 500 kg adult cows (1.1 animal-units). The patches create a model landscape that is 1663

x 1580 patches (1.66 km x 1.58 km, equal to 262.75 ha, of which 258.82 ha are within the pasture

under study and 3.93 ha are outside the fence line and thus inaccessible). This landscape aims to

replicate pasture 16 at the Colorado State University Research Foundation Maxwell Ranch, a

working cattle ranch in the Laramie Foothills ecoregion of north-central Colorado that is a transition

zone between the Rocky Mountains and the Great Plains. Several pastures on the ranch, including

pasture 16, have significant populations of *D. geyeri*, which generate ongoing management challenges

and have fatally poisoned cattle.

To make the model appropriately spatially explicit we included three sets of geographic data.

First, using data from the Worldview-2 satellite (8-band multispectral, resolution 2 m) from July 10,

2016, we created an index of non-tree/shrub vegetation distribution within the pasture using a soil-

adjusted vegetation index (SAVI) within ERDAS Imagine 2016 software at a resolution of 1 m

[35,36]. Second, as there are no developed watering locations in pasture 16, with ArcGIS Desktop

10.4 we digitized and rasterized (at 1 m) all locations of naturally occurring water as of July 2017

[37].

Lastly, in June and July of 2017 we mapped larkspur distribution and density in pasture 16

using a hybrid approach. We began by digitally dividing the pasture into 272 1-ha sampling plots.

Because we knew larkspur to be of patchy distribution, in each plot we first mapped all larkspur

patches (defined as areas with >1 larkspur plant • m^{-2}) using an iPad equipped with Collector for

ArcGIS 17.01 [38] and a Bad Elf Pro+ Bluetooth GPS receiver accurate to 2.5 m. To sample areas

outside of larkspur patches for larkspur density, we counted all living larkspur plants in a 6-m-wide

belt transect running horizontally across the plot, with the origin randomly assigned and any patches

excluded [39]. Using ArcGIS Desktop we then extended the belt-transect-derived larkspur density to

the rest of the plot (excluding patches), and both sets of data were integrated into a 1 m raster of

larkspur distribution.

The number of cows (individual agents) in the model varies according to the chosen

stocking density (SD, in AU • ha^{-1}). Cows are assigned the role of "leader" (5%), "follower" (85%),

or "independent" (10%) [16,40,41]. Each cow is also assigned a value for MSAL-tolerance and

larkspur-attraction. MSAL-tolerance determines the MSAL-level at which a cow will "die" and is

randomly assigned to create a normal distribution with 99.9% of values falling within 25% of the

mean ($\overline{x}$=4,000 mg, σ=333.33 mg) [42]. In this model, death does not result in the removal of a cow

from the herd; instead, in order to preserve herd and other model functions it is recorded as having

died, its MSAL-level is set to zero, and it continues to graze. Note that MSAL-tolerance can be

understood as modeling genetic, physiological, and situational susceptibility.

Larkspur-attraction determines how much larkspur the individual cow will consume when in

a patch with MSAL-content and is also randomly assigned to create a normal distribution with

99.9% of values falling within 25% of the mean ($\overline{x}$=1.0, σ=0.083). A value of 1.0 means that the

cow will consume larkspur at the same rate as other forage, while values greater or less than 1.0

cause the animal to, respectively, prefer or avoid larkspur. All functionally relevant state variables for

patches and cows, as well as global variables and inputs, are described in Table 1.

Table 1. Relevant model variables.

Entity	Variable	Description
Patches	forage-mass	Amount of currently available forage (g)
	n-forage-mass	Mean initial available forage in patches within a radius of 3 m (g)
	MSAL-content	Amount of toxic alkaloids currently in patch (mg)
	times-grazed	Number of times patch has been grazed
Cows	role	Role in the herd: leader, follower, or independent
	MSAL-level	Current amount of MSAL alkaloids in cow's body (mg); metabolized with a half-life of one grazing-day
	MSAL-tolerance	Level at which cow will be recorded as having died (MSAL-level>MSAL-tolerance); assigned randomly from a normal distribution ($\overline{x}$=4,000 mg, σ=333.33 mg)
	larkspur-attraction	Factor determining the relative amount of larkspur a cow will eat when in a patch with MSAL-content; assigned randomly from a normal distribution ($\overline{x}$=1, σ=0.083)
	herdmates	Agent-set consisting of nearest 20 cows
	mean-herd-distance	Mean distance to herdmates
	total-MSAL-intake	Total amount of MSAL alkaloids consumed during model run (mg)
	daily-MSAL-intake	Amount of MSAL alkaloids consumed during current day (mg)
	hydration	Hydration level, decreases to zero between visits to water
	ready-to-go	Used by leader cows only, a measure of their inclination to move on from an overgrazed site
Globals	waterers	Patch-set of all watering locations

	site-tolerance	Herd-size-dependent variable determining leader cows' tolerance for relatively overgrazed sites
	site-radius	Radius of site when choosing a new site; product of herd-cohesion-factor and herd size resulting in space per cow ranging from 10 m^2 to 1000 m^2
	herd-distance	Desired mean-herd-distance; product of herd-cohesion-factor resulting in range from 10 m to 100 m
Inputs	kgs-per-hectare	Mean amount of usable forage (kg • ha^{-1})
	mean-larkspur-mass	Mean mass of larkspur plants (g)
	MSAL-concentration	MSAL alkaloid concentration in larkspur plants (mg • g^{-1})
	herd-cohesion-factor (HCF)	Determines herd-distance and site-radius; range 1-10, increase leads to more cohesive herd
	stocking-density (SD)	Instantaneous stocking density (AU • ha^{-1})

Scales

The model simulates cow activities at multiple temporal and spatial scales. In each tick (one

cycle through the model code), each cow interacts with a single 1 m^2 patch (a feeding station) by

grazing (>99% of the time) or drinking water (twice per day) [13]. A tick does not represent time,

but rather the occurrence of this interaction. This is because the duration of this interaction will vary

depending on the amount of forage available, among other factors. Instead, time is represented by

consumption of forage. When the average consumption of the grazing herd is equal to the average

daily consumption of a 500 kg cow (12.5 kg), the model counts a grazing-day as having passed [43].

Total model run time is measured in animal-unit-months (AUMs) [44].

The narrowest scale of spatial interaction is the eating interaction occurring within a single

patch (1 m^2). When determining the next patch to graze, the cow's decision is based on a desire

either to move closer to its herdmates or to choose a nearby patch with maximum available forage.

This decision happens on the scale of 2-25 m. Finally, leader cows make decisions on the scale of

the entire pasture by deciding when it is time to visit water or time to move from the current feeding

site to a new site.

Thus, there are four programmed spatial scales (additional scales may be emergent) at which

the cows interact with the landscape: 1) the individual patch; 2) the scale of herd cohesion, set by the

user; 3) the current feeding site; and 4) other feeding sites, identifiable by leader cows. The number

of ticks that will pass before reaching a stopping point (say, 150 AUMs) depends on the number of

animals grazing, their herd cohesion, the amount and distribution of available forage, and stochastic

emergent properties of the model. For an expanded discussion of temporal and spatial scales of

foraging behavior of large herbivores, see Bailey and Provenza [13].

Process overview and scheduling

189   Fig 1 illustrates the model execution process for each tick. Each cow moves through each step of

190   the process, but only performs those steps linked to its role.

191   Fig 1. Pseudo-coded flow chart of model processes, with role of cows executing each process in parentheses. 1= leader,
192   2=follower, 3= independent.

Check hydration

194   Each leader cow checks it hydration level, which is tied to forage consumption such that it

195   depletes to zero twice per day. We chose two water visits per day based on personal communication

196   about GPS collar data for the region [D. Augustine, USDA ARS, pers. communication; see 45]. If an

197   individual leader detects its hydration level as less than or equal to zero, it initiates a movement to

198   water for the whole herd.

Go to water

The water source in pasture 16 is a stream that is intermittently below ground. The go-to-

water procedure directs each cow to go to the nearest waterer patch with two or fewer cows already

present. The hydration value for each cow is then set to maximum, and the value for ready-to-go for

leader cows is set to site-tolerance – 1. This reflects the understanding that cattle will quickly graze

and trample areas around water, rendering them unsuitable for grazing. Instead, they will pick

desirable foraging areas in proximity to but not directly surrounding a watering site, expanding

outward as these areas are grazed [13]. The model thus encourages a site change upon drinking

water, but only if the area surrounding the watering site meets the criteria for increasing ready-to-go

(explained below). A global variable ensures that no other processes occur during a tick when

watering occurs.

Check site change

This process is only executed by leader cows, each of which assesses the mean number of

times patches within a radius of 10 m have been grazed. If these patches have been grazed relatively

more (defined as >0.5 • mean times-grazed of all patches + 1.2) than the pasture as a whole, the

value of ready-to-go increases by one. If this value reaches a pre-defined threshold (which increases

with herd size), the individual then initiates a site change, but only if the individual's hydration value

is not approaching zero, in which case it instead initiates the go-to-water procedure. We arrived at

the threshold formula for increasing the value of ready-to-go by using visual debugging and

evaluation related to site change frequency, as well as theory on the optimization of grazing effort

[13,46].

If conditions for a site change are satisfied, the deciding leader cow first identifies the best

five available sites, using criteria of number of times-grazed, forage-mass, and n-forage-mass to

determine a centroid patch. The nearest of these patches is then used to create a new site at a radius

that is linked to the user selected herd-cohesion-factor and the size of the herd, resulting in 10-1,000

$m^2 \cdot cow^{-1}$ in the new site. The leader cow then initiates the change-site procedure for itself and all

other cows.

Change site

227   This procedure is initiated according to role, so that leader cows have first choice of their

228   location in the new site, followers second, and independents third. Within the allocated new site,

229   each cow chooses the patch with the most forage that has no cows on it or any of its four direct

230   neighbors.

Assess herd

In combination with the environmental-movement procedure, this process represents >99%

of cow actions in the model. Each cow first sets its herdmates as the nearest 20 other cows [47]. For

leader and follower cows, if the individual's mean distance to these herdmates is greater than herd-

distance, it "herds up". This is achieved by facing the centroid of the herdmates and moving to the

patch with maximum available forage that is 10-25 m in the direction of this centroid, within a cone

of vision of ±45 degrees [14]. For independent cows, the same process occurs but is only initiated if

the distance from herdmates is greater than 2.5 times the herd-distance of the other cows.

Independent cows are also repelled from the center of their herdmates by moving away by the same

procedure when they are within one-half of the herd-distance.

Environmental movement

If none of the above procedures are implemented, each cow will make a movement decision

based on local grazing conditions. If the patches within a radius of 10 m are relatively ungrazed

(mean times-grazed < 0.5) the cow will move to the patch with the most available forage within 2 m,

within a ±45 degree cone of vision [13]. If the same area is relatively well grazed (mean times-grazed

≥ 0.5), the cow then looks further afield, choosing the patch with the most available forage within

10 m, within a cone of vision of ±45 degrees.

Eat

The eat procedure is the core interaction between the cows and the forage, both non-

larkspur and larkspur. Behavior varies slightly depending on how many times the patch has

previously been grazed. If the current visit is the first time it has been grazed, the cow eats 40% of

the available forage [15,18]. If it is the second visit, it eats 50% of what remains. In the third and any

subsequent visits, it eats 60%. Each cow then increases its consumption-level by the same amount

and decreases its hydration value. If there is larkspur present (in the form of MSAL-content), that is

consumed according to the individual cow's larkspur-attraction value, increasing the MSAL-level of

the cow. The corresponding patch values are decreased to account for consumption. Lastly, times-

grazed in the patch is increased by one.

Assess toxicosis

This process is triggered at the end of each grazing-day for all cows in order to assess their

toxicosis status, which is measured as their MSAL-level relative to their MSAL-tolerance. Note that

MSAL-level is measured continuously throughout the model run, and has an elimination half-life of

one grazing-day [48]. If MSAL-level exceeds MSAL-tolerance, the count of deaths for the model run

is increased by one, MSAL-level is set to zero, and the cow continues. Numerous other data on

toxicosis status are recorded for all cows at this point. Lastly, the MSAL-level for each cow is

multiplied by 0.5.

Design concepts

Emergence

Because the actions of the cows are encoded via simple behavior-based processes, nearly all

model patterns can be considered emergent. These include the stochastic distribution of the herd

and subherds, forage consumption, larkspur consumption, grazing pressure and patterns, and site

changes. Assessment of these un-coded emergent properties and patterns was critical to establishing

the credibility of the model [20].

Adaptation, objectives, learning, and predictions

The cows adapt to the grazing environment as they and their fellow cows graze, continually

seeking their main model objective of maximizing forage consumption within behavioral limits [14].

There is no encoded learning or prediction, as the cows are programmed to be familiar with the

location of forage and water in the pasture. However, it may be that learning and prediction are

emergent, in that activities that we might consider to be evidence of those behaviors are visible in

the model as a result of the simple encoded behaviors.

Sensing and interaction

The cows sense each other and their environment at multiple spatial scales. Interaction

occurs with other cows whenever moving to a new patch, both via sensing if a patch is already

occupied and by seeking to herd up when too far from their herdmates.

Stochasticity

There is no environmental stochasticity in this model iteration, as we sought to make the

landscape as realistic as possible by incorporating relevant data from the real pasture 16. However,

cattle interactions with the forage and larkspur demonstrate moderate stochasticity.

Initialization

Landscape initialization begins by loading the SAVI layer and a user-input value for available

forage per ha (kgs-per-hectare). The model uses a nonlinear exponential formula to distribute forage

such that the patches with the least forage contain one-third of the mean forage, while the patches

with the most contain three times the mean forage. Next, the model incorporates the larkspur

distribution layer, using inputs of median larkspur mass (g) and mean MSAL concentration (mg • g^{-1})

to generate an MSAL alkaloid (hereafter simply "alkaloid") content for each patch. These values are

based on our unpublished data on *D. geyeri* mass and toxicity at the Maxwell Ranch such that

larkspur plants in areas of high SAVI were 50% larger than the median, and larkspur plants in areas

of low SAVI were 50% smaller than the median. Finally, the model incorporates the water location

layer. All other patch variables are derived from these inputs. Fig 2 shows the initialized landscape.

Fig 2. Model landscape, 1.66 km x 1.58 km. (a) Initialized full model landscape, with darker green indicating areas with
greater aboveground forage biomass. (b) Landscape with larkspur locations only, with darker purple representing higher
MSAL-content and with results of hybrid sampling method evident. (c) Landscape with watering locations only, pointed
out by arrows.

The final step in model initialization is to create the cows by using the input of stocking-

density multiplied by the area of the pasture. All cows are initially in the same random location in the

pasture. This location is largely irrelevant as the cows immediately go to water, but we did not want

it to be the same location each time because this would be unrealistic (pasture 16 has multiple

entrances for cattle) and would limit stochasticity. At this point, the model is fully initialized and is

executed following the processes laid out above.

Simulation

We used the BehaviorSpace tool in NetLogo to run a full factorial simulation of four

different levels of both herd-cohesion-factor (1, 4, 7, and 10) and stocking-density (0.25, 0.5, 1.0,

and 2.0 AU • ha^{-1}). We replicated each combination 30 times, for a total of 480 simulations. Input

median larkspur mass was 3.5 g and input MSAL alkaloid concentration was 3.0 mg • g^{-1}. We chose

these values to be representative of an excellent growing year with larkspur plants at bud stage, when

the alkaloid pool (total available mg) is highest—arguably the most dangerous possible conditions.

This is also a time of year that cattle grazing in larkspur habitat is frequently avoided, despite being a

highly desirable time for grazing [1,4,49]. Input value for kgs-per-hectare was 500 kg, based on

current ranch usage and typical values for the area.

Observation

Of primary importance were data related to alkaloid consumption, assessed according to

dose-response data from previous research [42]. Most interesting was the number of times in a

model run that any individual cow crossed the threshold into potentially lethal acute toxicosis,

during which they would be expected to be recumbent and unable to stand, with a high likelihood of

death [42]. To measure the number of such cases, the model counted cows whose MSAL-level

exceeded their MSAL-tolerance at the end of a grazing-day.

The model also recorded data underlying the trends found for lethal acute toxicosis, most

importantly data on daily, total, and maximum alkaloid intake. These data assisted in identifying

potential mechanisms for the role of herd cohesion and stocking density in influencing deaths.

Additional data, such as forage consumption, number of site changes, travel distance per day, and

evenness of grazing impact, provide additional insight and model output verification.

Statistical analysis

We used both JMP Pro 13.0.0 and R statistical software, version 3.3.3 for data analysis and

presentation [50,51]. Data for daily alkaloid intake, which amounted to 1.88 million data points, were

organized and cleaned using OpenRefine 2.8 [52]. We began by assessing the role and relative

influence of HCF and SD in generating lethal acute toxicosis, within two contexts: first, using their

16 combinations as "management levels" to explore overall trends in a management-relevant

manner; and, second, using HCF and SD as continuous variables within a regression framework to

provide more information on the relative influence of each. To regress the lethal acute toxicosis

count data we used a generalized linear model (GLM) with a negative binomial distribution and a

log-link function using the MASS package in R [53,54]. To confirm that the negative binomial

distribution was the correct choice, we compared it to a GLM with a Poisson distribution and a log-

link function. The GLM with the negative binomial distribution was superior, using residual

deviance and Aikaike's information criterion (AIC) as judgment criteria [55].

To identify mechanisms for how HCF and SD were influencing deaths, we used the same

negative binomial GLM approach to analyze the relationship between various intake data and lethal

acute toxicosis. We did so by first hypothesizing which factors were driving deaths, and then looked

at single-factor models for each, assessed using AIC values and model coefficients [55]. Because the

goal was to identify key mechanisms rather than determine the best predictive model, this provided

more insight than examining a global model or various permutations of factors.

Finally, we analyzed the relationship of HFC and SD to the identified mechanisms using

multiple linear regression (R base package). While there were some indications of heteroscedasticity

and outliers, we determined that linear regression was robust to those errors in these cases. We

confirmed this by also fitting alternate models within other regression frameworks (robust and non-

parametric), which returned very similar results. Interaction effects are shown when significant;

otherwise, they were excluded from the models.

Results

Model output verification

A core element in the evaluation of behavior-based mechanistic effect models is a

comparison between multiple emergent model patterns and observed patterns in the real system

[20]. In this case, this helped to establish that the modeled cows, coded for individual behaviors,

acted like real cows when interacting with one another and the landscape, at least in regard to

behaviors relevant to larkspur consumption. Toward this end, first we offer Fig 3 to illustrate how

varying HCF influences herding patterns, and to show how grazing was distributed across the

pasture in one model run.

Fig 3. The effect of varying herd-cohesion-factor (HCF) on herd patterns, displayed at different levels of pasture usage
(AUMs). Note that the cows depicted in these images are drawn 200 times larger than they really are to aid visualization,
which makes them appear closer to one another than they are. Pasture size is 1.66 km x 1.58 km, and stocking density
for all images is 1.0 AU • ha^{-1}. White cows are leaders, black followers, and gray independents. Yellow indicates patches
that have been grazed twice, red three times. (a) HCF=10, AUMs=14; (b) HCF=7, AUMs=68; (c) HCF=4, AUMs=119;
(d) HCF=1, AUMs=163. Typical usage for this pasture (258.82 ha) is 150 AUMs.

Decreasing HCF increases overall herd separation and leads to more wandering among the

independent cows and others. Note that in Fig 3a the cows have formed distinct subherds. This

appears to be an emergent property of cows grazing with high herd cohesion (herd-distance ≤ 20

m).

The cows initially graze the areas with high forage amounts (dark green) in relative proximity

to the water, and gradually extend their impact outward, targeting high productivity areas. By the end

of the grazing cycle (Fig 3d), they have visited the entire pasture, though areas furthest from water

have been grazed less [56]. Areas of initial high forage mass have been grazed two or more times,

while many areas of low forage mass have not been grazed at all. These results are in line with well-

established qualitative understanding of grazing patterns in large pastures [13,44].

The variation in forage consumption among individuals also aligned well with the variation

seen in real cows foraging native pasture. While a grazing-day for the whole herd was defined as

mean consumption of 2.5% of body weight (12.5 kg), the mean 99.9% daily range of consumption

for all model runs was 2.34-2.66% of body weight. This range of consumption aligns well with

common "rules of thumb" and predictive formulae [43,57,58].

The mean value for site changes per day for the 16 management levels varies from 2.3 for

few cows grazing very loosely (HCF=1, SD=0.25) to 6.0 for many cows grazing very cohesively

(HCF=10, SD=2.0). These values are in line with the estimate of 1-4 hours per feeding site by Bailey

and Provenza [13]. For runs with few cows grazing with little cohesion (HCF=1, SD=0.25), mean

daily travel was 4.16 km, while many cows grazing very cohesively (HCF=10, SD=2.0) traveled an

average of 7.40 km per day. These numbers and the positive trend also track well with data from

previous studies [59].

As a last point of output verification, we were interested to see if the number of modeled

cases of larkspur-induced lethal acute toxicosis would parallel numbers from the literature when we

modeled grazing to be similar to the current management scheme. When modeled to reflect current

management practices, with HCF=4, SD=0.5, and for 150 AUMs (removing approximately 45% of

available forage), we recorded a mean of 2.8 cases of lethal acute toxicosis across 30 model

iterations. This amounts to 2.4% of cows, which falls within the estimate of 2-5% in pastures with

dangerous amounts of larkspur [4]. Additionally, individual model runs of zero deaths occurred in all

but four of the management levels, which aligns with our anecdotal understanding of producer

experience.

Lethal acute toxicosis

On its own, increased herd cohesion demonstrated the potential to significantly reduce

deaths. For example, at a stocking density of 0.5 AU • ha^{-1}, mean deaths declined from 4.33 at

HCF=1 to 1.37 at HCF=10. Similarly, increased stocking density in the absence of changes in herd

cohesion also greatly reduced deaths, for example from a mean of 7.5 at SD=0.25 to 0.70 at SD=2

at a constant HCF of 4. Working together, increases in both herd cohesion and stocking density

from the minimum to the maximum achieved a 99.6% reduction in deaths (Fig 4). The mean value

for MSAL-tolerance among dead cows was 3,725.8 mg, while the mean value for larkspur-attraction

was a factor of 1.06. Of 1,132 total deaths in the simulation, 3.9% were among cows with the role of

leader, 78.7% were among followers, and 17.4% were among independents.

Fig 4. Box plots of distribution of counts of lethal acute toxicosis cases (MSAL-level ≥ MSAL-tolerance at end of
grazing-day). From 30 model runs for each combination of herd-cohesion factor (HCF) and stocking-density (SD),
ordered by median count of lethal acute toxicosis cases, with outliers as jittered circles.
415

The coefficient for HCF (Table 2), as a log odds ratio, indicates that an increase of one in

HCF resulted in a 13.5% decrease in occurrences of lethal acute toxicosis. The coefficient for SD

indicates that an increase of one in SD resulted in a 54.8% decrease. Lastly, the coefficient for the

interaction of HCF with SD indicates that an increase in either HCF or SD slightly increases the

effect of the other. The GLM β coefficients indicate that HCF had 91.8% of the influence of SD in

reducing deaths.

Table 2. Results of GLM with negative binomial distribution and log-link function for count of lethal acute toxicosis as
predicted by herd-cohesion-factor (HCF) and stocking-density (SD). β coefficients are from the same GLM without the
interaction present. GLM fit: Fisher scoring iterations=1; residual deviance=516.94 on 476 degrees of freedom;
AIC=1686.3.

Coefficient	Estimate	Std. error	p-value	β
Intercept	2.341	0.128	<0.001	
HCF	-0.145	0.024	<0.001	-0.225
SD	-0.793	0.136	<0.001	-0.245
HCF:SD	-0.079	0.029	0.007	

Identifying mechanisms

We hypothesized that five factors might explain how HCF and SD were reducing deaths:

mean individual daily alkaloid intake (the average single-day alkaloid intake in a model run), standard

deviation of individual daily alkaloid intake, mean maximum individual daily alkaloid intake (each

cow's worst day), standard deviation of maximum individual daily alkaloid intake, and the coefficient

of variation for individual total alkaloid intake. Results for the comparison of single-factor models

reveal varying influence on lethal acute toxicosis among these factors (Table 3).

Table 3. Results for comparison of single-factor negative binomial generalized linear models with a log-link function
using corrected Aikaike's information criterion. All values for quartiles are in mg, except for CV total, which is unitless.
Percent Δ deaths from Q_1 to Q_3 is observed percent change in lethal acute toxicosis count between quartile one and
three.

Mechanism	AICc	GLM coefficient	Quartile 1	Quartile 3	Pct. Δ deaths from Q_1 to Q_3
σ Maximum	1473.3	0.0072	410.3	591.2	130.70%
σ Daily	1510.1	0.0265	363.8	435.4	192.36%
Mean maximum	1671.2	0.0019	1275.0	1987.7	135.55%
Mean daily	1911.4	-0.0374	527.9	543.0	-55.58%
CV total	1930.2	-7.044	0.118	0.185	-6.69%

440 Because they had the most significant effect on lethal acute toxicosis, and were scored lowest for

441 AICc, we focused the rest of the analysis on examining the relationship of HCF and SD to standard

442 deviation of maximum individual daily alkaloid consumption, standard deviation of individual daily

443 alkaloid consumption, and mean maximum individual daily alkaloid consumption. A model for lethal

444 acute toxicosis count that contained these three mechanisms had an AICc score of 1368.3.

Daily alkaloid intake

Mean individual daily alkaloid intake represents the mean of every single-day alkaloid intake

for every cow, and ranged from a low of 525.1 mg (HCF=4, SD=0.25) to a high of 550.9 mg

(HCF=10, SD=0.25). Multiple linear regression results indicate that HCF and SD had limited

influence on mean daily intake (adj. R^2<0.19), with both associated with slight increases. On the

other hand, the standard deviation of daily alkaloid intake, which quantifies the spread of the

distribution of daily alkaloid intake values, differed significantly between management levels, from a

high mean of 460.5 mg (HCF=1, SD=0.25) to a low mean of 301.3 mg (HCF=10, SD=2). Multiple

linear regression results indicate that HCF and SD were strongly influential, with HCF exerting

93.0% more influence than SD (Table 4).

Table 4. Results of multiple linear regression for the standard deviation of individual daily alkaloid intake as predicted by
herd-cohesion-factor (HCF) and stocking-density (SD). Adj. R^2=0.76.

Coefficient	Estimate	Std. error	p-value	β
Intercept	487.79	2.61	<0.001	
HCF	-11.33	0.33	<0.001	-0.774
SD	-29.38	1.64	<0.001	-0.401

458   A box plot showing the distribution of all individual daily alkaloid intake values (n=1.88 • 10^5) at

459   each management level further illustrates these patterns (Fig 5).

460   Fig 5. Box plots of distribution of individual daily alkaloid intake (mg; n=1.88 • 10-5). From 30 model runs for each
461   combination of herd-cohesion factor (HCF) and stocking-density (SD), ordered by median standard deviation of daily
462   alkaloid intake, with outliers as jittered circles.

Maximum daily alkaloid intake

Mean maximum individual daily alkaloid intake quantifies the mean worst day for all cows

during a model run, and ranged from 1,045.6 mg (HCF=10, SD=2) to 2,450.2 mg (HCF=1,

SD=0.25). The standard deviation of maximum individual daily alkaloid intake quantifies how widely

dispersed this value was among the herd members, and ranged from 303.0 mg (HCF=10, SD=2) to

704.0 mg (HCF=4, SD=0.25). Regression results for both factors provide further insight into the

relationship of HCF and SD to lethal acute toxicosis (Tables 5-6).

Table 5. Results of multiple linear regression for the mean of maximum individual daily alkaloid intake as predicted by
herd-cohesion-factor (HCF) and stocking-density (SD). Adj. R^2=0.82. β coefficients are from the same model without
the interaction present.

Coefficient	Estimate	Std. error	p-value	β
Intercept	2547.56	28.58	<0.001	
HCF	-61.86	4.44	<0.001	-0.31
SD	-686.15	24.80	<0.001	-0.84
HCF:SD	22.75	3.85	<0.001	

475   Table 6. Results of multiple linear regression for the standard deviation of maximum individual daily alkaloid intake as
476   predicted by herd-cohesion-factor (HCF) and stocking-density (SD). Adj. R^2=0.47. No significant interaction was
477   present.

Coefficient	Estimate	Std. error	p-value	β
Intercept	718.57	11.33	<0.001	
HCF	-22.34	1.42	<0.001	-0.52
SD	-96.06	7.12	<0.001	-0.45

Distinct persistent subherds

Model outputs (Fig 6) suggested an apparent scalar behavioral discontinuity between

HCF=7 and HCF=10, which we believe results from the emergent property of distinct persistent

subherds.

Fig 6. Box plots of various model evaluation data demonstrating effect of distinct persistent subherds. (a) Mean
individual travel distance per grazing day (m) by herd cohesion factor (HCF); (b) Proportion of use of assess herd
procedure (versus environmental movement) to choose a new grazing patch, a measure of herd-based versus individual
optimization, by HCF; (c) Standard deviation of times-grazed count for all patches at end of model run, a measure of
grazing heterogeneity, by HCF; (d) Total model run length, an inverse indicator of grazing efficiency, by HCF at
stocking density=0.5.
489

Discussion

Research into best practices for grazing management in larkspur habitat has long focused on

either attempts to eliminate larkspur or on phenological avoidance (what we term "fight or flight").

Because elimination through herbicides or mowing is costly and often impractical [60], most

research and current recommendations focus on avoiding grazing in larkspur habitat at times of year

when it is considered most dangerous to cattle, exemplified by the toxic window concept [3,4,23].

While this approach has certainly helped many producers better understand larkspur toxicity

dynamics, there is no evidence that it has reduced the overall number of deaths. There are many

reasons for this, and interactions are complex and place-based, but we suggest that a reliance on a

static view of palatability is largely to blame.

An alternative to fight or flight is to manage grazing such that larkspur intake remains below

the threshold where there is an observable negative effect on the cattle. This study provides an

indication that this may be possible even in pastures with dangerous amounts of Geyer's larkspur.

For the first time, this model suggests that herd cohesion and stocking density are key drivers of

larkspur-induced toxicosis, and that management decisions that influence these factors hold

potential to limit deaths. Of crucial importance is the observation that herd cohesion, which has

received almost no consideration in the broader grazing management literature, is an important

determinant of risk of death from larkspur.

An essential point for understanding how increased herd cohesion and stocking density

reduced deaths is that Geyer's larkspur grows most densely in relatively productive areas, which are

thus desirable areas for foraging. Functionally, increased herd cohesion and stocking density lead to

increased competition for forage, making it more difficult for any individual to monopolize a

resource- and larkspur-rich area. Additionally, increased herd cohesion leads to less wandering

among individuals, making it less likely an individual cow will wander into a dense larkspur patch

alone. Evidence for the danger of wandering behavior is found in the disproportionate death rate of

cows with the role of independent. Lastly, increased stocking density does appear to lead to dilution,

but in the form of lowered maximum individual daily intake rather than lowered mean individual

daily intake.

Mechanistically, decreased risk of lethal acute toxicosis occurred through: 1) a narrowed

distribution of individual daily alkaloid intake, 2) lowered mean and narrowed distribution of outlier

alkaloid intake days. Herd cohesion played a stronger role in narrowing the distribution of daily

intake, stocking density was more influential in lowering the mean of outlier intake, and both played

a relatively equal role in narrowing the distribution of outlier intake events. Strong evidence for the

role of these as mechanisms is provided by the much lower AICc score for the model with the

mechanisms than for the model with HCF and SD (1386.3 vs. 1686.3). This suggests that other

management interventions that succeed in influencing these mechanisms would have similar success

in reducing deaths.

When we recognize that even in the worst-case scenario lethal acute toxicosis is a rare event

among thousands of grazing-days, it becomes clear why narrowing the distribution of individual

intake and reducing outliers is so important. With a mean MSAL-tolerance of 4,000 mg, an average

bad day in a herd with low herd cohesion and low stocking density would put an individual

(especially one with lower tolerance or higher attraction to larkspur) in danger. Meanwhile,

individuals grazing in a herd with high herd cohesion and at a high stocking density in the same

pasture, even those with low tolerance, would need at least a few upper-end intake days in a row to

risk death—an unlikely occurrence.

Note that we selected the bounds of herd cohesion and stocking density to align with what

we believe to be realistically achievable by managers in the western US. While stocking density is

easily understood, it may be worthwhile to describe how we think the various levels of herd-

cohesion-factor (HCF) could be achieved (reference Fig 3). We think of HCF values of 1 and 4 as

representative of most current extensive management, such that there is a small to moderate amount

of herding behavior but in which animals are often spread out across a large area. The difference

between these two might be accounted for by differences in breeding history, carnivore pressure, or

genetic drift. To achieve an HCF of 7, we think cattle would need to be selected for strong herding

instinct or be regularly, but not necessarily continually, herded. An HCF of 10 is comparable to

many herds of wild ungulates and is achievable through the continual presence of a herder or a

sustained effort at selecting for herding behavior.

There are two additional ways that a rapid increase in herd cohesion may be achieved. First,

a drastic increase in stocking density (via increased animal-units or subdivided pastures) to a level

that approaches "mob" grazing can forcibly increase cohesion. Second, the emerging technology of

virtual fencing holds tremendous promise for achieving rapid changes in grazing behavior, including

herd cohesion [61].

An unexpected emergent phenomenon occurred at HCF=1, in the form of distinct

persistent subherds (see Fig 3a). These subherds are small groups of >20 but usually <35 cows that

stick closely together for an entire inter-watering period, with some exchange of individuals or

combining when two groups meet. This does not occur at higher levels of HCF. Cows in distinct

persistent subherds traveled significantly greater distances, spent more time seeking to be closer to

herdmates rather than maximizing forage intake, and grazed more heterogeneously (Figs 7a-c).

Nevertheless, these cows reached 150 AUMs of forage consumption in 94.3% of the model run

time of cows at lower herd cohesion levels, suggesting higher grazing efficiency (Fig 7d). We believe

that these data are evidence of a scale-dependent behavioral discontinuity that may hold relevance to

other grazing management challenges [62].

Model parsimony and study limitations

Perhaps the most obvious omissions from the model are those behaviors that we determined

to hold little to no relevance to larkspur consumption, at least in this pasture. These include

response to slope, resting, and some inconsistently understood aspects of dominance behaviors.

While there is nothing preventing them from being included, we decided that in this case these

behaviors would introduce uncertainty while adding little realism to cattle-larkspur dynamics. The

model also excludes plant regrowth. For Geyer's larkspur, this is not an issue, as plants that are

clipped or grazed during the bud stage exhibit very little regrowth [K. Jablonski, pers. obs.]. For

other forage, we determined that regrowth in July in this semi-arid climate would not be substantial

enough within a single grazing period to warrant inclusion.

The occurrence and measurement of death in the model might strike some as unrealistic.

However, given that deaths in a herd would change herd behavior in unknown ways, and that the

owner of the cattle would likely intervene once one death-event had occurred, we believe that

counting the death and resetting the cow's MSAL-level is the most accurate way to assess risk at

different management levels.

Another potential limitation concerns the model used for alkaloid metabolism. While there

has been some effort at the generation of such a model [e.g., 48], these efforts have been limited to

highly controlled settings using hay and other stored feeds and periodic dosing with alkaloids.

Additionally, little to nothing is known about the role of other forage in exacerbating or mitigating

the effects of larkspur consumption. As such, we had no confidence that a continuous metabolic

model would be more useful than the simple daily half-life model that we used.

Despite these limitations, we are confident that we have realistically modeled cattle-larkspur

dynamics, that increased herd cohesion and stocking density lower the risk of lethal acute toxicosis,

and that variations in mean and maximum daily alkaloid intake are the predominant mechanism for

this reduction. However, the exact values for when risk approaches zero may be dependent on the

circumstances of this model iteration—that of *D. geyeri*, at the input values for mass and toxicity, on

a ranch in northern Colorado.

It is worth noting that dangerous levels of *D. geyeri* are typically found on a limited number

of a single operation's grazing units. This means that the inclusion of herding to increase herd

cohesion, for example, would usually only be necessary for a relatively brief period. In addition, it

means that any potential secondary effects of sub-lethal larkspur consumption, such as appetite

suppression or lethargy (whether and how these would occur is unclear), would be of similarly

limited duration. Nevertheless, in pastures with a dangerous amount of larkspur, negative sub-lethal

effects may be unavoidable even (or especially) when death is avoided.

As with any research where cattle lives and producer livelihoods are at stake, it is most

important to emphasize that producers should exercise caution when incorporating our findings into

their own management, including careful assessment of other potential effects of increased herd

cohesion or stocking density. Those with low amounts of Geyer's larkspur or with no history of

losses might find comfort in altering their grazing management to incorporate this study's findings.

Those with a great deal of larkspur (Geyer's or other species) or a history of losses should be more

careful.

Other model implications and future directions

There is a broad literature on the effect of stocking rate/stocking density on many outcomes

(though not larkspur-induced toxicosis) but very little on the effects of herd cohesion, nor on the

interaction of these factors [44]. This is likely due to the relative ease of varying cattle numbers

versus manipulating cattle behavior. Because this study provides evidence that it is not only the

number of animals but also how they behave that affect the likelihood of death by larkspur, we are

excited to explore the role of herd cohesion, particularly the emergent property of distinct persistent

subherds, in other aspects of grazing ecology. If herd cohesion is genetically encoded, matrilineally-

oriented, or management-determined (or a combination thereof), what role might it play in other

negative outcomes, such as overgrazing of riparian areas or exposure to predation by carnivores [63],

and how might we influence it in different scenarios? The evolving promise of affordable GPS tags

means that we may also start to be able to test this through direct observation of entire herds [64].

For cattle-larkspur dynamics, our next step is to place these modeling results in context with

ongoing plant experiments and producer surveys to better formulate management recommendations

that work. Additionally, we would like to improve our understanding of alkaloid metabolism and

tolerance, as well as the role of preference in larkspur intake. For alkaloid metabolism and tolerance,

this means building upon previous studies [e.g., 21], which have been undertaken in highly

controlled settings using periodic high dosing, to model the stochastic dosing in a dynamic

environment that occurs in reality. For larkspur preference, this means moving beyond the entirely

anecdotal evidence of bouts of larkspur consumption [e.g., 65] to a more sophisticated

understanding of the role of preference, diet mixing, and satiation in larkspur-induced toxicosis

[66,67].

A final next step for the model presented here is what Augusiak et al. [19] term model

output corroboration, wherein model outputs are compared to new, independent data and patterns.

As noted above, this is very difficult when cattle lives are at risk. However, the results presented here

have encourage us to start to think about how such corroborative data could be collected. This will

likely entail a combination of full-herd GPS with careful on-the-ground monitoring by a herder.

Though ABMs have some limitations, we believe they offer an exciting new tool for

understanding the grazing behavior of livestock. Indeed, the synergistic emergence of financially

viable GPS technology [64] and "virtual fencing" [61], along with the increasing power of desktop

computers, suggests that the time is right for a computational revolution in livestock grazing

management. We are excited that this study provides a first example of the potential of agent-based

models to contribute to this revolution.

Acknowledgements

We thank Joel Vaad, manager of the Maxwell Ranch, for access, assistance, and advice. Early

conversations with Michael A. Smith and the Sims family of McFadden, WY, provided crucial

insight into the relationship between grazing management and larkspur. Tanner Marshall assisted in

data collection and asked difficult questions that moved the work forward. We are grateful to

DigitalGlobe for a generous imagery grant that greatly improved our ability to assess the grazing

landscape, and to Hexagon Geospatial for providing the software to analyze the imagery.

References

1. Green B, Gardner D, Pfister J, Cook D. Larkspur poison weed: 100 years of delphinium research. Rangelands. 2009;31: 22–27. doi:10.2111/1551-501X-31.1.22

2. Pfister JA, Gardner DR, Panter KE, Manners GD, Ralphs MH, Stegelmeier BL, et al. Larkspur (Delphinium spp.) poisoning in livestock. J Nat Toxins. 1999;8: 81–94.

3. Welch KD, Cook D, Green BT, Gardner DR, Pfister JA, McDaneld TG, et al. Adverse effects of larkspur (Delphinium spp.) on cattle. Agriculture. 2015;5: 456–474. doi:10.3390/agriculture5030456

4. Pfister JA, Gardner DR, Price KW. Grazing risk on tall larkspur-infested ranges. Rangelands. 1997;19: 12–15. doi:10.2307/4001275

5. Darnhofer I, Gibbon D, Dedieu B. Farming Systems Research: an approach to inquiry. In: Darnhofer I, Gibbon D, Dedieu B, editors. Farming Systems Research into the 21st Century: The New Dynamic. Dordrecht: Springer Netherlands; 2012. pp. 3–31. Available: http://www.springerlink.com/index/10.1007/978-94-007-4503-2_1

6. Provenza F, Pringle H, Revell D, Bray N, Hines C, Teague R, et al. Complex Creative Systems. Rangelands. 2013;35: 6–13. doi:10.2111/RANGELANDS-D-13-00013.1

7. McLane AJ, Semeniuk C, McDermid GJ, Marceau DJ. The role of agent-based models in wildlife ecology and management. Ecol Model. 2011;222: 1544–1556. doi:10.1016/j.ecolmodel.2011.01.020

8. Grimm V. Ten years of individual-based modelling in ecology: what have we learned and what could we learn in the future? Ecol Model. 1999;115: 129–148. doi:10.1016/S0304-3800(98)00188-4

9. Grimm V, Revilla E, Berger U, Jeltsch F, Mooij WM, Railsback SF, et al. Pattern-Oriented Modeling of Agent-Based Complex Systems: Lessons from Ecology. Science. 2005;310: 987–991. doi:10.1126/science.1116681

10. Northrop RB. Introduction to Complexity and Complex Systems. CRC Press; 2010.

11. Dumont B, Hill DRC. Spatially explicit models of group foraging by herbivores: what can Agent-Based Models offer? Anim Res. 2004;53: 419–428. doi:10.1051/animres:2004028

12. Funtowicz SO, Ravetz JR. Science for the post-normal age. Futures. 1993;25: 739–755. doi:10.1016/0016-3287(93)90022-L

13. Bailey DW, Provenza FD. Mechanisms Determining Large-Herbivore Distribution. In: Prins HHT, Langevelde FV, editors. Resource Ecology. Springer Netherlands; 2008. pp. 7–28. doi:10.1007/978-1-4020-6850-8_2

675   14.   Distel RA, Laca EA, Griggs TC, Demment MW. Patch selection by cattle: maximization of
676   intake rate in horizontally heterogeneous pastures. Appl Anim Behav Sci. 1995;45: 11–21.
677   doi:10.1016/0168-1591(95)00593-H

678   15.   Laca EA, Distel RA, Criggs TC, Demment MW. Effects of Canopy Structure on Patch
679   Depression by Grazers. Ecology. 1994;75: 706–716. doi:10.2307/1941728

680   16.   Sato S. Leadership during actual grazing in a small herd of cattle. Appl Anim Ethol. 1982;8:
681   53–65. doi:10.1016/0304-3762(82)90132-8

682   17.   Shiyomi M. How are distances between grazing cows determined: A case study. Appl Entomol
683   Zool. 2004;39: 575–581. doi:10.1303/aez.2004.575

684   18.   WallisDeVries MF, Laca EA, Demment MW. From feeding station to patch: scaling up food
685   intake measurements in grazing cattle. Appl Anim Behav Sci. 1998;60: 301–315.
686   doi:10.1016/S0168-1591(98)00158-0

687   19.   Augusiak J, Van den Brink PJ, Grimm V. Merging validation and evaluation of ecological
688   models to 'evaludation': A review of terminology and a practical approach. Ecol Model.
689   2014;280: 117–128. doi:10.1016/j.ecolmodel.2013.11.009

690   20.   Grimm V, Martin BT. Mechanistic effect modeling for ecological risk assessment: Where to go
691   from here? Integr Environ Assess Manag. 2013;9: e58–e63. doi:10.1002/ieam.1423

692   21.   Green BT, Welch KD, Pfister JA, Chitko-McKown CG, Gardner DR, Panter KE. Mitigation
693   of Larkspur Poisoning on Rangelands Through the Selection of Cattle. Rangelands. 2014;36:
694   10–15. doi:10.2111/RANGELANDS-D-13-00031.1

695   22.   Pfister JA, Gardner DR, Panter KE. Consumption of Low Larkspur (Delphinium
696   nuttallianum) by Grazing Sheep. Rangel Ecol Manag. 2010;63: 263–266. doi:10.2111/REM-D-
697   09-00084.1

698   23.   Pfister JA, Manners GD, Ralphs MH, Hong ZX, Lane MA. Effects of Phenology, Site, and
699   Rumen Fill on Tall Larkspur Consumption by Cattle. J Range Manag. 1988;41: 509–514.
700   doi:10.2307/3899528

701   24.   Ralphs MH, Olsen JD, Pfister JA, Manners GD. Plant-animal interactions in larkspur
702   poisoning in cattle. J Anim Sci. 1988;66: 2334–2342.

703   25.   Pfister JA, Provenza FD, Panter KE, Stegelmeier BL, Launchbaugh KL. Risk Management to
704   Reduce Livestock Losses from Toxic Plants. J Range Manag. 2002;55: 291–300.
705   doi:10.2307/4003137

706   26.   Pfister JA, Provenza FD, Manners GD, Gardner DR, Ralphs MH. Tall Larkspur Ingestion:
707   Can Cattle Regulate Intake Below Toxic Levels? J Chem Ecol. 1997;23: 759–777.
708   doi:10.1023/B:JOEC.0000006409.20279.59

709   27.   Smith MA, Waggoner, Jr. JW, Sims S. Larkspur: Managing grazing to avoid poisoning cattle.
710   University of Wyoming Cooperative Extension; 2010 Jul. Report No.: MP-111.13.

28.   Panter KE, Manners GD, Stegelmeier BL, Lee S, Gardner DR, Ralphs MH, et al. Larkspur
poisoning: toxicology and alkaloid structure–activity relationships. Biochem Syst Ecol. 2002;30:
113–128. doi:10.1016/S0305-1978(01)00123-5

29.   Grina JA, Schroeder DR, Wydallis ET, Stermitz FR. Alkaloids from Delphinium geyeri. Three
new C20-diterpenoid alkaloids. J Org Chem. 1986;51: 390–394. doi:10.1021/jo00353a023

30.   Grimm V, Berger U, DeAngelis DL, Polhill JG, Giske J, Railsback SF. The ODD protocol: A
review and first update. Ecol Model. 2010;221: 2760–2768.
doi:10.1016/j.ecolmodel.2010.08.019

31.   Allen VG, Batello C, Berretta EJ, Hodgson J, Kothmann M, Li X, et al. An international
terminology for grazing lands and grazing animals. Grass Forage Sci. 2011;66: 2–28.
doi:10.1111/j.1365-2494.2010.00780.x

32.   Shiyomi M, Tsuiki M. Model for the spatial pattern formed by a small herd in grazing cattle.
Ecol Model. 1999;119: 231–238. doi:10.1016/S0304-3800(99)00059-9

33.   Wilensky U. Netlogo 6.01 [Internet]. Evanston, IL: Center for Connected Learning and
Computer-Based Modeling, Northwestern University; 1999. Available:
http://ccl.northwestern.edu/netlogo

34.   Grimm V. Visual Debugging: A Way of Analyzing, Understanding and Communicating
Bottom-up Simulation Models in Ecology. Nat Resour Model. 2002;15: 23–38.
doi:10.1111/j.1939-7445.2002.tb00078.x

35.   Hexagon Geospatial. ERDAS IMAGINE 2016. Madison, Alabama; 2016.

36.   Huete AR. A soil-adjusted vegetation index (SAVI). Remote Sens Environ. 1988;25: 295–309.
doi:10.1016/0034-4257(88)90106-X

37.   ESRI. ArcGIS Desktop 10.4. Redlands, CA: ESRI; 2015.

38.   ESRI. Collector for ArcGIS 17.01. Redlands, CA: ESRI; 2017.

39.   Bonham CD. Measurements for Terrestrial Vegetation. Wiley; 1989.

40.   Bailey DW. Daily selection of feeding areas by cattle in homogeneous and heterogeneous
environments. Appl Anim Behav Sci. 1995;45: 183–200. doi:10.1016/0168-1591(95)00586-H

41.   Harris NR, Johnson DE, McDougald NK, George MR. Social Associations and Dominance of
Individuals in Small Herds of Cattle. Rangel Ecol Manag. 2007;60: 339–349. doi:10.2111/1551-
5028(2007)60[339:SAADOI]2.0.CO;2

42.   Welch KD, Green BT, Gardner DR, Cook D, Pfister JA. The effect of administering multiple
doses of tall larkspur (Delphinium barbeyi) to cattle. J Anim Sci. 2015;93: 4181.
doi:10.2527/jas.2015-9101

43.   National Research Council. Nutrient Requirements of Beef Cattle: Seventh Revised Edition:
745   Update 2000 [Internet]. Washington, D.C.: National Academies Press; 2000.
746   doi:10.17226/9791

747   44.   Holechek JL, Pieper RD, Herbel CH. Range management: principles and practices. 6th ed.
748   Pearson Education, Inc.; 2011.

749   45.   Augustine DJ, Derner JD. Controls over the strength and timing of fire-grazer interactions in a
750   semi-arid rangeland. J Appl Ecol. 2014;51: 242–250. doi:10.1111/1365-2664.12186

751   46.   Senft RL, Coughenour MB, Bailey DW, Rittenhouse LR, Sala OE, Swift DM. Large Herbivore
752   Foraging and Ecological Hierarchies. BioScience. 1987;37: 789–799. doi:10.2307/1310545

753   47.   Lazo A. Social segregation and the maintenance of social stability in a feral cattle population.
754   Anim Behav. 1994;48: 1133–1141. doi:10.1006/anbe.1994.1346

755   48.   Green BT, Welch KD, Gardner DR, Stegelmeier BL, Davis TZ, Cook D, et al. Serum
756   elimination profiles of methyllycaconitine and deltaline in cattle following oral administration
757   of larkspur (Delphinium barbeyi). Am J Vet Res. 2009;70: 926–931. doi:10.2460/ajvr.70.7.926

758   49.   Gardner DR, Pfister JA. Toxic alkaloid concentrations in Delphinium nuttallianum,
759   Delphinium andersonii, and Delphinium geyeri in the Intermountain Region. Rangel Ecol
760   Manag. 2007;60: 441–446. doi:10.2111/1551-5028(2007)60[441:TACIDN]2.0.CO;2

761   50.   SAS Institute. JMP. Cary, NC: SAS Institute, Inc.; 2016.

762   51.   R Core Team. R: a language for statistical computing [Internet]. Vienna, Austria: R Foundation
763   for Statistical Computing; 2017. Available: https://www.R-project.org

764   52.   Google/Open source. OpenRefine 2.8 [Internet]. 2017. Available:
765   http://openrefine.org/index.html

766   53.   Faraway JJ. Extending the Linear Model with R: Generalized Linear, Mixed Effects and
767   Nonparametric Regression Models, Second Edition. CRC Press; 2016.

768   54.   Hilbe JM. Negative Binomial Regression. Cambridge University Press; 2011.

769   55.   Anderson DR. Model based inference in the life sciences: a primer on evidence. New York ;
770   London: Springer; 2008.

771   56.   Hart RH, Bissio J, Samuel MJ, J. W. Waggoner J. Grazing Systems, Pasture Size, and Cattle
772   Grazing Behavior, Distribution and Gains. J Range Manag. 1993;46: 81–87.
773   doi:10.2307/4002452

774   57.   Galyean ML, Gunter† SA. Predicting forage intake in extensive grazing systems. J Anim Sci.
775   2016;94: 26–43. doi:10.2527/jas.2016-0523

776   58.   Cordova FJ, Wallace JD, Pieper RD. Forage Intake by Grazing Livestock: A Review. J Range
777   Manag. 1978;31: 430–438. doi:10.2307/3897201

59. Walker JW, Heitschmidt RK, Dowhower SL. Evaluation of Pedometers for Measuring
Distance Traveled by Cattle on Two Grazing Systems. J Range Manag. 1985;38: 90–93.
doi:10.2307/3899343

60. Nielsen DB, Ralphs MH, Evans JO, Call CA. Economic feasibility of controlling tall larkspur
on rangelands. J Range Manag Arch. 1994;47: 369–372.

61. Anderson DM, Estell RE, Holechek JL, Ivey S, Smith GB. Virtual herding for flexible
livestock management – a review. Rangel J. 2014;36: 205–221. doi:10.1071/RJ13092

62. Allen CR, Holling CS. Discontinuities in Ecosystems and Other Complex Systems. Columbia
University Press; 2008.

63. Barnes M. Livestock Management for Coexistence with Large Carnivores, Healthy Land and
Productive Ranches. Keystone Conservation; 2015.

64. Bailey DW, Trotter MG, Knight CW, Thomas MG. Use of GPS tracking collars and
accelerometers for rangeland livestock production research. J Anim Sci. 2017;95: 360–360.
doi:10.2527/asasann.2017.740

65. Ralphs MH, Jensen DT, Pfister JA, Nielsen DB, James LF. Storms Influence Cattle to Graze
Larkspur: An Observation. J Range Manag. 1994;47: 275–278. doi:10.2307/4002547

66. Provenza FD. Twenty-five years of paradox in plant-herbivore interactions and" sustainable"
grazing management. Rangel Arch. 2003;25: 4–15.

67. Provenza FD, Villalba JJ, Dziba LE, Atwood SB, Banner RE. Linking herbivore experience,
varied diets, and plant biochemical diversity. Small Rumin Res. 2003;49: 257–274.
doi:10.1016/S0921-4488(03)00143-3

799