1   An agent-based model of cattle grazing toxic Geyer's larkspur
2

3   Kevin E. Jablonski[1*], Randall B. Boone[2], Paul J. Meiman[1]
4
5   [1] Department of Forest and Rangeland Stewardship, Colorado State University, Fort Collins,
6   Colorado, United States of America.
7   [2] Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, Colorado, United
8   States of America
9
10 * Corresponding author
11 Email: kevin.jablonski@colostate.edu

Abstract

By killing cattle and otherwise complicating management, the many species of larkspur

(*Delphinium* spp.) present a serious, intractable, and complex challenge to livestock grazing

management in the western United States. Among the many obstacles to improving our

understanding of cattle-larkspur dynamics has been the difficulty of testing different grazing

management strategies in the field, as the risk of dead animals is too great. Agent-based models

(ABMs) provide an effective method of realistically testing alternate management strategies without

risk to livestock. ABMs are especially useful for modeling complex systems such as livestock grazing

management, and allow for realistic bottom-up encoding of cattle behavior. Here, we introduce a

spatially-explicit, behavior-based ABM of cattle grazing in a pasture with a dangerous amount of

Geyer's larkspur (*D. geyeri*). This model indicates that herd cohesion and stocking density are key

drivers of larkspur-induced toxicosis, and that alteration of these factors within realistic bounds can

mitigate risk. Crucially, the model points to herd cohesion, which has received little attention in the

discipline, as playing an important role in lethal acute toxicosis. As the first ABM to simulate grazing

cattle at realistic scales, this model also demonstrates the tremendous potential of ABMs to

illuminate grazing management dynamics, including fundamental aspects of livestock behavior

amidst heterogeneity.

Introduction

The many species of larkspur (*Delphinium* spp. L.) present a serious, intractable, and complex

challenge to livestock grazing management in the western United States [1–3]. Larkspur plants

contain numerous norditerpinoid alkaloids, which are potent neuromuscular paralytics that, for

reasons that are not entirely understood, are particularly effective at killing cattle, with yearly herd

losses estimated at 2-5% for those grazing in larkspur habitat [3,4]. To avoid such losses, producers

will often abandon or delay grazing in pastures with larkspur, which creates a substantial opportunity

cost and an impediment to achieving management objectives [1,4].

Among the many challenges to improving our understanding of cattle-larkspur dynamics has

been the difficulty of testing different grazing management strategies in the field, as the risk of dead

animals is too great. Additionally, the complexity of livestock grazing management, especially when

considered across the wide range of habitats and management regimes in which larkspur is found,

suggests that results from individual field experiments would be unlikely to be broadly useful [5,6].

What is needed instead is a method of realistically testing grazing management strategies without risk

to livestock and with the flexibility to test multiple scenarios. Agent-based models (ABMs) provide

such a method.

ABMs are computational simulation tools that focus on the behavior of individual "agents"

as they interact with one another and the environment [7]. They differ from other types of

simulation models in being bottom-up (versus top-down) with group-level behaviors emerging from

(usually) realistic individual behaviors rather than deterministic formulae [8]. ABMs are thus

particularly useful in modeling complex systems, where the results of the interactions among system

elements are not easily predicted or understood [9,10]. Indeed, it has been suggested that bottom-

up-simulation may be the best way to increase our understanding of complex systems, which is one

of the most important challenges confronting modern science [9,11,12].

As noted by Dumont and Hill [11], ABMs are "particularly suited to simulate the behavior of

groups of herbivores foraging within a heterogeneous environment". The authors encourage the use

of ABMs in situations where experimentation is impractical, and those where comparison of

different management strategies is needed. Despite this encouragement, and despite the growing

enthusiasm for ABMs in other disciplines, they have been little used in livestock grazing

management research, despite the existence of relevant studies to parameterize such a model [e.g.,

13–18]

Previous research into the relationship between grazing management and larkspur toxicosis

has largely focused on timing of grazing, with some attention paid to mineral supplementation, pre-

grazing with sheep, and, increasingly, genetic susceptibility [3,4,19–22]. Some papers have suggested

that cattle behavior, influenced by management, can play a role in mitigating larkspur deaths [23,24],

but these ideas have received little empirical study. Only anecdotally has it been observed that,

regardless of timing of grazing, it may be possible to eliminate losses to larkspur by increasing

stocking density, due to a dilution effect (same amount of alkaloids, more cattle) or perhaps changes

in herd behavior [25].

In this paper, we introduce a spatially-explicit, behavior-based ABM of cattle grazing in a

pasture with a dangerous amount of Geyer's larkspur (*Delphinium geyeri* Green), in which MSAL-type

alkaloids are the dominant toxin [26,27]. This model provides significant management-relevant

insight for producers dealing with larkspur and demonstrates the great potential of ABMs to address

questions in livestock grazing management, including not only other discrete challenges but also

fundamental aspects of livestock behavior amidst heterogeneity.

Methods

The model description follows the updated Overview, Design Concepts, and Details (ODD)

protocol, an accepted method for standardizing published descriptions of ABMs [28].

Purpose

We designed this model to test the effect of covarying instantaneous stocking density [29]

and herd cohesion (also known as troop length) [30] on the likelihood of bovine alkaloid toxicosis

caused by *D. geyeri*. We developed and executed the model in NetLogo 6.01, using the

BehaviorSpace tool to implement simulations [31].

Basic principles

Behavior-based encoding of cattle activities was the guiding principle of model design. As

noted by Mclane et al. [7], "the behavior-based approach leads to a more complex web of decisions,

and the responses of the animal to stimuli are often more multifaceted". We add that the behavior-

based approach is also more likely to allow for instructive emergent properties. In practice, the

behavior-based approach means that at every step of the coding process we sought literature on

actual cattle behavior and then encoded that behavior as realistically as possible. When literature was

lacking we used our best estimates of cattle behavior from our years as livestock researchers and

managers. The behavior-based approach also found expression in model validation, when one mode

of validation was whether the cows in the model "act like cows". This was achieved through a

lengthy process of visual debugging [32].

A second core principle was parsimony. Because this is the first ABM that we know of to

incorporate cattle at the individual scale of interaction with the environment (1 m^2) and extended to

a realistic pasture size, we were initially tempted to include every cattle behavior we could. However,

our focus on parsimony to the question at hand meant that we instead included only those behaviors

relevant to the consumption of larkspur. A final guiding principle was that when a judgement call

was needed, we erred on the side of making the effects of alkaloid toxicosis more prominent. If the

model was to show an effect of grazing management on reducing larkspur-induced toxicosis, we

wanted to be sure that we had taken every precaution against preconditioning it to do so.

Entities and state variables

The model has two kinds of entities: pixels representing 1 m^2 patches of land and agents

representing 500 kg adult cows (1.1 animal-units). The patches create a model landscape that is 1663

x 1580 patches (262.75 ha, of which 258.82 ha are within the pasture under study and 3.93 ha are

outside the fence line and thus inaccessible). This landscape aims to replicate pasture 16 at the

Colorado State University Research Foundation Maxwell Ranch, a working cattle ranch in the

Laramie Foothills ecoregion of north-central Colorado that is a transition zone between the Rocky

Mountains and the Great Plains. Several pastures on the ranch, including pasture 16, have significant

populations of *D. geyeri*, which generate ongoing management challenges and have fatally poisoned

cattle.

To make the model appropriately spatially explicit we included three sets of geographic data.

First, using data from the Worldview-2 satellite (8-band multispectral, resolution 2 m) from July 10,

2016, we created an index of non-tree/shrub vegetation distribution within the pasture using a soil-

adjusted vegetation index (SAVI) within ERDAS Imagine 2016 software at a resolution of 1 m

[33,34]. Second, as there are no developed watering locations in pasture 16, with ArcGIS Desktop

10.4 we digitized and rasterized (at 1 m) all locations of naturally occurring water as of July 2017

[35].

Lastly, in June and July of 2017 we mapped larkspur distribution and density in pasture 16

using a hybrid approach. We began by digitally dividing the pasture into 272 1-ha sampling plots.

Because we knew larkspur to be of patchy distribution, in each plot we first mapped all larkspur

patches (defined as areas with >1 larkspur plant • m^{-2}) using an iPad equipped with Collector for

ArcGIS 17.01 [36] and a Bad Elf Pro+ Bluetooth GPS receiver accurate to 2.5 m. To sample areas

outside of larkspur patches for larkspur density, we counted all living larkspur plants in a 6-m-wide

belt transect running horizontally across the plot, with the origin randomly assigned and any patches

excluded [37]. Using ArcGIS Desktop we then extended the belt-transect-derived larkspur density to

the rest of the plot (excluding patches), and both sets of data were integrated into a 1 m raster of

larkspur distribution.

The number of cows (individual agents) in the model varies according to the chosen

stocking density (SD, in AU • ha^{-1}). Cows are assigned the role of "leader" (5%), "follower" (85%),

or "independent" (10%) [16,38,39]. Functionally relevant state variables for patches and cows, as

well as global variables and inputs, are described in Table 1.

Table 1. Relevant model variables.

Entity	Variable	Description
Patches	forage-mass	Amount of currently available forage (g)
	n-forage-mass	Mean initial available forage in patches within a radius of 3 m (g)
	MSAL-content	Amount of toxic alkaloids currently in patch (mg)
	times-grazed	Number of times patch has been grazed
Cows	role	Role in the herd: leader, follower, or independent
	herdmates	Agent-set consisting of nearest 20 cows
	mean-herd-distance	Mean distance to herdmates
	consumption-level	Total amount of forage consumed during model run (g)
	total-MSAL-intake	Total amount of MSAL alkaloids consumed during model run (mg)
	daily-MSAL-intake	Amount of MSAL alkaloids consumed during current day (mg)
	hydration	Hydration level, decreases to zero between visits to water
	ready-to-go	Used by leader cows only, a measure of their inclination to move on from an overgrazed site
Globals	waterers	Patch-set of all watering locations
	tolerance	Herd-size-dependent variable determining leader cows' tolerance for relatively overgrazed sites
	site-radius	Radius of site when choosing a new site; product of herd-cohesion-factor and herd size resulting in space per cow ranging from 10 m^2 to 1000 m^2
	herd-distance	Desired mean-herd-distance; product of herd-cohesion-factor resulting in range from 10 m to 100 m
Inputs	kgs-per-hectare	Mean amount of usable forage (kg • ha^{-1})
	mean-larkspur-mass	Mean mass of larkspur plants (g)
	MSAL-concentration	MSAL alkaloid concentration in larkspur plants (mg • g^{-1})
	herd-cohesion-factor (HCF)	Determines herd-distance and site-radius; range 1-10, increase leads to more cohesive herd
	stocking-density (SD)	Instantaneous stocking density (AU • ha^{-1})

Scales

The model simulates cow activities at multiple temporal and spatial scales. In each tick (one

cycle through the model code), each cow interacts with a single 1 m^2 patch (a feeding station) by

grazing (>99% of the time) or drinking water (twice per day) [13]. A tick does not represent time,

but rather the occurrence of this interaction. This is because the duration of this interaction will vary

depending on the amount of forage available, among other factors. Instead, time is represented by

consumption of forage. When the average consumption of the grazing herd is equal to the average

daily consumption of a 500 kg cow (12.5 kg), the model counts a grazing-day as having passed [40].

Total model run time is measured in animal-unit-months (AUMs) [41].

The narrowest scale of spatial interaction is the eating interaction occurring within a single

patch (1 m^2). When determining the next patch to graze, the cow's decision is based on a desire

either to move closer to its herdmates or to choose a nearby patch with maximum available forage.

This decision happens on the scale of 2-25 m. Finally, leader cows make decisions on the scale of

the entire pasture by deciding when it is time to visit water or time to move from the current feeding

site to a new site.

Thus, there are four programmed spatial scales (additional scales may be emergent) at which

the cows interact with the landscape: 1) the individual patch; 2) the scale of herd cohesion, set by the

user; 3) the current feeding site; and 4) other feeding sites, identifiable by leader cows. The number

of ticks that will pass before reaching a stopping point (say, 165 AUMs) depends on the number of

animals grazing, their herd cohesion, the amount and distribution of available forage, and stochastic

emergent properties of the model. For an expanded discussion of temporal and spatial scales of

foraging behavior of large herbivores, see Bailey and Provenza [13].

Process overview and scheduling

158   Fig 1 illustrates the model execution process for each tick. Each cow moves through each step of

159   the process, but only performs those steps linked to its role.

Fig 1. Pseudo-coded flow chart of model processes, with role of cows executing each process in parentheses. 1= leader, 2=follower, 3= independent.

Check hydration

Each leader cow checks it hydration level, which is tied to forage consumption such that it depletes to zero twice per day. If an individual leader detects its hydration level as less than or equal to zero, it initiates a movement to water for the whole herd.

Go to water

The water source in pasture 16 is a stream that is intermittently below ground. The go-to-water procedure directs each cow to go to the nearest waterer patch with two or fewer cows already present. The hydration value for each cow is then set to maximum, and the value for ready-to-go for leader cows is set to tolerance – 1. This ensures that the herd is intolerant of remaining near a waterer if the area has been grazed already, and encourages a site change after drinking water. A global variable ensures that no other processes occur during a tick when watering occurs.

Check site change

This process is only executed by leader cows, each of which assesses the mean number of

times patches within a radius of 10 m have been grazed. If these patches have been grazed relatively

more (defined as >0.5 • mean times-grazed of all patches + 1.2) than the pasture as a whole, the

value of ready-to-go increases by one. If this value reaches a pre-defined threshold (which increases

with herd size), the individual then initiates a site change, but only if the individual's hydration value

is not approaching zero, in which case it instead initiates the go-to-water procedure. We arrived at

the threshold formula for increasing the value of ready-to-go by using visual debugging and

validation related to site change frequency, as well as theory on the optimization of grazing effort

[13,42].

If conditions for a site change are satisfied, the deciding leader cow first identifies the best

five available sites, using criteria of number of times-grazed, forage-mass, and n-forage-mass to

determine a centroid patch. The nearest of these patches is then used to create a new site at a radius

that is linked to the user selected herd-cohesion-factor (HCF) and the size of the herd, resulting in

10-1,000 m^2 • cow^{-1} in the new site. The leader cow then initiates the change-site procedure for itself

and all other cows.

Change site

191 This procedure is initiated according to role, so that leader cows have first choice of their

192 location in the new site, followers second, and independents third. Within the allocated new site,

193 each cow chooses the patch with the most forage that has no cows on it or any of its four direct

194 neighbors.

Assess herd

In combination with the environmental-movement procedure, this process represents >99%

of cow actions in the model. Each cow first sets its herdmates as the nearest 20 other cows [43]. For

leader and follower cows, if the individual's mean distance to these herdmates is greater than herd-

distance, it "herds up". This is achieved by facing the centroid of the herdmates and moving to the

patch with maximum available forage that is 10-25 m in the direction of this centroid, within a cone

of vision of ±45 degrees [14]. For independent cows, the same process occurs but is only initiated if

the distance from herdmates is greater than 2.5 times the herd-distance of the other cows.

Independent cows are also repelled from the center of their herdmates by moving away by the same

procedure when they are within one-half of the herd-distance.

Environmental movement

If none of the above procedures are implemented, each cow will make a movement decision

based on local grazing conditions. If the patches within a radius of 10 m are relatively ungrazed

(mean times-grazed < 0.5) the cow will move to the patch with the most available forage within 2 m,

within a ±45 degree cone of vision. If the same area is relatively well grazed (mean times-grazed ≥

0.5), the cow then looks further afield, choosing the patch with the most available forage within 10

m, within a cone of vision of ±45 degrees.

Eat

The eat procedure is the core interaction between the cows and the forage and larkspur.

Behavior varies slightly depending on how many times the patch has previously been grazed. If the

current visit is the first time it has been grazed, the cow eats 40% of the available forage [15,18]. If it

is the second visit, it eats 50% of what remains. In the third and any subsequent visits, it eats 60%.

Each cow then increases its consumption-level by the same amount and decreases its hydration

value. If there is larkspur present (in the form of MSAL-content), that is consumed at the same level

as the rest of the forage, and MSAL-intake values are increased. The corresponding patch values are

decreased to account for consumption. Lastly, times-grazed in the patch is increased by one.

Design concepts

Emergence

Because the actions of the cows are encoded via simple behavior-based processes, nearly all

model results can be considered emergent properties. These include the stochastic distribution of

the herd and subherds, forage consumption, larkspur consumption, grazing pressure and patterns,

and site changes [44].

Adaptation, objectives, learning, and predictions

The cows adapt to the grazing environment as they and their fellow cows graze, continually

seeking their main model objective of maximizing forage consumption within behavioral limits [14].

There is no encoded learning or prediction, as the cows are programmed to be familiar with the

location of forage and water in the pasture. However, it may be that learning and prediction are

emergent, in that activities that we might consider to be evidence of those behaviors are visible in

the model as a result of the simple encoded behaviors.

Sensing and interaction

The cows sense each other and their environment at multiple spatial scales. Interaction

occurs with other cows whenever moving to a new patch, both via sensing if a patch is already

occupied and by seeking to herd up when too far from their herdmates.

Stochasticity

There is no environmental stochasticity in this model iteration, as we sought to make the

landscape as realistic as possible by incorporating relevant data from the real pasture 16. However,

cattle interactions with the forage and larkspur are highly stochastic, as most model processes

involve a degree of randomization.

Observation

Of primary importance are data related to forage intake and different levels of daily and two-

day MSAL alkaloid consumption, corresponding to dose-response studies in the literature [45]. Most

interesting was the number of times in a model run that any individual cow crossed the threshold

into potentially lethal acute toxicosis, during which they would be expected to be recumbent and

unable to stand, with a high chance of death [45]. To measure the number of such cases, the model

tracked the number of cows that received a dose of ≥ 8 mg $\bullet$ kg^{-1} of body weight over the course of

one to two days [45].

We were also interested whether the reduction in cases of acute toxicosis generated by

increasing stocking density and, especially, herd cohesion, was mirrored by corresponding changes in

the number of animals experiencing sub-lethal acute toxicosis. Specifically, we looked at two levels:

instances where a cow consumed ≥ 4 mg $\bullet$ kg^{-1} and <8 mg $\bullet$ kg^{-1} or ≥ 2 mg $\bullet$ kg^{-1} and <4 mg $\bullet$ kg^{-1} of

toxic alkaloids in one to two days. In the first case, while this level of consumption is unlikely to

cause death, we can still consider this a "danger zone". We can consider the latter case to be a

relatively safe outcome with a strong likelihood of "no observable adverse effect" and thus the "safe

zone" [3,45].

A final statistical observation of interest was the distribution of individual overall total and

maximum daily alkaloid intake. If cattle are going to consume larkspur with the aim of keeping

consumption at sub-lethal levels, it is important to determine if it is possible to decrease the variance

of alkaloid consumption through management. Lastly, the model also reported various validation

data such as the count of site changes and travel distance per day.

Initialization

Landscape initialization begins by loading the SAVI layer and a user-input value for available

forage per ha (kgs-per-hectare). The model uses a nonlinear exponential formula to distribute forage

such that the patches with the least forage contain one-third of the mean forage, while the patches

with the most contain three times the mean forage. Next, the model incorporates the larkspur

distribution layer, using inputs of median larkspur mass (g) and mean MSAL concentration (mg • g^{-1})

to generate an MSAL alkaloid (hereafter simply "alkaloid") content for each patch. These values are

based on our unpublished data on *D. geyeri* mass and toxicity at the Maxwell Ranch such that

larkspur plants in areas of high SAVI were 50% larger than the median, and larkspur plants in areas

of low SAVI were 50% smaller than the median. Finally, the model incorporates the water location

layer. All other patch variables are derived from these inputs. Fig 2 shows the initialized landscape.

275
Fig 2. Model landscape. (a) Initialized full model landscape, with darker green indicating areas with greater aboveground
forage biomass. (b) Landscape with larkspur locations only, with darker purple representing higher MSAL-content and
with results of hybrid sampling method evident. (c) Landscape with watering locations only, pointed out by arrows.

The final step in model initialization is to create the cows by using the input of stocking-

density multiplied by the area of the pasture. All cows are initially in the same random location in the

pasture. This location is largely irrelevant as the cows immediately go to water, but we did not want

it to be the same location each time because this would be unrealistic (pasture 16 has multiple

entrances for cattle) and would limit stochasticity. At this point, the model is fully initialized and is

executed following the processes laid out above.

Simulation

We used the BehaviorSpace tool in NetLogo to run a full factorial simulation of four

different levels of both herd-cohesion-factor (1, 4, 7, and 10) and stocking-density (0.25, 0.5, 1.0,

and 2.0 AU • ha^{-1}). We replicated each combination 30 times, for a total of 480 simulations. Input

median larkspur mass was 3.5 g and input MSAL alkaloid concentration was 3.0 mg • g^{-1}. We chose

these values to be representative of an excellent growing year with larkspur plants at middle bud

stage, when the alkaloid pool (total available mg) is highest—arguably the most dangerous possible

conditions. This is also a time of year that cattle grazing in larkspur habitat is frequently avoided,

despite being a highly desirable time for grazing [1,4,46]. Input value for kgs-per-hectare was 500 kg,

based on current ranch usage and typical values for the area.

Statistical analysis

We used both JMP Pro 13.0.0 and R statistical software, version 3.3.3 for analysis [47,48]. To

analyze count data from the model, we used a Generalized Linear Model (GLM) with a negative

binomial distribution and a log-link function using the MASS package in R [49,50]. To confirm that

the negative binomial distribution was the correct choice, we compared each to a GLM with a

Poisson distribution and a log-link function. In all cases, the GLM with the negative binomial

distribution was far superior, using residual deviance and Aikaike's information criterion (AIC) as

judgment criteria [51].

We further analyzed the count data using combinations of herd-cohesion-factor and

stocking-density as "treatments" with a non-parametric Games-Howell test for multiple comparison

of means (R package userfriendlyscience), which is robust to unequal variances [52]. For data on

total and maximum daily alkaloid intake, we used multiple linear regression (R base package) and a

Tukey HSD test (JMP) for comparison of "treatments" [52].

Results

Model function and validation

While a complete detailing of ABM function and validation for overall grazing behavior is

beyond the scope of this paper, it is nonetheless important to demonstrate that the modeled cows

"act like cows" insofar as larkspur consumption is concerned, and that their effect on the landscape

is plausibly that of real cows. Toward this end, we offer Fig 3 to illustrate how varying HCF

influences herding patterns, and to show how grazing was distributed across the pasture in one

model run.

Fig 3. The effect of varying herd-cohesion-factor (HCF) on herd patterns. Note that the cows depicted in these images are drawn 200 times larger than they really are to aid visualization, which makes them appear closer to one another than they are. Stocking density for all images is 1.0 AU • ha⁻¹. White cows are leaders, black followers, and gray independents. Yellow indicates patches that have been grazed twice, red three times. (a) HCF=10, AUMs=14; (b) HCF=7, AUMs=68; (c) HCF=4, AUMs=119; (d) HCF=1, AUMs=163. Typical usage for this pasture (258.82 ha) is 165 AUMs.

Decreasing HCF increases overall herd separation and leads to more wandering among the independent cows and others. Note that in Fig 3a the cows have formed distinct independent subherds. This appears to be an emergent property of cows grazing with high herd cohesion (herd-distance ≤ 20 m).

The cows initially graze the areas with high forage amounts (dark green) in proximity to the

water, and gradually extend their impact outward. By the end of the grazing cycle (Fig 3d), they have

visited the entire pasture, though areas further from water have been grazed less. Areas of initial

high forage mass have been grazed two or more times, while many areas of low forage mass have

not been grazed at all. These results are in line with well-established understanding of grazing

patterns in large pastures [41].

The mean value for site changes per day for the 16 different combinations of HCF and SD

varies from 2.5 for few cows grazing very loosely (HCF=1, SD=0.25) to 6.9 for many cows grazing

very cohesively (HCF=10, SD=2.0). These values are in line with the estimate of 1-4 hours per

feeding site by Bailey and Provenza [13]. For runs with few cows grazing with little cohesion

(HCF=1, SD=0.25), mean daily travel was 4.44 km, while many cows grazing very cohesively

(HCF=10, SD=2.0) traveled an average of 8.06 km per day. These numbers and the positive trend

also track well with data from previous studies [e.g., 53].

As a last point of model validation, we were interested to see if the number of modeled cases

of larkspur-induced lethal acute toxicosis would parallel numbers from the literature when we

modeled grazing to be similar to the current management scheme. When modeled to reflect current

management practices, with HCF=4, SD=0.5, and for 165 AUMs (removing approximately 45% of

available forage), we recorded a mean of 5.2 cases of lethal acute toxicosis across 30 model

iterations. This amounts to 4.0% of cows, which falls within the estimate in the literature of losses of

2-5% in pastures with dangerous amounts of larkspur [4,54].

Lethal acute toxicosis

The coefficient for HCF (Table 2), as a log odds ratio, indicates that an increase of 1 in HCF

resulted in an 19.0% decrease in occurrences of lethal acute toxicosis. The coefficient for SD

indicates that an increase of 1 in SD resulted in a 52.3% decrease. Lastly, the coefficient for the

interaction of HCF with SD indicates that an increase in either HCF or SD increases the effect of

the other.

Table 2. Results of GLM with negative binomial distribution and log-link function for count of lethal acute toxicosis as
predicted by herd-cohesion-factor (HCF) and stocking-density (SD). β coefficients are from the same GLM without the
interaction present. GLM fit: Fisher scoring iterations=1; residual deviance=519.7 on 476 degrees of freedom;
AIC=2086.3.

Coefficient	Estimate	Std. error	p-value	β
Intercept	3.161	0.142	<0.001	
HCF	-0.211	0.025	<0.001	-0.162
SD	-0.74	0.137	<0.001	-0.127
HCF:SD	0.086	0.029	0.003	

356   Because our goal is to provide management-relevant insight, we compared the 16

357   combinations of HCF and SD as treatments, which present a clearer parallel to grazing management

358   practices. Fig 4 shows box plots of the distributions for the 16 combinations of HCF and SD, and

359   Table 3 shows the results of a post-hoc multiple comparisons of means using a Games-Howell test.

Fig 4. Box plots of distribution of counts of lethal acute toxicosis (≥ 8 mg • kg⁻¹ in 1-2 days) cases. From 30 model runs for each combination of herd-cohesion factor (HCF) and stocking-density (SD), ordered by mean count of lethal acute toxicosis cases, with outliers as jittered circles.

Table 3. Connected letters table for the results of multiple comparisons of means using a Games-Howell test for mean count of lethal acute toxicosis (LAT) by herd-cohesion-factor (HCF) and stocking-density (SD). Levels not connected by the same letter are significantly different at α=0.05.

HCF:SD							Mean ct. of LAT
4:0.25	A						13.20
1:0.25	A						12.77
1:0.5	A						10.23
7:0.25	A	B					9.00
4:0.5		B	C				5.20
1:1			C				4.37
4:1			C	D			3.07
1:2			C	D	E		3.70
4:2*			C	D	E	F	G	1.90
7:0.5				D	E	F		1.73
7:1					E	F		1.13
7:2						F		0.70
10:0.25						F	G	0.70
10:0.5						F	G	0.63
10:1						F	G	0.40
10:2							G	0.00

*Single large outlier with otherwise tight distribution makes 4:2 difficult to interpret. See Fig 4.

Sub-lethal acute toxicosis

The coefficient for HCF (Table 4) indicates that an increase of 1 in HCF resulted in a 4.8% decrease in occurrences of sub-lethal acute toxicosis in the danger zone. The coefficient for SD indicates that an increase of 1 in SD resulted in a 12.2% decrease. The coefficient for the interaction of HCF with SD indicates an interaction such that an increase in either HCF or SD slightly increases the effect of the other. Table 5 shows the result of the post-hoc means comparison of occurrences of sub-lethal acute toxicosis in the danger zone for the 16 combinations of HCF and SD.

Table 4. Results of GLM with negative binomial distribution and log-link function for count of sub-lethal acute toxicosis in the danger zone (≥4 mg • kg⁻¹ and <8 mg • kg⁻¹ in 1-2 days) as predicted by herd-cohesion-factor (HCF) and stocking-density (SD). β coefficients are from the same GLM without the interaction present. GLM fit: Fisher scoring iterations=1; residual deviance=499.7 on 476 degrees of freedom; AIC=5127.9.

Coefficient	Estimate	Std. error	p-value	β

Intercept	6.036	0.029	<0.001	
HCF	-0.05	0.004	<0.001	$-2.88 \cdot 10^{-3}$
SD	-0.13	0.025	<0.001	$-1.95 \cdot 10^{-3}$
HCF:SD	-0.017	0.004	<0.001	

381 Table 5. Connected letters table of the results of post-hoc multiple comparisons of means using a Games-Howell test for
382 mean counts of sub-lethal acute toxicosis (SLAT) in the danger zone ($\geq$4 mg $\bullet$ kg^{-1} and <8 mg $\bullet$ kg^{-1} in 1-2 days) by
383 herd-cohesion-factor (HCF) and stocking-density (SD). Levels not connected by the same letter are significantly
384 different at α=0.05.

HCF:SD													Mean ct. of SLAT
1:0.25	A												361.3
4:0.25	A	B											352.2
1:0.5		B	C										336.2
7:0.25		B	C	D									327.4
4:0.5			C	D	E								319.3
1:1				D	E	F							309.5
4:1					E	F	G						293.9
7:0.5						F	G	H					289.5
1:2							G	H					275.3
7:1								H	I				258.4
4:2									I	J			241.7
10:0.25									I	J	K		220.9
7:2										J	K		215.3
10:0.5											K	L	184.1
10:1												L	165.1
10:2													M	122.0

The coefficient for HCF (Table 6) indicates that an increase of 1 in HCF resulted in an 3.3%

increase in occurrences of sub-lethal acute toxicosis in the safe zone. The coefficient for SD

indicates that an increase of 1 in SD resulted in a 1.5% decrease. Table 7 shows the result of the

post-hoc means comparison of occurrences of sub-lethal acute toxicosis in the safe zone for the 16

combinations of HCF and SD.

Table 6. Results of GLM with negative binomial distribution and log-link function for count of sub-lethal acute toxicosis
in the safe zone ($\geq$2 mg $\bullet$ kg^{-1} and <4 mg $\bullet$ kg^{-1} in 1-2 days) as predicted by herd-cohesion-factor (HCF) and stocking-
density (SD). As there was very little interaction between the terms, the results shown are for the GLM without the
interaction. GLM fit: Fisher scoring iterations=1; residual deviance=478.2 on 476 degrees of freedom; AIC=5947.4.

Coefficient	Estimate	Std. error	p-value	β

Intercept	7.28	0.007	<0.001	
HCF	0.033	0.005	<0.001	$-4.29 \cdot 10^{-5}$
SD	-0.015	0.001	<0.001	$-4.64 \cdot 10^{-4}$

396 Table 7. Connected letters table of the results of post-hoc multiple comparisons of means using a Games-Howell test for
397 mean counts of sub-lethal acute toxicosis (SLAT) in the safe zone (≥ 2 mg • kg^{-1} and <4 mg • kg^{-1} in 1-2 days) by herd-
398 cohesion-factor (HCF) and stocking-density (SD). Levels not connected by the same letter are significantly different at
399 $\alpha=0.05$.

HCF:SD									Mean ct. of SLAT
10:1	A								2164.5
10:0.5	A								2145.7
10:0.25	A								2133.0
10:2		B							1988.8
7:0.5			C						1828.6
7:1			C	D					1804.4
7:0.25				D	E				1725.0
7:2					E	F			1690.8
4:1					E	F	G		1661.5
4:2					E	F	G		1654.2
4:0.5						F	G		1643.4
4:0.25						F	G	H	1618.4
1:1							G	H	I	1590.3
1:0.5								H	I	1560.2
1:0.25									I	1545.6
1:2									I	1542.5

400
Distribution of total and maximum alkaloid consumption

While there were some indications of heteroscedasticity and outliers for both models, we determined

that linear regression was robust to those errors in this case. We confirmed this by also fitting

alternate models within other regression frameworks (robust and non-parametric), which returned

very similar results. Because interaction between HCF and SD was weak in both cases, we left this

term out of both models. Tables 8-11 show the results of multiple linear regression and the post-hoc

means comparison of the standard deviation of total MSAL alkaloid consumption and standard

deviation of maximum MSAL alkaloid consumption for the 16 combinations of HCF and SD.

Table 8. Results of multiple linear regression for the standard deviation of mean total MSAL alkaloid consumption as
predicted by herd-cohesion-factor (HCF) and stocking-density (SD). Adj. R^2=0.79.

Coefficient	Estimate	Std. error	p-value	β
Intercept	1989.548	34.578	<0.001	
HCF	-76.907	4.339	<0.001	-0.372
SD	-834.55	21.713	<0.001	-0.807

411
412 Table 9. Connected letters table of the results of post-hoc multiple comparisons of means using a Tukey HSD test for
413 the mean standard deviation (σ) of total MSAL alkaloid consumption (mg) by herd-cohesion-factor (HCF) and stocking-
414 density (SD). Levels not connected by the same letter are significantly different at α=0.05.

HCF:SD										Mean σ tot. alk. cons.
1:0.25	A									3034.5
4:0.25		B								2834.0
7:0.25			C							2451.8
1:0.5				D						2119.3
4:0.5				D						2001.2
7:0.5					E					1761.2
10:0.25					E					1742.9
1:1						F				1467.4
4:1						F	G			1409.1
10:0.5						F	G			1397.2
7:1							G			1290.3
1:2								H		1015.3
10:1								H		1015.2
4:2								H	I	973.5
7:2									I	889.5
10:2										J	680.0

415
416 Table 10. Results of multiple linear regression for the standard deviation of maximum daily alkaloid consumption as
417 predicted by herd-cohesion-factor (HCF) and stocking-density (SD). Adj. R^2=0.55.

Coefficient	Estimate	Std. error	p-value	β
Intercept	401.441	9.82	<0.001	
HCF	-25.042	1.232	<0.001	-0.621
SD	-82.856	6.167	<0.001	-0.411

418
419 Table 11. Connected letters table of the results of post-hoc multiple comparisons of means using a Tukey HSD test for
420 the mean standard deviation (σ) of maximum daily MSAL alkaloid consumption (mg) by herd-cohesion-factor (HCF)
421 and stocking-density (SD). Levels not connected by the same letter are significantly different at α=0.05.

HCF:SD				Mean σ max. daily alk. cons.
1:0.25	A			646.6
4:0.25	A	B		621.6
7:0.25	A	B		607.2
1:0.5		B	C	575.8

4:0.5	C	D								538.2
1:1	C	D	E							509.9
4:1		D	E	F						475.2
1:2			E	F						468.7
7:0.5			E	F						462.3
4:2				F	G					423.2
7:1				F	G	H				414.2
7:2					G	H	I			375.2
10:0.5						H	I			346.6
10:1							I	J		330.5
10:0.25							I	J		311.2
10:2								J		277.4

Discussion

Research into best practices for grazing management in larkspur habitat has long focused on either attempts to eliminate larkspur or on phenological avoidance (what we term "fight or flight"). Because elimination through herbicides or mowing is costly and often impractical, most research and current recommendations focus on avoiding grazing in larkspur habitat at times of year when it is considered most dangerous to cattle, exemplified by the toxic window concept [3,4,21]. While this approach has certainly helped many producers better understand larkspur toxicity dynamics, there is no evidence that it has reduced the overall number of deaths. There are many reasons for this, and interactions are complex and place-based, but we suggest that a reliance on a static view of palatability is largely to blame.

The alternative to fight or flight is to manage grazing such that larkspur intake remains below the threshold where there is an observable negative effect on the cattle. This study provides answers to the question of whether this may be possible, and under which circumstances. For the first time, this model suggests that herd cohesion and stocking density are key drivers of toxicosis, and that management decisions that influence these factors hold potential to limit the risk of death. Of crucial importance is the observation that herd cohesion, which has received almost no

439 consideration in the broader grazing management literature, may be the most important determinant

440 of risk of death.

Lethal acute toxicosis

We observed significant differences in the count of cases of lethal acute toxicosis. The

simplest interpretation is that management that results in few cows grazing very loosely is much

more dangerous to cattle than management that results in many cows grazing very cohesively.

However, the model results point to more nuance than that. The standardized coefficients (Table 2)

indicate that herd cohesion is 28% more influential than stocking density, and we can see that

influence in the ordering of Table 3. These results suggest that, regardless of stocking density, cattle

that graze very cohesively (especially subherds with mean inter-animal distance ≤ 10 m) significantly

lower the risk of death caused by *D. geyeri*. A combination of highly cohesive herd behavior with

high stocking density (2.0 animal units per ha) eliminated lethal acute toxicosis.

Sub-lethal acute toxicosis

Because we seek to manage cattle to consume larkspur without dying, the results for sub-

lethal acute toxicosis are also of interest. For sub-lethal acute toxicosis in the danger zone, similar

ordering between Tables 4 and 3 suggests similar management lessons to those of the lethal acute

toxicosis analysis. The GLM coefficients confirm this, with the standardized coefficients indicating

that herd cohesion is 48% more influential than stocking density in this case. In short, if we aim to

limit alkaloid consumption that approaches lethality, once again highly cohesive herds grazing at

high stocking density is our best approach. Note, however, that elimination of sub-lethal acute

toxicosis in the danger zone may not be possible when there are high levels of toxic Geyer's

larkspur.

Herd cohesion and stocking density have opposite effects on the number of cases of sub-

lethal acute toxicosis in the safe zone, unlike the previous two cases. This means that increased herd

cohesion actually increases the number of safe zone cases, while increased stocking density still

lowers them, with herd cohesion exerting an 8% stronger influence. This points to the dilution

effect of increased stocking density interacting with the smoothing effect of herd cohesion on the

distribution of consumption, which we explore further in the next section.

Distribution of total and maximum alkaloid consumption

Table 8 makes apparent that stocking density, which is strongly correlated with total alkaloid

consumption, also exerts a strong influence on its distribution, with 217% of the influence of herd

cohesion. This means that increased stocking density not only limits total alkaloid consumption

through dilution, but also, in combination with herd cohesion, plays a strong role in limiting the

likelihood of extreme total consumption for individual cows.

That the ordering of Tables 3, 5, and 11 are similar should not come as a surprise. Tables 10

and 11 tell us that herd cohesion and stocking density work together to narrow the distribution of

maximum daily alkaloid consumption, which represents each cow's worst day in a given model run,

with herd cohesion exerting 51% more influence than stocking density. This is the key to

preventing risky levels of alkaloid consumption—keep the distribution of maximum daily

consumption evenly distributed, avoiding outliers.

Parsimony and study limitations

Perhaps the most obvious omissions from our model are individual cattle preference and

stochastic consumption of larkspur. A growing body of evidence indicates that individuals do

develop preferences, and past anecdotes have suggested that cattle may engage in brief but intense

bouts of larkspur consumption [55,56]. Why not, then, include these factors? First, because past

evidence of bouts of larkspur consumption is entirely anecdotal, there are no data to use to

parameterize consumption. It is not even clear if this really occurs, or if the anecdotes are due to

selective observation.

Second, we are comfortable with preference and stochastic consumption remaining below

the minimum scale of the model. That this is the case is not a reason to discount the ability of the

model to address the question of larkspur consumption. Indeed, it is reasonable to understand the

model as though behaviors such as preference and diet mixing are happening. Because there is

heterogeneity in the distribution of forage and larkspur, and heterogeneity and stochasticity in the

cows' consumption, within the context of larkspur consumption we believe these behaviors are

sufficiently realized.

Aside from behaviors and environmental factors occurring below 1 m^2, the model also

excludes behaviors that we determined to hold little to no relevance to larkspur consumption, at

least in this pasture. These include response to slope, resting, and some inconsistently understood

aspects of dominance behaviors. Despite our decision to exclude these behaviors, it is important to

note that they could be incorporated if they become relevant.

The model also excludes plant regrowth. For larkspur, this is not an issue, as plants that are

clipped or grazed during the bud stage exhibit very little regrowth (K. Jablonski, pers. obs.). For

other forage, we determined that regrowth in July in this semi-arid climate would not be substantial

enough within a single grazing period to warrant inclusion. Regrowth of non-larkspur forage would

also serve to lessen the risk of larkspur-induced toxicosis.

Most important are the limitations to making direct management-relevant conclusions from

the model results. While we are confident in the fundamental conclusion that increased herd

cohesion and stocking density lower the risk of lethal acute toxicosis, the exact values for when that

risk approaches zero may be dependent on the circumstances of this model iteration—that of *D.*

geyeri, at the input values for mass and toxicity, on a ranch in northern Colorado. This model is

predictive only in that it can help us understand the range of potential outcomes, rather than predict

exactly what will occur. It is thus essential that every manager interpret these conclusions in context.

Other model implications and future directions

There is a broad literature on the effect of stocking rate/stocking density on many outcomes

(though not larkspur-induced toxicosis) but very little on the effects of herd cohesion, nor on the

interaction of these factors [41]. This is likely due to the relative ease of varying stocking rate versus

manipulating cattle behavior. Because this study provides evidence that it is not only the number of

animals but perhaps more importantly how they behave that affects the likelihood of death by

larkspur, we are interested in what other challenges this finding might apply to. The evolving

promise of affordable GPS tags means that we may also start to be able to test this through direct

observation of entire herds, which can be used to validate and improve grazing ABMs [58].

We are also excited to explore the emergent property of distinct persistent subherds that

appears to emerge when the desired mean inter-animal distance of the cows is < 20 m. If this desire

is either genetically encoded or management-determined (or both), and this subherd behavior is

important to reducing the risk of death by larkspur, how might it affect other negative outcomes,

such as overgrazing of riparian areas or exposure to predation by carnivores [59]?

Though ABMs have some limitations, we believe they offer an exciting new tool for

understanding the grazing behavior of livestock. Indeed, the synergistic emergence of financially

viable GPS technology [58] and "virtual fencing" [60], along with the increasing power of desktop

computers, suggests that the time is right for a modeling revolution in livestock grazing

management. We are excited that this study provides a first example of the potential of ABMs to

contribute to this revolution.

Management implications

The model results indicate that it may be possible to limit or eliminate the risk of fatal

poisoning of cattle by Geyer's larkspur by increasing herd cohesion and/or stocking density. This

provides context for the anecdotal report of Smith et al. [25] for the Sims Ranch in McFadden, WY,

where it was found that an increase in stocking density all but eliminated deaths due to *D. geyeri*.

However, an important caveat is that it is essential that managers understand their level of risk,

which we suggest is best represented by the total MSAL alkaloid pool of a pasture. Though we have

modeled a pasture that ranks among the densest (overall, 0.17 plants $\bullet$ m^{-2}) and most toxic

populations of *D. geyeri* that we have seen, it could be worse elsewhere.

We selected the bounds of herd cohesion and stocking density to align with what we believe

to be realistically achievable by managers in the western US. While stocking density is easily

understood, it may be worthwhile to describe how we think the various levels of herd cohesion

could be achieved (reference Fig 3). We think of HCF values of 1 and 4 as representative of most

current extensive management, such that there is a small to moderate amount of herding behavior

but in which animals are often spread out across a large area. The difference between these two

might be accounted for by differences in breeding history, carnivore pressure, or genetic drift. To

achieve an HCF of 7, we think cattle would need to be selected for strong herding instinct or be

actively, but not necessarily continually, herded. An HCF of 10 is comparable to many herds of wild

ungulates, and may be achievable through the continual presence of a herder or a sustained effort at

selecting for herding behavior.

It is worth noting that dangerous levels of *D. geyeri* are typically found on a limited number

of a single operation's grazing units. This means that the inclusion of herding, for example, would

usually only be necessary for a relatively brief period. In addition, it means that any potential

secondary effects of sub-lethal larkspur consumption, such as appetite suppression (whether and

how this would occur is unclear) would be of similarly limited duration.

There are two additional ways that a rapid increase in herd cohesion may be achieved. First,

a drastic increase in stocking density (via increased animal-units or subdivided pastures) to a level

that approaches "mob" grazing can forcibly increase cohesion. Second, the emerging technology of

virtual fencing holds tremendous promise for achieving rapid changes in grazing behavior, including

herd cohesion [60].

As with any research where cattle lives and producer livelihoods are at stake, it is most

important to emphasize that producers should exercise caution when incorporating our findings into

their own management. Those with low amounts of Geyer's larkspur and with no history of losses

might find comfort in altering their grazing management to incorporate this study's findings. Those

with a great deal of larkspur (Geyer's or other species) and a history of losses should be more

careful. As researchers, our next step is to place these modeling results in context with ongoing plant

experiments and producer surveys to better formulate management recommendations that work.

Acknowledgements

We thank Joel Vaad, manager of the Maxwell Ranch, for access, assistance, and advice. Early

conversations with Michael A. Smith and the Sims family of McFadden, WY, provided crucial

insight into the relationship between grazing management and larkspur. Tanner Marshall assisted in

data collection and asked difficult questions that moved the work forward. We are grateful to

DigitalGlobe for a generous imagery grant that greatly improved our ability to assess the grazing

landscape, and to Hexagon Geospatial for providing the software to analyze the imagery.

References

1. Green B, Gardner D, Pfister J, Cook D. Larkspur poison weed: 100 years of delphinium research. Rangelands. 2009;31: 22–27. doi:10.2111/1551-501X-31.1.22

2. Pfister JA, Gardner DR, Panter KE, Manners GD, Ralphs MH, Stegelmeier BL, et al. Larkspur (Delphinium spp.) poisoning in livestock. J Nat Toxins. 1999;8: 81–94.

3. Welch KD, Cook D, Green BT, Gardner DR, Pfister JA, McDaneld TG, et al. Adverse effects of larkspur (Delphinium spp.) on cattle. Agriculture. 2015;5: 456–474. doi:10.3390/agriculture5030456

4. Pfister JA, Gardner DR, Price KW. Grazing risk on tall larkspur-infested ranges. Rangelands. 1997;19: 12–15. doi:10.2307/4001275

5. Darnhofer I, Gibbon D, Dedieu B. Farming Systems Research: an approach to inquiry. In: Darnhofer I, Gibbon D, Dedieu B, editors. Farming Systems Research into the 21st Century: The New Dynamic. Dordrecht: Springer Netherlands; 2012. pp. 3–31. Available: http://www.springerlink.com/index/10.1007/978-94-007-4503-2_1

6. Provenza F, Pringle H, Revell D, Bray N, Hines C, Teague R, et al. Complex Creative Systems. Rangelands. 2013;35: 6–13. doi:10.2111/RANGELANDS-D-13-00013.1

7. McLane AJ, Semeniuk C, McDermid GJ, Marceau DJ. The role of agent-based models in wildlife ecology and management. Ecol Model. 2011;222: 1544–1556. doi:10.1016/j.ecolmodel.2011.01.020

8. Grimm V. Ten years of individual-based modelling in ecology: what have we learned and what could we learn in the future? Ecol Model. 1999;115: 129–148. doi:10.1016/S0304-3800(98)00188-4

9. Grimm V, Revilla E, Berger U, Jeltsch F, Mooij WM, Railsback SF, et al. Pattern-Oriented Modeling of Agent-Based Complex Systems: Lessons from Ecology. Science. 2005;310: 987–991. doi:10.1126/science.1116681

10. Northrop RB. Introduction to Complexity and Complex Systems. CRC Press; 2010.

11. Dumont B, Hill DRC. Spatially explicit models of group foraging by herbivores: what can Agent-Based Models offer? Anim Res. 2004;53: 419–428. doi:10.1051/animres:2004028

12. Funtowicz SO, Ravetz JR. Science for the post-normal age. Futures. 1993;25: 739–755. doi:10.1016/0016-3287(93)90022-L

13. Bailey DW, Provenza FD. Mechanisms Determining Large-Herbivore Distribution. In: Prins HHT, Langevelde FV, editors. Resource Ecology. Springer Netherlands; 2008. pp. 7–28. doi:10.1007/978-1-4020-6850-8_2

14.   Distel RA, Laca EA, Griggs TC, Demment MW. Patch selection by cattle: maximization of
intake rate in horizontally heterogeneous pastures. Appl Anim Behav Sci. 1995;45: 11–21.
doi:10.1016/0168-1591(95)00593-H

15.   Laca EA, Distel RA, Criggs TC, Demment MW. Effects of Canopy Structure on Patch
Depression by Grazers. Ecology. 1994;75: 706–716. doi:10.2307/1941728

16.   Sato S. Leadership during actual grazing in a small herd of cattle. Appl Anim Ethol. 1982;8:
53–65. doi:10.1016/0304-3762(82)90132-8

17.   Shiyomi M. How are distances between grazing cows determined: A case study. Appl Entomol
Zool. 2004;39: 575–581. doi:10.1303/aez.2004.575

18.   WallisDeVries MF, Laca EA, Demment MW. From feeding station to patch: scaling up food
intake measurements in grazing cattle. Appl Anim Behav Sci. 1998;60: 301–315.
doi:10.1016/S0168-1591(98)00158-0

19.   Green BT, Welch KD, Pfister JA, Chitko-McKown CG, Gardner DR, Panter KE. Mitigation
of Larkspur Poisoning on Rangelands Through the Selection of Cattle. Rangelands. 2014;36:
10–15. doi:10.2111/RANGELANDS-D-13-00031.1

20.   Pfister JA, Gardner DR, Panter KE. Consumption of Low Larkspur (Delphinium
nuttallianum) by Grazing Sheep. Rangel Ecol Manag. 2010;63: 263–266. doi:10.2111/REM-D-
09-00084.1

21.   Pfister JA, Manners GD, Ralphs MH, Hong ZX, Lane MA. Effects of Phenology, Site, and
Rumen Fill on Tall Larkspur Consumption by Cattle. J Range Manag. 1988;41: 509–514.
doi:10.2307/3899528

22.   Ralphs MH, Olsen JD, Pfister JA, Manners GD. Plant-animal interactions in larkspur
poisoning in cattle. J Anim Sci. 1988;66: 2334–2342.

23.   Pfister JA, Provenza FD, Panter KE, Stegelmeier BL, Launchbaugh KL. Risk Management to
Reduce Livestock Losses from Toxic Plants. J Range Manag. 2002;55: 291–300.
doi:10.2307/4003137

24.   Pfister JA, Provenza FD, Manners GD, Gardner DR, Ralphs MH. Tall Larkspur Ingestion:
Can Cattle Regulate Intake Below Toxic Levels? J Chem Ecol. 1997;23: 759–777.
doi:10.1023/B:JOEC.0000006409.20279.59

25.   Smith MA, Waggoner, Jr. JW, Sims S. Larkspur: Managing grazing to avoid poisoning cattle.
University of Wyoming Cooperative Extension; 2010 Jul. Report No.: MP-111.13.

26.   Panter KE, Manners GD, Stegelmeier BL, Lee S, Gardner DR, Ralphs MH, et al. Larkspur
poisoning: toxicology and alkaloid structure–activity relationships. Biochem Syst Ecol. 2002;30:
113–128. doi:10.1016/S0305-1978(01)00123-5

27.   Grina JA, Schroeder DR, Wydallis ET, Stermitz FR. Alkaloids from Delphinium geyeri. Three
new C20-diterpenoid alkaloids. J Org Chem. 1986;51: 390–394. doi:10.1021/jo00353a023

28.   Grimm V, Berger U, DeAngelis DL, Polhill JG, Giske J, Railsback SF. The ODD protocol: A
646   review and first update. Ecol Model. 2010;221: 2760–2768.
647   doi:10.1016/j.ecolmodel.2010.08.019

648   29.   Allen VG, Batello C, Berretta EJ, Hodgson J, Kothmann M, Li X, et al. An international
649   terminology for grazing lands and grazing animals. Grass Forage Sci. 2011;66: 2–28.
650   doi:10.1111/j.1365-2494.2010.00780.x

651   30.   Shiyomi M, Tsuiki M. Model for the spatial pattern formed by a small herd in grazing cattle.
652   Ecol Model. 1999;119: 231–238. doi:10.1016/S0304-3800(99)00059-9

653   31.   Wilensky U. Netlogo 6.01 [Internet]. Evanston, IL: Center for Connected Learning and
654   Computer-Based Modeling, Northwestern University; 1999. Available:
655   http://ccl.northwestern.edu/netlogo

656   32.   Grimm V. Visual Debugging: A Way of Analyzing, Understanding and Communicating
657   Bottom-up Simulation Models in Ecology. Nat Resour Model. 2002;15: 23–38.
658   doi:10.1111/j.1939-7445.2002.tb00078.x

659   33.   Hexagon Geospatial. ERDAS IMAGINE 2016. Madison, Alabama; 2016.

660   34.   Huete AR. A soil-adjusted vegetation index (SAVI). Remote Sens Environ. 1988;25: 295–309.
661   doi:10.1016/0034-4257(88)90106-X

662   35.   ESRI. ArcGIS Desktop 10.4. Redlands, CA: ESRI; 2015.

663   36.   ESRI. Collector for ArcGIS 17.01. Redlands, CA: ESRI; 2017.

664   37.   Bonham CD. Measurements for Terrestrial Vegetation. Wiley; 1989.

665   38.   Bailey DW. Daily selection of feeding areas by cattle in homogeneous and heterogeneous
666   environments. Appl Anim Behav Sci. 1995;45: 183–200. doi:10.1016/0168-1591(95)00586-H

667   39.   Harris NR, Johnson DE, McDougald NK, George MR. Social Associations and Dominance of
668   Individuals in Small Herds of Cattle. Rangel Ecol Manag. 2007;60: 339–349. doi:10.2111/1551-
669   5028(2007)60[339:SAADOI]2.0.CO;2

670   40.   National Research Council. Nutrient Requirements of Beef Cattle: Seventh Revised Edition:
671   Update 2000 [Internet]. Washington, D.C.: National Academies Press; 2000.
672   doi:10.17226/9791

673   41.   Holechek JL, Pieper RD, Herbel CH. Range management: principles and practices. 6th ed.
674   Pearson Education, Inc.; 2011.

675   42.   Senft RL, Coughenour MB, Bailey DW, Rittenhouse LR, Sala OE, Swift DM. Large Herbivore
676   Foraging and Ecological Hierarchies. BioScience. 1987;37: 789–799. doi:10.2307/1310545

677   43.   Lazo A. Social segregation and the maintenance of social stability in a feral cattle population.
678   Anim Behav. 1994;48: 1133–1141. doi:10.1006/anbe.1994.1346

44. Meuret M, Provenza F. The Art and Science of Shepherding: Tapping the Wisdom of French
Herders. Acres U.S.A., Incorporated; 2014.

45. Welch KD, Green BT, Gardner DR, Cook D, Pfister JA. The effect of administering multiple
doses of tall larkspur (Delphinium barbeyi) to cattle. J Anim Sci. 2015;93: 4181.
doi:10.2527/jas.2015-9101

46. Gardner DR, Pfister JA. Toxic alkaloid concentrations in Delphinium nuttallianum,
Delphinium andersonii, and Delphinium geyeri in the Intermountain Region. Rangel Ecol
Manag. 2007;60: 441–446. doi:10.2111/1551-5028(2007)60[441:TACIDN]2.0.CO;2

47. SAS Institute. JMP. Cary, NC: SAS Institute, Inc.; 2016.

48. R Core Team. R: a language for statistical computing [Internet]. Vienna, Austria: R Foundation
for Statistical Computing; 2017. Available: https://www.R-project.org

49. Faraway JJ. Extending the Linear Model with R: Generalized Linear, Mixed Effects and
Nonparametric Regression Models, Second Edition. CRC Press; 2016.

50. Hilbe JM. Negative Binomial Regression. Cambridge University Press; 2011.

51. Anderson DR. Model based inference in the life sciences: a primer on evidence. New York ;
London: Springer; 2008.

52. Rafter J, Abell M, Braselton J. Multiple Comparison Methods for Means. SIAM Rev. 2002;44:
259–278. doi:10.1137/S0036144501357233

53. Walker JW, Heitschmidt RK, Dowhower SL. Evaluation of Pedometers for Measuring
Distance Traveled by Cattle on Two Grazing Systems. J Range Manag. 1985;38: 90–93.
doi:10.2307/3899343

54. Nielsen DB, Ralphs MH, Evans JO, Call CA. Economic feasibility of controlling tall larkspur
on rangelands. J Range Manag Arch. 1994;47: 369–372.

55. Provenza FD, Villalba JJ, Dziba LE, Atwood SB, Banner RE. Linking herbivore experience,
varied diets, and plant biochemical diversity. Small Rumin Res. 2003;49: 257–274.
doi:10.1016/S0921-4488(03)00143-3

56. Ralphs MH, Jensen DT, Pfister JA, Nielsen DB, James LF. Storms Influence Cattle to Graze
Larkspur: An Observation. J Range Manag. 1994;47: 275–278. doi:10.2307/4002547

57. Laycock WA. Alkaloid Content of Duncecap Larkspur after Two Years of Clipping. J Range
Manag. 1975;28: 257–259. doi:10.2307/3897769

58. Bailey DW, Trotter MG, Knight CW, Thomas MG. Use of GPS tracking collars and
accelerometers for rangeland livestock production research. J Anim Sci. 2017;95: 360–360.
doi:10.2527/asasann.2017.740

59.   Barnes M. Livestock Management for Coexistence with Large Carnivores, Healthy Land and
Productive Ranches. Keystone Conservation; 2015.

60.   Anderson DM, Estell RE, Holechek JL, Ivey S, Smith GB. Virtual herding for flexible
livestock management – a review. Rangel J. 2014;36: 205–221. doi:10.1071/RJ13092

716