AQMM: Enabling Absolute Quantification of Metagenome and Metatranscriptome

Authors: Xiao-Tao Jiang, Ke Yu, Li-Guan Li, Xiao-Le, Yin, An-Dong, Li, Tong Zhang*

Affiliations: Environmental Biotechnology Laboratory, Department of Civil Engineering, University of Hong Kong

* Corresponding author Email: zhangt@hku.hk

November 12, 2017

Abstract

Metatranscriptome has become increasingly important along with the application of next generation sequencing in the studies of microbial functional gene activity in environmental samples. However, the quantification of target active gene is hindered by the current relative quantification methods, especially when tracking the sharp environmental change. Great needs are here for an easy-to-perform method to obtain the absolute quantification. By borrowing information from the parallel metagenome, an absolute quantification method for both metagenomic and metatranscriptomic data to per gene/cell/volume/gram level was developed. The effectiveness of AQMM was validated by simulated experiments and was demonstrated with a real experimental design of comparing activated sludge with and without foaming. Our method provides a novel bioinformatic approach to fast and accurately conduct absolute quantification of metagenome and metatranscriptome in environmental samples. The AQMM can be accessed from https://github.com/biofuture/aqmm.

Keywords: metagenome, metatranscriptome, absolute quantification, differential expression genes

Background

Shotgun metatranscriptomics is a powerful tool in identifying the overall expression of microorganisms in an environment (Alexander et al. 2015, Gifford et al. 2011, Shi et al. 2009, Turner et al. 2013, Yu and Zhang 2012), shedding light on discovering how microbes respond to environmental changes or diseases status (Jorth et al. 2014, Mason et al. 2012) and capturing gene expression patterns for functionally important bacteria in engineering systems (Oyserman et al. 2015, Stark et al. 2014). For these applications, accurate quantification is required to detect the true variations or differential expression genes (DEGs).

Traditionally, the abundance of a transcript in RNA-sequencing (RNA-seq) is thought to be influenced by the library size and inherent dependence on the expression levels of other transcripts as described in a comprehensive review (Rapaport et al. 2013). Following this idea, transcripts in RNA-seq was generally quantified by within-sample normalization. One of the most common quantification methods was RPKM (Mortazavi et al. 2008) (reads per kilobase of exon model per million mapped reads) which considered factors of both the length of gene and library size. Another improved within-sample normalization method was TPM (transcript per million) (Wagner et al. 2012) which only considered the transcript rather than the whole library size and respected the invariance of relative molar RNA concentration (rmc). The TPM was thought to be better fitted in sample comparison due to its unit-free characteristics. The FPKM (substitute the reads with fragments in RPKM) was an adaption of RPKM to pair-end reads. These above methods are all relative quantification (RQ) and suffer from the 'composition effects' (the increase of one transcript will decrease other unrelated transcript). To relieve this problem, Robinson and Oshlack proposed a new normalization method "TMM" (trimmed mean of

M-values) to detect the DEGs under the hypothesis that most of the genes are not

differentially expressed (Robinson and Oshlack 2010), which has been integrated into

popular DEGs detection R software edgeR (Robinson et al. 2010). The scaling factor

in edgeR for normalization is the TMM value. Another method was to compute the

median of the ratio as the scaling factor and it could be conducted by R software

DESeq/DESeq2 (Love et al. 2014). It is also based on the assumption that most genes

are not DEGs and this method then calculates the scaling factor (median of ratios)

associated with this sample to perform further normalization. In the two software, the

negative binomial distribution was applied to adjust the distribution of transcript

between different conditions to relieve the dispersion effects of deviation from

standard passion distribution (Rapaport et al. 2013). Although with these efforts in

optimizing the normalization process, these indices were all still RQ based and the

relationship could be distorted while performing comparative analysis across samples,

especially when borrowing these methods from traditional Eukaryote RNA-seq to

current Prokaryote metatranscriptome studies (Conesa et al. 2016). One feasible way

to solve the problem was to get the absolute quantification (AQ) of expression level

for each transcript. For example, the qRT-PCR has long been applied in RNA-seq or

microarray data for AQ (Becker-André and Hahlbrock 1989, Whelan et al. 2003). In

addition, there were methods by spiking in exterior/alien RNA in microarray to get the

per cell absolute quantification (Kanno et al. 2006) and internal standard approach to

estimate per liter expression in marine metatranscriptome (Gifford et al. 2011).

However, the experiment to perform spiking internal standard was difficult due to its

skill-demanding nature and for metatranscriptome data, factors like the time to add

spike-in material, the type and the amount of alien RNA required still needed to be

elaborately designed. Hence, it was not as popular as those RQ methods. The

quantification methods in the newly developed analyzing pipelines for

metatranscriptome like IMP (Narayanasamy et al. 2016), MetaTrans (Martinez et al.

2016), COMAN (Ni et al. 2016) and SAMSA (Westreich et al. 2016) were still all

based on RQ methods; this would result in accelerated spreading of the inaccurate

quantification in many studies.

To solve the problem of RQ and get an accurate quantification without performing

spike-in experiment, an AQ bioinformatics software package AQMM was developed

by combining metagenome and metatranscriptome data to achieve the goal of

accurate and comparable quantification. In this study, we firstly introduced the

AQMM algorithm flow, and then compared and validated it with RQ methods by

simulated metagenome and metatranscriptome data. Moreover, we further applied this

algorithm to a real combination of metagenome and metatranscriptome dataset in

quantifying genes and transcripts of resistome in six foaming activated sludge (FAS)

and non-foaming activated sludge (NFAS) samples.

Results

Overall view of AQMM algorithm

The AQMM **(Fig. 1)** was designed to perform AQ of parallel metagenome and

metatranscriptome dataset no matter whether spike-in experiment/internal standard

was initially added or not. The major aims were to obtain the AQ of

genes/transcripts/taxa in samples and to accurately detect DEGs in metatranscriptome

data. The assumptions under the algorithm include: 1) with the known extraction ratio

of DNA for a DNA extraction Kit for a type of sample, the total weight of DNA per

volume of the sample could be calculated. The weight of the sequenced library of

DNA could be estimated with the molecular weight and bases numbers of A, T, C and

G in the sample. Then, the ratio of sequenced DNA to total weight of DNA per

volume of the sample could be calculated. In addition, by utilizing the universal

single-copy phylogenetic marker genes (USCMGs)**,** the number of cells for a

metagenome library could be estimated accurately (Nayfach and Pollard 2015). With

the above information, cells per volume could be calculated for a metagenome data; 2)

Using the same volume of the sample contains the same number of cells for DNA and

RNA extraction, the cell number per volume to extract RNA was the same as the

parallel DNA sample; 3) With the known ratio of RNA extraction and the rRNA ratio

of total RNA, by a similar process, the sequenced RNA weight ratio could be

calculated, and then the equivalent cell numbers in a metatranscriptome could be

deduced accordingly. With the cell numbers included in the metagenome and

metatranscriptome data, the abundance of gene/transcript could be normalized to per

cell level. Moreover, as the number of cells per volume is available, per cell

quantification could be easily transformed into per volume quantification.

Comparing and validating AQMM with RQ methods using simulated

metatranscriptome demo

To reveal the problem of RQ methods like RPKM, edgeR and DESeq2 and to assess

the effectiveness of AQMM, simulated metatranscriptomic datasets comprised of

known community structure and expression levels were generated **(Fig. 2; Details in**

methods). The simulated data was with known ground truth absolute expression for

each gene. For simplicity, to focus on the quantification of metatranscriptome in

identifying DEGs, we assume the DNA content are not changed like what happens in

a reactor with a stable biomass concentration, however the gene expression under

condition A and B are significantly changed with fold of 2 or 16 in part of the bacteria

like what happens under sharp environmental change. In order to focus only on the of

influence normalization methods, in generation of the simulated metatranscriptome,

the base qualities of were all set with 50 and to eliminate the influence of mapping

process, the mapping criteria of bowtie2 was set to exactly match without gap and

mismatch allowed (bowtie2 parameters, -N 0 -L 31, --rdg 100,150 --rfg 100,150

--gbar 100,150). The result of DEGs detection was in **Table 1**. We can observe that

compared with ground truth, the RQ methods detect quite a large portion of false

positive higher gene expression under condition A. On the contrary, the AQMM

method which aims to obtain the AQ has limited errors detection even with a given

variance in RNA extraction efficiency **(Table 1)**. Noticeably, in real combination of

metagenome and metatranscriptome, the metagenome could also be totally different,

and in this case, the AQMM is still applicable.

Case study: AQ of activate resistome in FAS and NFAS

The AQMM was applied in the six metagenome and metatranscriptome dataset of

FAS and NFAS, the AQ of the sequenced cells generated by the pipeline were shown

in **Table 2**. In detail, the metagenome contained 8 to 11.8 GBs data and

metatranscriptome with a depth between 13 and 16 GBs for each sample. The "per

cell/volume" quantifying values were the fundamental of normalizing to cells or

volume in order to perform comparison among different studies. The cell number per

milliliter in literature was at 3.3E+09 using flow Cytometer to quantify(Foladori et al.

2010) and was from 2.1E+09 to 5.5E+09 using CFU and flow Cytometer (Manti et al.

2008) level for AS which was a bit lower than the obtained number in this study at the

magnitude of E+10 cells per milliliter**.** Overall number of mRNA molecules per cell

are 387.98 ± 102.86 and 235.21 ± 30.59 averagely for FAS and NFAS, respectively

(Table S1), which is consistent with previous observation of coastal bacterioplankton

by 142-238 mRNA molecules per Cell (Gifford et al. 2011, Moran et al. 2013).

As WWTPs become the hot-spot of antibiotic resistant genes (ARGs) to the receiving environment. Hence, the expressions of ARGs in the AS were in great concerns and further profiled. Overall, the abundance of ARGs per cell in FAS and NFAS were 0.0517 ± 0.0034 and 0.0483 ± 0.0041; and the transcript of ARGs per cell were 0.0140 ± 0.0039 and 0.0059 ± 0.0009, respectively **(Table S2 & S3)**. The overall transcription of ARGs was significantly higher in FAS compared with NFAS. At DNA level, only tetracycline resistance gene was higher in FAS and beta-lactam was higher in NFAS, other types were not significantly different. However, at transcript level, all the types were all significantly higher in FAS. Among the nine transcribed ARGs types, beta-lactam and sulfonamide resistance genes were the most abundant expressed ARG types in both FAS and NFAS. Per volume ARGs abundance and expression at type level were shown in **Fig. 3**. The overall ARGs abundance per milliliter AS in FAS and NFAS were $2.51E+09 \pm 2.44E+08$ and $2.66E+09 \pm 5.63E+08$; and the transcript of ARGs per milliliter were $9.83E+09 \pm 3.82E+08$ and $4.49E+09 \pm 5.10E+08$, respectively. With the AQ results, the transcripts per copy gene (TPCG), which represents of the transcribe rate could be further derived. The unclassified, quinolone, multidrug and beta-lactam were more active in FAS compared with NFAS in terms of TPCG, **(Table S4)**. For the detected ARGs, the host taxonomy was assigned by LCA algorithms using all the genes annotation in the same Contig. Thirteen orders were detected to carry ARGs and eleven of them were transcribed (**Fig. 3**). The most ARGs transcribed order was Enterobacteriales. The active ARGs in bacteria enclosed in foams of FAS posed potential threats for the public as ARGs carrying bacteria could spread into the air from the foams bubbles.

The co-expression of ARGs and MRGs was also studied to check whether there were

co-expression effects at the RNA level. Using this dataset, we observed co-expression

within ARGs, within MRGs, and between ARGs and MRGs (**Fig. 4**). Numerous types

of MRGs were detected in the metagenome and metatranscriptome. The most

abundant MRG was Cu resistant genes and for the ARGs, beta-lactam, tetracycline

and aminoglycoside were the most expressed types. The highest number of

co-expression within MRGs was Cr and Fe; while within ARGs was beta-lactam and

tetracycline. The most MRG and ARGs co-expression was Cr, which co-expression

with nine types of ARGs. This was the first transcript level evidence of the

co-expression of ARGs and MRGs in AS.

Discussion

Metatranscriptome enabled the study of whole metabolic pathways expression of the

system and many studies had already taken this advantage for different environments,

such as in marine (Mason et al. 2012), rhizosphere of the plant (Turner et al. 2013),

human oral disease (Jorth et al. 2014). Each study has specific method to integrate the

metagenome and metatranscriptome information to understand the microbes and their

activities in the system. The quantification of metatranscriptome was generally RQ

based methods. The RQ methods are problematic as they may not be able to reflect

the actual expression level of a population in the whole community. Due to the

relative characteristics, the RQ methods are always suffer from the so-called

composition effects, which indicates that the upgrade of one gene should definitely

make other genes downgrade. Additionally, the RQ methods are just a relative portion

rather than a value with biological implications. On the contrary, the AQ could be

more biological meaningful at per cell/volume unit. Hence, it was necessary to

conduct AQ to compare different samples. In this study, we proposed an AQ method

and developed a set of algorithms to conveniently calculate the absolute number of

sequenced cells for each RNA library by borrowing cell numbers from a

corresponding data set of DNA library of the same sample.

Noticeably, there were several hypotheses for the application of the proposed method.

Firstly, the sample used to extract DNA and RNA should contain the same cell

numbers per volume which could be easily met with sufficient mixing of samples.

Secondly, the DNA and RNA extraction efficiency should be estimated, as well as the

rRNA ratio in total RNA. This was likely difficult to achieve. However, for an

environmental sample, generally literature based data could be used for the extraction

kit, for example, to FastDNA SPIN Kit for Soil, the extract efficiency was estimated

as 28.4% (Mumy and Findlay 2004). Most importantly, as the parallel samples were

extracted under the same condition, the difference between samples was minimized

(DNA extraction data, unpublished). This AQMM method is capable of performing

absolute quantification of both metagenome and metatranscriptome without the

requirement to do complex spike-in experiments. Importantly, AQMM avoids the RQ

problems of composition effects and able to detect accurate DEGs. Hence, the

proposed AQMM is a method in between experimental spike-in based AQ methods

and those improved RQ methods of TMM based edgeR.

With AQ, a number of indices with various biological meaning were proposed in this

study (Methods), for example, the transcript per copy gene (TPCG) index is a

reflection of the transcribe rate of the gene, which could never be delivered by RQ

methods. It was demonstrated with simulating RNA-seq that the organism abundance

(community structure) was important at normalizing metatranscritptome data in

identifying DEGs (Klingenberg and Meinicke 2017). The gene per cell (GPC) and

transcript per cell (TPC) in AQMM are global level normalization indices and the

scaling factor is the total number of cells in the DNA or RNA library. This global

scaling factor could be easily transformed into taxa specific scaling factors with the

relative quantification of different taxa with indices of transcript of taxon A per cell

(TTPC). Hence, the normalization in AQMM is well fit for the factor of microbial

abundance in metatranscriptome data.

AS is important biological wastewater treatment process and this system is considered

as a hot spot for ARG dissemination into the receiving water. The foaming of AS

would result in spreading of foams with AS bacteria into the surrounding environment.

Understanding the active resistome and the host bacteria in foaming AS enables

engineers understanding the risk of sludge foaming incurred to the surrounding

environment. We observed a wide profile of active ARG types in the FAS, the

identification of opportunity pathogen bacteria *Pseudomonas* carrying active ARGs

alerts us the risk of spreading ARGs-carrying bacteria. Additionally, per cell mRNA

molecules is an important indication of the activity of the cell, generally natural

bacterial communities was observed to hold a lower inventory of transcripts (Moran et

al. 2013); and the absolute quantification obtained with AQMM was well-fitted with

previous observation.

Conclusions

In this study, we filled the gap of lacking a bioinformatic algorithm to perform AQ of

metatranscriptomic data. The developed AQMM was demonstrated to gain enhanced

performance at identifying DEGs compared with those RQ methods benchmarked

with simulated metagenomic and meatranscriptomic data. Additionally, with the

AQMM, the active resistome in foaming and normal activated sludge were quantified

to per cell/volume level and even down to the transcription per copy gene. The active

ARG host were quantified and the co-expression of MRGs and ARGs was revealed

for the first time in AS.

Materials and methods

Absolute quantification of gene abundance and transcript expression

We developed a package of scripts AQMM (absolute quantification of metagenome

/metatranscriptome) to perform comparative analysis.

The formula for cells per mL:

$$\boldsymbol{C} = N_c / \frac{L_{size} * 10^9 * \frac{(R_A * 313.2 + R_T * 304.2 + R_C * 289.18 + R_G * 329.21)}{6.022 * 10^{23}}}{X/\alpha} \qquad (1)$$

C is value of cell numbers per mL AS

N_c is the estimated cell numbers for the sequenced DNA library with USCMGs

L_{size} is the sequencing depth

R_A, R_T, R_C and R_G are ratios of A, T, C and G

X is the overall extracted weight (ng) of DNA for 1 mL AS

α is DNA extraction efficiency, for FAST DNA Kit for Soil, α is estimated as 28.2%

(Mumy and Findlay 2004).

The sequenced cells for RNA sequencing, for a RNA-seq with library size of L_{size}

after removing all ribosomal RNA, the equivalent sequenced cells for this sample is

$$\boldsymbol{E_c} = C * \frac{L_{size} * 10^9 * \frac{R_A * 329.2 + R_U * 306.2 + R_C * 305.2 + R_G * 345.2}{6.022 * 10^{23}}}{Y * \gamma/\beta} \qquad (2)$$

E_c is the estimated number of cells sequenced for this RNA library

C is value of cell numbers per mL AS

L_{size} is the sequencing depth

R_A, R_U, R_C and R_G are ratios of A, U, C and G, the value they multiplied are molecular

weight

Y is the overall extracted weight (ng) of RNA for 1 mL AS

β is RNA extraction efficiency, the estimated β is about 7.5% as used in this study.

This value was deduced from AS empirical data of proportion of RNA biomass by

engineering perspective and the extracted RNA biomass.

γ is non-ribosomal RNA ratio, for AS the estimated γ is about 0.03.

Based on the two AQ numbers of cells for each sample, the gene or transcript

abundance matrix could be further normalized into the following indices.

GPC (Gene per Cell): an indication of the overall abundance of the gene in system.

$\text{GPC} = \dfrac{N_{read}*L_{read}/L_{gene}}{N_c}$ (3)

TPC (Transcript per Cell): an indication of overall activity of the gene in system.

$\text{TPC} = \dfrac{N_{read}*L_{read}/L_{gene}}{E_c}$ (4)

TPCG (Transcript per copy gene): an indication of the absolute activity of one copy

gene in the system, equivalent to transcribe rate for each gene.

$\text{TPCG} = {\text{TPC}}/{GPC}$ (5)

GTPC (Gene of taxon A per Cell): an indication of the overall abundance of the taxon

in system averagely.

$\text{GTPC} = \sum_{i=1}^{n} GPC_i$ (6)

TTPC (Transcript of taxon A per Cell): an indication of overall activity of the taxon

in system averagely.

$\text{TTPC} = \sum_{i=1}^{n} TPC_i$ (7)

ATCT (Averagely transcript per copy gene of taxon A): indication of the averagely

absolute activity per copy expressed gene in taxon A

$\text{ATCT} = \dfrac{1}{n}\sum_{i=1}^{n} TPCG_i$ (8)

N_c is the estimated cell numbers for the sequenced DNA library,

N_{read} is the number of reads or transcript mapping to the target gene

L_{read} is the length of reads

L_{gene} is the length of the target gene

n is the number of genes affiliated to taxa A.

When the number of cells per mL was obtained, using the GPC, genes per mL could

be calculated.

Simulating metatranscriptome data

To validate our method and comparing with those RQ methods in identifying the

DEGs, simulated data was generated by workflow illustrated in **Fig. 2**. For simplicity,

the DNA was set unchanged to mimic the activated sludge community composition

with 16 strains from different phylogeny. The metatranscriptome data sets were

generated for two conditions A and B, each with three biological duplications; for the

condition A and B, there were part of the strains with folds of significantly changed

expression (**Table S5**). To only focus on the quantification method, all the system

errors caused by other factors like base qualities, cDNA synthesis, assembly, mapping

parameters were not considered.

Sampling

AS samples were collected in Shatin wastewater treatment plant at three locations

along the flow direction while serious foaming happened at 2016-04-08 and nearly no

foaming happened at 2016-04-25. Samples were collected on site by storing in liquid

nitrogen immediately and then transported to the laboratory for RNA extraction. The

DNA samples were mixed with 1:1 100% ethanol and AS and then stored at -20 °C

fridge. Totally six samples were collected for both DNA and RNA samples alongside

the segment aeration tank in three locations as depicted in **Fig. 5**.

Whole DNA, total RNA extraction, removal of ribosomal RNA, cDNA synthesis

and next generation sequencing

FAST DNA Kit was used to extract total DNA from 1 mL mixed AS samples. RNeasy
Mini was used to extract the total RNA from 0.5 mL AS stored in liquid nitrogen. The
extracted RNA was then processed by DNase I to eliminate the DNA in the RNA
samples. Then both Illumina Ribo-Zero rRNA removal KIT (Bacteria) and Ribo-Zero
rRNA removal KIT (Human/Mouse/Rat) was applied for each sample to remove
rRNA from Prokaryote and Eukaryote respectively in order to get the total clean
non-ribosomal RNA. Generally, metatranscriptome rRNA depletion was only used the
Ribo-Zero for Bacteria, in this study, the addition of Eukaryote rRNA removal was
due to a fact that by only using the Ribo-Zero Bacteria rRNA removal Kit for AS,
there was still over half of RNA were rRNA from Eukaryote (our previous experiment,
data unpublished). To get more non-rRNA, the Ribo-Zero rRNA Kit to remove
Eukaryote was also used. RNA then was fragmented into 170 bps library and was
reverse-transcribed to construct cDNA library for sequencing. The quality of DNA
and RNA were assessed with Agilent 2100 Bioanalyzer (Agilent Technologies, Palo
Alto, CA, USA). All the samples was sent to sequence, considering the complexity of
AS and the aims of this study to detect the expression of low abundance gene, we
gave each sample a very deep sequencing depth which doubled the sequencing depth
in previous studies. All the samples were sequenced with Hiseq 4000 in
BGI-ShenZhen. DNA samples with PE-150 with library size of 300 bps. And RNA
with PE101 of library size 170 bps.

Bioinformatics analysis

Quality filtering was firstly performed on DNA and RNA reads to keep only high
quality reads using trimmomatic v1.04 (Bolger et al. 2014). DNA datasets were

pooled together and assembled by CLC Genomics Workbench 6.5.3 (CLC Bio,

Aarhus, Denmark, https://www.qiagenbioinformatics.com/) with default parameters.

Finally, 1,430,611 contigs with length over 100 bps (N50, 2,416 bps; 2,457,704,443

bps length in total) were obtained and 74.5% of reads could be mapped back to these

Contigs. All these contigs were sent to predict genes with Prodigal (version 1.5)

(Hyatt et al. 2010) using `-meta` parameter and finally 3,234,330 genes were obtained.

By removing exactly the same genes using USEARCH (version 8.0.1623) (Edgar

2010) unique command (parameters -fastx_uniques), 3,234,246 million genes were

kept; this set was defined as 'unique gene set'. Reads were mapped back to the contig

set and 'unique gene set' to obtain reads coverage matrixes for contigs and genes. The

matrix of genes was finally normalized to cell numbers. For metatranscriptome

samples, after quality filtering, the SortMeRNAv1.9 was used to remove all the

possible ribosomal RNA by aligning to six databases of bacteria, archaea and

eukaryotic small and large subunits (Kopylova et al. 2012). RNA reads for each

sample were then mapped back to the `unique gene set` to get the transcript coverage

for each gene with CLC genomic workbench 6.5.3 using parameters of gap penalty 2,

gap extension 3, length fraction 0.8 and similarity at least 0.9.

Taxonomy composition of the metagenome was generated with MEGAN6 (Huson et

al. 2015). In detail, all genes were aligned to NCBI NR database (version 201603)

with diamondv1.09 (Buchfink et al. 2015) to find out the homology proteins. To each

gene, the local common ancestors (LCA) were applied using the taxonomy

information of the hit NR protein in NCBI taxonomy database (Acland et al. 2014)

and then this gene was annotated with the common ancestor taxonomy. We further

processed the NCBI taxonomy annotation results to remove those subdivisions and

subgroups to format the annotation to 7 levels from kingdom to species. Among total

3,234,246 unique genes predicted, 2,348,907 could be aligned to NR database. The

remaining 885,339 (27.3%) genes could not be annotated with the NR database. The

abundance of each taxon was a sum of all the annotated genes under that taxon in

every sample. Antibiotic resistant genes (ARGs) were annotated with SARG database

which contained a type-subtype structure annotation (Yang et al. 2016). Metal

resistance genes (MRGs) were detected by aligning the "unique gene set" to the MRG

database (Li et al. 2017). Absolute abundance and transcript was determined by

AQMM.

Declarations

Data availability

The metagenome and metatranscriptome raw data were deposited in NCBI SRA under

accession number XXX.

Analyzing document

The analyzing document for the whole data analysis and simulation process could be

accessed from

https://github.com/biofuture/aqmm/blob/master/Analysing_document.txt

Conflict of interest

386 The authors declare no conflict of interest

Acknowledgements

388 The authors would like to thank GRF of Hong Kong for financial support

389 (172099/14E). Xiaotao Jiang would like to thank The University of Hong Kong for

390 the postgraduate scholarship, and Dr. Ke Yu, Dr. Li-Guan, and Dr. Andong, Li, would

391 like to thank The University of Hong Kong for postdoc fellowship.

Contributions

T. Zhang and X.-T. Jiang design the study of quantification. X.-T. Jiang developed the software and performed the wet-lab and simulation experiments. X.-T. Jiang performed the bioinformatics analyses. X.-T. Jiang, A.D. Li and K. Y. did the DNA and RNA extraction experiment. L.-G. Li did the MRG analyses. T. Zhang and X.-T. Jiang wrote the manuscript. T. Zhang, X.-T. Jiang, A.D. Li, L.G. Li and X.L. Yin revised the manuscript.

Reference

Acland, A., Agarwala, R., Barrett, T., Beck, J., Benson, D.A., Bollin, C., Bolton, E., Bryant, S.H., Canese, K. and Church, D.M. (2014) Database resources of the national center for biotechnology information. Nucleic acids research 42(Database issue), D7.

Alexander, H., Jenkins, B.D., Rynearson, T.A. and Dyhrman, S.T. (2015) Metatranscriptome analyses indicate resource partitioning between diatoms in the field. Proceedings of the National Academy of Sciences 112(17), E2182-E2190.

Becker-André, M. and Hahlbrock, K. (1989) Absolute mRNA quantification using the polymerase chain reaction (PCR). A novel approach by a P CR aided t ranscipt t itration assay (PATTY). Nucleic acids research 17(22), 9437-9446.

Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics, btu170.

Buchfink, B., Xie, C. and Huson, D.H. (2015) Fast and sensitive protein alignment using DIAMOND. Nature Methods 12(1), 59-60.

Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A., Szcześniak, M.W., Gaffney, D.J., Elo, L.L. and Zhang, X. (2016) A survey of best practices for RNA-seq data analysis. Genome biology 17(1), 13.

Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26(19), 2460-2461.

Foladori, P., Bruni, L., Tamburini, S. and Ziglio, G. (2010) Direct quantification of bacterial biomass in influent, effluent and activated sludge of wastewater treatment plants by using flow cytometry. Water Research 44(13), 3807-3818.

Gifford, S.M., Sharma, S., Rinta-Kanto, J.M. and Moran, M.A. (2011) Quantitative analysis of a deeply sequenced marine microbial metatranscriptome. The ISME journal 5(3), 461-472.

Huson, D., Beier, S., Buchfink, B., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.-J. and Tappu, R. (2015) MEGAN6-Microbiome analysis involving hundreds

of samples and billions of reads, preparation.

Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W. and Hauser, L.J. (2010)
Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC
bioinformatics 11(1), 119.

Jorth, P., Turner, K.H., Gumus, P., Nizam, N., Buduneli, N. and Whiteley, M. (2014)
Metatranscriptomics of the human oral microbiome during health and disease. MBio 5(2),
e01012-01014.

Kanno, J., Aisaki, K.-i., Igarashi, K., Nakatsu, N., Ono, A., Kodama, Y. and Nagao, T. (2006)
" Per cell" normalization method for mRNA measurement by quantitative PCR and
microarrays. BMC genomics 7(1), 64.

Klingenberg, H. and Meinicke, P. (2017) How To Normalize Metatranscriptomic Count Data
For Differential Expression Analysis. bioRxiv, 134650.

Kopylova, E., Noé, L. and Touzet, H. (2012) SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics 28(24), 3211-3217.

Li, L.G., Xia, Y. and Zhang, T. (2017) Co-occurrence of antibiotic and metal resistance genes
revealed in complete genome collection. Isme Journal 11(3), 651-662.

Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome biology 15(12), 550.

Manti, A., Boi, P., Falcioni, T., Canonico, B., Ventura, A., Sisti, D., Pianetti, A., Balsamo, M.
and Papa, S. (2008) Bacterial cell monitoring in wastewater treatment plants by flow
cytometry. Water Environment Research 80(4), 346-354.

Martinez, X., Pozuelo, M., Pascal, V., Campos, D., Gut, I., Gut, M., Azpiroz, F., Guarner, F.
and Manichanh, C. (2016) MetaTrans: an open-source pipeline for metatranscriptomics.
Scientific reports 6, 26447.

Mason, O.U., Hazen, T.C., Borglin, S., Chain, P.S., Dubinsky, E.A., Fortney, J.L., Han, J.,
Holman, H.-Y.N., Hultman, J. and Lamendella, R. (2012) Metagenome, metatranscriptome
and single-cell sequencing reveal microbial response to Deepwater Horizon oil spill. The
ISME journal 6(9), 1715-1727.

Moran, M.A., Satinsky, B., Gifford, S.M., Luo, H., Rivers, A., Chan, L.-K., Meng, J., Durham,
B.P., Shen, C. and Varaljay, V.A. (2013) Sizing up metatranscriptomics. The ISME journal
7(2), 237.

Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L. and Wold, B. (2008) Mapping and
quantifying mammalian transcriptomes by RNA-Seq. Nature methods 5(7), 621-628.

Mumy, K.L. and Findlay, R.H. (2004) Convenient determination of DNA extraction
efficiency using an external DNA recovery standard and quantitative-competitive PCR.
Journal of Microbiological Methods 57(2), 259-268.

Narayanasamy, S., Jarosz, Y., Muller, E.E., Heintz-Buschart, A., Herold, M., Kaysen, A.,
Laczny, C.C., Pinel, N., May, P. and Wilmes, P. (2016) IMP: a pipeline for reproducible

reference-independent integrated metagenomic and metatranscriptomic analyses. Genome
biology 17(1), 260.

Nayfach, S. and Pollard, K.S. (2015) Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome. Genome
biology 16(1), 51.

Ni, Y., Li, J. and Panagiotou, G. (2016) COMAN: a web server for comprehensive
metatranscriptomics analysis. BMC genomics 17(1), 622.

Oyserman, B.O., Noguera, D.R., del Rio, T.G., Tringe, S.G. and McMahon, K.D. (2015)
Metatranscriptomic insights on gene expression and regulatory controls in Candidatus
Accumulibacter phosphatis. The ISME journal.

Rapaport, F., Khanin, R., Liang, Y., Pirun, M., Krek, A., Zumbo, P., Mason, C.E., Socci, N.D.
and Betel, D. (2013) Comprehensive evaluation of differential gene expression analysis
methods for RNA-seq data. Genome biology 14(9), 3158.

Robinson, M.D., McCarthy, D.J. and Smyth, G.K. (2010) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26(1),
139-140.

Robinson, M.D. and Oshlack, A. (2010) A scaling normalization method for differential
expression analysis of RNA-seq data. Genome biology 11(3), R25.

Shi, Y.M., Tyson, G.W. and DeLong, E.F. (2009) Metatranscriptomics reveals unique
microbial small RNAs in the ocean's water column. nature 459(7244), 266-U154.

Stark, L., Giersch, T. and Wünschiers, R. (2014) Efficiency of RNA extraction from selected
bacteria in the context of biogas production and metatranscriptomics. Anaerobe 29, 85-90.

Turner, T.R., Ramakrishnan, K., Walshaw, J., Heavens, D., Alston, M., Swarbreck, D.,
Osbourn, A., Grant, A. and Poole, P.S. (2013) Comparative metatranscriptomics reveals
kingdom level changes in the rhizosphere microbiome of plants. The ISME journal 7(12),
2248-2258.

Wagner, G.P., Kin, K. and Lynch, V.J. (2012) Measurement of mRNA abundance using
RNA-seq data: RPKM measure is inconsistent among samples. Theory in Biosciences 131(4),
281-285.

Westreich, S.T., Korf, I., Mills, D.A. and Lemay, D.G. (2016) SAMSA: a comprehensive
metatranscriptome analysis pipeline. BMC bioinformatics 17(1), 399.

Whelan, J.A., Russell, N.B. and Whelan, M.A. (2003) A method for the absolute
quantification of cDNA using real-time PCR. Journal of immunological methods 278(1),
261-269.

Yang, Y., Jiang, X.T., Chai, B.L., Ma, L.P., Li, B., Zhang, A.N., Cole, J.R., Tiedje, J.M. and
Zhang, T. (2016) ARGs-OAP: online analysis pipeline for antibiotic resistance genes
detection from metagenomic data using an integrated structured ARG-database.
Bioinformatics 32(15), 2346-2351.

Yu, K. and Zhang, T. (2012) Metagenomic and metatranscriptomic analysis of microbial

community structure and gene expression of activated sludge. Plos One 7(5), e38183.

Figures and legends

Fig. 1: Schematic flow diagram for absolute quantification of metagenome and

metatranscriptome to cell/volume level.

Fig. 2 Flowchart of the simulation datasets generation and analyzing process to get

the differential expression genes.

Fig. 3: Absolute quantification of type level ARGs abundance and transcription in

FAS and NFAS. ARGs-carry hosts abundance and expression. * represents significant

difference (*P*-value < 0.05).

Fig. 4: Co-expression of ARGs and MRGs in Shatin AS, a) was the network of ARGs and MRGs expression; b) was statistical of co-expression within MRGs; c) was statistical of co-expression with ARGs; d) was statistical of co-expression of ARGs and MRGs. Lines in the network represented Spearman association over 0.6, *P*-value 0.05 the *P*-value was adjusted with B-H method.

525 **Fig. 5** Samples were collected for foaming activated sludge at 08/04/2016 and

526 non-foaming activated sludge at 25/04/2016 alongside the bioreactor at Shatin

527 wastewater treatment plant.

529 **Table 1** Comparing relative quantification methods with AQMM on detection of

530 DEGs for simulated metatranscriptome data.

	# of genes Higher expression in B	No expression difference	# of genes Higher expression in A
Theoretical Ground Truth	**28524**	**36572**	**0**
RPKM+t-Test (P < 0.05)	16477	11558	**37062**
edgeR	18278	20778	**26040**
AQMM-5%-variation	28744.72 ± 143.53	35807.52 ± 48.08	**543.77 ± 129.72**
AQMM-10%-variation	28740.83 ± 298.43	35801 ± 188.31	**554.17 ± 256.81**
AQMM-20%-variation	28549.48 ± 1007.17	35941.86 ± 919.86	**604.66 ± 654.76**
AQMM-50%-variation	16673.93 ± 9394.27	47694.99 ± 9600.33	727.08 ± 1775.09

533 **Table 2:** Summary of sequencing outputs and absolute quantification of each sample

534 at cell level with AQMM.

Sample ID	Type	Library size (bps clean data)	Total extracted DNA and RNA (ng/mL)	Estimated sequenced cells *	Estimated cells per mL *
DNA1	Foaming AS	8,567,524,200	49,140	1,541	**6.11E+10**
DNA2	Foaming AS	11,786,228,700	54,600	2,179	**6.98E+10**
DNA3	Foaming AS	10,108,576,800	58,380	1,919	**7.66E+10**
DNA4	Normal AS	8,755,895,700	57,974	1,425	**6.52E+10**
DNA5	Normal AS	9,196,724,100	66,752	1,541	**7.73E+10**

DNA6	Normal AS	11,185,847,400	75,194	1,957	**9.09E+10**
RNA1	Foaming AS	14,894,959,100	12,270	**98,936**	
RNA2	Foaming AS	13,598,855,700	12,710	**99,744**	
RNA3	Foaming AS	15,551,044,400	20,290	**78,449**	
RNA4	Normal AS	15,376,790,700	8,350	**160,343**	
RNA5	Normal AS	16,156,607,900	8,790	**189,776**	
RNA6	Normal AS	13,700,741,100	10,735	**154,925**	

*: Estimated sequenced cells for DNA libraries was using MicrobeCensus and for RNA libraries using AQMM. The assumption for AQMM was that per ml sample used for DNA and RNA extraction contained the same number of cells.