MapZ forms a stable ring structure that acts as a track for FtsZ treadmilling in *Streptococcus mutans*

Yongliang Li[1], Shipeng Shao[2], Xiao Xu[1], Xiaodong Su[3], Yujie Sun[2*], Shicheng Wei[1*]

1. Central Laboratory/Department of Oral and Maxillofacial Surgery, School and Hospital of Stomatology, Peking University; National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing 100081, China

2. State Key Laboratory of Membrane Biology, Biodynamic Optical Imaging Center (BIOPIC), School of Life Sciences, Peking University, Beijing 100871, China

3. State Key Laboratory of Protein and Plant Gene Research, School of Life Sciences, Peking University, Beijing 100871, China.

*: Correspondence: Shicheng Wei[1] or Yujie Sun[2]

1. Central Laboratory/Department of Oral and Maxillofacial Surgery, School and Hospital of Stomatology, Peking University, 22 Zhonguancun South Road, Haidian District, Beijing 100081, China Tel: (+86)-10-82195771; E-mail address: sc-wei@pku.edu.cn

2. State Key Laboratory of Membrane Biology, Biodynamic Optical Imaging Center (BIOPIC), School of Life Sciences, Peking University, 5 Yiheyuan Road Beijing 100871, China Tel: (+86)-10-62744060; E-mail address: sun_yujie@pku.edu.cn

Abstract

Proper formation of the FtsZ ring at the site of division is crucial for initiation of bacterial

cell division. A number of systems have been reported to regulate the spatial localisation of the

FtsZ ring. MapZ, a recently identified FtsZ regulatory protein in *Streptococcaceae*, has been found

to mark the division site and position the FtsZ ring. However, it is unclear how MapZ regulates the

dynamics of the FtsZ ring in *Streptococcaceae*, which is important for understanding the underlying

mechanism of this system. Here, we studied how MapZ regulates the FtsZ ring in *Streptococcus*

mutans, and its effect on the directional motion of FtsZ filaments. The results show that FtsZ

filaments in the ring exhibited treadmilling motion in *S. mutans*, similar to that in *Escherichia coli*

and *Bacillus subtilis*. Interestingly, deletion of MapZ led to unconstrained movement of

treadmilling FtsZ filaments and a shorter lifetime of the constricting FtsZ ring. Furthermore, using

both super resolution imaging and single molecule tracking, we revealed that MapZ forms an

immobile ring-like structure at the division site. Taken together, this study suggests that MapZ

forms a stable ring that acts as a track for FtsZ treadmilling in *S. mutans*.

In bacteria, the tubulin homologue FtsZ is a core component of the division machinery. On

the initiation of cell division, FtsZ forms a ring structure called the Z ring, which localises to the

middle of a cell[1] and recruits other components to construct the division machinery[2]. Over the past

few decades, a number of studies have focused on the Z-ring as well as its regulatory system. The

Min system[3], which includes three proteins, MinC, MinD, and MinE, was the first division site

selection system discovered in *Escherichia coli*. Oscillation of the MinCD complex from pole to

pole defines a minimum concentration of MinC at the middle of the cell, which directly prevents

FtsZ polymerisation. The nucleoid occlusion (NO) system[4] was also identified, which contains the

special DNA-binding proteins SimA and Noc in *E.coli* and *Bacillus subtilis*, respectively. Although

both of these proteins have been found to inhibit FtsZ assembly via binding to DNA of the undivided chromosome, a recent study has demonstrated that in addition to these two systems, some other unidentified mechanisms should also play a role in the early stage of cell division in *B. subtilis*[5]. However, many bacteria actually lack both of Min and NO systems and they evolved other Z ring regulatory systems. For instance, MipZ[6] was discovered in *Caulobacter crescentus* and is conserved among *α proteobacteria.* It directly perturbs FtsZ polymerisation and forms a complex with the DNA-binding protein ParB, which binds to replication origin regions. The complex restricts FtsZ assembly forming a Z ring in the mid-cell, where MipZ has the lowest concentration. Distinct from the negative regulatory systems mentioned above, two positive control systems PomZ[7] and SsgA/SsgB[8] were identified in *Myxococcus xanthus* and *Streptomyces*, respectively. These two systems localise at the division site prior to FtsZ and directly recruit FtsZ to the mid-cell to form the Z ring.

Recently, another important positive regulatory system was discovered in *Streptococcus pneumoniae* and is conserved in *Streptococcaceae*, which lack the aforementioned FtsZ regulatory systems. The new system is mediated by the membrane protein MapZ (also called LocZ)[9, 10], which positions the Z ring at the division site and controls its constriction.

The extracellular domain of MapZ is composed of two sub-domains,[11] and the subdomain tandem to the transmembrane domain acts as pedestal for proper localisation of the C-terminal domain, which directly binds to peptidoglycan. This two-domain structure implies that the "pedestal" domain may play the role of a switch, thereby regulating the interaction of MapZ and nascent peptidoglycans. This study supports the view that MapZ binds to nascent peptidoglycans at the site of division and is then pushed toward the daughter cell equator by peptidoglycan elongation, thereby marked the future division site. However, a recent study claimed that MapZ is not the initial factor that determines the division site[12]. This study showed that the origin of chromosome replication (oriC), which is located at the site of division prior to the MapZ ring, is

crucial for cell division. Deletion of MapZ had no remarkable effect on oriC localisation. However,

perturbing the segregation of oriC led to mislocalisation of MapZ.

Although it has been shown that MapZ interacts with FtsZ to position the Z-ring, how MapZ

regulates the dynamics of the FtsZ ring is unclear. Here, we studied how MapZ regulates the

position and movement of the FtsZ ring. The results show that FtsZ filaments in the ring exhibit a

treadmilling motion in *Streptococcus mutans*, similar to that in *E. coli* and *B. subtilis*[13, 14]. Deletion

of MapZ showed an interesting impact on the movement of treadmilling FtsZ filaments, the plane

angle of the FtsZ ring at its early formation stage, and the lifetime of the constricting FtsZ ring.

Further, our data revealed a stable ring like structure of MapZ at the division site, suggesting that

MapZ ring serves as a track for treadmilling FtsZ filaments in *S. mutans*.

Results

Characterise MapZ homologue in *Streptococcus mutans*.

MapZ was identified in *S. pneumoniae* (spMapZ). Here, we identified a MapZ homologue in

S. mutans (smMapZ); consistent with a previous report[15], smMapZ shows 37% sequence identity

with spMapZ, with 32% and 50% sequence identity for the cytoplasmic and extracellular domains,

respectively. Strikingly, most of the first 40 amino acids of spMapZ, which were thought to be

sufficient for the interaction of MapZ and FtsZ, were deficient in smMapZ while the two

phosphorylation sites[9] (T67 and T78) and the amino acids[11] (R409, Y411, N428, Y430, Y450, F451

and N454) required for binding to pneumococcal peptidoglycans are conserved (Extended Data 1a).

Subsequent analysis of the secondary structure of the intracellular domain (Extended Data 1d) and

prediction of sequence disorder tendency (Extended Data 1c) and amino acid conservation

(Extended Data 1b) indicated that smMapZ forms a similar potential structure with spMapZ, and

both contain a predicted functional subdomain at the N-terminus of the cytoplasmic domain and

two subdomains of the extracellular domain.

Deletion of *smmapZ* (Δ*smmapZ*) led to shorter, irregularly shaped cells and mini-cells (Figure

1a, b), which is similar to the phenotype in *S. pneumoniae*. To assess the localisation pattern of

smMapZ, we fused enhanced green fluorescent protein (EGFP) to the N-terminus of smMapZ

under the control of the native promoter. Time-lapse imaging of EGFP-smMapZ (Figure 1c)

indicated that smMapZ formed a ring at the mid-cell and a pair of rings appeared as the cell

elongated. The distance between the two smMapZ rings increased as a function of cell length while

the distance between the smMapZ rings and their nearest cell pole did not change remarkably

(Figure 1d), suggesting that the smMapZ ring at the mid-cell split into two rings, which moved to

the cell equator of the new daughter cells. Interestingly, smMapZ with cytoplasmic domain deletion

(*smmapZΔcyto*) resulted in loss of the ability to position at the constricting septum, resulting in

distribution throughout the cell membrane, whereas the extracellular domain null mutation

(*smmapZΔextra*) retained a septum localisation pattern (Figure 1e). This result differs from that of

spMapZ. Further multi-domain truncation analysis revealed that the predicted subdomain at the N-

terminus of the cytoplasmic domain and the subdomain tandem to the transmembrane domain

respond to such localisation patterns (Extended Data 2). Evolutionary tree analysis revealed that

smMapZ and spMapZ belong to different evolutionary branches (Figure s1).

Deletion of *smmapZ* led to mislocalisation of the FtsZ ring.

We next assessed the effect of smMapZ on the localisation of FtsZ. The red fluorescence

protein mCherry was used to label smMapZ and EGFP was fused to the C-terminus of FtsZ under

the control of an inducible promoter. Consistent with a previous report, FtsZ and smMapZ were

colocalised at the mid-cell in a newborn cell. Moreover, smMapZ was required for correct

localisation of the FtsZ ring, as 64% of FtsZ rings were unable to localise at the mid-cell in the

Δ*smmapZ* strain, while only 22.7% of FtsZ rings in the wild type (WT) strain were mislocalised

(Figure 2b). As the cell grew, the smMapZ ring split into a pair, which moved toward the cell

equator of the daughter cells before the FtsZ ring. A third smMapZ ring emerged at the division

site, constricting with the FtsZ ring. Subsequently, new FtsZ rings assembled, colocalising with the

smMapZ rings of daughter cells (Figure 2a).

Deletion of *smmapZ* led to unconstrained treadmilling of FtsZ filaments in the Z-ring.

As a tubulin homologue in prokaryotic, the treadmilling motion of FtsZ filaments was

confirmed recently[13, 14]. MapZ did not affect the polymerisation and GTPase activity of FtsZ[9], but

deletion of MapZ did impact FtsZ ring localisation and constriction. Thus, we proposed that MapZ

might control the treadmilling of FtsZ filaments. To confirm this hypothesis, we labelled FtsZ with

a brighter green fluorescent protein, mNeongreen, and used total internal reflection fluorescence

microscopy (TIRFM) to study the dynamics of FtsZ rings. The results revealed a periodic intensity

fluctuation of the FtsZ ring (Figure 3a, Extended Data 3a). With kymography, we observed

directional movement of the FtsZ ring (Figure 3b, Movie s1). In contrast with *E. coli,* in which only

one FtsZ ring forms during a cell cycle, three FtsZ rings could be observed in the majority of *S.*

mutans cells at a later stage of the cell cycle. Surprisingly, we observed translocation of FtsZ

filaments from the middle (old) FtsZ ring to the nearby (new) FtsZ rings (Figure 3d, Movie s3),

and the speed of translocation was similar to that of treadmilling inside the ring.

Interestingly, deletion of *smmapZ* led to highly unconstrained movement of FtsZ filaments

(Figure 4a, Movie s5, s6), but did not perturb the treadmilling and translocation of FtsZ filaments

(Figure 3a, b, d, Movie s2, s4) or alter the speed of treadmilling (Figure 3c), which suggests a

crucial role of smMapZ in guiding the treadmilling movement of FtsZ filaments. Although the

unconstrained FtsZ filaments could still form a ring-like structure as the cell grew, the angle of the

FtsZ ring plane changed back and forth during FtsZ ring formation, which might ultimately lead to

aberrant localisation of the Z ring. Nevertheless, once the Z ring was formed, the angle of the Z

ring plane would no longer change (Extended Data 3b, Movie s5). We also noticed that deletion of

smmapZ resulted in a decrease in chance to observe cells with three FtsZ rings (Extended Data 3c).

By monitoring the complete process of emergence and disappearance of the three FtsZ ring

structure, we discovered that the middle FtsZ ring in the Δ*smmapZ* strain only remained 5-6 min

after new FtsZ ring formation, exhibiting a shorter lifetime than the life time (>10 min) in the WT

strain (Figure 4b, Movie s7, s8). Analysis of the intensity trace also indicated that deleting *smmapZ*

led to a faster fluorescence decrease of the middle FtsZ ring than that in WT (Figure 4c). This

suggested that smMapZ might play a role in guiding the treadmilling and stabilising the constricting

FtsZ ring.

smMapZ ring exhibits discontinuous and stable structure

To understand how smMapZ guides the treadmilling of FtsZ filaments, we performed three-

dimensional structured illumination microscopy (3D-SIM) to study the structure of a Halotag-fused

smMapZ (Halo$_{JF549}$-smMapZ[16]) ring. The smMapZ ring exhibited a heterogeneous ring-like

structure at the division site (Figure 5a-c, Movie s11), similar to previously studied FtsZ, EzrA,

Pbp2[17], FtsA and ZipA[18]. Such discontinuous ring structures allow for direct observation of their

directional movements using single molecule TIRF imaging[13, 14]. Here, to study the dynamics of

smMapZ ring, we labelled smMapZ with mNeongreen for TIRF imaging. Interestingly, no

remarkable periodic motion of smMapZ was observed by intensity traces or kymograph analysis

(Figure 5d, Extended Data 4a, Movie s9), suggesting that the smMapZ ring is immobile. We further

observed the relative movement between smMapZ and FtsZ using a dual color assay. By labeling

smMapZ and FtsZ with EGFP and mScarlet-I, respectively, we revealed that the FtsZ ring exhibited

directional movement while the smMapZ ring showed no remarkable movement (Figure 5e).

Single molecule tracking allowed to measure the spatially specific dynamic features of targeted

molecules. Here, we labelled Halotag fused smMapZ with low concentrations (62.5 pM) of

HaloLigand-JF$_{646}$ to monitor the motion of single smMapZ molecules. The results revealed two

populations of smMapZ molecules: (i) stationary and (ii) diffusive (Figure 5f, Movie s10). In a few

cases, we also observed one single smMapZ molecule switched between the two modes (Extended

Data 4b). We analysed the localisation pattern of stationary smMapZ molecules in 62 cells. In 38

cells that exhibited newborn cell length or three FtsZ ring structures, the stationary smMapZ

molecules co-localised with the FtsZ ring. In the rest 24 growing cells, the stationary smMapZ

molecules localised at the cell equator of the future daughter cells (Figure 5g, Extended Data 4c).

The low mobility of smMapZ molecules supports that smMapZ forms a stable ring structure.

However, as most single molecules were bleached in 2-3 min, single molecule trajectories were

insufficient to study the stability of the smMapZ ring over a longer period of time. Therefore, we

labelled smMapZ with mMaple3, a photoconvertable fluorescent protein that converts from green

fluorescence to red fluorescence under purple illumination[19]. Using TIRF illumination, we

photoconverted mMaple3-fused smMapZ molecules in the bottom portion of the ring using a 405

nm laser and monitored their movement every 3 min. The results showed that over a total duration

of 21 min, neither the position nor the intensity of the photoconverted smMapZ molecules changed

remarkably, indicating that the smMapZ ring is a stable structure and the exchange of smMapZ

molecules occurs at a very slow rate (Figure 5h, Extended Data 4d).

Discussion

Recently a membrane protein MapZ mediated FtsZ regulatory system was uncovered in *S.*

pneumoniae, which was shown to be conserved in *Streptococcaceae*. As a positive FtsZ regulator,

MapZ first binds to the division site forming a ring structure and then positions the FtsZ ring[9, 10].

Two recent studies[11, 12] gained insight into how MapZ marks the division site, but little work has

been focused on how MapZ controls the FtsZ ring structure. Here, our data suggested that MapZ

forms a stable heterogeneous ring structure at the division site, which may act as a track for the

treadmilling of FtsZ filaments (Figure 6). This is primarily demonstrated by the characterisation of

the smMapZ ring structure and FtsZ treadmilling in *S. mutans*.

smMapZ is essential for regular cell shape formation and proper localistion of the constricting

septum in *S. mutans*; it marks the division site prior to the FtsZ ring (Figure 2a). Deletion of

smmapZ results in a shorter, irregularly shaped cell, and a mislocalised FtsZ ring (Figure 1a, b,

Figure 2b).

To understand the mechanism by which the smMapZ ring affects the FtsZ ring localisation,

we focused on the treadmilling of FtsZ, which was confirmed recently. Similar to that in *E. coli*

and *B. subtilis,* the FtsZ ring in *S. mutans* demonstrated periodic fluctuations of intensity and a

directional motion in high temporal resolution imaging (Figure 3a, b, c, Extended Data 3a, Movie

s1). Interestingly, different from that in *E. coli* and *B. subtilis*, we observed translocation of FtsZ

from the middle ring to the newly formed FtsZ ring (Figure 3d). The translocation velocity was

similar to the treadmilling vecolity of FtsZ filaments inside the ring. Considering that the

translocated FtsZ would participate in new ring formation, it is reasonable to note that such kind of

FtsZ translocation is a functional process rather than abnormal localisation. The translocation of

FtsZ also supports the view that the FtsZ ring consists of multiple protofilaments, thereby

demonstrating a bead-like structure or discontinuous clusters[17, 18, 20, 21].

Further, we revealed that deletion of *smmapZ* resulted in unconstrained movement of FtsZ

filaments and an unstable plane of the FtsZ ring (Figure 4a; Extended Data 3b), suggesting that the

smMapZ ring may play a role in restricting and guiding FtsZ treadmilling. This can also explain

why deletion of MapZ leads to an irregular cell shape and minicell formation. Consistent with a

previous report[9], we observed a lower proportion of cells harbouring the FtsZ dumbbell structure

in the Δ*smmapZ* strain compared to the WT cells (Extended Data 3c). This was probably due to the

shorter lifetime of the middle FtsZ ring in the Δ*smmapZ* strain (Figure 4b, c). We speculate that the

shorter lifetime was caused by an increase in FtsZ translocation (Movie s5); however, we cannot

exclude the possibility that it may have been due to premature constriction of the FtsZ ring, as

previously described[9].

We characterised the smMapZ ring structure to understand how it guides the treadmilling of

FtsZ filaments. The discontinued ring-like structure of smMapZ was uncovered by 3D-SIM (Figure

5a-c). However, unlike FtsZ, FtsA and Pbp2B[14], the smMapZ ring or single smMapZ molecules

did not show remarkable directional movement around the short cell axis (Figure 5d and Extended

Data 3a). Thus, we hypothesise that smMapZ localisates at the division site prior to FtsZ and forms

a stable heterogeneous ring-like structure (Figure 5g, h; Extended Data 3c, d). This stable ring

structure acts as a track for FtsZ treadmilling (Figure 5e). However, we noticed that FtsZ still

formed a Z ring in the Δ*smmapZ* strain. The change of plane angle of the FtsZ ring often happened

at the early stage of FtsZ ring formation but not in the constricting stage. Thus, we propose that the

track role of MapZ only functions at the early stage of FtsZ ring formation, and other proteins also

participate in the stablisation of FtsZ treadmilling after Z ring formation, possibly including ErZA,

a spectrin like protein[22], which has been reported to interact with FtsZ and is required for correct

localisation of the FtsZ ring[23].

In summary, our study revealed the "track" role of smMapZ in FtsZ treadmilling, providing

new insight into the mechanism of the MapZ-mediated FtsZ regulatory system. Considering the

similar functions of smMapZ with that of spMapZ, we believe this does not only exist in *S. mutans*,

but also in other MapZ-conserved bacteria.

Material and methods

Bacterial strains, plasmid and mutation construction, and culture conditions

S. mutans standard strain UA159 and all derived strains used in this study are listed in Table

1. To generate the *S. mutans* mutations (gene deletion and fluorescent protein fusion), we used a

counter-selectable system called IFDC2 (provided by Zhoujie Xie). The transformation procedure

was performed as previously described[24]. Briefly, the overnight culture grown in Todd–Hewitt

medium supplemented with 0.3% yeast extract (THY; QinDao Wei si, China) was diluted (1:20)

in the same medium and incubated for 2–3 h (OD_{600} 0.2–0.3). Overlapping PCR reaction (5 µl) and

0.5 µl of CSP (1 µg/µl stock) were added to 500 µl of culture and incubated for 2 h. To select

antibiotic-resistant colonies, brain heart infusion (BHI; Difco Laboratories, Detroit, MI, USA) agar

plates were supplemented with 12-µg/ml erythromycin (Sigma-Aldrich, St. Louis, MO, USA).

Selection of the second transformation was performed on BHI plates supplemented with 4-mg/ml

p-cl-Phe (Sigma-Aldrich). For snapshot imaging, *S. mutans* strains were aerobically cultured in

THY at 37°C. For time-lapse imaging, the strains were cultured in a chemically defined medium[25]

supplemented with 0.3% yeast extract (C + Y). A final concentration of 10 mg/ml xylose was added

to the culture medium to induce the expression of Pzx9 and Pzx10[26] (provided by Jiezhou Xie).

Sample preparation for snapshot imaging

An overnight culture grown in THY medium was diluted (1:100) in the same medium, and

incubated at 37°C aerobically to OD_{600} 0.4–0.5 (7 h). The culture was diluted again (1:100) in THY

and grown at 37°C aerobically to OD_{600} 0.1–0.2 (5 h). Cells were harvested by centrifugation at

$1,500 \times g$ for 3 min, washed three times with $1\times$ phosphate-buffered saline (PBS), and resuspended

in $1\times$ PBS. The bacteria were loaded between a gel pad ($1\times$ PBS supplemented with 1.5% low

melting agarose) and coverslip (Fisher 24×50, No. 1). For FtsZ imaging, spectinomycin was added

to all cultures at a final concentration of 1 mg/ml. D (+) xylose (Sigma-Aldrich) was added before

the last round of dilution at 10 mg/ml.

Sample preparation for 3D-SIM imaging

JF_{549} dye (500 nM; provided by Luke Lavis) was added into the culture at OD_{600} 0.1–0.2,

incubated for 15 min at 37°C, washed once with $1\times$ PBS, fixed in 4% paraformaldehyde for 15 min

at room temperature, and washed twice with $1\times$ PBS. Resuspended cells were loaded between the

coverslip (Fisher 24×50, No. 1.5) and gel pad.

Sample preparation for time-lapse imaging

EGFP_smMapZ and mMaple3_smMapZ. An overnight culture grown in THY medium was

diluted (1:100) in C + Y medium. After two dilutions, the culture at OD_{600} 0.1–0.2 was added to a

gel pad (C + Y medium supplemented with 1.5% low melting agarose) on a glass slide, and a

coverslip was added. Rectangular 3M double-sided tape was used to seal the coverslip and glass

slide.

HaloJF₆₄₆_smMapZ. JF_{646} dye (62.5 pM) was added into the culture at OD_{600} 0.1–0.2,

incubated for 15 min at 37°C and washed once with pre-warmed fresh C + Y medium. Resuspended

cells were loaded between the coverslip and the gel pad.

Imaging of FtsZ_mNeongreen. Spectinomycin was added to all cultures at a final

concentration of 1 mg/ml. D (+) xylose was added at 10 mg/ml before the last round of dilution.

Slide preparation

Coverslips were cleaned with Piranha solution (30% H_2O_2: 98% H_2SO_4 v:v = 1:3 at 90°C for 30

min), washed with Milli-Q water, stored in filtered Milli-Q water, and dried with high purity

nitrogen before use

Snapshot imaging of smMapZ and FtsZ

The samples were visualised on an N-STORM (Nikon, Tokyo, Japan) system equipped with

a 100× oil TIRF objective (Nikon PLAN APO, 1.49 NA), Andor-897 EMCCD (Andor, Belfast,

Northern Ireland), the perfect focus system (Nikon), a laser source (405, 488, 561, and 647 nm)

and 1.5× magnification optics. Images were collected with Nis-Elements AR software (Nikon). For

EGFP_smMapZ and mCherry_smMapZ, 300 frames were collected with a 100 ms exposure time.

For FtsZ_EGFP, 50 frames were collected with a 100 ms exposure time.

Time-lapse imaging of EGFP_smMapZ and mMaple3_smMapZ

Time-lapse observation of EGFP_smMapZ was performed with a DeltaVision OMX SR (GE

Healthcare, Little Chalfont, UK) system equipped with a 1.42 NA 60× oil objective (Olympus,

Tokyo, Japan), laser source (405, 488, 561, and 642 nm), four sCOMS, and an environment control

system. Using the OMX Acquisition software, the images were collected with a 60 ms exposure

time, with 5 min intervals at 37°C.

Time-lapse imaging of mMaple3_smMapZ was performed with an N-STORM microscope

equipped with a live cell instrument. A laser at 405 nm was used to photoconvert the mMaple3 to

the red state with an initial exposure time of 300 ms. Then, a 561 nm laser was used to excite the

red state mMaple3 with a 400 ms exposure time, and a 488 nm laser was used for the green state

with a 300 ms exposure time.

3D-SIM imaging of HaloJF$_{549}$_smMapZ

Super-resolution 3D-SIM imaging was performed with a DeltaVision OMX SR system, with

a 10 ms exposure time and 50% transmission. The 3D-SIM raw data were reconstructed with

SoftWoRx 6.0 (Applied Precision, Issaquah, WA, USA) using a Wiener filter setting of 0.001 and

a channel that specifically measured optical transfer functions. The immersion oil was optimised

to 1.518.

Imaging of the dynamics of FtsZ_mNeongreen, mNeongreen_smMapZ and

HaloJF646_smMapZ

The images were collected by near-TIRF illumination on an N-STORM microscope.

Exposure time was 0.165 s, interval time was 0.495 s, and mNeongreen was excited by the 488 nm

laser. The images were captured at room temperature. For the single molecular track of HaloJF646-

smMapZ, exposure time was 0.1 s, and the interval time was 0.5 s. The 647 nm laser was used to

excite the JF646 dye.

Data analysis

Localisation of smMapZ. The averaged image stacks were obtained, and Snapshot analysis

accomplished, with ImageJ software (NIH, Bethesda, MD, USA). Cell outlines and corresponding

fluorescence signals were manually detected and analysed using MicrobeTracker[27] and custom

MATLAB (MathWorks, Natick, MA, USA) codes. Information regarding the cell length of WT

and ΔsmMapZ was extracted and the graph was generated in Microsoft Excel software (Microsoft

Corp., Redmond, WA, USA). To compare the distance between the two outer rings, and between

the smMapZ ring and the cell pole, we manually eliminated the cell in which only one ring structure

could be observed, then the data were calculated with a custom MATLAB script and plotted in

Excel. To sort the fluorescence intensity of the smMapZ truncations as a function of cell length,

averaged images were deconvoluted using Huygens and subsequently analysed using Coli-

Inspector running under the ImageJ plugin ObjectJ (http://simon.bio.uva.nl/objectj/), as previously

described[11].

The analysis of periodic intensity fluctuations of the Z ring and treadmilling were performed

in Fiji (http://fiji.sc/). First, drift correction of the time-lapse imaging was performed using series

registration based on the Fiji plugin Descriptor. We selected a medium detection brightness and

six-pixel for detection size; the type of detection was set as minima and maxima, and the rest were

set as default values. After drift correction, a 3×3 pixel (~320 nm) region of interest was chosen

inside the Z ring, and the Z-axis profile was plotted with Fiji to show the periodic intensity

fluctuations. For analysis of FtsZ treadmilling, the images were resized to 26 nm/pixel with the

bicubic method, and a segmented line (line width is 11) was drawn inside the Z ring, parallel to its

long axis, and a multiple kymograph plugin was used to demonstrate movement. The velocity of

treadmilling was calculated based on the kymograph using the macro "read velocities from tsp" as

described in the manual.

Single particle tracking. The Fiji plugin TrackMate (http://imagej.net/TrackMate) was used

to analyse the trajectory of single molecules of smMapZ. We selected the LoG detector and set 0.4

μm as the estimated blob diameter. The LAP Tracker function, with a 0.5 μm maximum distance

for frame-to-frame linking, was employed to complete the tracking. The trajectory data were

exported; further analysis of logD was carried out in R studio (R Development Core Team, Vienna,

Austria) with a custom script. The two-status analysis of single trajectory was performed in

MATLAB with the HMM-Bayes procedure[28] , as previously described.

smMapZ sequence analysis. To identify smMapZ, we selected spMapZ as the query sequence

and used BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi) to search the genome of *S. mutans*. The

two sequences were aligned with SnapGene software (GSL Biotech LLC, Chicago, IL, USA). The

transmembrane domain was predicted using TMHMM Server v. 2.0

(http://www.cbs.dtu.dk/services/TMHMM/). Network Protein Sequence Analysis[29] (http://npsa-

pbil.ibcp.fr) was used to predict the secondary structure of the cytoplasmic domain of smMapZ as

previously described. In total, 150 protein sequences were selected to determine amino acid

conservation using the Consurf web server[30] (http://consurf.tau.ac.il/2016/). Prediction of sequence

disordered tendency was performed with IUPred[31] (http://iupred.enzim.hu/).

ACKNOWLEDGEMENTS

This work was funded by Peking University's 985 Grant and the National Natural Science

Foundation of China (No. 30973317, 31271423, and 21390412). We thank Zhoujie Xie (Institute

of Microbiology, CAS) for his assistance in the construction of the mutations and Chunyan Shan

(The Core Imaging Facility of Peking University) for her help on DeltaVision OMX. We thank He

Yu, Professor Can Xie and ShuJin Luo (School of Life Sciences, Peking University) for their help

with analysis of evolution tree.

1.   Bi EF, Lutkenhaus J. FtsZ ring structure associated with division in Escherichia coli. *Nature* 1991, **354**(6349)**:** 161-164.

2.   Adams DW, Errington J. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. *Nature reviews Microbiology* 2009, **7**(9)**:** 642-653.

3.   Rowlett VW, Margolin W. The bacterial Min system. *Current biology : CB* 2013, **23**(13)**:** R553-556.

4.   Wu LJ, Errington J. Nucleoid occlusion and bacterial cell division. *Nature reviews Microbiology* 2011, **10**(1)**:** 8-12.

5.   Rodrigues CDA, Harry EJ. The Min System and Nucleoid Occlusion Are Not Required for Identifying the Division Site in Bacillus subtilis but Ensure Its Efficient Utilization. *Plos Genet* 2012, **8**(3).

6.   Thanbichler M, Shapiro L. MipZ, a spatial regulator coordinating chromosome segregation with cell division in Caulobacter. *Cell* 2006, **126**(1)**:** 147-162.

390   7.   Treuner-Lange A, Aguiluz K, van der Does C, Gomez-Santos N, Harms A, Schumacher D,
391   *et al.* PomZ, a ParA-like protein, regulates Z-ring formation and cell division in
392   Myxococcus xanthus. *Mol Microbiol* 2013, **87**(2)**:** 235-253.

393
394   8.   Willemse J, Borst JW, de Waal E, Bisseling T, van Wezel GP. Positive control of cell
395   division: FtsZ is recruited by SsgB during sporulation of Streptomyces. *Genes &*
396   *development* 2011, **25**(1)**:** 89-99.

397
398   9.   Fleurie A, Lesterlin C, Manuse S, Zhao C, Cluzel C, Lavergne JP*, et al.* MapZ marks the
399   division sites and positions FtsZ rings in Streptococcus pneumoniae. *Nature* 2014,
400   **516**(7530)**:** 259-262.

401
402   10.   Holeckova N, Doubravova L, Massidda O, Molle V, Buriankova K, Benada O*, et al.* LocZ is
403   a new cell division protein involved in proper septum placement in Streptococcus
404   pneumoniae. *mBio* 2014, **6**(1)**:** e01700-01714.

405
406   11.   Manuse S, Jean NL. Structure-function analysis of the extracellular domain of the
407   pneumococcal cell division site positioning protein MapZ. *Nature communications* 2016,
408   **7:** 12071.

409
410   12.   van Raaphorst R, Kjos M, Veening JW. Chromosome segregation drives division site
411   selection in Streptococcus pneumoniae. *Proc Natl Acad Sci U S A* 2017, **114**(29)**:** E5959-
412   e5968.

413
414   13.   Yang X, Lyu Z. GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ
415   organizes septal cell wall synthesis. *Science* 2017, **355**(6326)**:** 744-747.

416
417   14.   Bisson-Filho AW, Hsu YP. Treadmilling by FtsZ filaments drives peptidoglycan synthesis
418   and bacterial cell division. *Science* 2017, **355**(6326)**:** 739-743.

419
420   15.   Garcia PS, Simorre JP, Brochier-Armanet C, Grangeasse C. Cell division of Streptococcus
421   pneumoniae: think positive! *Current opinion in microbiology* 2016, **34:** 18-23.

422
423   16.   Grimm JB, English BP, Chen J, Slaughter JP, Zhang Z, Revyakin A*, et al.* A general method
424   to improve fluorophores for live-cell and single-molecule microscopy. *Nature methods*
425   2015, **12**(3)**:** 244-250, 243 p following 250.

426
427   17.   Strauss MP, Liew ATF, Turnbull L, Whitchurch CB, Monahan LG, Harry EJ. 3D-SIM Super
428   Resolution Microscopy Reveals a Bead-Like Arrangement for FtsZ and the Division
429   Machinery: Implications for Triggering Cytokinesis. *Plos Biology* 2012, **10**(9).

431   18.   Rowlett VW, Margolin W. 3D-SIM Super-resolution of FtsZ and Its Membrane Tethers in
432   Escherichia coli Cells. *Biophysical Journal* 2014, **107**(8)**:** L17-L20.

433
434   19.   Wang S, Moffitt JR, Dempsey GT, Xie XS, Zhuang X. Characterization and development of
435   photoactivatable fluorescent proteins for single-molecule-based superresolution
436   imaging. *Proc Natl Acad Sci U S A* 2014, **111**(23)**:** 8452-8457.

437
438   20.   Guo F, Tao H, Buss J, Coltharp C, Hensel Z, Jie X. In Vivo Structure of the E. coli FtsZ-ring
439   Revealed by Photoactivated Localization Microscopy (PALM). *PloS one* 2010, **5**(9).

440
441   21.   Buss J, Coltharp C, Huang T, Pohlmeyer C, Wang S-C, Hatem C*, et al.* In vivo organization
442   of the FtsZ-ring by ZapA and ZapB revealed by quantitative super-resolution microscopy.
443   *Molecular Microbiology* 2013, **89**(6)**:** 1099-1120.

444
445   22.   Cleverley RM, Barrett JR, Basle A, Bui NK, Hewitt L, Solovyova A*, et al.* Structure and
446   function of a spectrin-like regulator of bacterial cytokinesis. *Nature communications*
447   2014, **5:** 5421.

448
449   23.   Jorge AM, Hoiczyk E, Gomes JP, Pinho MG. EzrA contributes to the regulation of cell size
450   in Staphylococcus aureus. *PloS one* 2011, **6**(11)**:** e27542.

451
452   24.   Xie Z, Okinaga T, Qi F, Zhang Z, Merritt J. Cloning-independent and counterselectable
453   markerless mutagenesis system in Streptococcus mutans. *Applied and environmental*
454   *microbiology* 2011, **77**(22)**:** 8025-8033.

455
456   25.   van de Rijn I, Kessler RE. Growth characteristics of group A streptococci in a new
457   chemically defined medium. *Infection and immunity* 1980, **27**(2)**:** 444-448.

458
459   26.   Xie Z, Qi F, Merritt J. Development of a tunable wide-range gene induction system useful
460   for the study of streptococcal toxin-antitoxin systems. *Applied and environmental*
461   *microbiology* 2013, **79**(20)**:** 6375-6384.

462
463   27.   Sliusarenko O, Heinritz J, Emonet T, Jacobs-Wagner C. High-throughput, subpixel
464   precision analysis of bacterial morphogenesis and intracellular spatio-temporal
465   dynamics. *Mol Microbiol* 2011, **80**(3)**:** 612-627.

466
467   28.   Monnier N, Barry Z, Park HY, Su KC, Katz Z, English BP*, et al.* Inferring transient particle
468   transport dynamics in live cells. *Nature methods* 2015, **12**(9)**:** 838-840.

469
29.   Combet C, Blanchet C, Geourjon C, Deleage G. NPS@: network protein sequence
analysis. *Trends in biochemical sciences* 2000, **25**(3)**:** 147-150.

30.   Landau M, Mayrose I, Rosenberg Y, Glaser F, Martz E, Pupko T*, et al.* ConSurf 2005: the
projection of evolutionary conservation scores of residues on protein structures. *Nucleic*
acids research 2005, **33**(Web Server issue)**:** W299-302.

31.   Dosztanyi Z, Csizmok V, Tompa P, Simon I. IUPred: web server for the prediction of
intrinsically unstructured regions of proteins based on estimated energy content.
Bioinformatics 2005, **21**(16)**:** 3433-3434.

Figure legends

492 Figure 1. **Characterisation of a MapZ homologue in *S. mutans* (smMapZ)**. (a) Cell shape of

493 wild type (WT) and deletion of *smmapZ* (Δ*smmapZ*). Membrane was stained with FM4-64. A

minicell is indicated by the white arrow. Scale bar, 1 μm. (b) Analysis of cell length of WT and

Δ*smmapZ*. (c) Live cell imaging of enhanced green fluorescent protein (EGFP) smMapZ. White

arrows indicate the split smMapZ rings, and the red arrow indicates the third smMapZ ring. (d)

Analysis of the distance between the two outer rings, and between the ring and the nearest pole.

(e) Localisation pattern of WT smMapZ, smMapZ with cytoplasmic domain deletion

(smMapZΔcyto), and the extracellular domain null mutation (smMapZΔextra). A schematic of

the mutations is shown on the left, cell shape and EGFP signals in the middle, and corresponding

maps of fluorescence intensity (sorted by cell length) on the right.

506 Extended Data 2. **Localisation of smMapZ truncations.** The potential subdomains of the

507 cytoplasmic domain were predicted based on its secondary structure. The potential subdomains of

508 the extracellular domain were predicted based on amino acids conservation and disorder tendency

509 analysis.

514 Figure 2. **Effect of smMapZ on FtsZ localisation**. (a) Dual colour imaging of smMapZ and FtsZ

515 in different stages of the cell cycle. (b) Localisation of FtsZ in WT and Δ*smmapZ*. Cell shape

516   observed by phase contrast microscopy and fluorescence of EGFP are shown on the left, and

517   statistical analysis of FtsZ ring position is shown on the right.

Figure 3. **FtsZ shows directional movement on smMapZ**. (a) Typical periodic fluctuation of the

FtsZ ring in WT and Δ*smmapZ*. (b) Typical direct movement of FtsZ in WT and Δ*smmapZ*. (c)

Distribution of velocity of FtsZ movement inside the rings. (d) Translocation of the FtsZ filament.

Schematics are shown above. "O" refers to the old ring and "N" refers to newly formed rings.

Fluorescence images are shown below. The red line indicates cell shape, and yellow arrows mark

the initiation of translocation.

Figure 4. **Deletion of *smmapZ* lead to unconstrained FtsZ treadmilling**. (a) Unconstrained

movement of the FtsZ filaments in Δ*smmapZ*. Diagrams of the first four snapshots are shown in

the upper panel, and the cell shape in bright field, marked by a dotted line, is shown in the upper

left. In the diagrams, the dark blue arrows indicate the position and direction of movement of

FtsZ filaments, and light blue arrows indicate the movement path of FtsZ. In snapshots of the

time-lapse imaging shown below, yellow arrows mark the position of FtsZ and the change of

plane of the FtsZ ring. (b) Snapshots of the time-lapse imaging of three FtsZ ring structures in

WT and Δ*smmapZ*. White arrows mark the newly formed FtsZ ring position and yellow arrows

show the position of the old FtsZ ring. (c) Fluorescence intensity traces of the old FtsZ ring as a

function of time.

Extended Data 3. **Additional results of the effect of smMapZ on FtsZ treadmilling.** (a) Two

representative fluorescence intensity fluctuations of FtsZ_mNeongreen in WT and Δ*smmapZ*. (b)

Abnormal FtsZ ring plane and division septum. (c) Proportion of three FtsZ ring structures in WT

and Δ*smmapZ*.

Figure 5. **Characterisation of the smMapZ ring**. (a) Three-dimensional structured illumination microscopy of HaloJF$_{549}$_smMapZ. (b) Heterogeneous ring structure of smMapZ in the box region of (a). (c) Cartoon of the heterogeneous ring structure in (b). (d) Intensity trace and kymograph analysis of mNeongreen_smMapZ. (e) Dual colour analysis of the movement of FtsZ and smMapZ. (f) Distribution of logD of single molecule tracking of smMapZ. D is the diffusion coefficient. The Gaussian fitting curve, the average D, and area of the two populations are shown. (g) Kymograph analysis of representative constrained molecules (left) and the localised relationship between the FtsZ ring in the newborn (upper) and the grown (lower) cells (right). (h) Snapshots of the time-lapse imaging of mMaple3_smMapZ. The smMapZ ring structure is marked by yellow arrows.

Extended Data 4. **Additional characterisation of the smMapZ ring.** (a) Additional intensity

traces of mNeongreen_smMapZ. (b) Hidden Markov model-Bayes analysis of the trajectory with

a combination of immobilisation and diffusion. Two diffusive statuses (D-D) were predicted, and

blue and pink annotations represent the higher and lower diffusion coefficient status, respectively.

(c) Typical images of the relationship between smMapZ molecules and FtsZ rings. The average

of the first 50 frames of Halo$_{JF646}$_smMapZ is shown. White arrows mark the cells in which

immobilised smMapZ molecules co-localise with the FtsZ ring. Yellow arrows mark the cells in

which immobilised smMapZ molecules do not co-localise with the FtsZ rings. Blue arrows mark

the cells in which smMapZ molecules are diffuse.

Figure 6. **Model of smMapZ guiding the treadmilling of FtsZ**. smMapZ forms a stable

heterogeneous ring-like structure at the division site. The stable ring acts as a track for the

treadmilling of FtsZ and, as the cell grows, this stable smMapZ ring moves to the future division

site of daughter cells before the FtsZ ring. Newly formed FtsZ will move along the stable ring,

and the translocated FtsZ filaments will also participate in new ring formation.

597 Table 1

S. mutans strains	Genotype and description	reference
UA159	S. mutans str UA159	Gift from Yifei Zhang, Central Laboratory of School and Hospital of Stomatology, Peking University,China
△smmapZ	UA159; △Smu473	This study
EGFP_smMapZ	UA159; Smu473::EGFP-Smu473	This study
mCherry_smMapZ	UA159; Smu473::mCherry-Smu473	This study
Halo_smMapZ	UA159; Smu473::Halo tag-Smu473	This study
mNeongreen_smMapZ	UA159; Smu473::mNeongreen-Smu473	This study
EGFP_smMapZΔcyto	UA159; Smu473::EGFP-Smu473-Δcytoplasmic domain	This study
EGFP_smMapZΔextra	UA159; Smu473::EGFP-Smu473-Δextracellular domain	This study
EGFP_smMapZΔcyto(1-40)	UA159; Smu473::EGFP-Smu473-ΔM1-F40	This study
EGFP_smMapZΔcyto(41-104)	UA159; Smu473::EGFP-Smu473-ΔD41-A104	This study
EGFP_smMapZΔcyto(105-175)	UA159; Smu473::EGFP-Smu473-ΔP105-K175	This study
EGFP_smMapZΔcyto(41-175)	UA159; Smu473::EGFP-Smu473-ΔD41-K175	This study
EGFP_smMapZΔextra(198-401)	UA159; Smu473::EGFP-Smu473-ΔQ198-T401	This study
EGFP_smMapZΔextra(402-521)	UA159; Smu473::EGFP-Smu473-ΔP402-Y521	This study
FtsZ-EGFP	UA159; Pzx10::FtsZ-EGFP, Str^R	This study
FtsZ-mNeongreen	UA159; Pzx9::FtsZ-mNeongreen, Str^R	This study
FtsZ-mScarlet	UA159; Pzx9::FtsZ-mScarlet-I, Str^R	This study
△smmapZ FtsZ-EGFP	UA159; △Smu473 Pzx10::FtsZ-EGFP, Str^R	This study
mCherry_smMapZ FtsZ-EGFP	UA159; Smu473::mCherry-Smu473 Pzx10::FtsZ-EGFP, Str^R	This study
EGFP_smMapZ FtsZ-mScarlet	UA159; Smu473::EGFP-Smu473 Pzx9::FtsZ-mScarlet-I, Str^R	This study
Halo_smMapZ FtsZ-mNeongreen	UA159; Smu473::Halo tag-Smu473 FtsZ-mNeongreen, Str^R	This study

Supplementary

Figure s1: Evolutionary tree analysis of smMapZ and spMapZ in *Streptococcaceae.*

Movie s1. Directional movement of FtsZ inside the Z ring in WT cells.

Movie s2. Directional movement of FtsZ inside the Z ring in △*smmapZ* cells.

Movie s3. Translocation of FtsZ in WT cells.

Movie s4. translocation of FtsZ in △*smmapZ* cells.

Movie s5. Unconstrained movement of FtsZ in △*smmapZ* cells. The bright field image was

added via image calculator in Fiji.

Movie s6. Unconstrained movement of FtsZ in △*smmapZ* cells.

Movie s7. Lifetime of constricting FtsZ ring in △*smmapZ* cells.

Movie s8. Lifetime of constricting FtsZ ring in WT cells.

Movie s9. Timelapse of mNeongreen_smMapZ in WT cells.

Movie s10. Single molecule tracking of smMapZ.

Movie s11. Ring structure of Halo$_{JF549}$_smMapZ.