Quantitative imaging of chromatin decompaction in living cells

Elisa Dultz*†, Roberta Mancini*, Guido Polles[1], Pascal Vallotton*, Frank Alber[1] and Karsten Weis*

* Institute of Biochemistry, Department of Biology, ETH Zurich, Zurich, Switzerland

[1] Molecular and Computational Biology, University of Southern California, Los Angeles, USA

† corresponding author: elisa.dultz@bc.biol.ethz.ch

Running head: chromatin decompaction live

Abbreviations:

        TSA – trichostatin A

        ORF – open reading frame

        qPCR – quantitative PCR

        SE – standard error

        SPB – spindle pole body

        GST – Glutathione S transferase

        smFISH – single molecule fluorescence *in situ* hybridization

        Pol II – RNA polymerase II

Abstract

Chromatin organization is highly dynamic and regulates transcription. Upon transcriptional activation, chromatin is remodeled and referred to as "open", but quantitative and dynamic data of this decompaction process are lacking. Here, we have developed a quantitative superresolution microscopy assay in living yeast cells to visualize chromatin decompaction using the *GAL7-10-1* locus as a model system. Upon transcriptional activation of these three clustered genes, we detect an increase of the mean distance across this locus by ~100 nm. This decompaction is linked to active transcription but is not sensitive to the histone deacetylase inhibitor trichostatin A or to deletion of the histone acetyl transferase Gcn5. By contrast, the deletion of *SNF2* (encoding the ATPase of the SWI/SNF chromatin remodeling complex) or the deactivation of the histone chaperone complex FACT lead to a strongly reduced decompaction. Our assay quantifies and visualizes for the first time the chromatin conformation changes brought about by transcriptional activation and suggests that nucleosome remodeling and eviction activities are major contributors to chromatin reorganization.

Introduction

DNA in the cell nucleus is present as chromatin in a tight complex with histones and other proteins. This complex is central to spatially organize the DNA strand by balancing the negative charges of the phosphate backbone and is also crucial for gene regulation. The three-dimensional chromatin conformation is highly dynamic and remodeled continuously as cells change their physiological state or their transcriptional programs. This remodeling is orchestrated by histone modifiers and chromatin remodelers, which change the interaction between nucleosomes as well as the interaction between histones, DNA and the protein complement present at a chromatin site, thereby affecting the spatial packing of nucleosomes, their location, mobility and density.

Activation of transcription typically leads to a change in chromatin conformation manifested in higher accessibility of the DNA to digestion or transposon integration (Tsompana and Buck, 2014). Although the changes in histone occupancy and accessibility have been studied extensively, the quantitative structural changes of chromatin on a single cell level remain poorly understood, and it is still largely unclear how the activities of chromatin remodeling complexes are spatially and temporally integrated in living cells. The chromatin 'opening' associated with transcriptional activation consists of two distinct changes in chromatin structure: spatial decompaction by changes in nucleosome-nucleosome interactions, and linear decompaction by changes in nucleosome density (Even-Faitelson *et al.*, 2016). Both of these processes are thought to be tightly linked to the posttranslational modifications of histones, which characterize transcriptionally active and transcriptionally silent chromatin. For example, histone tail acetylation is generally associated with active chromatin (Li *et al.*, 2007). *In vitro*, specific histone modifications directly influence the spatial organization of chromatin by mediating or restricting nucleosome-nucleosome interactions. For example, the acetylation of histone H4 at lysine 16 prevents the interaction between the acetylated tail and a neighboring nucleosome, lowering the propensity to form a compact 30 nm fiber (Shogren-Knaak *et al.*, 2006). However, no effect on linear chromatin compaction was observed on autosomes in mammalian cells and *C. elegans* upon loss of H4K16 modification during differentiation (Taylor *et al.*, 2013; Lau *et al.*, 2017). In addition, evidence is accumulating that extended regular higher order chromatin structures like a 30 nm fiber do not form *in vivo,* and that chromatin fibers are present mostly in dispersed states in living cells (Fussner *et al.*, 2012; Hsieh *et al.*, 2015; Chen *et al.*, 2016). This is corroborated by recent visualizations of chromatin chains in intact mammalian nuclei, showing that the locally appearance of the chromatin fiber is similar in regions of eu- and heterochromatin (Ou *et al.*, 2017). It is therefore unclear how local chromatin folding is influenced by histone modifications *in vivo*.

In addition to their potential to mediate inter- and intrachromosomal interactions and thus to shape
the longer-range organization of the chromatin fiber, nucleosomes constitute direct obstacles to the
passage of RNA Polymerase II (Pol II) during transcription. Histones and entire nucleosomes are
therefore evicted from the DNA during transcription by the concerted action of chromatin
remodelers like the SWI/SNF and SWR1 complexes and histone chaperones like Asf1 and FACT
(reviewed in Venkatesh and Workman, 2015). This nucleosome eviction (leading to a lower
nucleosome density) would be expected to cause an increase in the effective length of the chromatin
fiber, but this has not been investigated *in vivo*.

In order to study chromatin dynamics and organization *in vivo*, we took advantage of the budding
yeast *GAL* locus, a highly regulated gene cluster which has served as a paradigm for inducible gene
expression. The *GAL* locus comprises the *GAL7*, *GAL10* and *GAL1* genes located next to each other on
chromosome II. These three genes, encoding enzymes required for the metabolism of galactose, are
highly regulated depending on the carbon sources present in the growth medium. The genes are
repressed in the presence of glucose, active in the presence of galactose (but the absence of glucose)
and 'derepressed' in the absence of both glucose and galactose (e.g. in raffinose). Intricate
regulation of carbon metabolic genes allows *S. cerevisiae* to adapt to and successfully compete with
other organisms for various sugars present in the environment (New *et al.*, 2014).

GAL gene activation involves the recruitment of several histone modifying enzymes (Carrozza *et al.*,
2002; Wang *et al.*, 2002; Govind *et al.*, 2007) and the nucleosome occupancy at the locus is
dramatically reduced (Schwabish and Struhl, 2004; Govind *et al.*, 2007; Bryant *et al.*, 2008). How this
affects the chromatin conformation *in vivo* and to which extent chromatin decompaction and
transcriptional activation are interdependent remains unclear. To address these important questions,
we developed an assay to visualize chromatin compaction in living cells. Monitoring the distance
between two chromosomal loci on either side of the *GAL* locus revealed a drastic linear
decompaction upon activation of the *GAL* gene cluster. This decompaction was tightly coupled to
transcriptional activity. Furthermore, the observed opening was not regulated by histone acetylation
but depended on the activity of nucleosome-evicting chromatin remodelers.

Results

An assay to quantitatively analyze transcription-induced chromatin decompaction in living cells

Results
An assay to quantitatively analyze transcription-induced chromatin decompaction in living
cells
To probe chromatin decompaction during transcription in a quantitative manner in living cells, we
developed a microscopy-based assay to follow chromatin conformation in *S. cerevisiae* over time. We
chose the *GAL7-GAL10-GAL1* gene cluster as a model system, since it is very well studied and the
presence of three co-regulated genes spanning ~ 5.8 kb is expected to give a clear decompaction
response. LacO and TetO repeats were introduced on either side of the *GAL* gene cluster or in a
control region and visualized with LacI-GFP and TetR-mCherry (Figure 1A). Bright green and red dots
were readily detected in all cells (Figure 1A), and their distance was analyzed in cells grown in glucose
(repressed), raffinose (derepressed) or galactose (active). In a control region adjacent to but not
spanning the *GAL* locus (18 kb between LacO- and TetO-repeat insertion sites), no change in the
distribution of measured distances was observed between the different carbon sources (Figure 1B).
In contrast, in strains with a distance of 14 or 31 kb between the insertion points of the repeats
across the *GAL* locus, the distribution clearly shifted towards larger distances in activated cultures
(Figure 1B and C) increasing from a mean distance of 255±13 nm in glucose to 310±7 nm in galactose
in the 14 kb reporter strain or from 327±12 nm to 391±14 nm in the 31 kb reporter strain (2D
distances measured on single imaging plates; values are means of three biological replicates with
standard error of the mean). As expected, the deletion of the transcriptional activator Gal4 abolished

the decompaction response, while deletion of the transcriptional repressor Gal80 led to
decompaction already in raffinose-grown cells (Supplemental Figure 1). Thus, we have developed a
microscopy assay that can readily detect chromatin decompaction upon transcriptional activation in
living cells.

To analyze the dynamics of decompaction, we acutely induced the *GAL* genes by adding galactose to
cells growing in raffinose. This led to an increase in the population mean distance over the course of
20 minutes (Figure 1D) mimicking the kinetics of transcriptional activation as seen e.g. by
quantitative PCR (qPCR) analysis (Green *et al.*, 2012). Similarly, the addition of glucose to cells
growing in galactose resulted in fast compaction kinetics (Figure 1E) corresponding to the rapid shut-
down of transcription due to glucose-induced repression. Decompaction could also be induced with
faster kinetic within less than five minutes if cells were pregrown in galactose and then repressed
with glucose for only 1 h before re-induction with galactose (Figure 1F). In this case, the presence of
elevated levels of Gal1 and Gal3 proteins induce a 'transcriptional memory' and lead to faster
activation (Kundu and Peterson, 2010). Thus, the kinetics of chromatin compaction closely follows
transcriptional activity.

Active transcription correlates with chromatin decompaction in single cells
To directly correlate transcriptional activity with distance measurements on a single cell level, we
applied single molecule fluorescent *in situ* hybridization (smFISH) to visualize transcripts as they are
produced from the *GAL* locus in our distance reporter strains. We simultaneously used FISH probes
for all three *GAL* genes *GAL1*, *GAL10* and *GAL7*, all labeled with the same fluorophore (Qasar670).
While cytoplasmic mRNAs are visible as individual spots, or hazy signal due to the large number of
molecules in induced cells, the transcription site is visible as a bright focus in the nucleus close to the
repeat-marked gene loci (Figure 2A). Transcription spots and cytoplasmic mRNA foci are absent in
cells grown in raffinose (Figure 2A, first panel). Upon induction with galactose, the fraction of cells
with transcription spots increased over the course of 30 min paralleling the increase in the mean
distance in the population after induction (Figure 2B). After 60 minutes or longer, a transcription
spot can be seen in virtually every cell (Figure 2A), but since our automatic detection routine does
not detect all transcription spots, the maximum measured percentage of cells with transcription spot
is ~ 80 % (Figure 2B).

To correlate transcription with distance, we analyzed early time points after induction, when only a
fraction of the cells shows a transcription spot (Figure 2B). We separately determined the distance
between the LacO- and TetO-repeat markers in cells with or without a detectable transcription spot.
Importantly, distances were measured in 3D on image stacks to provide additional resolution, since
fixation and FISH procedures led to an overall shrinkage of the sample. While only few cells with a
clear transcription spot were observed after 5 min and decompaction was still minimal, after 10 and
30 min the distances in cells with a transcription spots were significantly larger than in cells without a
transcription spot (Figure 2C and D). Thus, decompaction correlates with transcription also on a
single cell level.

Histone acetylation is dispensable for chromatin decompaction at the *GAL* locus
The increased distance distribution observed in our assay, which is indicative of decompaction of
chromatin during transcription, could result from reduced internucleosomal interactions, from the
eviction of nucleosomes, or both. Internucleosomal interactions are often mediated by histone tails
and are thought to be regulated by posttranslational modification. Histones H3 and H4 contribute
most to direct internucleosomal interactions but are also required for the recruitment of
transcriptional activators. Acetylation of H3 *in vivo* occurs predominantly through Gcn5 (Grant *et al.*,
1997). This histone acetyl transferase is a component of the SAGA complex, which is recruited to the

GAL locus by the transcriptional activator Gal4 (Carrozza *et al.*, 2002; Govind *et al.*, 2007). However,
the deletion of neither the SAGA histone acetyl transferase Gcn5 nor its activating subunit Ada2
influenced chromatin decompaction across the *GAL* locus (Figure 3A). These mutations also do not
affect pre-initiation complex formation at the *GAL* promoter (Bhaumik and Green, 2001) or mRNA
production (Stafford and Morse, 2001; Green *et al.*, 2012), suggesting that acetylation is dispensable
for transcriptional activation and chromatin decompaction induced by Gal4. While dispensable for
activation, Gcn5 activity is crucial for the association of the *GAL* locus with the nuclear periphery
upon activation (Dultz *et al.*, 2016). Therefore, our results rule out that the observed decompaction
response is a consequence of relocalization of the *GAL* locus to the nuclear pore complex, which
might, for example, generate pulling forces that could lead to stretching of the chromatin fiber.

We also tested whether the inhibition of histone deacetylases using trichostatin A (TSA) would affect
chromatin decompaction. TSA treatment neither changed the steady state distance distributions
(Figure 3B) nor did it affect induction or repression kinetics (Figure 3C). Since both the deletion of
GCN5 and the inhibition of histone deacetylases targeted by TSA have been shown to globally affect
acetylation levels of histones (Vogelauer *et al.*, 2000), these results indicate that the global
acetylation state of histones *per se* does not play a major role in regulating chromatin compaction at
the *GAL* locus.

Mutants that affect transcription also affect decompaction
Although Gcn5 was dispensable for chromatin decompaction, deletion of the SAGA component
Spt20, which is required for the stability of the complex (Grant *et al.*, 1997) and for its recruitment by
Gal4 (Bhaumik and Green, 2001; Larschan and Winston, 2001; Bryant and Ptashne, 2003), led to
strongly reduced decompaction of the *GAL* locus (Figure 4 A and C). As previously reported (Bhaumik
and Green, 2001), Spt20 mutants also show strongly reduced *GAL* gene transcription (Figure 4B).

Whereas SAGA is thought to be required to stabilize the general transcriptional coactivator Mediator
at the *GAL* promoter, Gal4 is also able to directly recruit this complex (Traven *et al.*, 2006). Indeed,
cells with a deletion of *GAL11,* a Mediator component that mediates direct interaction with Gal4, or
expressing a temperature sensitive mutant of the Mediator protein Med7 (*med7-163*) phenocopied
spt20Δ cells by exhibiting reduced overall decompaction in galactose as well as slowed decompaction
kinetics (Figure 4 A and C and Supplemental Figure 2).

To analyze the effect of *SPT20* or *GAL11* deletions on a single cell level, we carried out single
molecule FISH. After overnight growth in galactose, most cells displayed cytoplasmic mRNA signals,
but the signal was strongly reduced compared to wildtype cells (Figure 4D). In addition, the intensity
of transcription spots, which were still present in most cells, were reduced on average to ~ 30 % of
the wildtype intensity (Figure 4D and E). Thus, also in the mutant cells, reduced transcription
correlates with reduced decompaction of the *GAL* locus. Our results reveal a close linkage between
transcriptional activation and decompaction, but they are not sufficient to establish a causality or
temporal order between the two processes.

Nucleosome remodelers are required for decompaction
Next, we examined the function of nucleosome remodelers in our decompaction assay. As
demonstrated by chromatin immunoprecipitation experiments, histone occupancy in the promoter
of transcribed genes – but also in the open reading frame (ORF) of highly transcribed Pol II genes
including the *GAL* genes – is reduced several fold upon activation (Lee *et al.*, 2004; Schwabish and
Struhl, 2004). Nucleosome eviction is mediated by the SWI/SNF chromatin remodeling complex and
the histone chaperones Asf1 (Antisilencing function 1) and FACT (FAcilitates Chromatin
Transcription), a heterodimeric complex consisting of Pob3 and Spt16 (Schwabish and Struhl, 2006,
2007; Venkatesh and Workman, 2015). If the observed decompaction across the *GAL* locus is due to

nucleosome eviction, depletion of either of these factors should lead to a reduction of
decompaction.

SWI/SNF is a chromatin remodeling complex with nucleosome sliding and nucleosome eviction
activities. At the *GAL* locus, it is required for rapid Pol II recruitment and activation (Kundu and
Peterson, 2010). We found that deletion of *SNF2*, encoding the ATPase subunit of the complex,
slowed down the kinetics of decompaction and strongly reduced the degree of final decompaction as
well (Figure 4A and C) consistent with a reduction in nucleosome eviction across the locus
(Schwabish and Struhl, 2006). Although inactivation of SWI/SNF has previously been reported not to
alter steady state *GAL* gene expression (Lemieux and Gaudreau, 2004), we observed strongly reduced
mRNA levels both by qPCR and sm FISH in snf2∆ cells (Figure 4B and D). In contrast to the SWI/SNF
complex, the chromatin remodeling complexes INO80 (represented by arp6∆) and SWR1
(represented by swr1∆) were not required for chromatin decompaction at the *GAL* locus (Figure 4A).

Histone chaperones play an important role in nucleosome eviction and deposition. Therefore, we
tested the involvement of the histone chaperones Asf1 and FACT, which have both been implicated
in the regulation of the *GAL* genes (Schwabish and Struhl, 2006; Xin *et al.*, 2009). Deletion of *ASF1* did
not prevent chromatin decompaction in our assay (Figure 5A), which is consistent with previous
findings that showed only a small decrease in histone H3 (but not H2B) removal at *GAL* promoters
and ORFs (Schwabish and Struhl, 2006). In contrast, cells carrying temperature sensitive alleles for
either component of the FACT complex (pob3-7 or spt16ts) did not show decompaction one hour
after a shift to activating conditions at the restrictive temperature (Figure 5B). Together, these
findings indicate that the eviction of nucleosomes mediated by SWI/SNF and FACT is the major
source of decompaction observed in our assay.

Decompaction of the *GAL* locus is transcription dependent
If the observed decompaction is due to the removal of nucleosomes at the promoter and the ORF,
the degree of decompaction will be expected to scale with the length of the ORF. To test this
prediction, we introduced *GAL* promoter driven reporter genes at a different genomic site ~ 10 kb
telomeric of the native *GAL* locus and measured changes in chromatin distance in the presence or
absence of galactose (Figure 6A). The *GAL* promoter immediately followed by a terminator sequence
did not exhibit a significant decompaction response. In contrast, robust decompaction was observed
in a strain where the entire *GAL1* gene including promoter and downstream sequences was inserted.
Decompaction, although to a weaker extent than for the entire *GAL1* gene, was also observed in a
strain where we introduced a *GAL1* promoter-driven ORF coding for Glutathione S transferase (GST).
Since the ORF of a single *GST* is shorter than the one of *GAL1* (700 bp versus 1 kb), we also examined
the response in a reporter strain carrying two consecutive copies of the *GST* ORF (*2xGST*). However,
the presence of a second *GST* ORF only slightly increased the decompaction, indicating that the *GAL1*
ORF or 3'UTR have specific properties that enhance the observed decompaction response. However,
the *2xGST* construct also contained MS2 stem loop sequences in the 5'UTR that might affect
transcription (Heinrich *et al.*, 2017) and thus decompaction.

Decompaction leads to an untypical open chromatin state at the *GAL* locus
Our data provide the first quantitative measurement of transcription-dependent decompaction in
living cells. To understand the changes in chromatin structure underlying the observed decompaction
response, we more closely analyzed the distance distributions of the populations. The 3D distance of
genomic loci on the same chromosome has been shown to scale with the distance in base pairs (van
den Engh *et al.*, 1992). This is true also on chromosome II of budding yeast in both glucose and
galactose grown cells as shown previously by us (Dultz *et al.*, 2016 and Figure 6B grey and black
datapoints). One explanation for the observed changes in compaction at the *GAL* locus could be that

the *GAL* locus – compared to bulk chromatin – is *hyper*-compacted in the repressed state. However,
this is not the case: in the repressed or derepressed state (glucose or raffinose growth), the median
distances of TetO and LacO marker pairs across the *GAL* locus (Figure 6B open red circles, data for
raffinose grown cells not shown) are in line with the distances of other pairs of loci on chromosome
II. This indicates that the compaction state of the locus is comparable to other regions on the same
chromosome. In contrast, in galactose-grown cells, the median distance increased far above the
distance expected from the linear distance on the chromosome (Figure 6B, filled red circles), showing
that the *GAL* locus is strongly *hypo*-compacted in its active state. This is consistent with the
interpretation that linear decompaction at the *GAL* locus upon transcriptional activation is due to the
eviction of nucleosomes, and that the high transcriptional activity over three clustered genes at the
GAL locus leads to a very low histone density and thus hypo-compaction.

Since our data provide quantitative measurements of this hypo-compaction, we applied
computational modeling to predict the level of histone loss that could lead to such a distance
increase. To this end, we adapted a previously developed polymer model of the yeast genome, which
accurately recapitulates several experimentally determined features (Tjong *et al.*, 2012; Dultz *et al.*,
2016), by increasing the resolution of the model in the proximity of the *GAL* locus. In this model, the
compaction of the chromatin fiber in the *GAL* locus had to be reduced by ~60% in order to
recapitulate the observed increase in distances (Figure 6C and D), which corresponds to a 1.7-fold
increase in linear extension. Since the wrapping of DNA on nucleosomes leads to an approximately 6-
fold compaction, this suggests that about 30 % of the nucleosomes at the *GAL* locus are evicted upon
transcriptional activation. This is in accordance with nucleosome occupancy measurements
(Schwabish and Struhl, 2006) and, at a density of 162bp/nucleosome (Horz and Zachau, 1980),
corresponds to the eviction of approximately 11 of 36 nucleosomes across the locus.

Discussion
Recently, the development of chromosome conformation capture techniques has revolutionized the
characterization of three-dimensional interaction landscapes of chromatin and has led to many
important discoveries and new models of chromatin organization (reviewed in Pombo and Dillon,
2015; Bonev and Cavalli, 2016). However, these techniques require the fixation of cells and are
usually applied to cell populations. Furthermore, it is not possible to directly convert the interactions
detected by chromosome conformation capture into 3D distances in the cell nucleus. In fact, a direct
comparison of chromosome capture analyses with DNA fluorescence *in situ* hybridisation results has
revealed that the two techniques do not always result in congruent findings (Williamson *et al.*, 2014).
It is therefore crucial to analyze chromatin conformation *in situ* and also *in vivo*.

Chromatin decompaction upon transcriptional activation has been visualized in mammalian cells
using large repeat arrays (Tumbar *et al.*, 1999; Dietzel *et al.*, 2004; Verschure *et al.*, 2005; Hu *et al.*,
2009). FISH analysis has also been employed to obtain distance measurements at activated loci
(Chambeyron and Bickmore, 2004; Chambeyron *et al.*, 2005; Morey *et al.*, 2007). Although making
important contributions to our understanding of changes in chromatin conformation upon
transcriptional activation, quantitative interpretation is limited because fixation and DNA FISH
protocols introduce modifications to the cell ultrastructure and repeat arrays may themselves
influence the local chromatin environment. Therefore, we have used the budding yeast
Saccharomyces cerevisiae to generate a system where live cell analysis is possible in a quantitative
manner at an endogenous gene locus. Although we initially use a 3-gene system to establish our
assay, we also show that this assay is sensitive enough to visualize decompaction of a single activated
gene. Furthermore, the use of budding yeast allows us to study the basic level of chromatin
decompaction in the absence of confounding changes in chromatin conformation like enhancer-
promoter looping or looping out from a chromosome territory. Thus, our data yield quantitative

measurements of chromatin decompaction upon transcriptional activation and allow us to model the
changes in chromatin properties at an activated locus.

The role of acetylation

Interestingly, we found that deletion of the lysine acetyl transferase gene *GCN5* or inhibition of lysine
deacetylation by trichostatin A had no observable effect on the distance distribution in our
compaction assay. In contrast, the treatment of mammalian cells with TSA was previously reported
to affect chromatin organization globally and to lead to decompaction of heterochromatic regions
(Toth *et al.*, 2004; Lleres *et al.*, 2009). However, these studies used global histone signals (intensity or
FRET-FLIM) to assess compaction levels. These analyses thus do not give sequence-specific readout
and report on spatial rather than on linear compaction. In agreement with our findings, DNA FISH
analysis in mouse cells, which also tested linear compaction between two loci, showed decompaction
in polycomb silenced regions but not elsewhere (Eskeland *et al.*, 2010). Thus, very low acetylation
levels may be crucial for the formation of certain subnuclear compartments in heterochromatic
regions of the nucleus but may play a less prominent role in linear decompaction in the relatively
open bulk yeast chromatin. However, it cannot be excluded that specific acetylation added by acetyl
transferases other than Gcn5 or removed by non-TSA sensitive deacetylases contributes to linear
chromatin decompaction. Furthermore, acetylation could facilitate the binding of factors to specific
DNA sequences by rendering the interactions between histones and DNA less tight. This could allow
nucleosomes to slide more easily or create binding sites for bromodomain-containing chromatin
effector proteins (Filippakopoulos and Knapp, 2014). In contrast to the *GAL* genes, other budding
yeast genes were shown to fully depend on the HAT activity of Gcn5 for transcriptional activation.
For example, this is true for a heterologous β-estradiol responsive transcriptional activator based on
the viral VP16 transactivation domain (Stafford and Morse, 2001). Intriguingly, we found that upon
activation via a β-estradiol-inducible VP16 fusion protein (Louvion *et al.*, 1993), Gcn5 is essential for
decompaction (Supplemental Figure 3) and neither *gcn5Δ* nor *spt20Δ* cells were able to decompact
the *GAL* locus upon addition of β-estradiol (Supplemental Figure 3). This observation also
corroborates the conclusion that transcriptional activation precedes decompaction.

The interplay of transcription and decompaction

Our mutant analysis shows that transcriptional activation and chromatin decompaction are tightly
coupled. To directly address the role of Pol II in decompaction, we carried out experiments with cells
expressing a temperature-sensitive mutant allele of the largest subunit of RNA Pol II, *rpb1-1*. *rpb1-1*
cells were induced with galactose one hour after a shift to the restrictive temperature. Both
expression of *GAL10* mRNA and decompaction were reduced but not abolished (Supplemental Figure
4A and B). Thus, it appears that induction at this locus is so strong that the temperature-sensitive
rpb1-1 mutant is not fully effective to suppress decompaction and transcription. Nevertheless,
together with the observation that an ectopic *GAL* promoter in the absence of an ORF is not
sufficient to confer decompaction (Figure 6A), our results suggest that ORF transcription by Pol II is
required for decompaction of the chromatin at the *GAL* locus. This is consistent with the observation
in mammalian cells that only initial decompaction is mediated by transcriptional activation but for
complete decompaction elongation by RNA polymerase is necessary (Hu *et al.*, 2009). We
hypothesize that the removal of nucleosomes at the promoter presents a first stage of
decompaction, but elongation and the concomitant eviction of nucleosomes along the ORF
constitutes the major contribution to chromatin opening. This interpretation is in line with the
observation that opening of the *GAL* locus is not detected when both glucose and galactose are
present (Figure 4C Galactose+Glucose condition), a condition where SWI/SNF activity has been
reported to evict nucleosomes at the promoter (Bryant *et al.*, 2008). However, in addition to the
length of the ORF, the strength of activation and possibly the composition of the 3' and 5' UTRs may

also contribute to the level of nucleosome eviction and thus decompaction. Importantly, the absence
of a decompaction response from the promoter alone also indicates that longer range interactions
mediated by the promoter, for example promoter-enhancer loops, do not play a role in our system
allowing us to focus on the local changes in chromatin organization. Such effects or the release from
higher spatial compaction states may have contributed to the transcription-independent
decompaction that were observed previously (Hu *et al.*, 2009).

After transcriptional repression by the addition of glucose, the signal of *GAL* mRNAs in the cytoplasm
quickly diminishes. Although the nuclear FISH signal at the site of transcription is reduced as well, a
weak spot can still be detected in many cells even 60 minutes after glucose addition (Supplemental
Figure 5A and C). This most likely corresponds to polyadenylated *GAL* mRNAs previously reported to
persist at the *GAL* genes upon transcriptional shut-off (Abruzzi *et al.*, 2006). Interestingly, cells with a
brighter RNA dots showed a distance distribution shifted to longer distances compared to cells with a
weaker or absent RNA dot signal (Supplemental Figure 5B). This could indicate that transcription is
still ongoing in those cells and thus the brighter RNA signal would stem from nascent transcripts.
Alternatively, long-lived, chromatin-associated RNA dots might contribute to keeping chromatin in a
partially decompacted state.

The role of nucleosome eviction
Histones are evicted during transcriptional elongation but repositioned after the passage of Pol II.
Both eviction and deposition require the activity of histone chaperones like FACT. In highly
transcribed genes like the *GAL* genes, individual transcription events follow closely after each other
(Lenstra *et al.*, 2015). In this case, deposition of histones between rapid rounds of transcription may
not be possible, and thus a net loss of histones and nucleosomes occurs. Such a response would
explain the dramatic unfolding of chromatin across the *GAL* locus observed in our assay. This is
consistent with our observation that mutants with reduced transcriptional activity display a reduced
decompaction. Here, initiation and polymerase II progression occurs at a lower frequency, which
could allow for more time to deposit histones after the passage of Pol II. This interpretation is also in
agreement with the observed correlation between ORF length and decompaction. However, for the
moment it remains unclear why the *GAL1* ORF and its native 3'UTR confer higher levels of
decompaction compared to a GST ORF terminated by an ADH1 terminator. Of note, the presence of
MS2 stem loops in our 2xGST construct may present a serious obstacle to Pol II (Heinrich *et al.*, 2017)
and therefore could lead to reduced Pol II density on the body of the gene.

Outlook
Transcription initiation displays large cell-to-cell variability, and the same can be expected also for
decompaction. Unfortunately, we were not able to reliably measure decompaction kinetics on a
single cell level, due to the large fluctuations in distances measured over time even for cells in steady
state growth conditions. Therefore, further developments of reporter systems with reduced
biological noise in combination with improved superresolution microscopy approaches will be
needed to enable the analysis of decompaction kinetics on the single cell level. For example, the
contribution of DNA sequences that are not involved in the observed response could be minimized
by placing fluorescent reporters directly adjacent to the gene of interest. Adaptions of our system
thus have the potential to shed further light on the dynamics of chromatin decompaction by allowing
to study this process quantitatively in single cells under physiological conditions. The data from live
cell microscopy experiments are highly complementary to data obtained from chromosome capture
techniques and are very valuable to better understand chromatin organization and its dynamics *in*
vivo. Furthermore, using CRISPR/Cas technologies, our assay could be readily transferred to higher
eukaryotic systems as well. Optimized cassettes for the integration of LacO repeats by CRISPR/Cas
have recently been created (Tasan *et al.*, 2017) and the targeting of labelled CIRSPR/Cas complexes

(Chen *et al.*, 2013; Chen and Huang, 2014; Ma *et al.*, 2016) will in the future provide a powerful and
flexible tool to dissect the contribution of individual players to chromatin compaction in various
model systems.

Methods

Plasmid construction

Plasmid were constructed using standard molecular biology techniques. pKW1008 was constructed
by ligating annealed oligos UC586/867 into pFA6-GFP(s65t)-kan (Longtine *et al.*, 1998) using PacI/AscI
to replace GFP with a FLAG-tag. pKW2695, pKW2704 and pKW3264 were constructed by ligating PCR
products of primers CH4505/CH4506, CH4513/CH4514 or CH6/CH106 on genomic DNA into
pKW1689 (Green *et al.*, 2012) after cutting vector and insert with XhoI and SacI. pKW3035 was
constructed from a PCR product of UC5681/UC5682 on pKW3010 (Backlund *et al.*, 2014) cut with
DraIII and ligated into pAFS135 (Straight *et al.*, 1996) also cut with DraIII. pKW3681 and pKW3682
were constructed by inserting fragments generated by primers CH1211/CH1213 or CH1211/CH1177
on genomic DNA into the pFA6a backbone containing a NatMx resistance cassette (cut with
PacI/XhoI). pKW3683 was constructed by inserting into the same vector a stitched PCR product of the
GAL1 promoter (CH1211/CH1218) and GST (CH1214/CH1215) with PacI/XhoI. pKW3999 was
constructed in multiple steps. The final construct contains homology regions for a region on
chromosome II which were amplified from genomic DNA using primers CH1795/CH1796 and
CH1797/CH1798. Between the homology sites, the *GAL1* promoter amplified by CH1211/CH2043
drives a construct with 24 MS2 stem loops (Grunwald and Singer, 2010) in the 5' UTR and encoding
2xGST (CH1789/CH1794 and CH1791/CH1794) tagged with V5. It also contains the NatNT2 resistance
cassette for selection and can be integrated after cutting NotI/AscI. The Gal4DBD-ER-VP16 activation
construct was transferred from the construct published in (Louvion *et al.*, 1993) into a YIplac204
backbone with an additional NatMx cassette added for selection in TRP+ strains (pKW3504). All
plasmids used in this study are listed in Supplementary Table 1. Primers are listed in Supplementary
Table 2.

Yeast strain construction

S. cerevisiae strains were constructed in the background of BY4741 and BY4742 (Brachmann *et al.*,
1998) using standard yeast genetic techniques either by transformation of a linearized plasmid or of
a PCR amplification product with homology to the target site (Baudin *et al.*, 1993). The mutants
spt20Δ, gal80Δ, gcn5Δ, asf1Δ, snf2Δ, gal11Δ, pob3-7, spt16ts, med7-163, rpb1-1, gal4Δ, swr1Δ and
arp6Δ were constructed by mating of the wildtype strains with the corresponding strains from the
MATa, MATα deletion collections (Winzeler *et al.*, 1999) or the MATa temperature sensitive
collection (Li *et al.*, 2011) followed by sporulation and tetrad dissection. *ADA2* was deleted by PCR-
directed mutagenesis using primers CH395/CH396 on pKW1008 to generate KWY5104. Reporter
strains for the experiments shown in Figure 6A were generated by transformation of KWY4067 with
the PCR product of CH1272/CH1273 on pKW3681 (=> KWY6245), pKW3682 (=> KWY6247) or
pKW3683 (=> KWY6252) respectively. Genotypes were confirmed by PCR. All yeast strains used in
this study are listed in Supplementary Table 3.

Yeast culture conditions

Yeast cells were cultured in complete synthetic medium with 2 % of either glucose, galactose or
raffinose as indicated at 30 ˚C. Temperature sensitive mutants were grown at 25 ˚C. Some kinetic
experiments with mutants showing significant growth defects were also carried out at 25 ˚C. All
experiments were carried out with cells in exponential growth phase. For microscopy experiments,
cells were usually inoculated from saturated cultures into fresh medium and grown overnight to OD
0.5-0.8 and then imaged. Alternatively, cells were diluted in the morning and grown for additional 2-

3 cell cycles before the start of the experiment. Trichostatin A (Sigma Catalog #:T8552) was used at
50 μM.

Microscopy

Cells were pregrown in raffinose-containing medium and inoculated in 1 ml in 24 well plates
overnight so that they reached an OD of 0.5-0.8 in the morning. Cells were then transferred to
concanavalin A coated 384 well plates (Matrical) and imaged on a temperature controlled Nikon Ti
Eclipse equipped with a Spectra LED lamp using a Apochromat VC 100x objective NA 1.4 (Nikon)
(filters: Spectra emission filters 475/28 & 542/27 and DAPI/FITC/Cy3/Cy5 Quad HC Filterset with
410/504/582/669 HC Quad dichroic and a 440/521/607/700 HC quadband filter (Semrock)) with
exposure times of 50-200 ms. For timelapse experiments, induction with different sugars or
trichostatin A was carried out on stage after the first imaging timepoint by mixing with an equal
volume of trichostatin or sugar containing medium. Different fields of view were imaged at each
timepoint to prevent negative effects of phototoxicity or bleaching on the analysis. 3D microscopy of
living cells was carried out on a SpinningDisk microscope (Yokogawa Confocal Scanner Unit CSU-W1-
T2) built on a Nikon TiE body and controlled with the VisiVIEW software using dual camera
acquisition mode and a 100x NA 1.49 CFI Apo TIRF objective. Cameras were Orca Flash 4.0 V2,
excitation lasers were a DPSS 488 nm (200mW) and a diode 561 nm (200 mW) laser. Z scanning was
performed in streaming mode with a LUDL BioPrecision2 Piezo Stage using 2-by-2 binning with 100
ms exposure times per frame. Filters were: Dichroic quad-band DAPI/GFP/RFP/CY5, splitting filter to
camera ports: 561LP, emission filters GFP/ET525/50 and mCherry ET630/75 respectively.

Single molecule fluorescence in situ hybridization

Single molecule fluorescence *in situ* hybridization was carried out according to (Mugler *et al.*, 2016)
with slight adaptations. The indicated strains were inoculated in synthetic medium containing 2 %
raffinose and grown overnight to saturation. The next day, cells were diluted into fresh synthetic
complete media containing 2 % raffinose or 2 % galactose and grown overnight at 30 °C to
exponential growth phase (OD600 = 0.6-0.8). Cells were then induced by addition of glucose or
galactose to final concentration of 2 % and fixed after the indicated timepoints for 15 min at 30 °C
and for 15 min at 25 °C with 4 % paraformaldehyde (EM grade 32 % paraformaldehyde acqueous
solution electron Microscopy Sciences 15714), washed with buffer B (1.2 M sorbitol, 100 mM $KHPO_4$
at pH 7.5, 4 °C) and stored at 4 °C overnight. Cells were then spheroplasted for 20 min using 1 % 20T
zymolyase in 1.2 M sorbitol, 100 mM $KHPO_4$ at pH 7.5, 20 mM vanadyl ribonuclease complex and
20 μM β-mercaptoethanol, washed with buffer B to stop the spheroplasting reaction and then
washed into 10 % formamide (Merck Millipore S4117) in 2x SSC.

Mixtures of DNA probes coupled to Quasar670 (Stellaris, LGC Biosearch, Novato, CA; probes were
synthesized by BioCat, Heidelberg, Germany) were used for smFISH, targeting the *GAL1*, *GAL7* and
GAL10 ORFs (Supplementary Table 4). Per sample, 0.5 μl of each probe mix (stock 25 uM) was mixed
with 2 μl of salmon-sperm DNA (10 mg ml−1, Life Technologies, 15632-011) and 2 μl yeast transfer
RNA (10 mg/ml, Life Technologies, AM7119). The probe mix was denatured in 50 μl per sample of
hybridization buffer F (20 % formamide, 10 mM $NaHPO_4$ at pH 7.0) for 3 min at 95 °C and then mixed
with 50 μl per sample hybridization buffer H (4× SSC, 4 mg/ml BSA (acetylated) and 20 mM vanadyl
ribonuclease complex). Cells (approximately corresponding to pellet of 5 ml initial culture) were
resuspended in the hybridization mix and incubated for 8-12 hours at 37 °C. After four washing steps
(10 % formamide/2× SSC; 0.1 % Triton/2x SSC; 2x SSC/DAPI; 2x SSC), cells were stored at 4 °C. Cells
were imaged in concanavaline A coated 384 wells. Microscopy was performed with an inverted epi-
fluorescence microscope (Nikon Ti) equipped with a Spectra X LED light source and a Hamamatsu
Flash 4.0 sCMOS camera using a PlanApo 100 x NA 1.4 oil-immersion objective and the NIS Elements
software. 31 z planes were acquired. The stack of the Qasar670 channel was acquired first due to

significant bleaching. mCherry and GFP channels were acquired plane by plane to minimize shift between channels.

Image analysis

Images were processed using FIJI (NIH ImageJ 1.51p and previous) and Diatrack (Vallotton *et al.*, 2017). For 2D analysis of distances, position of chromosome location marked with LacI or TetR were detected by a 2D Gaussian fit using custom written scripts in Matlab (Dultz *et al.*, 2016). Distances between detected spots were calculated. Due to negligible shift between the two channels in these datasets, correction was not required. For 3D analysis of distances (data acquired on dual camera spinning disk microscope), positions of chromosome location marked with LacI or TetR were detected in Diatrack in 3D using across-color tracking to pair corresponding spots from the same cell (parameters used: 'exclude blurred' 0.07-0.09, 'exclude dim' 20-200, full width half maximum for Gaussian fitting 2.5). Subsequently, positions were corrected for shift between the two channels using bead images (1 μm TetraSpeck fluorescent microspheres, ThermoFisher Scientific) acquired under identical imaging conditions. For shift correction, the field of view was subdivided into 150x150 pixel large squares and the mean shift for beads acquired in this region was applied to correct the positions of chromosome locations acquired in this region of the camera. The distances between corrected positions were calculated and plotted in Matlab as boxplots. For FISH analysis, position of chromosome location marked with LacI or TetR were detected in Diatrack in 3D using across-color tracking to pair corresponding spots from the same cell. Positions of transcriptions spots were detected on maximum intensity projections of the Qasar670 channel in Diatrack and used to categorize cells as having or not having a transcription spot for separate analysis of the two populations. For analysis of transcription spot intensities, transcription spots were detected on 3D stacks in Diatrack and intensity values were obtained from the Diatrack session file. Only the intensity of those transcription spots for which corresponding chromosome locations marked with LacI or TetR could be detected in Diatrack were included in the analysis. Bootstrapping was used to obtain 95 % confidence intervals of the medians and mean. Typically 200-400 cells (but at least 90) were analyzed per condition in each experiment.

Quantitative real-time PCR

qPCR was performed as described previously (Dultz *et al.*, 2016): 1 ml of cells at OD600 0.8-1 was harvested by centrifugation and snap frozen in liquid nitrogen. RNA was extracted using the RNeasy kit from Qiagen via mechanical disruption. 300 ng of total extracted RNA was used for reverse transcription. The RNA was first treated with DNase I using the DNA-free kit from Ambion according to the protocol of the manufacturer. Reverse transcription was performed according to the protocol of the manufacturer using Superscript II reverse transcriptase (Invitrogen) with random hexamer primers. Quantitative Real Time PCR was performed on a StepOnePlus Instrument (Invitrogen) using Absolute Blue QPCR Mix with SYBR Green and ROX (ThermoFisher Scientific). All experiments were carried out in three technical replicates and three biological replicates. Data was analyzed by the comparative CT method using *ACT1* as endogenous control. Primers used for qPCR are listed in Supplementary Table 2.

Computational modeling

Average spatial distances were calculated from ensembles containing 1000 structure models of the entire haploid yeast genome. Ensembles were generated for each of the two *GAL* locus activation states (active/galactose and inactive/glucose), as described in (Tjong *et al.*, 2012; Dultz *et al.*, 2016). All chromosomes were modeled as chains of connecting beads subject to a number of spatial restraints. In particular, all chromosomes were confined inside a nucleus with a radius of 1 μm; the nucleolus and spindle pole body (SPB) are placed on opposite sides of the nucleus along the central nuclear axis; all non-ribosomal-DNA gene regions are excluded from the nucleolar volume;

centromeric regions are proximal to the SPB, whereas telomeric regions are tethered to the nuclear envelope (allowing a maximal distance between telomeres and the nuclear envelope of 50 nm). Each bead accommodated ~3.2kb of genome sequence as described in (Tjong *et al.*, 2012).

To reproduce the experimental data, we modeled a gradual decrease in chromatin compaction in the proximity of the *GAL* locus. For the 60 kb region starting at the position of the *GAL* locus, the compaction ratio was set to 1.6 kb per bead. We then determined the optimal chromatin compaction for the 6.2 kb *GAL* locus so that the models reproduce most closely the experimentally observed 3D fluorophore distances in both activation states. The *GAL* locus compaction in the inactive glucose state was found to be ~375 bp/bead whereas in the active galactose state it was set to ~200 bp/bead.

Each ensemble of genome structures was generated from 1000 independent simulations, each starting from random configurations. The optimization procedure consisted of a simulated annealing Molecular Dynamics run followed by Conjugate Gradients score minimization, both performed using IMP (Russel *et al.*, 2012).

Author contributions

Conception and design of the work: ED and KW; Data collection: ED and RM; Data analysis and interpretation: ED and RM; development of code for image analysis: ED and PV; modelling: GP & FA Drafting the article: ED; critical revision of the article: ED, RM, PV, GP, FA & KW.

Acknowledgements

We are grateful to members of the Weis laboratory for critical discussions and suggestions on the manuscript. We thank the Scientific Center for Optical and Electron Microscopy of ETHZ (ScopeM) for support with microscopy. This work was supported by the Swiss National Fond (SNF 159731) and NIH (U54DK107981 to F.A).

References

Abruzzi, K.C., Belostotsky, D.A., Chekanova, J.A., Dower, K., and Rosbash, M. (2006). 3'-end formation signals modulate the association of genes with the nuclear periphery as well as mRNP dot formation. The EMBO journal *25*, 4253-4262.

Backlund, M.P., Joyner, R., Weis, K., and Moerner, W.E. (2014). Correlations of three-dimensional motion of chromosomal loci in yeast revealed by the double-helix point spread function microscope. Molecular biology of the cell *25*, 3619-3629.

Baudin, A., Ozier-Kalogeropoulos, O., Denouel, A., Lacroute, F., and Cullin, C. (1993). A simple and efficient method for direct gene deletion in Saccharomyces cerevisiae. Nucleic acids research *21*, 3329-3330.

Bhaumik, S.R., and Green, M.R. (2001). SAGA is an essential in vivo target of the yeast acidic activator Gal4p. Genes & development *15*, 1935-1945.

Bonev, B., and Cavalli, G. (2016). Organization and function of the 3D genome. Nature reviews. Genetics *17*, 661-678.

Brachmann, C.B., Davies, A., Cost, G.J., Caputo, E., Li, J., Hieter, P., and Boeke, J.D. (1998). Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. Yeast *14*, 115-132.

Bryant, G.O., Prabhu, V., Floer, M., Wang, X., Spagna, D., Schreiber, D., and Ptashne, M. (2008). Activator control of nucleosome occupancy in activation and repression of transcription. PLoS biology *6*, 2928-2939.

Bryant, G.O., and Ptashne, M. (2003). Independent recruitment in vivo by Gal4 of two complexes required for transcription. Molecular cell *11*, 1301-1309.

Carrozza, M.J., John, S., Sil, A.K., Hopper, J.E., and Workman, J.L. (2002). Gal80 confers specificity on HAT complex interactions with activators. The Journal of biological chemistry *277*, 24648-24652.

Chambeyron, S., and Bickmore, W.A. (2004). Chromatin decondensation and nuclear reorganization of the HoxB locus upon induction of transcription. Genes & development *18*, 1119-1130.

Chambeyron, S., Da Silva, N.R., Lawson, K.A., and Bickmore, W.A. (2005). Nuclear re-organisation of the Hoxb complex during mouse embryonic development. Development *132*, 2215-2223.

Chen, B., Gilbert, L.A., Cimini, B.A., Schnitzbauer, J., Zhang, W., Li, G.W., Park, J., Blackburn, E.H., Weissman, J.S., Qi, L.S., and Huang, B. (2013). Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. Cell *155*, 1479-1491.

Chen, B., and Huang, B. (2014). Imaging Genomic Elements in Living Cells Using CRISPR/Cas9. Methods in enzymology *546*, 337-354.

Chen, C., Lim, H.H., Shi, J., Tamura, S., Maeshima, K., Surana, U., and Gan, L. (2016). Budding yeast chromatin is dispersed in a crowded nucleoplasm in vivo. Molecular biology of the cell.

Dietzel, S., Zolghadr, K., Hepperger, C., and Belmont, A.S. (2004). Differential large-scale chromatin compaction and intranuclear positioning of transcribed versus non-transcribed transgene arrays containing beta-globin regulatory sequences. Journal of cell science *117*, 4603-4614.

Dultz, E., Tjong, H., Weider, E., Herzog, M., Young, B., Brune, C., Müllner, D., Loewen, C., Alber, F.,
and Weis, K. (2016). Global reorganization of budding yeast chromosome conformation in different
physiological conditions. The Journal of cell biology, jcb.201507069.

Eskeland, R., Freyer, E., Leeb, M., Wutz, A., and Bickmore, W.A. (2010). Histone acetylation and the
maintenance of chromatin compaction by Polycomb repressive complexes. Cold Spring Harbor
symposia on quantitative biology *75*, 71-78.

Even-Faitelson, L., Hassan-Zadeh, V., Baghestani, Z., and Bazett-Jones, D.P. (2016). Coming to terms
with chromatin structure. Chromosoma *125*, 95-110.

Filippakopoulos, P., and Knapp, S. (2014). Targeting bromodomains: epigenetic readers of lysine
acetylation. Nature reviews. Drug discovery *13*, 337-356.

Fussner, E., Strauss, M., Djuric, U., Li, R., Ahmed, K., Hart, M., Ellis, J., and Bazett-Jones, D.P. (2012).
Open and closed domains in the mouse genome are configured as 10-nm chromatin fibres. EMBO
reports *13*, 992-996.

Govind, C.K., Zhang, F., Qiu, H., Hofmeyer, K., and Hinnebusch, A.G. (2007). Gcn5 promotes
acetylation, eviction, and methylation of nucleosomes in transcribed coding regions. Molecular cell
25, 31-42.

Grant, P.A., Duggan, L., Cote, J., Roberts, S.M., Brownell, J.E., Candau, R., Ohba, R., Owen-Hughes, T.,
Allis, C.D., Winston, F., Berger, S.L., and Workman, J.L. (1997). Yeast Gcn5 functions in two
multisubunit complexes to acetylate nucleosomal histones: characterization of an Ada complex and
the SAGA (Spt/Ada) complex. Genes & development *11*, 1640-1650.

Green, E.M., Jiang, Y., Joyner, R., and Weis, K. (2012). A negative feedback loop at the nuclear
periphery regulates GAL gene expression. Molecular biology of the cell *23*, 1367-1375.

Grunwald, D., and Singer, R.H. (2010). In vivo imaging of labelled endogenous beta-actin mRNA
during nucleocytoplasmic transport. Nature *467*, 604-607.

Heinrich, S., Sidler, C.L., Azzalin, C.M., and Weis, K. (2017). Stem-loop RNA labeling can affect nuclear
and cytoplasmic mRNA processing. Rna *23*, 134-141.

Horz, W., and Zachau, H.G. (1980). Deoxyribonuclease II as a probe for chromatin structure. I.
Location of cleavage sites. J Mol Biol *144*, 305-327.

Hsieh, T.H., Weiner, A., Lajoie, B., Dekker, J., Friedman, N., and Rando, O.J. (2015). Mapping
Nucleosome Resolution Chromosome Folding in Yeast by Micro-C. Cell *162*, 108-119.

Hu, Y., Kireev, I., Plutz, M., Ashourian, N., and Belmont, A.S. (2009). Large-scale chromatin structure
of inducible genes: transcription on a condensed, linear template. The Journal of cell biology *185*, 87-
100.

Kundu, S., and Peterson, C.L. (2010). Dominant role for signal transduction in the transcriptional
memory of yeast GAL genes. Molecular and cellular biology *30*, 2330-2340.

Larschan, E., and Winston, F. (2001). The S. cerevisiae SAGA complex functions in vivo as a
coactivator for transcriptional activation by Gal4. Genes & development *15*, 1946-1956.

Lau, M.S., Schwartz, M.G., Kundu, S., Savol, A.J., Wang, P.I., Marr, S.K., Grau, D.J., Schorderet, P.,
Sadreyev, R.I., Tabin, C.J., and Kingston, R.E. (2017). Mutation of a nucleosome compaction region
disrupts Polycomb-mediated axial patterning. Science *355*, 1081-1084.

1 Lee, C.K., Shibata, Y., Rao, B., Strahl, B.D., and Lieb, J.D. (2004). Evidence for nucleosome depletion at
2 active regulatory regions genome-wide. Nature genetics *36*, 900-905.

3 Lemieux, K., and Gaudreau, L. (2004). Targeting of Swi/Snf to the yeast GAL1 UAS G requires the
4 Mediator, TAF IIs, and RNA polymerase II. The EMBO journal *23*, 4040-4050.

5 Lenstra, T.L., Coulon, A., Chow, C.C., and Larson, D.R. (2015). Single-Molecule Imaging Reveals a
6 Switch between Spurious and Functional ncRNA Transcription. Molecular cell *60*, 597-610.

7 Li, B., Carey, M., and Workman, J.L. (2007). The role of chromatin during transcription. Cell *128*, 707-
8 719.

9 Li, Z., Vizeacoumar, F.J., Bahr, S., Li, J., Warringer, J., Vizeacoumar, F.S., Min, R., Vandersluis, B.,
10 Bellay, J., Devit, M., Fleming, J.A., Stephens, A., Haase, J., Lin, Z.Y., Baryshnikova, A., Lu, H., Yan, Z.,
11 Jin, K., Barker, S., Datti, A., Giaever, G., Nislow, C., Bulawa, C., Myers, C.L., Costanzo, M., Gingras, A.C.,
12 Zhang, Z., Blomberg, A., Bloom, K., Andrews, B., and Boone, C. (2011). Systematic exploration of
13 essential yeast gene function with temperature-sensitive mutants. Nature biotechnology *29*, 361-
14 367.

15 Lleres, D., James, J., Swift, S., Norman, D.G., and Lamond, A.I. (2009). Quantitative analysis of
16 chromatin compaction in living cells using FLIM-FRET. The Journal of cell biology *187*, 481-496.

17 Longtine, M.S., McKenzie, A., 3rd, Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and
18 Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and
19 modification in Saccharomyces cerevisiae. Yeast *14*, 953-961.

20 Louvion, J.F., Havaux-Copf, B., and Picard, D. (1993). Fusion of GAL4-VP16 to a steroid-binding
21 domain provides a tool for gratuitous induction of galactose-responsive genes in yeast. Gene *131*,
22 129-134.

23 Ma, H., Tu, L.C., Naseri, A., Huisman, M., Zhang, S., Grunwald, D., and Pederson, T. (2016).
24 Multiplexed labeling of genomic loci with dCas9 and engineered sgRNAs using CRISPRainbow. Nature
25 biotechnology *34*, 528-530.

26 Morey, C., Da Silva, N.R., Perry, P., and Bickmore, W.A. (2007). Nuclear reorganisation and chromatin
27 decondensation are conserved, but distinct, mechanisms linked to Hox gene activation. Development
28 *134*, 909-919.

29 Mugler, C.F., Hondele, M., Heinrich, S., Sachdev, R., Vallotton, P., Koek, A.Y., Chan, L.Y., and Weis, K.
30 (2016). ATPase activity of the DEAD-box protein Dhh1 controls processing body formation. eLife *5*.

31 New, A.M., Cerulus, B., Govers, S.K., Perez-Samper, G., Zhu, B., Boogmans, S., Xavier, J.B., and
32 Verstrepen, K.J. (2014). Different levels of catabolite repression optimize growth in stable and
33 variable environments. PLoS biology *12*, e1001764.

34 Ou, H.D., Phan, S., Deerinck, T.J., Thor, A., Ellisman, M.H., and O'Shea, C.C. (2017). ChromEMT:
35 Visualizing 3D chromatin structure and compaction in interphase and mitotic cells. Science *357*.

36 Pombo, A., and Dillon, N. (2015). Three-dimensional genome architecture: players and mechanisms.
37 Nature reviews. Molecular cell biology *16*, 245-257.

38 Russel, D., Lasker, K., Webb, B., Velazquez-Muriel, J., Tjioe, E., Schneidman-Duhovny, D., Peterson, B.,
39 and Sali, A. (2012). Putting the pieces together: integrative modeling platform software for structure
40 determination of macromolecular assemblies. PLoS biology *10*, e1001244.

Schwabish, M.A., and Struhl, K. (2004). Evidence for eviction and rapid deposition of histones upon
transcriptional elongation by RNA polymerase II. Molecular and cellular biology *24*, 10111-10117.

Schwabish, M.A., and Struhl, K. (2006). Asf1 mediates histone eviction and deposition during
elongation by RNA polymerase II. Molecular cell *22*, 415-422.

Schwabish, M.A., and Struhl, K. (2007). The Swi/Snf complex is important for histone eviction during
transcriptional activation and RNA polymerase II elongation in vivo. Molecular and cellular biology
27, 6987-6995.

Shogren-Knaak, M., Ishii, H., Sun, J.M., Pazin, M.J., Davie, J.R., and Peterson, C.L. (2006). Histone H4-
K16 acetylation controls chromatin structure and protein interactions. Science *311*, 844-847.

Stafford, G.A., and Morse, R.H. (2001). GCN5 dependence of chromatin remodeling and
transcriptional activation by the GAL4 and VP16 activation domains in budding yeast. Molecular and
cellular biology *21*, 4568-4578.

Straight, A.F., Belmont, A.S., Robinett, C.C., and Murray, A.W. (1996). GFP tagging of budding yeast
chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion.
Current biology : CB *6*, 1599-1608.

Tasan, I., Sustackova, G., Zhang, L., Kim, J., Sivaguru, M., HamediRad, M., Wang, Y., Genova, J., Ma, J.,
Belmont, A.S., and Zhao, H. (2017). CRISPR/Cas9-mediated Knock-in of an Optimized TetO Repeat for
Live Cell Imaging of Endogenous Loci. bioRxiv.

Taylor, G.C., Eskeland, R., Hekimoglu-Balkan, B., Pradeepa, M.M., and Bickmore, W.A. (2013). H4K16
acetylation marks active genes and enhancers of embryonic stem cells, but does not alter chromatin
compaction. Genome research *23*, 2053-2065.

Tjong, H., Gong, K., Chen, L., and Alber, F. (2012). Physical tethering and volume exclusion determine
higher-order genome organization in budding yeast. Genome research *22*, 1295-1305.

Toth, K.F., Knoch, T.A., Wachsmuth, M., Frank-Stohr, M., Stohr, M., Bacher, C.P., Muller, G., and
Rippe, K. (2004). Trichostatin A-induced histone acetylation causes decondensation of interphase
chromatin. Journal of cell science *117*, 4277-4287.

Traven, A., Jelicic, B., and Sopta, M. (2006). Yeast Gal4: a transcriptional paradigm revisited. EMBO
reports *7*, 496-499.

Tsompana, M., and Buck, M.J. (2014). Chromatin accessibility: a window into the genome.
Epigenetics & chromatin *7*, 33.

Tumbar, T., Sudlow, G., and Belmont, A.S. (1999). Large-scale chromatin unfolding and remodeling
induced by VP16 acidic activation domain. The Journal of cell biology *145*, 1341-1354.

Vallotton, P., van Oijen, A.M., Whitchurch, C.B., Gelfand, V., Yeo, L., Tsiavaliaris, G., Heinrich, S.,
Dultz, E., Weis, K., and Grunwald, D. (2017). Diatrack particle tracking software: review of
applications and performance evaluation. Traffic.

van den Engh, G., Sachs, R., and Trask, B.J. (1992). Estimating genomic distance from DNA sequence
location in cell nuclei by a random walk model. Science *257*, 1410-1412.

Venkatesh, S., and Workman, J.L. (2015). Histone exchange, chromatin structure and the regulation
of transcription. Nature reviews. Molecular cell biology *16*, 178-189.

Verschure, P.J., van der Kraan, I., de Leeuw, W., van der Vlag, J., Carpenter, A.E., Belmont, A.S., and van Driel, R. (2005). In vivo HP1 targeting causes large-scale chromatin condensation and enhanced histone lysine methylation. Molecular and cellular biology *25*, 4552-4564.

Vogelauer, M., Wu, J., Suka, N., and Grunstein, M. (2000). Global histone acetylation and deacetylation in yeast. Nature *408*, 495-498.

Wang, A., Kurdistani, S.K., and Grunstein, M. (2002). Requirement of Hos2 histone deacetylase for gene activity in yeast. Science *298*, 1412-1414.

Williamson, I., Berlivet, S., Eskeland, R., Boyle, S., Illingworth, R.S., Paquette, D., Dostie, J., and Bickmore, W.A. (2014). Spatial genome organization: contrasting views from chromosome conformation capture and fluorescence in situ hybridization. Genes & development *28*, 2778-2791.

Winzeler, E.A., Shoemaker, D.D., Astromoff, A., Liang, H., Anderson, K., Andre, B., Bangham, R., Benito, R., Boeke, J.D., Bussey, H., Chu, A.M., Connelly, C., Davis, K., Dietrich, F., Dow, S.W., El Bakkoury, M., Foury, F., Friend, S.H., Gentalen, E., Giaever, G., Hegemann, J.H., Jones, T., Laub, M., Liao, H., Liebundguth, N., Lockhart, D.J., Lucau-Danila, A., Lussier, M., M'Rabet, N., Menard, P., Mittmann, M., Pai, C., Rebischung, C., Revuelta, J.L., Riles, L., Roberts, C.J., Ross-MacDonald, P., Scherens, B., Snyder, M., Sookhai-Mahadeo, S., Storms, R.K., Veronneau, S., Voet, M., Volckaert, G., Ward, T.R., Wysocki, R., Yen, G.S., Yu, K., Zimmermann, K., Philippsen, P., Johnston, M., and Davis, R.W. (1999). Functional characterization of the S. cerevisiae genome by gene deletion and parallel analysis. Science *285*, 901-906.

Xin, H., Takahata, S., Blanksma, M., McCullough, L., Stillman, D.J., and Formosa, T. (2009). yFACT induces global accessibility of nucleosomal DNA without H2A-H2B displacement. Molecular cell *35*, 365-376.

Figure legends

Figure 1: Measuring transcription-dependent chromatin decompaction *in vivo*. (A) Chromosome
labeling scheme with LacO and TetO repeats on either side of the *GAL7-10-1* locus and example
image of cells expressing the reporter constructs. (B) Distribution of distances between LacO and
TetO repeats in cell populations constitutively grown in media with glucose, galactose or raffinose as
a carbon source. Boxes indicate 25 and 75 percentiles, blue line indicates the median of the
distribution. Outliers are shown as grey crosses. (C) Histogram of distance distribution for TetR and
LacI dots for 31 kb reporter strain in constitutive growth in glucose (black) or galactose (red). (D)
Mean distance of cell population during addition of galactose to cells grown in raffinose (or
continuous growth in either galactose or raffinose). Shaded areas show the standard error (SE) of
four independent experiments. (E) Mean distance of cell population during addition of glucose to
cells grown in galactose (or continuous growth in either galactose or raffinose). Shaded areas show
the SE of four independent experiments. (F) Mean distance of cell populations (number of cells per
time point approx. 200) during medium switch from glucose to galactose. Cells were pre-induced by
overnight growth in galactose and then grown in glucose for one hour. Black curve shows
fluorescence of Dextran-Alexa Fluor 680 (3000 MW, anionic) present in the glucose medium to
monitor the medium switch in the microfluidics setup.

Figure 2: Opening correlates with transcription on single cell level. (A) Individual slices of microscope
images showing single molecule FISH signal (bottom) in cells expressing genomic markers (green and
red) and stained with DAPI (blue) grown in raffinose (0 min) or after different times of induction with
galactose. The cells were hybridized with probes against *GAL1*, *GAL7* and *GAL10* mRNAs
simultaneously, all labeled with the far-red dye Qasar670. In the image of the 10 min time point, two
transcription spots in cells with little cytoplasmic mRNA signal are indicated by white arrowheads. A
cell with many cytoplasmic mRNA foci is indicated with a red arrow. (B) Quantification of percentage
of cells in which a bright transcription spot could be detected (Mean of three biological replicates,
error bars indicate standard deviation). (C) Distribution of distances across the GAL locus for all cells
(dark grey) and cells without (light grey) or with (green) transcription spot. Stars indicate that median
of the distances in cells classified as carrying a transcription spot is outside the 95 % confidence
interval of the median of the distances in cells classified as not carrying a transcription spot and vice
versa. n: number of cells analyzed. Histograms below show comparison of distance distributions at
10 min and 30 min after induction. Shown is the data of one representative experiment of three
biological replicates. (D) Means of median distances in cells classified for absence or presence of a
transcription spot from three biological replicates (Error bars represent standard deviation).

Figure 3: The role of acetylation in chromatin opening. (A) Boxplot showing the distance distribution
in wildtype cells or the indicated mutants. Cells were grown in raffinose to exponential phase and
then glucose, galactose or raffinose were added to a final concentration of 2 % (raffinose 2 %+ 2 %)
and cells imaged one hour after sugar treatment. Blue lines represent medians. Dashed lines show
median distance in the wildtype in glucose (grey) and galactose (green) with the shaded area
representing the 95 % confidence interval as determined by bootstrapping. Outliers are indicated by
grey crosses. The boxes extend to the 25th and 75th quartile. Data shown is of one representative
experiment of at least three biological replicates. (B) Boxplot showing the distance distribution in
cells treated with TSA or DMSO as solvent control for 16 hours. (C) Opening and closing kinetics after
treatment with TSA or DMSO for 16 hours. Mean of three biological replicates is shown, error bars
represent standard error of the mean.

Figure 4: Decompaction correlates with transcriptional activation. (A) 3D distance distributions in the
indicated mutants measured one hour after addition of the indicated sugars. Blue lines indicate
median of the distribution, grey crosses represent outliers. Dashed lines show median distance in the

wildtype in glucose (grey) and galactose (green) with the shaded area representing the 95 %
confidence interval as determined by bootstrapping. Each distribution is from one representative
experiment of at least three biological replicates and encompasses > 200 cells. (B) Fold expression of
GAL10 mRNA in different mutants relative to the wildtype determined by quantitative PCR
(normalized for *ACT1* mRNA as endogenous control). Means of three or more biological replicates are
shown. Error bars represent standard deviations. (C) Kinetics of chromatin opening and closing in
wildtype and mutants cells. Cells were grown overnight in medium with 2 % raffinose or 2 %
galactose. Galactose or glucose respectively were added or not added to 2 % final concentration at
time point 0. Shown are the means of three or more biological replicates with the standard error
represented by the shaded areas. (D) smFISH for *GAL1*, *GAL7* and *GAL10* in the indicated strains
growing at steady state in galactose. Insets show enhanced contrast for indicated cells. Images are
maximum intensity projections of z stacks encompassing the entire cells. (E) Boxplot showing the
intensities of transcriptions spots in the different strains (distribution from one representative
experiment of at least three biological replicates).

Figure 5: Nucleosome chaperones are crucial for chromatin opening. (A) Boxplots showing the
distance distribution in wildtype and *asf1Δ* cells grown in raffinose and induced with the indicated
carbon sources for one hour. (B) Boxplots showing the distance distribution in wildtype cells and cells
carrying temperature sensitive mutations of the FACT components Pob3 or Spt16. Cells were grown
in raffinose at 25 °C, shifted to 37 °C for 1 hour and then induced with the indicated carbon sources
for one hour at 37 °C. On all boxplots, blue lines indicate the median of the distribution, grey crosses
represent outliers. Dashed lines show median distance in the wildtype in glucose (grey) and galactose
(green) with the shaded area representing the 95 % confidence interval as determined by
bootstrapping. Each distribution is from a representative experiment of at least three biological
replicates and encompasses > 90 cells.

Figure 6: Dependence of 3D distance on ORF length and linear distance on chromosome. (A) Distance
distributions in cells harboring the indicated inducible reporter constructs (pro: GAL1-10 promoter
followed directly by the *ADH1* terminator). Stars indicate that median of distribution in galactose is
outside of the 95 % CI interval of the median in raffinose and vice versa (95 % CI determined by
bootstrapping). Representative experiment is shown. (B) Experimentally determined mean 3D
distance in various strains carrying the chromatin location markers TetO and LacO at different
distances from each other on chromosome II. Each datapoint is the mean of three or more medians
from different biological replicates. Grey and black datapoints are plotted from data published in
Dultz et al. 2016 Figure 4B. (C) Average 3D distance vs. genomic distance curves obtained from
simulated structures at two different values of compaction for the GAL locus. The markers
correspond to the experimental averages in the active and inactive conditions. In order to reproduce
the experimentally observed 3D distances, the compaction of GAL locus was set to 200 and 375
bp/bead respectively in the active and inactive conditions. (D) Two randomly selected snapshots of
the simulated GAL locus in the inactive (Gluc, left panel) and active (Gal, right panel) conditions.

Supplemental Figure 1: Transcription activation mutants also affect opening of the *GAL* locus. 3D
distance distribution of indicated strains. Strains were grown to exponential phase in raffinose
containing medium. The indicated sugars were added to a concentration of 2 % final and
measurements were taken after 1h of incubation.

Supplemental Figure 2: Kinetics of opening in *med7-163* temperature sensitive mutant at restrictive
temperatures. Cells were grown overnight in medium with 2 % raffinose or 2 % galactose at 25 °C
and shifted to 37 °C one hour before timepoint 0. Galactose was added or not added to 2 % final

concentration at timepoint 0. Shown are the means of three biological replicates with the standard
error represented by the shaded areas.

Supplemental Figure 3: Decompaction mediated by VP16 depends on SAGA's HAT activity. (A)
Schematic representation of the VP16 induction system. The Gal4 DNA binding domain (Gal4DBD) is
fused to the activation domain of VP16 linked by the hormone binding domain of human estrogen
receptor. Upon addition of β-estradiol to the cells, the fusion protein is released from its interaction
with a heat shock protein and can activate transcription from promoters containing a Gal4
recognition motif. (B-E) Decompaction kinetics in response to activation via galactose or β-estradiol.
Cells were grown to exponential phase in medium with 2 % raffinose. Induction was carried out on
the microscope stage at timepoint 0 with galactose (2 % final concentration) or β-estradiol (1 mM
final concentration). Shown are the means of three biological replicates with the standard error
represented by the error bars. (B) Wildtype. (C) Wildtype expressing β-estradiol responsive VP16
activator. (D) *gcn5Δ* expressing β-estradiol responsive VP16 activator. (E) *spt20Δ* expressing β-
estradiol responsive VP16 activator.

Supplemental Figure 4: Transcription correlates with opening. (A) Kinetics of opening and closing in
rpb1-1 temperature sensitive mutant at permissive and restrictive temperatures. Cells were grown
overnight in medium with 2 % raffinose or 2 % galactose at 25 °C and shifted to 37 °C were indicated
one hour before timepoint 0. Galactose or glucose respectively were added or not added to 2 % final
concentration at timepoint 0. Shown are the means of three biological replicates with the standard
error represented by the shaded areas. (B) Fold expression of *GAL10* mRNA relative to growth at 25
°C in raffinose (normalized for *ACT1* mRNA as endogenous control). Cells were grown in 2 % raffinose
into exponential phase. Cells were then shifted to the indicated temperature for one hour and
subsequently induced with galactose (+gal) or not for one more hour. Means of three biological
replicates are shown. Error bars represent standard deviations. Inset shows blow-up for uninduced
(raffinose) growth.

Supplemental Figure 5: RNA dot persists after glucose repression and correlates with more open
chromatin. (A) Maximum projections of smFISH for *GAL1*, *GAL7* and *GAL10* mRNAs detected by oligos
labelled with Qasar670 in cells grown to exponential phase in galactose and then repressed by the
addition of glucose to a final concentration of 2%. Images from all conditions are scaled the same.
Bottom panels show same images with higher contrast. Images were filtered with a Gaussian filter
for presentation purposes. Scale bar represents 2 um. (B) Boxplot showing the distance distribution
in cells grown either in galactose or raffinose continuously or repressed by addition of glucose to
galactose grown cells. Distributions of cells with or without transcription spot are analyzed separately
10 and 60 min after glucose repression. Data from one representative experiment of at least three
biological replicates is shown. (C) Mean % of cells with nuclear RNA dot from three biological
replicates.

Figure 1

Figure 1: Measuring transcription-dependent chromatin decompaction in vivo. (A) Chromosome labeling scheme with LacO and TetO repeats on either side of the GAL7-10-1 locus and example image of cells expressing the reporter constructs. (B) Distribution of distances between LacO and TetO repeats in cell populations constitutively grown in media with glucose, galactose or raffinose as a carbon source. Boxes indicate 25 and 75 percentiles, blue line indicates the median of the distribution. Outliers are shown as grey crosses. (C) Histogram of distance distribution for TetR and LacI dots for 31 kb reporter strain in constitutive growth in glucose (black) or galactose (red). (D) Mean distance of cell population during addition of galactose to cells grown in raffinose (or continuous growth in either galactose or raffinose). Shaded areas show the standard error (SE) of four independent experiments. (E) Mean distance of cell population during addition of glucose to cells grown in galactose (or continuous growth in either galactose or raffinose). Shaded areas show the SE of four independent experiments. (F) Mean distance of cell populations (number of cells per time point approx. 200) during medium switch from glucose to galactose. Cells were pre-induced by overnight growth in galactose and then grown in glucose for one hour. Black curve shows fluorescence of Dextran-Alexa Fluor 680 (3000 MW, anionic) present in the glucose medium to monitor the medium switch in the microfluidics setup.

Figure 2

Figure 2: Opening correlates with transcription on single cell level. (A) Individual slices of microscope images showing single molecule FISH signal (bottom) in cells expressing genomic markers (green and red) and stained with DAPI (blue) grown in raffinose (0 min) or after different times of induction with galactose. The cells were hybridized with probes against GAL1, GAL7 and GAL10 mRNAs simultaneously, all labeled with the far-red dye Qasar670. In the image of the 10 min time point, two transcription spots in cells with little cytoplasmic mRNA signal are indicated by white arrowheads. A cell with many cytoplasmic mRNA foci is indicated with a red arrow. (B) Quantification of percentage of cells in which a bright transcription spot could be detected (Mean of three biological replicates, error bars indicate standard deviation). (C) Distribution of distances across the GAL locus for all cells (dark grey) and cells without (light grey) or with (green) transcription spot. Stars indicate that median of the distances in cells classified as carrying a transcription spot is outside the 95 % confidence interval of the median of the distances in cells classified as not carrying a transcription spot and vice versa. n: number of cells analyzed. Histograms below show comparison of distance distributions at 10 min and 30 min after induction. Shown is the data of one representative experiment of three biological replicates. (D) Means of median distances in cells classified for absence or presence of a transcription spot from three biological replicates (Error bars represent standard deviation).

Figure 3

Figure 3: The role of acetylation in chromatin opening. (A) Boxplot showing the distance distribution in wildtype cells or the indicated mutants. Cells were grown in raffinose to exponential phase and then glucose, galactose or raffinose were added to a final concentration of 2 % (raffinose 2 %+ 2 %) and cells imaged one hour after sugar treatment. Blue lines represent medians. Dashed lines show median distance in the wildtype in glucose (grey) and galactose (green) with the shaded area representing the 95 % confidence interval as determined by bootstrapping. Outliers are indicated by grey crosses. The boxes extend to the 25th and 75th quartile. Data shown is of one representative experiment of at least three biological replicates. (B) Boxplot showing the distance distribution in cells treated with TSA or DMSO as solvent control for 16 hours. (C) Opening and closing kinetics after treatment with TSA or DMSO for 16 hours. Mean of three biological replicates is shown, error bars represent standard error of the mean.

Figure 4

Figure 4: Decompaction correlates with transcriptional activation. (A) 3D distance distributions in the indicated mutants measured one hour after addition of the indicated sugars. Blue lines indicate median of the distribution, grey crosses represent outliers. Dashed lines show median distance in the wildtype in glucose (grey) and galactose (green) with the shaded area representing the 95 % confidence interval as determined by bootstrapping. Each distribution is from one representative experiment of at least three biological replicates and encompasses > 200 cells. (B) Fold expression of GAL10 mRNA in different mutants relative to the wildtype determined by quantitative PCR (normalized for ACT1 mRNA as endogenous control). Means of three or more biological replicates are shown. Error bars represent standard deviations. (C) Kinetics of chromatin opening and closing in wildtype and mutants cells. Cells were grown overnight in medium with 2 % raffinose or 2 % galactose. Galactose or glucose respectively were added or not added to 2 % final concentration at time point 0. Shown are the means of three or more biological replicates with the standard error represented by the shaded areas. (D) smFISH for GAL1, GAL7 and GAL10 in the indicated strains growing at steady state in galactose. Insets show enhanced contrast for indicated cells. Images are maximum intensity projections of z stacks encompassing the entire cells. (E) Boxplot showing the intensities of transcriptions spots in the different strains (distribution from one representative experiment of at least three biological replicates).

Figure 5

Figure 5: Nucleosome chaperones are crucial for chromatin opening. (A) Boxplots showing the distance distribution in wildtype and asf1Δ cells grown in raffinose and induced with the indicated carbon sources for one hour. (B) Boxplots showing the distance distribution in wildtype cells and cells carrying temperature sensitive mutations of the FACT components Pob3 or Spt16. Cells were grown in raffinose at 25 °C, shifted to 37 °C for 1 hour and then induced with the indicated carbon sources for one hour at 37 °C. On all boxplots, blue lines indicate the median of the distribution, grey crosses represent outliers. Dashed lines show median distance in the wildtype in glucose (grey) and galactose (green) with the shaded area representing the 95 % confidence interval as determined by bootstrapping. Each distribution is from a representative experiment of at least three biological replicates and encompasses > 90 cells.

Figure 6

Figure 6: Dependence of 3D distance on ORF length and linear distance on chromosome. (A) Distance distributions in cells harboring the indicated inducible reporter constructs (pro: GAL1-10 promoter followed directly by the ADH1 terminator). Stars indicate that median of distribution in galactose is outside of the 95 % CI interval of the median in raffinose and vice versa (95 % CI determined by bootstrapping). Representative experiment is shown. (B) Experimentally determined mean 3D distance in various strains carrying the chromatin location markers TetO and LacO at different distances from each other on chromosome II. Each datapoint is the mean of three or more medians from different biological replicates. Grey and black datapoints are plotted from data published in Dultz et al. 2016 Figure 4B. (C) Average 3D distance vs. genomic distance curves obtained from simulated structures at two different values of compaction for the GAL locus. The markers correspond to the experimental averages in the active and inactive conditions. In order to reproduce the experimentally observed 3D distances, the compaction of GAL locus was set to 200 and 375 bp/bead respectively in the active and inactive conditions. (D) Two randomly selected snapshots of the simulated GAL locus in the inactive (Gluc, left panel) and active (Gal, right panel) conditions.

Supplemental Figure 1

Supplemental Figure 1: Transcription activation
mutants also affect opening of the GAL locus. 3D
distance distribution of indicated strains. Strains
were grown to exponential phase in raffinose
containing medium. The indicated sugars were
added to a concentration of 2 % final and
measurements were taken after 1h of incubation.

Supplemental Figure 2

Supplemental Figure 2: Kinetics of opening in med7-163 temperature sensitive mutant at restrictive temperatures. Cells were grown overnight in medium with 2 % raffinose or 2 % galactose at 25 °C and shifted to 37 °C one hour before timepoint 0. Galactose was added or not added to 2 % final concentration at timepoint 0. Shown are the means of three biological replicates with the standard error represented by the shaded areas.

Supplementary Figure 3

Supplemental Figure 3: Decompaction mediated by VP16 depends on SAGA's HAT activity. (A) Schematic representation of the VP16 induction system. The Gal4 DNA binding domain (Gal4DBD) is fused to the activation domain of VP16 linked by the hormone binding domain of human estrogen receptor. Upon addition of β-estradiol to the cells, the fusion protein is released from its interaction with a heat shock protein and can activate transcription from promoters containing a Gal4 recognition motif. (B-E) Decompaction kinetics in response to activation via galactose or β-estradiol. Cells were grown to exponential phase in medium with 2 % raffinose. Induction was carried out on the microscope stage at timepoint 0 with galactose (2 % final concentration) or β-estradiol (1 mM final concentration). Shown are the means of three biological replicates with the standard error represented by the error bars. (B) Wildtype. (C) Wildtype expressing β-estradiol responsive VP16 activator. (D) gcn5Δ expressing β-estradiol responsive VP16 activator. (E) spt20Δ expressing β-estradiol responsive VP16 activator.

Supplemental Figure 4

Supplemental Figure 4: Transcription correlates with opening. (A) Kinetics of opening and closing in rpb1-1 temperature sensitive mutant at permissive and restrictive temperatures. Cells were grown overnight in medium with 2 % raffinose or 2 % galactose at 25 °C and shifted to 37 °C were indicated one hour before timepoint 0. Galactose or glucose respectively were added or not added to 2 % final concentration at timepoint 0. Shown are the means of three biological replicates with the standard error represented by the shaded areas. (B) Fold expression of GAL10 mRNA relative to growth at 25 °C in raffinose (normalized for ACT1 mRNA as endogenous control). Cells were grown in 2 % raffinose into exponential phase. Cells were then shifted to the indicated temperature for one hour and subsequently induced with galactose (+gal) or not for one more hour. Means of three biological replicates are shown. Error bars represent standard deviations. Inset shows blow-up for uninduced (raffinose) growth.

Supplemental Figure 5: RNA dot persists after glucose repression and correlates with more open chromatin. (A) Maximum projections of smFISH for GAL1, GAL7 and GAL10 mRNAs detected by oligos labelled with Qasar670 in cells grown to exponential phase in galactose and then repressed by the addition of glucose to a final concentration of 2%. Images from all conditions are scaled the same. Bottom panels show same images with higher contrast. Images were filtered with a Gaussian filter for presentation purposes. Scale bar represents 2 um. (B) Boxplot showing the distance distribution in cells grown either in galactose or raffinose continuously or repressed by addition of glucose to galactose grown cells. Distributions of cells with or without transcription spot are analyzed separately 10 and 60 min after glucose repression. Data from one representative experiment of at least three biological replicates is shown. (C) Mean % of cells with nuclear RNA dot from three biological replicates.