A comparison of tests for quantifying sensory eye dominance

Manuela Bossi[a], Lisa Hamm[b,c], Annegret Dahlmann-Noor[a,d], Steven C. Dakin[a,b,c,‡]

5   [a] *UCL Institute of Ophthalmology, University College London, Bath St, London EC1V 9EL, UK*

[b] *School of Optometry and Vision Science and* [c] *New Zealand National Eye Centre, University of Auckland, Auckland 1142, New Zealand*

[d] *Department of Paediatric Ophthalmology, NIHR Biomedical Research Centre at Moorfields Eye Hospital, City Road, London EC1V 2PD, UK*

10   [‡]Corresponding author, email: s.dakin@auckland.ac.nz

Running head: *Measuring sensory eye dominance*

Abstract

Purpose Clinicians rely heavily on stereoacuity to measure binocular visual function, but stereo-vision represents only one aspect of binocularity. Lab-based tests of *sensory eye dominance* (SED)
15   are commonplace, but have not been translated to wider clinical practice. Here we compare several methods of quantifying SED in a format suitable for clinical use. **Methods** We tested 30 participants with ostensibly normal vision on 8 tests. Seven tests (#1-7) were designed to quantify SED in the form of an interocular *balance-point* (BP). In tests #1-6, we estimated a *contrast-BP*, the interocular difference in contrast required for observers to be equally likely to base their judgement on either
20   eye, whereas in test #7 we measured binocular rivalry (interocular ratio of sensory dominance duration). We compare test-retest reliability (intra-observer consistency) and test-validity (inter-observer discriminatory power) and compare BP to stereoacuity (test #8). **Results** The test that best preserved inter-observer differences in contrast balance while maintaining good test-retest reliability was a polarity judgement using superimposed opposite-contrast polarity same-identity
25   optotypes. **Conclusions** A reliable and valid measure of SED can be obtained rapidly (20 trials) using a simple contrast-polarity judgement. Tests that use polarity-rivalrous stimuli elicit more reliable judgments than those that do not.

Significance Statement

Although sensory eye dominance is central to understanding normal and disordered binocular
30   vision, there is currently no consensus as to the best way to measure it. Here we compare several candidate measures of sensory eye dominance and conclude that a reliable measure of SED can be achieved rapidly using a judgement of stimulus contrast-polarity.

Introduction

35 *Background*

When combining signals from the two eyes into a coherent 'cyclopean' percept, observers rely (to differing extents) on one eye more than the other. Such *eye dominance* is important in a number of clinical settings. First, imbalances between the eyes in childhood can lead to severe loss of acuity in the weaker eye, a condition called amblyopia (Birch 2013). Treating amblyopia (e.g. by patching) attempts to
40 "rebalance" vision, by improving acuity in the weaker eye. Thus the relative contribution of each eye to cyclopean vision a key measure for understanding amblyopia. This is particularly the case as there is debate as to whether suppression of the weaker eye is a cause (e.g. Wong 2012, Birch 2013, Hess and Thompson 2015) or a consequence (e.g. Vedamurthy, Nahum et al. 2015, Kehrein, Kohnen et al. 2016, Bossi, Tailor et al. 2017) of the condition. Second, presbyopia (an age-related refractive error) can be
45 corrected using monovision, which enhances intermediate-distance vision in the dominant eye only, and so necessitates a decision on eye dominance. Finally, loss of visual function in age-related macular-disease leads to subtle changes in patients' reliance on their two eyes ("binocular balance"). A simple method for quantifying changes in sensory eye dominance may therefore have diagnostic value for this and other conditions (Wiecek, Lashkari et al. 2015).

50 A variety of methods are available to measure the relative contribution of each eye in a clinical setting. Clinical measures of eye dominance tend to be intuitive, for example assessing with which eye a patient can more easily wink (Miles 1930). Building on the notion that eye dominance is linked to motor control, comparisons of eye-alignment are still routinely used. For instance, in the *Miles test* and *Porta test,* near and distant targets are aligned binocularly, and then each eye is opened and closed in succession to
55 reveal which eye gives the more accurate estimate of position. Such tests of 'sighting' dominance are binary (left or right) and tend to elicit variable results (Johansson, Seimyr et al. 2015), that are dependent on specific test and viewing conditions (Rice, Leske et al. 2008).

Sensory eye dominance (SED), on the other hand, refers to the relative *perceptual* contribution of each eye to the cyclopean percept under dichoptic conditions (Ooi and He 2001). In the clinic,
60 *Worth's four dot test* produces a qualitative estimate of SED. This task has the observer wear red-green anaglyph glasses and report the perceived number and colour of 4 illuminated circles (1 red, 2 green and 1 white). The only *quantitative* direct clinical test of SED is the *neutral-density filter bar* (a version using red filters is the *Sbisa bar*), that quantifies the minimum filter-density that must be placed over one eye to induce the use of the other eye (McCormick, Bhola et al. 2002). However,
65 such tests can be challenging for children which likely contributes to their only moderate test-retest reliability (Crawford and Griffiths 2015, Piano and Newsham 2015) and their infrequent use for paediatric screening (Tailor, Balduzzi et al. 2014).

Neither sighting-dominance nor sensory dominance assess the functional advantage of having two eyes. When each eye makes a relatively equal contribution, observers experience binocular summation
70 (increased contrast sensitivity when using both eyes; Baker, Meese et al. 2007, Baker, Meese et al. 2007, Baker, Meese et al. 2008) and stereopsis (the perception of three-dimensional structure based on retinal disparity; Cumming and DeAngelis 2001, O'Connor, Birch et al. 2010). Although there is no standard clinical test of summation, stereoacuity is the most widely accepted clinical measure of binocular vision. Stereoacuity tests quantify the minimum retinal disparity supporting reliable discrimination of surface-
75 depth (Julesz 1971). Popular variants include the *TNO*, the *Randot* test (Birch, Williams et al. 2008) and the *Frisby* test (Frisby, Davis et al. 1996). Stereoacuity tests are simple to administer and can easily be explained to children. However, they have drawbacks; some contain monocular cues (Fricke and Siderov 1997), they can be difficult to administer in the presence of strabismus (McKee, Levi et al. 2003) and they

are not necessarily indicative of clinical conditions (since 1-14% of the general population are stereo-blind; Bosten, Goodbourn et al. 2015) which limits their utility for vision screening (Cotter, Cyert et al. 2015). For example, despite being stereo-blind patients with strabismic amblyopia can demonstrate normal binocular summation when balancing the visibility of dichoptic stimuli (Baker, Meese et al. 2007).

Outside the clinic, where stereoacuity is the foundation of binocular assessment, vision researchers have developed diverse, innovative and quantitative measures of sensory eye dominance. Rather than directly asking observers to make a judgement about contrast similarity (as required for e.g. the Sbisa bar), psychometric assessment tends to rely on forced-choice methods (for a study that linked traditional clinical methods of SED to a common psychophysical measure see Li, Lam et al. 2010). Pairing robust psychophysical judgements with carefully-designed stimuli allows assessment of the relative contribution of each eye to visual processing in different stages of the cortical hierarchy.

Here we consider SED tests first according to whether they do or do not preclude a coherent cyclopean percept (i.e. are intrinsically *rivalrous*). One test which poses no cyclopean conflict presents observers with dichoptic sine-wave gratings that differ only in phase and contrast (Figure 1, test #4). Observers are asked to indicate the location of the middle dark stripe of the phase-shifted grating that results from binocular summation (Ding and Sperling 2006, Huang, Zhou et al. 2009, Huang, Zhou et al. 2010, Zhou, Thompson et al. 2013, Kwon, Lu et al. 2014). Reported position allows one to determine the interocular contrast difference that supports equal contribution from each eye, and it is this contrast difference that quantifies "binocular balance". Another test embeds contrast manipulation within a global processing task (Mansouri, Thompson et al. 2008, Li, Lam et al. 2010, Black, Thompson et al. 2011, Li, Hess et al. 2013, Li, Hess et al. 2013, Li, Thompson et al. 2013, Hamm, Chen et al. 2017). In motion coherence paradigms (Li, Lam et al. 2010), the observers are required to discriminate the direction of a pattern of moving dots, comprised of signal-dots (moving in one direction) and noise-dots (moving in random directions). The most commonly used version of this approach for quantifying SED requires two phases: (1) determining the proportion of signal to noise dots supporting reliable direction discrimination and (2) determining the contrast of the signal-dots required to maintain performance under dichoptic conditions (Black, Thompson et al. 2011). This paradigm has also been used with global orientation tasks (Zhou, Huang et al. 2013). In all such tests, observers perceive a coherent cyclopean percept and the relative contrast supporting optimum performance is termed the interocular 'balance point' and quantifies SED.

A second category of tests use rivalrous stimuli to quantify SED. When different monocular images fall on corresponding retinal locations of the two eyes this can lead either to an experience of *binocular rivalry* (an alternation of percept between the stimuli presented to the two eyes; Wheatstone 1838), or to *diplopia* (double vision). The nature and extent of conflict between the information from the two eyes is a measure of the degree of SED. Bossi *et al* (2017) quantified this using dichoptic stimuli where each eye was presented with opposite contrast-polarity versions of the same symbol (Figure 1, test #3), whereas Kwon et al presented observers with spatially overlapping rivalrous letter pairs of differing contrast (Figure 1, task 2; Kwon, Wiecek et al. 2015). In both cases, the objective was to measure the 'contrast balance point', or the contrast mixture required for observers' percept to be equally likely to be driven by either eye's view. This balance point (like the non-rivalrous tests) quantifies SED.

Although all tests described use centrally presented stimuli, SED has been measured at multiple locations in the visual field. Xu et al mapped SED over 17 locations (at the fovea, 2 and 4 deg. eccentricity) and report gradual variation across the field (Xu, He et al. 2011). Hess and colleagues have measured contrast balance in patients with strabismus and report that patients exhibited stronger suppression across the field than controls (Babu, Clavagnier et al. 2017). Development of this, and many of the tests described, has been driven by research exploring the role of interocular suppression in amblyopia (Baker, Meese et al. 2008, Mansouri, Thompson et al. 2008, Huang, Zhou

125   et al. 2009). Several studies have examined the use of training (or *perceptual learning*) to shift balance towards the amblyopic eye as a treatment for amblyopia (Ooi, Su et al. 2013, Birch, Li et al. 2015). Although the benefits of such training for acuity in the amblyopic eye are clear, it is less clear if these benefits result from a shift in SED (Vedamurthy, Nahum et al. 2015, Bossi, Tailor et al. 2017) as has been reported(Black, Hess et al. 2012). Although the functional significance of changes in SED

130   is debatable, it is the case that such changes can arise from short term manipulation of binocular sensory experience (e.g. through patching) in both amblyopes (Zhou, Thompson et al. 2013) and controls (Zhou, Clavagnier et al. 2013, Zhou, Reynaud et al. 2014).

There have been several efforts to translate lab-based measure of SED to the clinic. Typically, research groups have adapted their own method for clinical use - making tests more

135   convenient/shorter/simpler (e.g. Black, Thompson et al. 2011, Kwon, Lu et al. 2014). Here we compared a variety of dichoptic stimuli using a common psychophysical protocol. We assessed resulting SED scores based on 1) *Reliability* (consistency between two measurements made on the same participant), and 2) *Validity* (the ability of the test to capture individual differences).

We selected (or created) tests suitable for a single, rapid psychophysical protocol (effectively

140   precluding 2-stage paradigms and tasks involving multiple stimulus locations). We ensured tasks probed a range of visual processes; from low-level (e.g. phase; Ding and Sperling 2006), to high-level processing of spatial form (e.g. letter identity; Kwon, Wiecek et al. 2015), as well as motion perception. We also set out to quantify the impact of rivalry on our measures, focusing on stimuli involving no-rivalry (Ding and Sperling 2006), contrast-polarity rivalry (Bossi, Tailor et al. 2017) and

145   spatial-form rivalry (Kwon, Wiecek et al. 2015).

Methods

Participants

We recruited thirty adult participants (18 female; 22-55yrs old) through the University of Auckland-Optometry clinic via email and poster advertisements. Based on pre-test screening, none had

150   significant disorders of vision and all wore optical correction as necessary. Three of the observers are authors; others were not informed as to the purpose of this study. Experimental protocols complied with the Declaration of Helsinki and were approved by the University of Auckland Human Research Ethics Committee.

Apparatus

155   Stimuli were presented on a linearised LG 21" LCD 3D monitor, with a 1920x1080 pixel resolution operating at 120Hz. Stimuli were generated in Matlab (Mathworks Ltd) using the Psychophysics Toolbox (Brainard, 1997; Pelli, 1997; Kleiner et al, 2007). During testing, stimuli were viewed through wireless LCD shutter glasses (nVidia Corp., Santa Clara, CA) allowing independent control of the image presented to each eye (effective framerate was 60Hz per eye). Shutter glasses were worn

160   over optical correction when necessary. A Minolta LS110 photometer (Konica-Minolta Ltd) was used to calibrate the monitor-luminance, using measurements made through the shutter glasses. Participants viewed the screen at a viewing distance of 100 cm to produce a pixel density of 55.9 pixel/deg.

 Stimuli

Red-green anaglyph versions of the stimuli are depicted in Figure 1, and described in Table 1. Stimuli appeared against a mid-grey background, and were surrounded by a "vergence-lock" frame (visible to both eyes) made of black and white alternating bars; this promoted fusion.

All tests involved presenting a pair of images independently to each eye. **Test #1-2.** Stimuli
170 comprised a pair of different letters, selected from 10 ETDRS Sloan letters (Ferris, Kassoff et al. 1982). The width and height of letters was 2.75° and the letters were the same (positive) contrast polarity. In test #1 letters were spatially separated (minimising rivalry) whereas in test #2 they were superimposed, promoting rivalry between the two different letters. **Test #3** Letter font, size and separation were identical to test #1. However, each of the two letters was comprised of
175 superimposed pair of opposite contrast polarity letters (with the same identity) presented separately to the two eyes. In one stimulus-letter the light component went to the left eye, the dark component to the right, and in the other stimulus-letter, vice-versa. This led to strong rivalry, driven by the conflicting contrast-polarity of the components of each letter. **Tests #4-5.** Stimuli were similar to Ding & Sperling (2006) and were comprised of two horizontal sine-wave gratings with the same
180 spatial frequency (SF; 0.52cpd), presented within a 2.94° square window. The two gratings were presented separately to each eye and had phases of ±45° (test #4, 90° phase difference, Figures 1-4) or ±90° (test #5, 180° phase difference, Figures 1-5), where a 0° phase grating would present a horizontal dark bar centrally positioned (on the y-axis) bisecting the square window. On a given trial the positive-phase grating was assigned randomly to one eye, the negative-phase grating to the
185 other. In the 90° condition (Test 4) addition of gratings led to a phase shifted cyclopean percept without rivalry, whereas in the 180° condition gratings resulted in rivalry (driven by a polarity conflict similar to Test 3). **Tests #6-7.** Stimuli were comprised of two 13.75° square filtered-noise patterns drifting in opposite directions (upwards or downwards). Stimuli were generated by filtering Gaussian random noise with a log Gabor filter which passed all orientations but a range of SFs (filter had a log-
190 Gaussian profile: mean SF of 6cpd, bandwidth 0.5 octaves). **Test #8.** Stimuli were stereograms - stereo-defined Sloan letters (5.5° square) - defined by interocular horizontal shifts of the pixels within carrier images (435 X 435 black-white pixel arrays which each patch subtending 7.8°). Stimuli were shifted with sub-pixel accuracy using bilinear interpolation.

195   **Figure 1.** (Grey panels) Examples of stimuli used in the eight tests (for the purpose of demonstration, stimuli are presented in red-green anaglyph versions). (Inset, lower right) Relative sizes of stimuli.

Test #	Test name	Symbol	Cue	Rivalry? (type)	Question	Measure	Primary reference
1	*Letter-strength (non-overlapping)*		*form*	✘	*Which letter appears stronger?*	Contrast-BP	Kwon, Wiecek et al. (2015)
2	*Letter-strength (overlapping)*		*form*	✔ *(form)*	*Which letter appears stronger?*	Contrast-BP	Kwon, Wiecek et al. (2015)
3	*Letter-polarity*		*form*	✔ *(polarity)*	*Which letter appears whiter?*	Contrast-BP	Bossi, Tailor et al. (2017)
4	*Phase combination*		*position*	✘	*Does middle dark bar fall above or below horizontal ref. line?*	Contrast-BP	Ding and Sperling (2006)
5	*Phase suppression*		*position*	✔ *(polarity)*	*Does middle dark bar fall above or below horizontal ref. line?*	Contrast-BP	Ding and Sperling (2006)
6	*Motion strength (contrast)*		*motion*	✔ *(motion)*	*Does the pattern move up or down?*	Contrast-BP	Black, Thompson et al. (2011)
7	*Motion rivalry (duration)*		*motion*	✔ *(motion)*	*Hold ↑ key when pattern moves ↑ & ↓ key when it moves ↓*	Time-ratio BP	Dieter, Sy et al. (2016)
8	*Stereoacuity*		*form*	✘	*Identify the letter you see*	Threshold disparity	Frisby, Davis et al. (1996)

Table 1. Summary of eight tests. BP= Balance Point (either contrast-BP in tests #1-6 or time-ratio BP in test #7).

 Procedure

We ran participants through eight tests, twice (to allow us to estimate test-retest reliability). Tests were always administered in the same order (#1-8). In all tests except #7 (motion rivalry) stimuli appeared for a maximum of 4.5 s, after which the display switched to showing the response-choices for a maximum of 10s. Observers could respond before the display of response options, which
205   triggered presentation of the next trial. Participants signalled their response verbally, which the experimenter recorded using the computer keyboard. Note that no feedback was provided in any test (since we are estimating bias in all but test #8). Each test took approximately 90s to administer. The experimenter instructed the participant before each test as highlighted in Table 1. Total duration of the experiment - incorporating 8 tests x 2 repeats, instructions and breaks - was around
210   50 min. Tests #1-7 measured a Balance Point (BP) from 0 (participant used their left eye only), to 1 (right eye only). For tests #1-6 BP was based on contrast, and for task #7 it was based on a ratio of time spent experiencing the stimuli presented to each eye.

Tests #1-6 quantified the ratio between the contrast-level applied to each dichoptic-image that leads to either of the two images (called A, B below) being equally likely to be chosen. This contrast-BP
215   was determined in 20 trials using an adaptive staircase algorithm (QUEST; Watson and Pelli 1983) which, for a stimulus comprised of the mixture of images *A* and *B*, converged on a threshold (α) producing 50% identification of (tests #1,2,4-6) image *A* or (Test 3) the "whiter" stimulus. The values QUEST produced were clamped in the range 0.0-1.0 (where 0.0 is exclusive presentation of the cue to the left eye, 0.5 equal presentation to left and right eye and 1.0 exclusive presentation to the
220   right eye). QUEST's guess-rate (γ) was set to 0.5 (2 AFC), the lapse rate (λ) to 0.01 and the slope-estimate (β) to 3.5. The initial guess for threshold or contrast-BP was 0.5 (i.e. balance) with an associated standard deviation of 0.7. For tests #1,2,4,5 and 6, QUEST set the contrast (*C*) of the right eye component (C_{right}) and C_{left} was set to 1-C_{right}, for test #3 contrast manipulation is described in the next section. Note that QUEST values were determined from a Monte Carlo simulation on a
225   population of ideal observers whose simulated-balances spanned the range 0.05-0.95 and whose other psychometric characteristics (β, λ) were taken from Kwon, Wiecek et al. (2015). **Test #7** consisted of 1 minute of exposure to rivalrous motion. Either upward or downward motion was randomly assigned to the left and right eyes for the initial 30s test period and then direction was switched across eyes for the remaining 30s (to counter-balance any bias for a stimulus in a given
230   direction rather than from a given eye). During stimulus presentation, observers pressed and held either the U ("up") or D ("down") button on the computer keyboard to indicate their dominant percept. In case of uncertainty, e.g. due to a "patchy" percept, the participant was instructed to report the more dominant direction. **Test #8** estimated a stereo-threshold (not bias) for letter-identification, using a 20-trial QUEST staircase. QUEST's guess-rate (γ) was set to 0.1 (10 AFC), the
235   lapse rate (λ) to 0.01 and the slope (β) to 3.5. The initial guess for threshold was set to 3'20".

Details of contrast manipulation for test #3 For test #3, the QUEST value set the luminance of the components of the opposite contrast polarity left/right-eye letters forming each of the two stimulus letters. First, the highest increment was randomly assigned to the stimulus-letter *A* or *B*. Then, the luminance of the light component-letter in one stimulus-letter and the dark component-letter in the
240   other stimulus-letter were set to be equal increments and decrements (range ±50cd/m2; example in Figure 2 of Bossi, Tailor et al. 2017). Thus, for a background luminance of 50 cd/m^2 if QUEST produced a value of 0.8 then the light component letter in, for example, stimulus letter A was 50+0.8*50=90 cd/m^2 and so the dark component letter in stimulus letter B was 50-0.8*50=10 cd/m^2. The dark component letter in stimulus letter A was 50-(1.0-0.8)*50=40 cd/m^2 and the light
245   component letter in the stimulus letter B was 50+(1.0-0.8)*50=60 cd/m^2. In the extreme case of

QUEST producing the limit value of 0.0, then the light component letter in, for example, stimulus letter A was 50+0.0*50=50 cd/m^2 and so the dark component letter in stimulus letter B was also 50-0.0*50=50 cd/m^2. The dark component letter in stimulus letter A was 50-1.0*50=0 cd/m^2 and the light component letter in the stimulus letter B was 50+1.0 *50=100cd/m^2. Conversely, when QUEST converged to 1.0, the dark component of A and the light component of B were matched (at 50 cd/m^2) while A had a light component at 100 cd/m^2 and B a dark component at 0 cd/m^2.

Analysis

For tests #1-6, BP was derived by fitting the 20 trials of a single QUEST-controlled run with a cumulative normal function using the Palamedes toolbox (Prins and Kingdom 2009). In order to quantify confidence in our fit parameters we pooled stimulus-levels (and their associated responses) into four bins, assumed binomially distributed errors at each level and bootstrapped fits to these binned data. Although this is not an exact estimate of error (since we cannot bootstrap the continuous data used to derive the contrast-BP) it is nonetheless a useful indicator of confidence in these estimates.

For test #7 we quantified BP as the proportion of left-eye dominance, using the proportion of frames (within a given trial) when the participants' response was consistent with their relying on the left-eye-view during exposure to rivalrous stimuli. We excluded responses occurring within 4.2 s (i.e. 500 frames) of the beginning, and the middle (when directions switched between eyes) of the 60s sequence. Responses composed of simultaneous keypresses were excluded from analysis.

We first describe BP for each individual and each test by averaging run 1 and 2 together, and also calculate the mean BP for each participant across all tests. To estimate test-reliability we calculated Bland Altman 95% limits of agreement, or Coefficient of Repeatability (CoR - lower values indicate higher repeatability; Vaz, Falkmer et al. 2013) as well as the Intraclass Correlation Coefficient (ICC – higher values indicate higher repeatability), based on a mean rating (k=2), absolute agreement, 1-way random-effects model (McGraw and Wong 1996). We examine how CoR and ICCs are influenced by the number of trials run (by re-calculating BP with Palamedes refits for fewer trials), and investigate the reliability of tests by comparing measures to one another. We note that highly reliable outcomes can arise from a measure which has low test-validity, being insensitive to differences in SED between individual (in other words, a test which always elicited the same outcome would be highly reliable, but not very useful). The absence of a gold standard measure of SED means we cannot directly assess which test is the most accurate by comparing it to such a standard. We therefore relied on Mean Average Precision (MAP) and Fractional Rank Precision (FRP) to incorporate both reliability *and* validity (Dorr, Elze et al. 2017). Finally we use regression to compare SED data to stereoacuity. FRP employs an information retrieval approach and evaluates a test by quantifying how identifiable a participant is from their set of test-scores.

Results

Tests #1-7 estimated observers Balance-Point (BP): #1-6 estimated observers' contrast Balance-Point, #7 quantified rivalry; test #8 measured stereoacuity. We assessed tests according to their 1) *reliability* (consistency between two measurements of the same test made on the same participant), and 2) *validity* (the ability of the test to capture individual differences amongst participants). Reliability was quantified using the Bland Altman Coefficient of Repeatability (CoR; the variance of the difference between runs 1 and run 2). The other metrics in table 2 provide statistical estimates of both reliability and validity.

Are observers more reliant on their left or right eye? Figure 2a plots the mean BPs (of the two runs) of tests #1-7 for all observers. Participants IDs are ordered along the abscissa by BP (averaged across the two runs). The overall mean-BP (for all participants; Green symbols), obtained by averaging mean contrast-BP, for tests #1-6 ,*and* mean proportion of instances when the percept was determined by the left eye (test #7) was 0.52 (σ=0.04). Mean contrast-BP (excluding test #7) was 0.52 (σ=0.05). Note the clustering of BP results around 0.5, indicating near-perfect binocular balance; as one would expect for observers with normal binocular vision. In line with previous studies, we report a prevalence of right eye-dominance of 21/30 (mean BP=0.54, σ=0.03). 9/30 participants were mainly left-eye dominant (mean BP=0.47; σ=0.02). Sixteen participants showed a marked eye-preference in all balance tests (only five had a left-eye preference). Tests #1, 3 4, 5 and 7 elicited BPs more similar to one another (and to the overall mean balance) compared to tests #2 and 6 (Figure 2a). Figure 2b plots a sample of data from three participants performing two runs of the seven balance tests. The dashed line denotes perfect binocular balance (BP=0.5). The green line indicates the mean estimated balance-level across the seven tests (with exact values provided in the green box above each plot). We found a generally high level of agreement across runs of the same test: the absolute difference was 0.08±0.09 between participants (averaging, for each participant, the difference across tests) and 0.05±0.07 between tests (averaging, for each test, the difference across participants). We observe consistency of balance estimates made using different tests, as indicated by the correlation between the mean balance on a certain test and the estimate across the other tests (tests #1-7: mean R= 0.564±0.003, *p*=0.102±0.004).

Figure 2. (a) Estimated balance point (BP) for all participants performing tests #1-7, ordered by mean BP across tests (green triangles; error bars are 95% confidence intervals). Here, BP=0.0 would indicate complete reliance on the left eye, 1.0 complete reliance on the right eye and 0.5 equal reliance on each eye (binocular balance). Tests are colour coded according to the legend. **(b)**

315 Illustrative results from three participants, chose from the left, central and right batches of IDs on the abscissa in 2a. Blue and red symbols show data from the first and second run respectively, and error bars are 95% confidence intervals on estimates. Mean BP across tasks (green line) is the green-boxed figure in each panel.

320 *Which is the "best" test?* We assessed 'best' in terms of (a) how reliable the tests are (across two runs) and (b) how well they capture individual variation in SED across our group. To visualise this, Figure 3 shows Bland Altman plots of data from the seven balance tests as well as test #8: stereoacuity (n.b. data are plot on different axes to other sub-plots). It is clear that some tests such as #6: motion strength, lead to a wide range of balance estimates across observers but also to high variability of balance-estimates

325 across runs (see also Figure 2). Conversely, tests such as letter-strength (Figure 3, task #1) elicit both a narrower range of balance estimates and much lower variability across runs. However, the high degree of repeatability of this test arises from it yielding a BP estimate of 0.5 for almost all participants, suggesting it is unable to differentiate subtle difference in SED (i.e. it has poor test-validity). We note that it is the closest test we have to the *Sbisa bar*, in that there is not spatially overlapping information

330 and participants are required to make a judgment of contrast.

Assuming that SED varies within the population tested, the best test maximises the range of measures across observers but minimises the range of measures across runs. A variety of statistics quantify this

and a selection are given in Table 2. Test #3 (letter-polarity) maximises intra class correlation (ICC), *F*, mean average precision (MAP), and fractional rank precision (FRP). The Coefficient of Reliability (CoR) is

335   lowest for test #1 - the letter-strength (non-overlapping) task. However, as indicated above, it would appear that this reliability comes at the expense of failing to capture individual variation in SED.

Based on ICCs and corresponding F values, tests #2 and #3 appear particularly useful. However, ICCs are driven by values at the ends of the measured range, making them susceptible to outliers. In our data set, participant ID3 reported quite unbalanced but reliable scores on tests #2, 3 and 5. This

340   individual may be an outlier, inflating ICC scores for these tests. MAP and the related FRP estimates circumvent this issue by scoring on rank rather than absolute value. This difference in method impacts test #6 the most, as it has a poor ICC (and F), but fair MAP and FRR values.

Figure 3. Bland-Altman plots of results from the eight tests. (#1-6) Mean contrast balance across runs is

345   plotted against the difference in contrast balance. (#7) Relative time spent in each possible rivalrous state plot against the different in this measure across runs. (#8) Mean stereoacuity plot against the difference in stereoacuity across runs. Shaded regions denote the 95% confidence intervals across estimates with error bars on these estimates calculated using the procedure given by Carkeet (2015). Note high repeatability in (#1) but restricted range of estimated balance across participants. Motion test (#6), in contrast, produced a broader

350   range of estimated balance but at the expense of repeatability. (#3) The letter-polarity test combines high repeatability with a broad range of estimated contrast-balance.

	ICC	F (p)	CoR	MAP	FRP
1. Letter-strength (non-overlapping)	0.603	4.035 (0.00014)	**0.058***	0.174	0.638
2. Letter-strength (overlapping)	0.727	6.336 (<0.00001)	0.224	0.197	0.648
3. Letter-polarity (overlapping)	**0.803***	**9.152* (<0.00001)**	0.105	**0.299***	**0.776***
4. Phase-comb (90°)	0.230	1.597 (0.10425)	0.169	0.170	0.592
5. Phase-suppression (180°)	0.629	4.388 (0.00006)	0.124	0.198	0.711
6. Motion strength (contrast)	0.316	1.924 (0.03988)	0.494	0.261	0.752
7. Motion rivalry (duration)	0.201	1.504 (0.13607)	0.208	0.141	0.539
8. Stereo	0.392	2.29 (0.013)	40.2	0.222	0.688

Table 2. Summary statistics for performance of the eight tests. *Indicates best performance for a given metric. *ICC*:intraclass correlation, *F*: ANOVA *F* statistic, *CoR*: coefficient of repeatability (lower is better), *MAP*: mean average precision (higher is better), *FRP*: fractional rank precision (higher is better; Dorr, Elze et al. 2017). Note Note CoR measures for Stereo are unique being expressed in units of arc sec.\

Figure 4 shows the variation of the CoR and ICC over run-length. Here we analysed from 25 to 100% of data (i.e. either trials 5-20, for tests #1-6 and 8 or from 4.2-30s and 34.2-60s in test #7). For each test (colour coding given in the legend), CoR-coefficient of repeatability (Figure 4a) and ICC-interclass correlation (Figure 4b) indexes are obtained by comparing first and second run across all observers. Predictably, there is better agreement towards the end of runs, as evidenced by the descending (Figure 4a) or ascending (Figure 4b) trend of the plotted lines. Figure 4c visually represents CoR (x-axis; lower is better) against ICC (y-axis; higher is better) indexes for data from the whole run for all of the tasks tested. Note the clustering of results. Results from the two motion- and the non-rivalrous phase-combination tests were substantially poorer than the other tests.

Figure 4. Plot of the evolution of (a) coefficient of repeatability (lower is better) and (b) interclass correlation (higher is better) over run-length (expressed as a percentage of the run analysed, e.g. 100%=20 trials) for the seven balance tests. The tests are listed in legend-panel (a). (c) Plot of intra-class correlation against coefficient of repeatability.

We next assessed whether increased binocular imbalance was associated with poorer stereoacuity and/or with more marked eye-dominance in rivalry. As shown in Figure 5a, we fitted a regression line to the data points corresponding to (y-axis) the magnitude of SED and (x-axis) the stereoacuity estimate from the same participant (mean of the two runs). The mean magnitude of SED was

obtained by averaging contrast-BP indexes across tests #1-6 and computing the absolute difference of this value from 0.5. The fit was consistent with only a marginally significant association between measures (R=0.30; *p*= 0.104). Figure 5b plots SED magnitude against the mean duration of instances of perceptual dominance, as measured in the rivalry test (#7). These data show a non-significant correlation of R=0.25, *p*=0.191. Participants rivalrous percepts lasted for an average of 1.67seconds (σ=±1.02s).

Figure 5 Data from each numbered participant (colour coding of tests is as Figure 3) showing either (a) stereoacuity (task #8) or (b) the duration of perceptual dominance (task #7) plot against magnitude of SED (based on the mean contrast-Balance-Point averaged across test #1-6). All data are averaged across two runs. Magnitude of SED is |SED-0.5|, i.e. 0.5=balanced vision, and 1=complete dominance of one eye. On each graph, a regression line has been fit to data, and correlation indexes (*R*) with associated *p*-values are reported.

Discussion

We compared eight tests that use dichoptic stimuli to quantify binocular visual function. We measured sensory-eye-dominance (SED) in tests #1-7, and stereoacuity in test #8 in 30 individuals with ostensibly normal vision. All SED tests involved estimation of a *Balance-Point* (BP): an estimate of the extent to which participants relied equally on the two components of a dichoptic stimulus-pair. Specifically, pairs of *contrast*-modulated stimuli were used in tests #1-6, comprising Sloan letters (adapted from Kwon, Wiecek et al. 2015 -tests #1-2, Bossi, Tailor et al. 2017 -test #3), reciprocally-shifted gratings (from Ding and Sperling 2006 -tests #4-5) or up/down drifting noise-patterns(loosely based on the motion task by Black, Thompson et al. 2011 -test #6). Test #7 (inspired by Dieter, Sy et al. 2016) measured rivalry, as the proportion of dominant-eye instances (presenting the same pairs of stimuli used in # 6 but at a fixed contrast). In addition, test #8 measured stereoacuity, using stereo-defined Sloan letters (variant of Frisby, Davis et al. 1996).

We observed a high level of consistency in BPs across tests for each participant. The mean magnitude of SED, obtained by averaging absolute BPs, was 0.55, σ=±0.03, with the majority of participants (22/30; mean BP=0.5; σ=0.03) showing right-eye sensory dominance, in line with previous results (e.g. Ehrenstein, Arnold-Schulz-Gahmen et al. 2005, Pointer 2012, Johansson, Seimyr et al. 2015). Overall, regardless of the statistics used, test #3 showed the best performance in term of validity and reliability; while tests #7 and #4 were the poorest (see Table 2). We found no correlation between stereoacuity and SED measures: better binocular balance (i.e. BPs around 0.5) was not associated with better stereoacuity (Figure 5a).

Why are some tests better than others? In terms of test-validity an ideal test captures a range of BP estimates across participants. We note that binocular imbalance manifests most robustly for stimuli presented at the same location so that estimates of BP from test #1 (non-overlapping letter strength) converge on 0.5. This fundamentally limits the validity of this test for capturing variation in SED within a population of people with normal vision.

In terms of test-reliability, one might expect that tests involving rivalrous stimuli would elicit less reliable responses from participants as e.g. rivalrous percepts typically fluctuate in time. We presented stimuli for relatively short durations, to limit such perceptual alternation. Nevertheless, our expectation - that these stimuli would still elicit more random response – was not borne out by the reliability of tests involving rivalrous stimuli. Rather, our finding of high levels of reliability on these tests suggests that the SED may selectively drive the early stages of a rivalrous percept. This is consistent with earlier reports of participants' first percept of a rivalrous display predicting their eye dominance (e.g. quantified using percept frequency) calculated for stimuli that alternated over dozens of seconds (Xu, He et al. 2011, Dieter, Sy et al. 2016).

Among the contrast-balance tests, those which rival in motion (#6) and form (#2) show the poorest test-retest reliability (CoR). By contrast, tests #3 and #5, which only rival in polarity, exhibit better reliability. We believe that this difference arises from polarity-rivalry removing the need for participants to explicitly judge relative stimulus strength. For two rivalrous, same-polarity but different-identity targets (e.g. test #2), observers likely judge relative stimulus-strength by comparing two independent estimates of contrast. In this case *binocular imbalance manifests as a difference in magnitude of the two cues* (i.e. one will be "stronger/whiter/brighter" than the other). In contrast, the appearance of two rivalrous, opposite-polarity but same-identity targets (test #3), alternates over time with the *target-pair* appearing as either left-black/right-white or left-white/right-black. In this scenario *binocular imbalance manifests as a difference in probability that one state will dominate over the other.* In other words, for test #2 the observer is faced with a judgement of a subtle difference in perceived luminance, whereas in test #3 they are always judging which of two (apparently) black and white letters is white. This task is straightforward for our adult participants and sufficiently simple for children that we can reliably administer a very similar test using an unsupervised measurement system, as part of a home-based binocular amblyopia therapy (Bossi, Tailor et al. 2017).

How could new tests fit into clinical practice? There is an unmet need for accessible, valid and reliable methods to measure binocularity in standard care. In the clinic, binocularity is frequently assessed measuring stereoacuity. However, 1-14% of the population is stereo-blind (Bosten, Goodbourn et al. 2015), and binocular function is necessary but not sufficient for stereoacuity. Moreover, standard stereo-tests (e.g. Frisby, randdot) result in non-measurable stereoacuity in cases where compensation for visual deficits (e.g. balancing stimulus visibility across the eyes) allow patients to see stereo (Tytla, Lewis et al. 1993), or achieve binocular summation (Baker, Meese et al. 2007). We therefore would anticipate SED being able to (a) quantify subtle binocular deficits in people with normal stereoacuity and (b) reveal the presence of binocular capacity in the absence of measurable stereoacuity. Our current results speak to (a) – we observe a range of SEDs in our test group – albeit centred on 0.5 or balanced vision. Differences between observers are stable, both in terms of test-retest reliability and in terms of agreement amongst different tests of SED. The current study does not speak to (b) since we have not targeted individuals with abnormal binocular vision. Inclusion of participants with poor binocular vision (the subject of an ongoing study) may influence the outcome of our evaluation For example, the excellent test-retest reliability of test #1 may outweigh its poor test validity as patients are more likely to exhibit a wider range of larger BPs which may drive this test well but could lead to ceiling effects in more sensitive tests.

455 Other clinical tests are problematic for their own reasons. For example, tests of "sighting-dominance" (e.g. Miles or Porta tests) deliver binary "left" or "right" measures of eye-dominance. Further, sighting-dominance can be inconsistent, influenced by gaze direction (Khan and Crawford 2001), and by the test used (Rice, Leske et al. 2008). The only continuous estimate of eye-dominance available to clinicians comes from Sbisa bars which rely on the clinician's judgement of what stimulus level induces a patient to

460 report diplopia. Our procedures have much in common with the Sbisa bar but automate stimulus selection, have the patient perform a forced-choice task, and use all of the response-information to calculate the balance point. This we believe will contribute to superior test reliability and better compliance from patients.

It is important to note that the sight-dominant eye does not necessarily support better visual acuity

465 (Pointer 2007), and laterality - measured with either SED or sight-dominance - only agrees in 50% of cases (Pointer 2012). Further, tests of SED (Suttle, Alexander et al. 2009, Pointer 2012) as well as those of other forms of binocularity, such as sight dominance (Rice, Leske et al. 2008) and stereoacuity (Heron and Lages 2012), produce results that differ depending on the particular visual task and test-conditions. Clearly, both sensory and motor aspects are involved in binocularity. Thus, a complete assessment of

470 functional binocular vision requires more than one type of binocular measurement. The lack of correlation between our (reliable) measures of SED and stereo-acuity suggests these tests are complementary. Future research could more directly focus on determining the sub-processes that support complete binocular vision, and developing a more complete battery of tests to probe them.

SED and amblyopia There is increasing interest in the feasibility of treating amblyopia in older children

475 and adults, either with occlusion therapy (Erdem, Cinar et al. 2011, Kishimoto, Fujii et al. 2014) or other monocular (Evans, Yu et al. 2011) and/or binocular methodologies, such as dichoptic training and videogame play (Tsirlin, Colpa et al. 2015). Based on the idea that interocular suppression is a cause not a symptom of amblyopia it has been proposed that binocular treatments achieve their results by reducing interocular suppression. However, a few studies now suggest that therapeutic benefit cannot be

480 underpinned solely by a reduction in interocular suppression (Vedamurthy, Nahum et al. 2015, Bossi, Tailor et al. 2017). There a number of current hypothesis about the mechanism supporting visual improvement but importantly there are also a large number of different measures of SED in use across different labs. The assumption that all tests of SED tap the same mechanism may not be true and this may hamper comparative evaluation of the outcomes of different binocular therapies. Having a standard

485 set of tests to measure binocularity could be useful in clarifying how and in which sense interocular suppression, as quantified by the magnitude of imbalance in SED, affects the therapeutic outcome for, and the mechanism of, amblyopia.

Conclusion

We compared several tests for rapidly quantifying sensory-eye dominance and a test of stereoacuity.

490 A judgement of which of two dichoptically-superimposed opposite contrast-polarity patterns dominated the participant's percept (test #3) supports a reliable and sensitive measure of sensory-eye dominance in only 20 trials. Practical and reliable measure of SED have application in amblyopia research and as part of a more thorough assessment of binocularity in the clinic.

Acknowledgements

495 We thank all the participants and Steven Lim and Young Oh for assistance with data collection. This work was funded by a UCL Impact award to SCD and by a grant from Cure Kids, NZ to SCD & LH.

References

500 Babu, R. J., S. Clavagnier, W. R. Bobier, B. Thompson and R. F. Hess (2017). "Regional Extent of Peripheral Suppression in Amblyopia." Invest Ophthalmol Vis Sci **58**(4): 2329-2340.

Baker, D. H., T. S. Meese and R. F. Hess (2008). "Contrast masking in strabismic amblyopia: attenuation, noise, interocular suppression and binocular summation." Vision Res **48**(15): 1625-1640.

Baker, D. H., T. S. Meese, B. Mansouri and R. F. Hess (2007). "Binocular summation of
505 contrast remains intact in strabismic amblyopia." Invest Ophthalmol Vis Sci **48**(11): 5332-5338.

Baker, D. H., T. S. Meese and R. J. Summers (2007). "Psychophysical evidence for two routes to suppression before binocular summation of signals in human vision." Neuroscience **146**(1): 435-448.

Birch, E., C. Williams, J. Drover, V. Fu, C. Cheng, K. Northstone, M. Courage and R. Adams
510 (2008). "Randot Preschool Stereoacuity Test: normative data and validity." J aapos **12**(1): 23-26.

Birch, E. E. (2013). "Amblyopia and binocular vision." Progress in Retinal and Eye Research **33**(0): 67-84.

Birch, E. E., S. L. Li, R. M. Jost, S. E. Morale, A. De La Cruz, D. Stager, Jr., L. Dao and D. R. Stager, Sr. (2015). "Binocular iPad treatment for amblyopia in preschool children." J AAPOS **19**(1): 6-
515 11.

Black, J. M., R. F. Hess, J. R. Cooperstock, L. To and B. Thompson (2012). "The measurement and treatment of suppression in amblyopia." J Vis Exp(70): e3927.

Black, J. M., B. Thompson, G. Maehara and R. F. Hess (2011). "A compact clinical instrument for quantifying suppression." Optom Vis Sci **88**(2): E334-343.
520 Bossi, M., V. K. Tailor, E. J. Anderson, P. J. Bex, J. A. Greenwood, A. Dahlmann-Noor and S. C. Dakin (2017). "Binocular Therapy for Childhood Amblyopia Improves Vision Without Breaking Interocular Suppression." Invest Ophthalmol Vis Sci **58**(7): 3031-3043.

Bosten, J. M., P. T. Goodbourn, A. J. Lawrance-Owen, G. Bargary, R. E. Hogg and J. D. Mollon (2015). "A population study of binocular function." Vision Res **110**(Pt A): 34-50.
525 Carkeet, A. (2015). "Exact parametric confidence intervals for Bland-Altman limits of agreement." Optom Vis Sci **92**(3): e71-80.

Cotter, S. A., L. A. Cyert, J. M. Miller and G. E. Quinn (2015). "Vision screening for children 36 to <72 months: recommended practices." Optom Vis Sci **92**(1): 6-16.

Crawford, L. C. J. and H. J. Griffiths (2015). "The repeatability of the Sbisa bar for testing
530 density of suppression." British and Irish Orthoptic Journal **12**: 35-40.

Cumming, B. G. and G. C. DeAngelis (2001). "The physiology of stereopsis." Annu Rev Neurosci **24**: 203-238.

Dieter, K. C., J. L. Sy and R. Blake (2016). "Individual differences in sensory eye dominance reflected in the dynamics of binocular rivalry." Vision Res.
535 Ding, J. and G. Sperling (2006). "A gain-control theory of binocular combination." Proceedings of the National Academy of Sciences of the United States of America **103**(4): 1141-1146.

Dorr, M., T. Elze, H. Wang, Z. L. Lu, P. J. Bex and L. Lesmes (2017). New precision metrics for contrast sensitivity testing.

Ehrenstein, W. H., B. E. Arnold-Schulz-Gahmen and W. Jaschinski (2005). "Eye preference
540 within the context of binocular functions." Graefe's Archive for Clinical and Experimental Ophthalmology **243**(9): 926-932.

Erdem, E., G. Y. Cinar, D. Somer, N. Demir, A. Burcu and F. Ornek (2011). "Eye patching as a treatment for amblyopia in children aged 10-16 years." Jpn J Ophthalmol **55**(4): 389-395.

Evans, B. J. W., C. S. Yu, E. Massa and J. E. Mathews (2011). "Randomised controlled trial of
545 intermittent photopic stimulation for treating amblyopia in older children and adults." Ophthalmic Physiol Opt **31**: 56-68.

Ferris, F. L., 3rd, A. Kassoff, G. H. Bresnick and I. Bailey (1982). "New visual acuity charts for clinical research." Am J Ophthalmol **94**(1): 91-96.

Fricke, T. R. and J. Siderov (1997). "Stereopsis, stereotests, and their relation to vision screening and clinical practice." Clinical and Experimental Optometry **80**(5): 165-172.

Frisby, J. P., H. Davis and K. McMorrow (1996). "An improved training procedure as a precursor to testing young children with the Frisby Stereotest." Eye (Lond) **10 (Pt 2)**: 286-290.

Hamm, L., Z. Chen, J. Li, J. Black, S. Dai, J. Yuan, M. Yu and B. Thompson (2017). "Interocular suppression in children with deprivation amblyopia." Vision Res **133**: 112-120.

Hess, R. F. and B. Thompson (2015). "Amblyopia and the binocular approach to its therapy." Vision Res.

Huang, C.-B., J. Zhou, Y. Zhou and Z.-L. Lu (2010). "Contrast and phase combination in binocular vision." PLoS One **5**(12): e15075.

Huang, C. B., J. Zhou, Z. L. Lu, L. Feng and Y. Zhou (2009). "Binocular combination in anisometropic amblyopia." J Vis **9**(3): 17 11-16.

Johansson, J., G. Ö. Seimyr and T. Pansell (2015). "Eye dominance in binocular viewing conditions." Journal of vision **15**(9): 21-21.

Julesz, B. (1971). Foundations of Cyclopean Perception, University of Chicago Press.

Kehrein, S., T. Kohnen and M. Fronius (2016). "Dynamics of Interocular Suppression in Amblyopic Children during Electronically Monitored Occlusion Therapy: First Insight." Strabismus **24**(2): 51-62.

Khan, A. Z. and J. D. Crawford (2001). "Ocular dominance reverses as a function of horizontal gaze angle." Vision Research **41**(14): 1743-1748.

Kishimoto, F., C. Fujii, Y. Shira, K. Hasebe, I. Hamasaki and H. Ohtsuki (2014). "Outcome of conventional treatment for adult amblyopia." Jpn J Ophthalmol **58**(1): 26-32.

Kwon, M., Z. L. Lu, A. Miller, M. Kazlas, D. G. Hunter and P. J. Bex (2014). "Assessing binocular interaction in amblyopia and its clinical feasibility." PLoS One **9**(6): e100156.

Kwon, M., E. Wiecek, S. C. Dakin and P. J. Bex (2015). "Spatial-frequency dependent binocular imbalance in amblyopia." Sci Rep **5**: 17181.

Li, J., R. F. Hess, L. Y. Chan, D. Deng, X. Chen, M. Yu and B. S. Thompson (2013). "How best to assess suppression in patients with high anisometropia." Optom Vis Sci **90**(2): e47-52.

Li, J., R. F. Hess, L. Y. Chan, D. Deng, X. Yang, X. Chen, M. Yu and B. Thompson (2013). "Quantitative measurement of interocular suppression in anisometropic amblyopia: a case-control study." Ophthalmology **120**(8): 1672-1680.

Li, J., C. S. Lam, M. Yu, R. F. Hess, L. Y. Chan, G. Maehara, G. C. Woo and B. Thompson (2010). "Quantifying sensory eye dominance in the normal visual system: a new technique and insights into variation across traditional tests." Invest Ophthalmol Vis Sci **51**(12): 6875-6881.

Li, J., B. Thompson, D. Deng, L. Y. Chan, M. Yu and R. F. Hess (2013). "Dichoptic training enables the adult amblyopic brain to learn." Curr Biol **23**(8): R308-309.

Mansouri, B., B. Thompson and R. F. Hess (2008). "Measurement of suprathreshold binocular interactions in amblyopia." Vision Res **48**(28): 2775-2784.

McCormick, A., R. Bhola, L. Brown, D. Squirrel, J. Giles and I. Pepper (2002). "Quantifying relative afferent pupillary defects using a Sbisa bar." Br J Ophthalmol **86**(9): 985-987.

McGraw, K. O. and S. P. Wong (1996). "Forming inferences about some intraclass correlation coefficients." Psychological methods **1**(1): 30.

McKee, S. P., D. M. Levi and J. A. Movshon (2003). "The pattern of visual deficit in amblyopia." Journal of Vision **3**(5): 380-405.

Miles, W. R. (1930). "Ocular Dominance in Human Adults." The Journal of General Psychology **3**(3): 412-430.

O'Connor, A. R., E. E. Birch, S. Anderson and H. Draper (2010). "The functional significance of stereopsis." Invest Ophthalmol Vis Sci **51**(4): 2019-2023.

Ooi, T. L. and Z. J. He (2001). "Sensory eye dominance." Optometry **72**(3): 168-178.

Ooi, T. L., Y. R. Su, D. M. Natale and Z. J. He (2013). "A push-pull treatment for strengthening the 'lazy eye' in amblyopia." Curr Biol **23**(8): R309-310.

600   Piano, M. and D. Newsham (2015). "A pilot study examining density of suppression measurement in strabismus." Strabismus **23**(1): 14-21.

Pointer, J. S. (2012). "Sighting versus sensory ocular dominance." Journal of Optometry **5**(2): 52-55.

Prins, N. and F. A. A. Kingdom. (2009). "Palamedes: Matlab routines for analyzing
605   psychophysical data.", from http://www.palamedestoolbox.org

Rice, M. L., D. A. Leske, C. E. Smestad and J. M. Holmes (2008). "Results of Ocular Dominance Testing Depend on Assessment Method." Journal of AAPOS : the official publication of the American Association for Pediatric Ophthalmology and Strabismus / American Association for Pediatric Ophthalmology and Strabismus **12**(4): 365-369.

610   Tailor, V., S. Balduzzi, S. Hull, J. Rahi, C. Schmucker, G. Virgili and A. Dahlmann-Noor (2014). "Tests for detecting strabismus in children age 1 to 6 years in the community." Cochrane Database of Systematic Reviews(7).

Tsirlin, I., L. Colpa, H. C. Goltz and A. M. Wong (2015). "Behavioral Training as New Treatment for Adult Amblyopia: A Meta-Analysis and Systematic Review." Invest Ophthalmol Vis Sci
615   **56**(6): 4061-4075.

Tytla, M. E., T. L. Lewis, D. Maurer and H. P. Brent (1993). "Stereopsis after congenital cataract." Invest Ophthalmol Vis Sci **34**(5): 1767-1773.

Vaz, S., T. Falkmer, A. E. Passmore, R. Parsons and P. Andreou (2013). "The Case for Using the Repeatability Coefficient When Calculating Test–Retest Reliability." PLoS ONE **8**(9): e73990.

620   Vedamurthy, I., M. Nahum, D. Bavelier and D. M. Levi (2015). "Mechanisms of recovery of visual function in adult amblyopia through a tailored action video game." Sci Rep **5**: 8482.

Vedamurthy, I., M. Nahum, S. J. Huang, F. Zheng, J. Bayliss, D. Bavelier and D. M. Levi (2015). "A dichoptic custom-made action video game as a treatment for adult amblyopia." Vision Res **114**: 173-187.

625   Watson, A. B. and D. G. Pelli (1983). "QUEST: a Bayesian adaptive psychometric method." Percept Psychophys **33**(2): 113-120.

Wheatstone, C. (1838). "Contributions to the Physiology of Vision.--Part the First. On Some Remarkable, and Hitherto Unobserved, Phenomena of Binocular Vision." Philosophical transactions of the Royal Society of London **128**: 371-394.

630   Wiecek, E., K. Lashkari, S. C. Dakin and P. Bex (2015). "Metamorphopsia and interocular suppression in monocular and binocular maculopathy." Acta Ophthalmol **93**(4): e318-320.

Wong, A. M. F. (2012). "New concepts concerning the neural mechanism of amblyopia and their clinical implications." Can J Ophthalmol **47**(5): 399-409.

Xu, J. P., Z. J. He and T. L. Ooi (2011). "A binocular perimetry study of the causes and
635   implications of sensory eye dominance." Vision research **51**(23-24): 2386-2397.

Zhou, J., S. Clavagnier and R. F. Hess (2013). "Short-term monocular deprivation strengthens the patched eye's contribution to binocular combination." J Vis **13**(5).

Zhou, J., P. C. Huang and R. F. Hess (2013). "Interocular suppression in amblyopia for global orientation processing." J Vis **13**(5): 19.

640   Zhou, J., A. Reynaud and R. F. Hess (2014). "Real-time modulation of perceptual eye dominance in humans." Proc Biol Sci **281**(1795).

Zhou, J., B. Thompson and R. F. Hess (2013). "A new form of rapid binocular plasticity in adult with amblyopia." Sci Rep **3**: 2638.