Information synergy: adding unambiguous quality information rescues social information use in ants

Tomer J. Czaczkes [1,3] and John J. Beckwith [1,2]

1) Institute of Zoology, Universität Regensburg, 93053 Regensburg, Germany
2) Department of Zoology, University of Oxford, Oxford OX13PS, UK
3) Corresponding author, email: tomer.czaczkes@ur.de

Abstract

Animals have access to many alternative information sources when making decisions, such as private information (e.g. memory) and social information. Social insects make extensive use of social information. However, when intentional social information (e.g. pheromone trails in ants) conflicts with private information (e.g. route memories), insects often follow their private information. Why is this? We propose that it is an asymmetry in information content between these two information sources that drives the neglect of social information. In ants, workers with certain information about the quality of a food source (memory) ignore valuable social information (pheromone trails) because the pheromone trails encode only a very ambiguous measure of food quality. This leads to a testable hypothesis; the addition of unambiguous quality information should rescue social information following. To test this, we trained ants to a poor quality (0.25M) food source, and then provided an alternative path along with either 1) no information, 2) a pheromone trail, 3) a 0.2µl droplet of 1.5M sugar solution, providing unambiguous quality information, or 4) both a trail and a droplet. When either no or only one information source was provided (1-3), most ants (60-75%) continued following their own memory. However, the addition of unambiguous quality information to a trail (4) rescued trail following: when both a pheromone trail and a sugar droplet were provided, 75% of ants followed the trail. We propose that information content – what type of information a source can convey, and what is its ambiguity, is a strong driver of which type of information is attended to.

Introduction

Animals have access to many types of information. These can be broadly separated into private information, which is available only to the individual animal, and public information, which is available to other animals as well [1]. Private information includes genetic information and non-genetic information such as internal states and, importantly, memories. Public information includes all observable cues, such as the colour of a flower. An important subset of public information is social information, which is information related in some way to other animals. These may be incidental cues, such as the presence of conspecifics at a resource, or intentionally produced signals, such as the waggle-dance of a honey bee.

Multiple information sources may provide information about the same thing. For example, both the colour and the odour of a flower patch, and whether other bees are dancing for that patch, might inform a bee about whether and where food is available in the environment [2,3]. These multiple information sources may interact in a variety of ways. Information may be used hierarchically, with one being used whenever it is available, and if it is not the next on the list is used [4]. When information sources agree, they may act additively or synergistically to improve behaviour. For example, ants which have private information about a food source (memory) which agrees with social information (a pheromone trail leading in the same direction) walk 25% faster than ants with either only one or none of these information sources [5]. These information sources also act additively by causing more ants to follow a pheromone trail [6]. Bumblebees which have experienced a novel flower odour in the nest (public information) are more likely to exploit the novel flowers, but only once they have tasted a flower scented with this novel odour (private information) [7].

A particularly interesting situation is when information sources conflict. One option in such situations is to weight information from different sources and produce and an intermediate value [6,8,9]. An alternative is to rely on one type of information and ignore the others. In some situations, such as choosing a path at a trail bifurcation, an intermediate option is not available. Different strategies might be undertaken due to different ecological conditions or the state of individual animals, and understanding what strategies animals use to decide which information to follow is a very active field of research [10–14]. Information use strategies can be broadly divided in to *when* strategies (when to use which information type) and *who* strategies (whose information to use) [15].

Evidence supporting various information use strategies has been described in wide array of taxa, including insects, fish, lizards, rats, and humans [16–19]. Information use strategies have been especially well researched in social insects such as ants and bees, as in these groups social

---

[1] Our definition of private information deviates somewhat from that of Danchin et al. [1], who consider public information more-or-less synonymous with social information. We consider social information a subset of public information.

information has the potential to play a very large role [see 11,20 and references therein]. Social
insects also represent a special case, since there are not usually expected to be any conflicts of
interest in many situations requiring communication, such as foraging and nest-site selection. This
should, at face value, strengthen the role of explicit social signals such as pheromone trails or
waggle-dances. However, somewhat surprisingly, in most cases in which such intentionally produced
social information is conflicted with private information, the ants and bees predominantly follow
their own memories [21–29]. This was found not to be the case in only a few instances [23,30]. This
preference could have strong implications for colony level behaviour, preventing colonies from
optimally exploiting their environment.

Why do ants and bees so often underweight or ignore the directional information provided by
their sisters? Explanations of information use strategies have been made in terms of the reliability of
an information source – how often using one type of information is associated with a positive
outcome, or how outdated the information is (e.g. [14,31]), or the cost of acquiring new information
[32–34]. However, what has not been considered is the *information content* of different types of
information.

Economists, and later behavioural ecologists, distinguish between two types of situations with an
unknown outcome: risk and ambiguity (or uncertainty) [35–37]. The term 'risk' refers to a situation
where the mean result or probabilities of a situation are known, but the precise value at any one
time is not. A gamble on a six-sided die is thus a risky proposition: the average score is 3, the
probability of any one side is 1/6, but the precise value on each roll is unknown. By contrast,
'ambiguity' or 'uncertainty' refers to a situation where neither the mean nor the variation is known; a
gamble on a die with an unknown number of sides is thus gambling under ambiguity.

When faced with ambiguity, humans have been found to behave in a manner inconsistent with
the predictions of expected utility theory, the standard economic theory of human behaviour.
Specifically, humans are found to display *ambiguity aversion.* Given a choice between a risky and an
ambiguous proposition, humans overwhelmingly prefer the risky proposition, although there is no *a*
priori rational reason for them to do so [38,39], (but see [37] for a Baysian perspective). This was
famously described by the Ellsberg paradox [38]. He gives the example of a single choice, in which
you are offered to pull one ball out of one of two urns. One urn contains an equal mix of red and
black balls, the other contains an unknown ratio of red to black balls. Pulling out a red ball is
rewarded with $100, but the black ball is unrewarded. The urns are thus a risky and an ambiguous
proposition, respectively. Most people choose the urn with the known distribution. Related to this,
Kahneman and Tversky, in their famous Prospect Theory paper [40], describe *the Certainty Effect.*
Here, subjects overweight certainty, and are willing to pay a premium to move from a risky to a

certain situation. For example, people usually prefer $3000 with certainty to an 80% chance of

$4000, although the expected payoff of the gamble is $3200. Effectively, subjects are willing to pay

$200 for certainty. Ambiguity aversion has also been documented in non-human primates [41,42].

A similar situation may be happening in social insects experiencing an information conflict.

Imagine an ant which has found a poor quality food source. It is shuttling back and forth to this food

source, and knows the quality of this resource with certainty. It then encounters a pheromone trail

leading in a different direction. This pheromone trail was deposited by another ant, which had

discovered a good food source. It would be in both of the ants' interests for the ant exploiting the

poor food to follow the pheromone trail. However, this often does not happen, even if the

recruitment to the alternative direction is very strong [21–29]. Why is this? We propose this is due

the (lack of) information content of the social communication signal. The pheromone trail provides

only very noisy, imprecise, and ambiguous information about the quality of a food source.

Pheromone trail strength varies with resource quality, time since discovery, recruitment rates, and

the individual pheromone depositions behaviour of ants: A strong trail may lead to a good food

source exploited by a few ants, or to a poor one exploited by many ants; a weak trail may lead to a

poor food source, or to a good one which has been unproductive for a while, or to a newly

discovered food source. Importantly, there is a very large variation in the amount of pheromone

deposited by individual ants to food sources of the same quality [43]. With no strong evidence that

the new food source is better, the forager avoids paying the costs of attempting to find an advertised

food source [32,44,45], and falls back on exploiting the food source she knows about.

This hypothesis leads to a testable prediction: If a worker foraging on a poor food source can be

given unambiguous information about the quality of a better food source a pheromone trail

advertises, she should follow the trail pheromone. Here, we set out to test this prediction.

Materials and methods

Study species and maintenance

We used 8 queenless colony fragments of the black garden ant, *Lasius niger* (Linnaeus),

collected from eight different colonies on the University of Regensburg campus. Colonies were

housed in a plastic box (40×30×20cm) with a layer of plaster on the bottom. Each box contained a

circular plaster nest (14cm diameter, 2cm high). Colonies contained c. 1000 workers and small

amounts of brood. The ants were fed *ad libitum* on 1M sucrose solution supplemented with

Drosophila melanogaster fruit flies. Colonies were deprived of food for four days prior to each trial to

give high and consistent motivation for foraging and pheromone deposition. Water was provided *ad*

libitum.

Experimental procedure

10 different experimental treatments were carried out in total, coded A-J. An overview of the

different treatments is provided in table S4 in the supplement. All treatments were variations on a

central design. The aim of the experiments was to test whether ants with a well-established memory

of finding food on one arm of a Y maze can be induced to search a newly-presented alternate arm by

the provision of various information sources. The information sources provided could be a

pheromone trail, a small droplet of sucrose (simulating trophallaxis), a novel scent in either the

droplet or on the newly presented path or both, or a combination of these. The scent treatments

were added to test whether similar droplet and pathway odours cause a further increase in trail

follow, as there could thus be an associative link between the quality of the droplet and the odour of

the new runway. However, as there was no effect of the various odour treatments (see supplement

S1 for details) we pooled our data into treatments groups for the final analysis, depending on the

types of information presented to the ants, and the quality of the original feeder to which it was

trained. This resulted in 5 treatment groups: i) no information (treatments A and B), ii) only a droplet

of 1.5M sucrose just before the bifurcation (treatments C-E), iii) only a pheromone trail on the new

path (treatment F), vi) both a pheromone trail and a droplet (treatments G-I), and finally v) a

pheromone and a droplet for ants trained to 1.5M sucrose (treatment J).

A trial began by allowing an ant onto the apparatus via a drawbridge. The drawbridge led to a

Y maze with one arm drawn out just out of reach of the ants, to form an L maze (Figure 1). The arm

that was out of reach (left or right) was systematically varied. The arms of the maze were 10cm long

and 1cm wide, narrowing to 2mm wide at the junction. The arms of the maze were covered with

paper overlays. The overlay on the stem of the maze was unscented, while the overlay on the arm of

the maze was scented with either rosemary or lemon essential oil – the odour used was

systematically varied between trials. Odour impregnation was achieved by storing the paper overlays

in a sealed plastic container containing one 0.1ml drop of essential oil on a glass petri dish for at least

24 hours. An acetate sheet affixed to the end of the L maze arm acted as a feeder. A large drop of

0.25M sucrose solution (1.5M in treatment J), flavoured similarly to the L maze arm leading to it, was

placed onto the feeder. Sucrose solutions were flavoured by adding 10µl essential oil per 100ml

sucrose solution, following Czaczkes *et al.* [46]. Once the ant found the feeder it was marked with a

dot of acrylic paint on the abdomen, and allowed to return to the nest. While the ant was in the nest,

unloading her sucrose load, the paper overlays on the stem and arm were replaced by fresh overlays,
to remove any trail pheromone the ant might have laid. The ant was then allowed to make 3 further
return visits to the same feeder, resulting in 4 visits to the original feeder in total. This is sufficient to
ensure that, even given the relatively low quality of food, most ants will return to this arm of the
maze given a choice (Oberhauser, Koch and Czaczkes, submitted, see also data below).

While the ant was in the nest prior to its fifth return to the feeder, the feeder from the
original arm of the Y maze was removed, and access to the second arm was given. The new arm was
either marked with a pheromone trail (treatments F - J) or not (treatments A-E). This pheromone trail
was produced by immersing 8 worker hindgut glands in 2ml of dichloromethane (DCM), following
von Thienen *et al*. [47]. 5.6µl of this mixture was applied in an even line along the paper overlay
covering the arm, using a capillary tube (Servoprax GmbH, Germany). This amount was calculated to
produce a pheromone trail of a realistic strength [47], and in control trials elicited a trail following
accuracy of 82% (594 / 715), which is indistinguishable from those recorded from naïve ants
following a reasonably strong naturally-deposited trails as reported by Czaczkes et al. [48] (82-83%
accuracy). See supplement S1 for detailed methods and results regarding the artificial pheromone
trail.

0.2ul droplet of 1.5M sucrose. May be flavoured to match the right arm

100mm

10mm

flavoured 0.25M sucrose feeder (not present on visit 5)

Bridge to nest

Access to second Y maze arm given after 4 visits to the first arm.

Paper overlay may be marked with trail pheromone.

Paper overlay may be scented with a novel scent.

Figure 1 – experimental setup. Ants are first trained for 4 visits on a Y maze with access to only one arm (here the top arm, in yellow in the figure but not coloured in the experiment), leading to a 0.25M sucrose feeder flavoured with lemon or rosemary (1.5M in treatment J). On the 5[th] outward journey the second arm of the Y maze (here bottom, green) is added, and the feeder removed. The new arm may be marked with a pheromone trail and/or scented with a novel scent (lemon or rosemary). Additionally, a small (0.2μl) droplet of 1.5M sucrose may be placed after a narrowing of the stem before the bifurcation, which the ant can drink but will not become satiated from. The droplet may be flavoured to match the new Y maze arm scent or may be unflavoured.

The new arm was also either scented (treatments B, D, E, H, I, J) with a different scent to that of the original arm, or was not (treatments A, C, F, G). Lastly, a small (c. 0.2μl) droplet of sucrose either was (treatments C, D, E, G, H, I, J) or was not (treatments A, B, F) placed on the stem of the Y maze, before the bifurcation but just after a narrowing of the stem. This ensured that the ant contacted the droplet as it walked towards the bifurcation. This droplet was either flavoured similarly to the scent on the newly presented runway (treatments E, I, J) or was unscented (treatments C, D,

G, H). The droplet was large enough for the ant to detect and drink, but not enough to satiate the ant

[49], which proceeded onwards after drinking the droplet. This droplet was designed to simulate

trophallaxis. Trophallaxis is used in the nest to unload food, and also provides ants with information

about the food available in the environment [50–53]. This information is attended to very strongly

[51,54]. *L. niger* ants can often be observed performing trophallaxis away from the nest, and

recipients of these trophallactic interactions usually continue on their outwards journey afterwards

(TJC, personal observation). Unfortunately, we could not reliably achieve such on-trail trophallaxis

with the trained ant under controlled experimental conditions.

We then noted which arm of the Y maze the ant chose. We took two choice measurements:

the initial decision, as defined by the ant crossing a line 2cm from the bifurcation, and the final

decision, as defined by the antennae of the ant reaching the end of the Y maze arm. As the ant

reached the end of the Y maze, it was allowed to walk onto a piece of paper and replaced on the

path leading to the Y maze, before the location of the sugar droplet. This allowed us to make 10

repeated measures of each ant. After 10 such measurements the ant was permanently removed

from the colony. The number of ants tested in each treatment is given in table S7 in the supplement.

Statistical analysis

Since the initial decision (crossing a line 2cm from the bifurcation) and the final decision (reaching

the end of the arm) rarely differed (<5% of cases), we analysed only the final decision. While we

collected 10 data points per ant to increase the potential power of our experiments, we finally

decided to perform a conservative analysis by only considering the first decision made by the ants.

This avoids the possibility of ants becoming frustrated in later trials. Initial data exploration revealed

a leftward side bias, which is not uncommon in the behaviour of ants or other animals [55], and so

training side was also added to the model. Finally, colony identity was added as a random effect. This

resulted in the following model:

Decision ~ treatment.group + training.side

223   + (colony as a random effect)

As the data is binomial we used a binomial error structure. Data were analysed in R v.3.4.1 [56] using

generalised linear mixed models from the LME4 package [57]. Model fit was checked using the

DHARMa package [58].

We also tested whether the decisions of each group differed significantly from random using two

tailed exact binomial tests.

Following Benjamin et al. [59] we set significance level for hypothesis testing at p = 0.005, terming p-
values greater than that but smaller than 0.05 'suggestive'. However, due to the limitations of using
p-values alone, we provide estimates and confidence intervals for all effects.

Results

The type of information available to a forager strongly influenced her decision of whether or
not to follow her memory. For ants trained to 0.25M sucrose, given just an unmarked novel path on
the Y maze, most ants (75%) choose to follow their own memory (figure 2), significantly more than
chance (two tailed binomial exact test, 46/61 followed their memory, mean = 0.75, 95% C.I. range =
0.63 – 0.86, p < 0.0001). The addition of a small 1.5M sucrose droplet does not significantly change
the choices made by the ants, who still preferentially follow their memory (binomial test, 60/87,
mean = 0.69, 95% C.I. = 0.58 – 0.78, P = 0.0005). The addition of a pheromone trail on the novel arm
still resulted in most ants (60%) following their memory, but this result was not different from
random choice (binomial test, 18/30, mean = 0.6, 95% C.I. = 0.41 – 0.77, p = 0.36). These three
groups do not differ significantly from each other (GLMM, z ≤ 1.90, p ≥ 0.057, for details of all
pairwise comparisons, including estimates for each comparison, see table S1 in supplement S1).

However, a striking synergistic effect of presenting a pheromone trail and a sucrose droplet
was found. If both were presented, only 25% of ants chose to follow their memory, with 75%
choosing to follow the novel path. This is significantly fewer than chance (binomial test, 23/92, mean
= 0.25, 95% C.I. = 0.17 – 0.35, p < 0.0001) and differs significantly from all other groups provided with
less information (GLMM, Z ≥ 2.97, P ≤ 0.0029).

The effect of presenting both information sources depends on the relative quality of the
droplet presented and the food quality the ant was trained on. If ants were trained on 1.5M sucrose,
53% of ants chose to follow their memory, which does not differ from random choice (binomial test,
16/30, mean = 0.53, 95% C.I. = 0.34-0.72, p = 0.86). This is a non-significantly, but suggestively,
greater than for the ants receiving the same information, but trained on poorer food (GLMM, z =
0.23, p = 0.020), and also suggestively less than the ants receiving no information (GLMM, z = -2.56, p
= 0.011). It is not significantly different from the ants receiving either one information source (GLMM
vs only droplet, z = 1.95, p = 0.051, vs only pheromone z = 0.60, p = 0.55).

The entire dataset on which these results are based in provided in online supplement S2.

Figure 2 – proportion of ants following their memory down the original Y-maze arm as a

function of what other information was available to them. Ants may have had either no

information about the new Y-maze arm, a pheromone trail on the new arm, a small 1.5M

sucrose droplet before the bifurcation, or a pheromone trail and a droplet. In most groups ants

were trained to 0.25M sucrose, but in one group they were trained to 1.5M sucrose. Whiskers

are 95% C.I. for the mean. Letters indicate groups that are significantly (P < 0.005) different

from each other. "***" and "n.s." signify groups that are either significantly different from

0.5 (exact binomial test, P < 0.0001), or not different from 0.5 (P > 0.05), respectively.

Sample sizes per group are show in the bars. Detailed statistical analysis results are presented

in tables S1 and S2 in online supplement S1.

Discussion

By providing unambiguous information about the quality of other resources in the environment the information asymmetry in a communication system was rebalanced. This rescued pheromone following behaviour in our experiment. When ants had no information about alternative food sources in the environment, they predominantly (75%) chose to continue exploiting their private information, returning to the sure but poor quality (0.25M) feeder. When ants were provided with the information that better food exists in the environment (a small 1.5M sucrose droplet), they nonetheless predominantly (69%) returned to the poor quality feeder they knew about. Even when a strong pheromone trail led in the opposite direction, 60% of the ants continued to exploit the known, poor quality feeder. However, when social information (a pheromone trail) and unambiguous quality information (the sugar droplet) was provided, the behaviour of the ants completely reversed, with 75% of choices made for the unknown, but advertised, route.

Information use strategies have been divided into 'when' and 'who' strategies – when should information be used, and who should one listen to [15]. Here, we propose considering information use in terms of a 'what' strategy – what is the information content of an information type, and what is its' level of ambiguity? Social information sources such as pheromone trails and honey bee waggle-dances do not provide unambiguous quality information, while private sensory information does. While pheromone trails and waggle-dances do encode some information about food quality, this information is extremely imprecise and noisy, both due to individual variation in responses to fixed food quality [43,60] and due to the effect of time and foraging patterns on these information sources (see introduction for details). We propose that is it this ambiguity in social information, or the certainty of the private information, which causes otherwise valuable social information to be neglected. Our data supports this claim.

Understanding information use strategies in terms of information asymmetry can explain previously puzzling findings. Leadbeater and Florent [7] unexpectedly found that bumblebees do not rate social information above personal experience, even when their personal experience becomes outdated. Bumblebees which had experienced a novel odour in the nest (social information) were just as unlikely to sample novel flowers scented with this odour as bees which did not experience the novel odour in the nest. Both groups continued choosing known-odour flowers, even when those flowers became unproductive. However, once the bees with the social information did, eventually, sample the novel flowers, they were much more likely to switch to the novel flowers than the group without social information. This is directly analogous to the situation we report: neither private sensory information providing reliable quality information (sampling the novel flowers), nor social information (smelling the novel odour in the nest) is sufficient to make bees change their choices.

However, when both are provided together (here, after the first sampling of the novel flower), bees
are willing to switch.

Similarly, understanding information use in terms of information quality can explain situations in
which social information does override personal information. This is because in such cases, the social
information is actually based on personal, sensory information, which provides unambiguous quality
information. For example, Dunlap et al. [14] and Smolla et al. [61] find that social information alone
can drive decision making during flower choice in bumblebees, and that social information even
outweighs private information when the two compete [14]. This result could be understood through
the psychological concept of blocking [20]. However, differences in information content between the
two information sources can also explain these results: in these studies the social information is
actually private information, based on making associations between a social cue (the presence of a
conspecific) and the direct personal experience of the forager with the flower [20]. There is thus no
quality ambiguity. Indeed, if ambiguity exists, it is lower in the social information: the forager not
only has her own experience of finding the flower acceptable, but also the presence of a conspecific
suggests that the conspecific also finds the flower acceptable, resulting in more weight of
information about quality for the socially advertised flower. Likewise, Josens et al. [54] describe how
trophallactic interactions with a forager providing scented, untainted food can cause foragers to
accept similarly scented food tainted with poison, which they would otherwise avoid. Here again, the
social information component is based in direct sensory experience of the food (trophallaxis), and
the reason the ants overweigh this information over their own later sensory information are similar
to those described for the bumblebee examples above.

Also supportive of the information content hypothesis is the response of ants whose receive
unambiguous information that alternative food qualities are *not* better – the ants trained to a high
quality 1.5M feeder in our experiment. Both paths (apparently) led to equally good food sources, so
there was no particular reason to prefer either, and indeed ants chose both feeders equally often.

Finding drops of liquid honeydew lying around on a trail is not a common situation in nature.
However, we argue that our results are analogous to an ant receiving a trophallactic interaction,
either on the trail or in the nest. Returning ants often perform trophallaxis with multiple partners on
returning to the nest, and these partners are then themselves likely to go out to forage. Similarly,
dancing honeybees often engage in trophallaxis with followers. Ants receiving trophallaxis may even
respond as if they have themselves successfully foraged, depositing trail pheromone when otherwise
only successful ants do so [62]. Ants can also occasionally be seen to perform trophallactic
interactions on the trail, with receivers continuing on their outwards journey afterwards (TC, pers.
obs.). Information gained during trophallactic interactions seems to be particularly well attended to –

even more so than information gained while foraging [51,54]. This information, gained in the nest or

on the trail, can then drive information use. Indeed, provision of unambiguous quality information,

may well be a major role of most trophallactic interactions, especially those occurring on a trail.

Provision of information about food associated cues such, as the odour of the shared food [50,54], is

also no doubt important, although we could not find any role for food odour in the current

experiment (see supplement S1).

Trophallactic networks within a nest are complex [63]. Social insect colonies contain a high

proportion of apparently taskless individuals ([64] and references therein), which can be divided into

inactives and interactives [65]. Interactives have a more central place in interaction networks within

the colony [64,65]. If they also receive trophallactic interactions from many foragers and food

receivers, they may act to homogenise the various food qualities being returned to the nest,

providing a reading of the average food being brought into the colony. Foragers interacting with

these ants may then compare this reading to their current exploited food source. If they find their

own food source to be worse, they may begin to follow social information. It has already been shown

that ants which receive high quality food via trophallaxis are less willing to accept medium quality

food, and vice versa for ants receiving poor quality food [Wendt & Czaczkes, in prep]. It is also known

that ants which have experienced higher quality food source avoid exploiting poorer but otherwise

acceptable food source discovered en-route [66]. Using newly developed imaging technologies

allows the flow of food via trophallaxis in a colony to be tracked and quantified [67], opening the

possibility to empirically test these suggestions. Such an information use pattern would correspond

to the strategy "copy if better", a strategy normally ignored, as direct quality comparison is not

usually considered possible.

Finally, we note that while our results are consistent with ambiguity aversion or the Certainty

Effect playing a role in insect behaviour, they do not constitute a direct test of either of these related,

but distinct, effects. The memory of the ants can be considered certain, while the pheromone trails

ambiguous. Thus, the ants may be following their memory due to its certainty, or avoiding the

pheromone trail due to its ambiguity. Moreover, our different treatments likely affected the

subjective probabilities of receiving a particular reward each ant assigned to each path. Only by

repeatedly testing the same ant in both a risky and an ambiguous situation could one demonstrate

that the behaviour of the ants truly deviates from standard economic behaviour, as described by

expected utility theory. It is intriguing that the results of this study are in line with the 'Certainty

Effect' from Prospect theory, while ants also perceive value relative to a reference point – another

key aspect of Prospect theory [Wendt & Czaczkes, in prep].

Here, we explored how three different information sources, one private and two public, interacted. In doing so, we found evidence that the information content of an information source is a key driver in whether it will be heeded or not. Uncertainty avoidance or the Certainty Effect [38,40] could be the driver behind the repeated findings that social insects often preferentially follow their own memory over the communications of their nestmates. We propose that as well as considering 'when' and 'who' strategies for information use, 'what' is an important question to ask: what type of information can be provided. Social insects, and animals in general, demonstrably use complex mechanisms to decide which information to attend to, and integrate information from a wide variety of sources in order to do so.

Acknowledgements

Thanks to Wolfhard von Thienen for advice on producing an artificial pheromone trail of a realistic strength, to Ellouise Leadbeater and Christoph Grüter for comments on previous versions of this manuscript, and to Andreas Roider and Nathan Carroll for comments on the economic background of this work.

References

1. Danchin E, Giraldeau LA, Valone TJ, Wagner RH. 2004 Public information: from nosy neighbors to cultural evolution. *Science* **305**, 487.

2. von Frisch K. 1967 *The dance language and orientation of bees*. Harvard University Press.

3. Reinhard J, Srinivasan MV, Guez D, Zhang SW. 2004 Floral scents induce recall of navigational and visual memories in honeybees. *J. Exp. Biol.* **207**, 4371–4381. (doi:10.1242/jeb.01306)

4. Dyer FC, Could JL. 1983 Honey Bee Navigation: The honey bee's ability to find its way depends on a hierarchy of sophisticated orientation mechanisms. *Am. Sci.* **71**, 587–597.

5. Czaczkes TJ, Grüter C, Jones SM, Ratnieks FLW. 2011 Synergy between social and private information increases foraging efficiency in ants. *Biol. Lett.* **7**, 521–524. (doi:10.1098/rsbl.2011.0067)

6. von Thienen W, Metzler D, Witte V. 2016 How memory and motivation modulate the responses to trail pheromones in three ant species. *Behav. Ecol. Sociobiol.* **70**, 393–407. (doi:10.1007/s00265-016-2059-5)

7. Leadbeater E, Florent C. 2014 Foraging bumblebees do not rate social information above personal experience. *Behav. Ecol. Sociobiol.* **68**, 1145–1150. (doi:10.1007/s00265-014-1725-8)

8.   Wystrach A, Mangan M, Webb B. 2015 Optimal cue integration in ants. *Proc R Soc B* **282**,
20151484. (doi:10.1098/rspb.2015.1484)

9.   Wehner R, Hoinville T, Cruse H, Cheng K. 2016 Steering intermediate courses: desert ants
combine information from various navigational routines. *J. Comp. Physiol. A* **202**, 459–472.
(doi:10.1007/s00359-016-1094-z)

10.   Rendell L *et al.* 2010 Why copy others? Insights from the social learning strategies tournament.
Science **328**, 208–213. (doi:10.1126/science.1184719)

11.   Grüter C, Leadbeater E. 2014 Insights from insects about adaptive social information use. *Trends*
Ecol. Evol. **29**, 177–184.

12.   Dall SRX, Giraldeau L-A, Olsson O, McNamara JM, Stephens DW. 2005 Information and its use by
animals in evolutionary ecology. *Trends Ecol. Evol.* **20**, 187–193. (doi:10.1016/j.tree.2005.01.010)

13.   Kendal RL, Coolen I, van Bergen Y, Laland KN. 2005 Trade-Offs in the Adaptive Use of Social and
Asocial Learning. *Adv. Study Behav.* **Volume 35**, 333–379.

14.   Dunlap AS, Nielsen ME, Dornhaus A, Papaj DR. 2016 Foraging Bumble Bees Weigh the Reliability
of Personal and Social Information. *Curr. Biol.* **26**, 1195–1199. (doi:10.1016/j.cub.2016.03.009)

15.   Laland KN. 2004 Social learning strategies. *Learn. Behav.* **32**, 4–14. (doi:VL - 32)

16.   Coolen I, Dangles O, Casas J. 2005 Social Learning in Noncolonial Insects? *Curr. Biol.* **15**, 1931–
1935. (doi:10.1016/j.cub.2005.09.015)

17.   Coolen I, Bergen YV, Day RL, Laland KN. 2003 Species difference in adaptive use of public
information in sticklebacks. *Proc. R. Soc. Lond. B Biol. Sci.* **270**, 2413–2419.
(doi:10.1098/rspb.2003.2525)

18.   Cote J, Clobert J. 2007 Social information and emigration: lessons from immigrants. *Ecol. Lett.* **10**,
411–417.

19.   Morgan TJH, Rendell LE, Ehn M, Hoppitt W, Laland KN. 2012 The evolutionary basis of human
social learning. *Proc. R. Soc. Lond. B Biol. Sci.* **279**, 653–662. (doi:10.1098/rspb.2011.1172)

20.   Leadbeater E, Dawson EH. 2017 A social insect perspective on the evolution of social learning
mechanisms. *Proc. Natl. Acad. Sci.* **114**, 7838–7845. (doi:10.1073/pnas.1620744114)

21.   Harrison JF, Fewell JH, Stiller TM, Breed MD. 1989 Effects of experience on use of orientation
cues in the giant tropical ant. *Anim. Behav.* **37**, 869–871. (doi:10.1016/0003-3472(89)90076-6)

22.   Rosengren R, Fortelius W. 1986 Ortstreue in foraging ants of the Formica rufa group — Hierarchy
of orienting cues and long-term memory. *Insectes Sociaux* **33**, 306–337.
(doi:10.1007/BF02224248)

23.   Aron S, Beckers R, Deneubourg J, Pasteels JM. 1993 Memory and chemical communication the
orientation of two mass-recruiting ant species. *Insectes Sociaux* **40**, 369–380.
(doi:10.1007/BF01253900)

24.   Grüter C, Czaczkes TJ, Ratnieks FLW. 2011 Decision making in ant foragers (Lasius niger) facing
conflicting private and social information. *Behav. Ecol. Sociobiol.* **64**, 141–148.
(doi:10.1007/s00265-010-1020-2)

25. Grüter C, Balbuena MS, Farina WM. 2008 Informational conflicts created by the waggle dance.
Proc. R. Soc. B Biol. Sci. **275**, 1321–1327. (doi:10.1098/rspb.2008.0186)

26. Quinet Y, Pasteels JM. 1996 Spatial specialization of the foragers and foraging strategy in Lasius
fuliginosus (Latreille) (Hymenoptera, Formicidae). *Insectes Sociaux* **43**, 333–346.
(doi:10.1007/BF01258407)

27. Traniello JFA. 1989 Chemical trail systems, orientation, and territorial interactions in the ant
Lasius neoniger. *J. Insect Behav.* **2**, 339–354. (doi:10.1007/BF01068060)

28. Fourcassie V, Beugnon G. 1988 How do red wood ants orient when foraging in a three
dimensional system? I. Laboratory experiments. *Insectes Sociaux* **35**, 92–105.
(doi:10.1007/BF02224141)

29. Stroeymeyt N, Franks NR, Giurfa M. 2011 Knowledgeable individuals lead collective decisions in
ants. *J. Exp. Biol.* **214**, 3046–3054. (doi:10.1242/jeb.059188)

30. Vilela EF, Jaffé K, Howse PE. 1987 Orientation in leaf-cutting ants (Formicidae: Attini). *Anim.*
Behav. **35**, 1443–1453. (doi:10.1016/S0003-3472(87)80017-9)

31. van Bergen Y, Coolen I, Laland KN. 2004 Nine-spined sticklebacks exploit the most reliable source
when public and private information conflict. *Proc. R. Soc. B Biol. Sci.* **271**, 957.

32. Wray MK, Klein BA, Seeley TD. 2012 Honey bees use social information in waggle dances more
fully when foraging errors are more costly. *Behav. Ecol.* **23**, 125–131.
(doi:10.1093/beheco/arr165)

33. Grüter C, Segers FH, Ratnieks FL. 2013 Social learning strategies in honeybee foragers: do the
costs of using private information affect the use of social information? *Anim. Behav.* **85**, 1443–
1449.

34. Webster MM, Laland KN. 2008 Social learning strategies and predation risk: minnows copy only
when using private information would be costly. *Proc. R. Soc. Lond. B Biol. Sci.* **275**, 2869–2876.

35. Knight FH. 1921 *Risk, Uncertainty and Profit*. New York: Houghton Mifflin Company.

36. Keynes JM. 1921 *A Treatise on Probability*. London: Macmillan.

37. Trimmer PC, Houston AI, Marshall JAR, Mendl MT, Paul ES, McNamara JM. 2011 Decision-making
under uncertainty: biases and Bayesians. *Anim. Cogn.* **14**, 465–476. (doi:10.1007/s10071-011-
0387-4)

38. Ellsberg D. 1961 Risk, Ambiguity, and the Savage Axioms. *Q. J. Econ.* **75**, 643–669.
(doi:10.2307/1884324)

39. Halevy Y. 2007 Ellsberg Revisited: An Experimental Study. *Econometrica* **75**, 503–536.
(doi:10.2307/4501998)

40. Kahneman D, Tversky A. 1979 Prospect Theory: An Analysis of Decision under Risk. *Econometrica*
47, 263. (doi:10.2307/1914185)

41. Hayden BY, Heilbronner SR, Platt ML. 2010 Ambiguity Aversion in Rhesus Macaques. *Front.*
Neurosci. **4**. (doi:10.3389/fnins.2010.00166)

42. Rosati AG, Hare B. 2011 Chimpanzees and bonobos distinguish between risk and ambiguity. *Biol.*
Lett. **7**, 15–18. (doi:10.1098/rsbl.2010.0927)

43. Beckers R, Deneubourg JL, Goss S. 1993 Modulation of trail laying in the ant Lasius niger
(Hymenoptera: Formicidae) and its role in the collective selection of a food source. *J. Insect*
Behav. **6**, 751–759. (doi:10.1007/BF01201674)

44. Verhaeghe J. 1982 Food recruitment in Tetramorium impurum (Hymenoptera: Formicidae).
Insectes Sociaux **29**, 67–85. (doi:10.1007/BF02224528)

45. Mautz D. 1971 Der Kommunikationseffekt der Schwanzeltanze bei Apis mellifica carnica (Pollm.).
J. Comp. Physiol. A **72**, 197–220. (doi:10.1007/BF00297822)

46. Czaczkes TJ, Schlosser L, Heinze J, Witte V. 2014 Ants use directionless odour cues to recall
odour-associated locations. *Behav. Ecol. Sociobiol.* **68**, 981–988. (doi:10.1007/s00265-014-1710-
2)

47. von Thienen W, Metzler D, Choe D-H, Witte V. 2014 Pheromone communication in ants: a
detailed analysis of concentration-dependent decisions in three species. *Behav. Ecol. Sociobiol.*
68, 1611–1627. (doi:10.1007/s00265-014-1770-3)

48. Czaczkes TJ, Castorena M, Schürch R, Heinze J. 2017 Pheromone trail following in the ant Lasius
niger: high accuracy and variability but no effect of task state. *Physiol. Entomol.* **42**, 91–97.
(doi:10.1111/phen.12174)

49. Mailleux A-C, Deneubourg J-L, Detrain C. 2000 How do ants assess food volume? *Anim. Behav.*
59, 1061–1069. (doi:10.1006/anbe.2000.1396)

50. Farina WM, Grüter C. 2009 Trophallaxis: a mechanism of information transfer. In *Food*
Exploitation by Social Insects: Ecological, Behavioral, and Theoretical Approaches, pp. 173–187.
Boca Raton, Florida.

51. Provecho Y, Josens R. 2009 Olfactory memory established during trophallaxis affects food search
behaviour in ants. *J. Exp. Biol.* **212**, 3221–3227. (doi:10.1242/jeb.033506)

52. Gil M, De Marco RJ. 2005 Olfactory learning by means of trophallaxis in Apis mellifera. *J. Exp.*
Biol. **208**, 671–680. (doi:10.1242/jeb.01474)

53. Hart AG, Ratnieks FLW. 2001 Why do honey-bee (Apis millifera) foragers transfer nectar to
several receivers? Information improvement through multiple sampling in a biological system.
Behav. Ecol. Sociobiol. **49**, 244–250. (doi:10.1007/s002650000306)

54. Josens R, Mattiacci A, Lois-Milevicich J, Giacometti A. 2016 Food information acquired socially
overrides individual food assessment in ants. *Behav. Ecol. Sociobiol.* , 1–12. (doi:10.1007/s00265-
016-2216-x)

55. Hunt ER, O'Shea-Wheller T, Albery GF, Bridger TH, Gumn M, Franks NR. 2014 Ants show a
leftward turning bias when exploring unknown nest sites. *Biol. Lett.* **10**, 20140945.
(doi:10.1098/rsbl.2014.0945)

56. R Core Team. 2012 *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
Foundation for Statistical Computing. See http://www.R-project.org.

57. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4. *J.*
Stat. Softw. **67**. (doi:10.18637/jss.v067.i01)

58. Hartig F. 2016 DHARMa - Residual Diagnostics for HierArchical (Multi-level / Mixed) Regression
Models. R package Version 0.1.3. See https://CRAN.R-project.org/package=DHARMa (accessed
on 18 December 2016).

59. Benjamin DJ, Burger J, Johannesson M, et al. 2017 Redefine statistical significance. *PsyArXiv*
(doi:10.17605/OSF.IO/MKY9J)

60. Seeley TD, Mikheyev AS, Pagano GJ. 2000 Dancing bees tune both duration and rate of waggle-
run production in relation to nectar-source profitability. *J. Comp. Physiol. A* **186**, 813–819.
(doi:10.1007/s003590000134)

61. Smolla M, Alem S, Chittka L, Shultz S. 2016 Copy-when-uncertain: bumblebees rely on social
information when rewards are highly variable. *Biol. Lett.* **12**, 20160188.
(doi:10.1098/rsbl.2016.0188)

62. Mailleux A-C, Buffin A, Detrain C, Deneubourg J-L. 2011 Recruitment in starved nests: the role of
direct and indirect interactions between scouts and nestmates in the ant Lasius niger. *Insectes*
Sociaux **58**, 559–567. (doi:10.1007/s00040-011-0177-7)

63. Sendova-Franks AB, Hayward RK, Wulf B, Klimek T, James R, Planque R, Britton NF, Franks NR.
2010 Emergency networking: famine relief in ant colonies. *Anim. Behav.* **79**, 473–485.

64. Charbonneau D, Poff C, Nguyen H, Shin MC, Kierstead K, Dornhaus A. 2017 Who Are the "Lazy"
Ants? The Function of Inactivity in Social Insects and a Possible Role of Constraint: Inactive Ants
Are Corpulent and May Be Young and/or Selfish. *Integr. Comp. Biol.* **icx029**.
(doi:10.1093/icb/icx029)

65. Charbonneau D, Dornhaus A. 2015 Workers 'specialized' on inactivity: Behavioral consistency of
inactive workers and their role in task allocation. *Behav. Ecol. Sociobiol.* **69**, 1459–1472.
(doi:10.1007/s00265-015-1958-1)

66. Wendt S, Czaczkes TJ. 2017 Individual ant workers show self-control. *Biol. Lett.* , 20170450.

67. Greenwald E, Segre E, Feinerman O. 2015 Ant trophallactic networks: simultaneous
measurement of interaction patterns and food dissemination. *Sci. Rep.* **5**.
(doi:10.1038/srep12496)

549