ABCF ATPases involved in protein synthesis, ribosome assembly and antibiotic

resistance: structural and functional diversification across the tree of life

Victoriia Murina*[1,2], Marje Kasari*[1], Hiraku Takada[1,2], Mariliis Hinnu[3], Chayan Kumar Saha[1], James

W. Grimshaw[4], Takahiro Seki[5], Michael Reith[1], Marta Putrinš[3], Tanel Tenson[3], Henrik Strahl[4], Vasili

Hauryliuk[1,2,3] and Gemma Catherine Atkinson[†1]

[1]Department of Molecular Biology, Umeå University, 901 87, Umeå, Sweden

[2]Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, 901 87, Umeå,

Sweden

[3]University of Tartu, Institute of Technology, Nooruse 1, 50411 Tartu, Estonia

[4]Centre for Bacterial Cell Biology, Institute for Cell and Molecular Biosciences Newcastle

University, Richardson Road, Newcastle upon Tyne, NE2 4AX, United Kingdom

[5]Department of Applied Chemistry and Biotechnology, Faculty of Engineering, Chiba University,

263-8522, Chiba, Japan

* These authors contributed equally to this work

[†]To whom correspondence should be addressed

<u>Abstract</u>

Within the larger ABC superfamily of ATPases, ABCF family members eEF3 in *Saccharomyces*

cerevisiae and EttA in *Escherichia coli* have been found to function as ribosomal translation factors.

Several other ABCFs including biochemically characterised VgaA, LsaA and MsrE confer resistance

to antibiotics that target the peptidyl transferase centre and exit tunnel of the ribosome. However,

the diversity of ABCF subfamilies, the relationships among subfamilies and the evolution of

antibiotic resistance factors from other ABCFs have not been explored. To address this, we

analysed the presence of ABCFs and their domain architectures in 4505 genomes across the tree of

life. We find 45 distinct subfamilies of ABCFs that are widespread across bacterial and eukaryotic

phyla, suggesting they were present in the last common ancestor of both. Surprisingly, currently

known antibiotic resistance (ARE) ABCFs are not confined to a distinct lineage of the ABCF family

tree. This suggests that either antibiotic resistance is a pervasive feature of bacterial ABCFs, or it is relatively easy to evolve antibiotic resistance from other ABCF functions. Our data suggest there are a number of previously unidentified ARE ABCFs in antibiotic producers and important human pathogens. We also find that ATPase-deficient mutants of all four *E. coli* ABCFs (EttA, YbiT, YheS and Uup) inhibit protein synthesis, indicative of their ribosomal function, and demonstrate a genetic interaction of ABCFs Uup and YheS with translational GTPase BipA involved in assembly of the 50S ribosome subunit. Finally, we show that *Bacillus subtilis* VmlR is a ribosome-binding resistance factor localised to the cytoplasm.

Author summary

Isolated members of the ABCF protein family of ATP-hydrolysing enzymes have been found to have important roles in protein synthesis and antibiotic resistance. However, their full diversity across the tree of life, and their evolutionary histories have never been examined. Therefore, we analysed the presence of ABCFs and their constituent domains in genomes across the tree of life, discovering 45 distinct subfamilies of ABCFs that are widespread across bacterial and eukaryotic phyla. This includes several subfamilies that we predict comprise novel antibiotic resistance (ARE) ABCFs, present in antibiotic producers and important human pathogens. There are significant gaps in our knowledge about the functional capabilities of different ABCF families. To address this, we have made ATPase domain mutants of all four *Escherichia coli* ABCFs, showing that they inhibit protein synthesis and indicating a role on the ribosome. Furthermore, we demonstrate a genetic interaction of two *E. coli* ABCFs with the GTPase BipA, involved in ribosome assembly. Finally, we show that *Bacillus subtilis* VmlR in the ARE2 subfamily is a ribosome-binding resistance factor localised to the cytoplasm. As more is discovered about the function of individual ABCFs, the more it will be possible to predict functions of uncharacterised members, using the ABCF family tree as a framework.

Introduction

Protein biosynthesis – translation – is the reading and deciphering of information coded in genes to produce proteins. It is one of the most ancient and central cellular processes, and control of the various stages of translation is achieved via an intricate interplay of multiple molecular interactions. For many years, enzymatic control of the ribosomal cycle was thought to be mainly orchestrated mainly by translational GTPases (trGTPases). That view of translation has been nuanced by the identification of multiple ATPases in the ABC superfamily that have important roles in translational regulation on the ribosome. The ABC protein eEF3 (eukaryotic Elongation Factor 3) is an essential factor for polypeptide elongation in *Saccharomyces cerevisiae* [1] with proposed roles in E-site tRNA release and ribosome recycling [2-4]. This fungi-specific translational ABC ATPase appeared to be an exception to the tenet that trGTPases are the enzymatic rulers of the ribosome, until ABCE1 (also known as Rli1), a highly conserved protein in eukaryotes and archaea, was identified as another ribosome recycling factor [5-7].

The ABC ATPases together comprise one of the most ancient superfamilies of proteins, evolving well before the last common ancestor of life [8]. The superfamily contains members with wide varieties of functions, but is best known for its membrane transporters [9]. Families of proteins within the ABC superfamily are named alphabetically ABCA to ABCH, following the nomenclature of the human proteins [10]. While most ABCs carry membrane-spanning domains (MSDs), these are lacking in ABCE and ABCF families [11]. ABCF proteins of eukaryotes include eEF3, and also other ribosome-associated proteins: Gcn20 is involved in sensing starvation by the presence of uncharged tRNAs on the eukaryotic ribosome [12]; ABC50 (ABCF1) promotes translation initiation in eukaryotes [13]; and both Arb1 (ABCF2) and New1 have been proposed to be involved in biogenesis of the eukaryotic ribosome [14, 15]. Ribosome-binding by ABCF proteins seemed to be limited to eukaryotes until characterisation of ABCF member EttA (energy-dependent translational throttle A) found in diverse bacteria. *Escherichia coli* EttA binds to the E-site of the ribosome where it is proposed to 'throttle' the elongation stage of translation in response to change in the

intercellular ATP/ADP ratio [16, 17]. EttA is one of four *E. coli* ABCFs, the others being YheS, Uup

and YbiT, none of which have yet been shown to operate on the ribosome [16].

Bacterial ABCF family members have been found to confer resistance to ribosome-inhibiting

antibiotics widely used in clinical practise, such as ketolides [18], lincosamides [19-22], macrolides

[23, 24], oxazolidinones [25], phenicols [25], pleuromutilins [22] and streptogramins A [22, 26]

and B [24]. These <u>a</u>ntibiotic <u>re</u>sistance (AREs) ABCFs have been identified in antibiotic-resistant

clinical isolates of *Staphylococcus*, *Streptomyces* and *Enterococcus* among others [27]. This includes

the so-called ESKAPE pathogens *Enterococcus faecium* and *Staphylococcus aureus* that contribute

to a substantial proportion of hospital-acquired multidrug-resistant infections [28]. As some efflux

pumps carry the ABC ATPase domain, it was originally thought that ARE ABCFs similarly confer

resistance by expelling antibiotics. However, as they do not carry the necessary transmembrane

domains, this is unlikely ~~[29, 30]~~. In support of this, it was recently shown that *Staphylococcus*

aureus ARE VgaA protects protein synthesis activity in cell lysates from antibiotic inhibition and

that *Enterococcus faecalis* ARE LsaA displaces radioactive lincomycin from *S. aureus* ribosomes

[31]. Using a reconstituted biochemical system, we have shown that VgaA and LsaA directly

protect the ribosome peptidyl transferase centre (PTC) from antibiotics in an ATP-dependent

manner [32]. Recent cryo-EM structures of AREs *Pseudomonas aeruginosa* MsrE and *Bacillus*

subtilis VmlR on the ribosome show that like EttA, these ABCFs bind to the E-site of the ribosome,

with extended inter-ABC domain linkers protruding into the PTC [33, 34]. The ARE ABCFs

therefore appear to either physically interact with the drug to displace it from the ribosome, or

allosterically induce a change in conformation of the ribosome that ultimately leads to drug drop-

off [35-37].

Here, we carry out an in-depth survey of the diversity of ABCFs across all species with sequenced

genomes. We find 45 groups (15 in eukaryotes and 30 in bacteria), including 7 groups of AREs. So-

called EQ_2 mutations, double glutamic acid to glutamine substitutions in the two ATPase active

sites of EttA lock the enzyme on ribosome in an ATP-bound conformation, inhibiting protein

synthesis and cellular growth [16, 17]. We have tested the effect of equivalent mutations in the other three *E. coli* ABCFs – YbiT, YheS and Uup – as well as the *Bacillus subtilis* ARE VmlR. We establish genetic associations of *E. coli* ABCFs YheS and Uup with the translational GTPase BipA (also known as TypA), and through microscopy and polysome profile analyses, confirm that VmlR does not confer lincomycin resistance through acting as a membrane-bound pump, but via direct interaction with cytoplasmic ribosomes.

Results

ABCFs are widespread among bacteria and eukaryotes

To identify candidate subfamilies of ABCFs and refine the classifications an iterative bioinformatic protocol of sequence searching and phylogenetic analysis was applied. Sequence searching was carried out against a local database of 4505 genomes from across the tree of life. To first get an overview of the breadth of diversity of ABCFs across life, sequence searching began with a local BlastP search against a translated coding sequence database limited by taxonomy to one representative per class, or order if there was no information on class for that species in the NCBI taxonomy database. From phylogenetic analysis of the hits, preliminary groups were identified and extracted to make Hidden Markov Models (HMMs) for further sequence searching and classification. Additional sequences from known ABCF AREs were included in phylogenetic analyses in order to identify groups of ARE-like ABCFs. HMM searching was carried out at the genus level followed by phylogenetic analysis to refine subfamily identification, with final predictions made at the species level. The resulting classification of 16848 homologous sequences comprises 45 subfamilies, 15 in eukaryotes and 30 in bacteria. Phylogenetic analysis of representatives across the diversity of ABCFs shows a roughly bipartite structure, with most eukaryotic sequences being excluded from those of bacteria with strong support (fig. 1). Five eukaryotic groups that fall in the bacterial part of the tree are likely to be endosymbiotic in origin (see the section *Bacteria-like eukaryotic ABCFs*, below).

ABCFs are widespread among bacteria and eukaryotes; there are on average four ABCFs per

bacterial genome, and five per eukaryotic genome. However, there is considerable variation in how

widespread each subfamily is (table 1). The presence of all subfamilies in each genome considered

here is shown in table S1, with the full set of sequence IDs and domain composition recorded in

table S2. Domain coordinates by amino acid position can be found in table S3. In bacteria,

Actinobacteria, and Firmicutes are the phyla with the largest numbers of ABCFs (up to 11 per

genome), due to expansions in ARE and potential novel ARE ABCF subfamilies. Among eukaryotes

plants and algae encode the most subfamilies, probably due in part to gene acquisition from

endosymbiosis events. The diatom *Fragilariopsis cylindrus* has 30 ABCFs and the Haptophyte

Emiliania huxleyi has 26. Bacterial contamination can sometimes inflate the number of genes in

eukaryotic genomes as noted previously for trGTPases [35]. However, as all the *Fragilariopsis* and

Emiliania sequences belong to typically eukaryotic subgroups, they do not appear to be the result

of bacterial contamination. The Tibetan antelope *Pantholops hodgsonii*, on the other hand has 25

ABCFs, 20 of which belong to bacterial subgroups. The genome is known to be contaminated by

Bradyrhizobium sequences [38], and thus the bacteria-like hits from *P. hodgsonii* are most likely

artifacts.

Table 1 | The subfamilies of the ABCF family, and the numbers (N) of phyla and species in which they are encoded

Subfamily	N Phyla	N Species	Notes on function, relationships and taxonomic distribution
YdiF	42	1852	Broad distribution in bacteria; polyphyletic
Uup [a]	24	3104	Broad distribution in bacteria; paraphyletic to EttA. Inc. P resistance TaeA [39]
EttA	18	2337	Broad distribution in bacteria; translation factor
YbiT	15	1874	Broad distribution in bacteria; potential translation factor
BAF2	7	305	Proteobacteria, Planctomycetes, Spirochaetes, Actinobacteria, Bacteroidetes, Gemmatimonadetes, Cyanobacteria
ARE1 [a]	6	269	M,L,S,P,K resistance, inc. VgaA [21], MsrA [18], MsrE [33]; Firmicutes, Actinobacteria, Spirochaetes, Bacteroidetes, Proteobacteria, Tenericutes
ARE3 [a]	5	261	L resistance inc LsaA [20]; Firmicutes, Spirochaetes, Proteobacteria, Fusobacteria, Actinobacteria
DAF1	5	58	Spirochaetes, Proteobacteria, Deferribacteres, Fibrobacteres, Chlamydiae
YfmM	5	587	Firmicutes, Tenericutes, Proteobacteria, Fusobacteria, Bacteroidetes
YheS	5	1234	Proteobacteria, Bacteroidetes, Cyanobacteria, Arthropoda, Elusimicrobia
BAF3	4	111	Bacteroidetes, Proteobacteria, Cyanobacteria, Elusimicrobia
DAF2	4	24	Proteobacteria, Planctomycetes, Spirochaetes, Omnitrophica
ARE2 [a]	2	54	Antibiotic resistance inc. VmlR[19]; Firmicutes, Tenericutes
ARE4 [a]	2	173	M,M16 resistance inc. CarA [40], srmB [40], tlrC [40]; Actinobacteria, Chloroflexi
ARE5 [a]	2	408	L,S resistance inc. VarM [26], LmrC [41]; Actinobacteria, Proteobacteria
BAF1	2	20	Firmicutes, Actinobacteria
PAF1	1	138	Proteobacteria
AAF1	1	313	Actinobacteria
AAF2	1	301	Actinobacteria
AAF4	1	219	Actinobacteria
AAF6	1	649	Actinobacteria
AAF3	1	8	Actinobacteria
AAF5	1	25	Actinobacteria
ARE6 [a]	1	8	L,S resistance SalA [22]; Firmicutes
ARE7 [a]	1	35	Oxazolidinone resistance OptrA [25], Firmicutes
BdAF1	1	176	Bacteroidetes
DAF3	1	36	Proteobacteria
FAF1	1	37	Firmicutes
FAF2	1	42	Firmicutes
SAF1	1	66	Firmicutes
ABCF2	27	560	Arb1 ribosome biogenesis factor; broadly distributed but lacking in Apicomplexa and Microsporidia
ABCF1	23	376	ABC50 translation initiation factor; found in plants, diverse algae, and opisthokonts excluding fungi
ABCF7	16	131	Found in plants, diverse algae, Alveolata, Excavata and Microsporidia
ABCF3	13	382	Gcn20 starvation response; Opisthokonts
eEF3L	9	83	Diverse algae, Chytridiomycota and choanoflagellates
ABCF4	7	37	Diverse algae and Filozoa
ABCF5	7	105	Diverse algae and fungi
cpYdiF	7	85	Diverse algae
algAF1	6	25	Diverse algae
cpEttA	6	18	Chloroplast targeting peptides predicted; plants and diverse algae
algUup	5	17	Found in diverse algae
ABCF6	5	17	Found in diverse algae and Amoebozoa
mEttA	3	14	mitochondrial targeting peptides predicted; diverse algae, Amoebozoa
New1	3	127	Fungi
eEF3	2	138	Translation factor; fungi

Notes - [a] Subfamilies containing known AREs; resistance to antibiotic class in notes column is as follows: M, 14- and 15-membered ring macrolides; M16, 16-membered ring macrolides; L, lincosamides; S, streptogramins; K, ketolides; P, pleuromutilins

154
155

156
157

The wide distribution and multi-copy nature of ABCFs suggests an importance of these proteins.

However they are not completely universal, and are absent in almost all Archaea. The

euryarcheaotes *Candidatus Methanomassiliicoccus intestinalis*, *Methanomethylophilus alvus*,

Methanomassiliicoccus luminyensis, and *Thermoplasmatales archaeon BRNA1* are the only archaea

found to encode ABCFs, in each case YdiF. ABCFs are also lacking in 214 bacterial species from

various phyla, including many endosymbionts. However, the only phylum that is totally lacking

ABCFs is Aquificae. ABCFs are almost universal in Eukaryotes; the only genomes where they were

not detected were those of Basidomycete *Postia placenta*, Microsporidium *Enterocytozoon bieneusi*,

and apicomplexan genera *Theileria, Babesia* and *Eimeria*.

Many of the phylogenetic relationships among subfamilies of ABCFs are poorly resolved (fig. 1).

This is not surprising, since these represent very deep bacterial relationships that predate the

diversification of major phyla and include gene duplication, and differential diversification and

loss, and likely combine both vertical inheritance and horizontal gene transfer. Although

relationships can not be resolved among all subfamilies, some deep relationships do have strong

support (i.e. maximum likelihood bootstrap percentage (MLB) of more than 85% and Bayesian

inference posterior probability (BIPP) of 1.0); EttA and Uup share a common ancestor to the

exclusion of other ABCFs with full support (100% MLB, 100% UFB and 1.0 BIPP fig. 1, fig. S1) and

YheS is the closest bacterial group to the eukaryotic ABCFs with strong support (94% MLB, 98%

UFB, and 1.0 BIPP; (fig. 1, fig. S1, text S1). This latter observation suggests that eukaryotic-like

ABCFs evolved from within the diversity of bacterial ABCFs. However, this depends on the root of

the ABCF family tree. To address this, phylogenetic analysis was carried out of all ABCFs from *E.*

coli, *Homo sapiens*, *S. cerevisiae* and *B. subtilis*, along with ABCE family sequences from the UniProt

database [42]. Rooting with ABCE does not provide statistical support for a particular group at the

base of the tree, but does support the eukaryotic subgroups being nested within bacteria, with

YheS as the closest bacterial group to the eukaryotic types (fig. 2). It also shows that eEF3 and

New1 are nested within the rest of the ABCF family, thus confirming their identity as ABCFs,

despite their unusual domain structure. To address the possibility that due to recombination the

two ABC domains of the ABCF family may have had different evolutionary histories, we repeated

our phylogenetic analysis of representative sequences with the ABC domains uncoupled and

aligned to each other (fig. 1, Text S1). With this very short alignment (204 positions) containing a

larger proportion of almost invariant active site residues, there is even less statistical support for

relationships among subgroups. Nevertheless, we still retain the branches that are well supported

in our full-length analyses (fig. 1), and thus there is no evidence for recombination.

Domain architectures in the ABCF family are variable

To assess the conservation of domains across the family tree of ABCFs, we extracted the domain

regions from subfamily alignments, made HMMs representing each domain region and scanned

every sequence in our database. We find the most common domain and subdomain structure is an

N-terminal ABC1 nucleotide binding domain (NBD) containing an internal Arm subdomain,

followed by the Linker region joining to the ABC2 NBD (fig. 3A). Variations on this basic structure

include deletions in the Arm and Linker regions, insertion of a Chromo subdomain in the ABC2

NBD, and extension of N and C termini by sequence extensions (fig. 3A). The domain structures of

eukaryotic ABCFs are more diverse than those of bacteria, with greater capacity for extensions of

the N-terminal regions to create new domains (fig. 3A). An increased propensity to evolve

extensions, especially at the N terminus is also seen with eukaryotic members of the trGTPase

family [35]. In bacteria, terminal extensions of ABCFs tend to be at the C terminus (fig. 3A-B).

Cryo-electron microscopy structures show that bacterial ABCFs bind to the same site of the

ribosome, which is different from that of eEF3 [2, 17, 33, 34], and this is reflected in their domain

architectures. eEF3 carries additional N-terminal domains and does not have the Arm subdomain

that in EttA binds the L1 stalk and ribosomal protein L1. Instead it carries a Chromo (Chromatin

Organization Modifier) subdomain in ABC2 that – from a different orientation to the Arm domain

of EttA - interacts with, and stabilises the conformation of the L1 stalk [2]. The Arm subdomain is

also missing in eEF3-like close relatives New1 and eEF3L. Although the Arm is widespread in

bacterial ABCFs, it is not universal; it is greatly reduced in a number of subfamilies, with the most

drastic loss seen in ARE1 and 2 (figs. 3A, and see *Putative AREs* section, below). Subdomain

composition can even vary within subfamilies; Uup has lost its arm independently in multiple

lineages (fig. 1, fig. S1).

Arms, linkers and CTD extensions are poorly conserved at the primary sequence level, but are

similar in terms of composition, all being rich in charged amino acids, particularly arginine and

lysine. Their variable presence and length suggests they can be readily extended or reduced during

evolution. The CTD of Uup forms a coiled coil structure that is capable of binding DNA [43].

However, whether DNA binding is its primary function is unclear. The CTD of YheS has significant

sequence similarity (E value 3.58e-03) to the tRNA binding CTD of Valine-tRNA synthetase (NCBI

conserved domains database accession cl11104). *In silico* coiled coil prediction suggests there is a

propensity of all these regions to form coiled coil structures (fig. 3B). Extensions and truncations of

the arms, linkers and CTD extensions possibly modulate the length of coiled coil protrusions that

extend from the globular mass of the protein.

Eukaryotic ABCFs comprise 15 subfamilies

eEF3, New1 and eEF3L

eEF3, eEF3L and New1 group together and are particularly distinct members of the ABCF family.

The eEF3 subfamily represents the classical fungal proteins, while eEF3L is a more divergent group

found mainly in protists (see below). eEF3 has a recent paralogue in *S. cerevisiae* (Hef3 (fig. 2),

YEF3B) that apparently arose as a result of the whole genome duplication in yeast [44, 45]. The

conservation of domain structure in eEF3/New1/eEF3L suggests they bind the ribosome similarly

(fig. 3A). Ribosome binding by eEF3 involves the Chromo subdomain, and the HEAT domain [2],

which in addition to New1 and eEF3L, is also found in the protein Gcn1, a binding partner of Gcn20

ABCF [46] (see the section *ABCF1-7*, below). eEF3 has a distinct C-terminal extension (fig. 3A),

through which it interacts with eEF1A [47]. It has been suggested that this leads to the recruitment

of eEF1A to *S. cerevisiae* ribosomes [47, 48]. The New1 CTD contains a region of sequence similarity to the eEF3 CTD; both contain a polylysine/arginine-rich tract of 20-25 amino acids (fig. S2). In *S. cerevisiae* eEF3, this is at positions 1009-1031, which falls within the eEF1A-binding site. Thus eEF1A binding may be a common feature of eEF3, New1, and also eEF3L, which commonly includes an eEF3-like C-terminal extension (table S2).

We find eEF3-like (eEF3L) factors in a range of eukaryotes including choanoflagellates, haptophytes, heterokonts, dinoflagellates, cryptophytes and red and green algae. This suggests the progenitor of eEF3 was an ancient protein within eukaryotes, and has been lost in a number of taxonomic lineages. Alternatively, eEF3L may have been horizontally transferred in eukaryotes. eEF3 is found in *Chlorella* viruses [49] and *Phaeocystis* viruses (fig. S3), suggesting this may be a medium of transfer. There is no "smoking gun" for viral-mediated transfer in the phylogenies, in that eukaryotic eEF3L sequences do not nest within viral sequence clades (fig. S3). However, given the close association of viral and protist eEF3L, this still remains a possibility. Curiously, the taxonomic distribution of eEF3L in diverse and distantly related protists is similar to that of the unusual elongation factor 1 (eEF1A) paralogue EFL [50] (table S4). The propensity for eEF3/eEF3L/New1 to be present in EFL-encoding organisms and absent in eEF1A-encoding organisms is significant at the level of $P<0.0001$ with Fisher's exact test. Like eEF3L, EFL can be found in viruses such as *Aureococcus anophagefferens* virus (NCBI protein accession YP_009052194.1). As eEF3 interacts with eEF1A [48], the equivalents eEF3L and EFL may also interact in the organisms that encode them.

Like eEF3, New1 is found across the fungal tree of life (table S1, fig. S3). *S. cerevisiae* New1 has previously been reported to carry a prion-like Y/N/Q/G repetitive region in the N-terminal region before the HEAT domain [51]. We find that this region is limited in taxonomic distribution to Saccharomycetale yeast (table S2, fig. 3A and fig. S2). Thus it is not found in the N-terminal region of *Schizosaccharomyces pombe* New1 (also known as Elf1).

ABCF1-7

ABCF1-7 comprise the "ancestral-type" eukaryotic ABCFs, in that they have the typical ABC domain

structure that is seen in bacterial ABCFs, and they lack the Chromo and HEAT domains that are

found in eEF3, New1 and eEF3L (fig. 3A). All of the terminal extensions found in ABCF subfamilies

are biased towards charged amino acids, often present as repeated motifs. The ABCF1 (ABC50)

NTD (which interacts with eIF2 [52]) is, due to its length and number of repeats, one of the most

striking (fig. S2). It contains multiple tracts of poly-lysine/arginine, poly–glutamic acid/aspartic

acid and – in animals - poly-glutamine. ABCF1 and ABCF2 have moderate support as sister groups

(fig. 1), and both have representatives in all eukaryotic superphyla, but are not universal. Notable

absences of ABCF1 are fungi, amoebozoa and most Aves (birds) (table 1, table S1).

ABCF2 (Arb1) is essential in yeast and its disruption leads to abnormal ribosome assembly [14],

and human ABCF2 can complement an Arb1 deletion, suggesting conservation of function [53]. The

protein is broadly distributed across eukaryotes, with the notable exception of the *Alveolata*

superphylum (Table S1). Like other ABCF terminal extensions, the ABCF2 N-terminal domain is

rich in lysine, in this case lysine and alanine repeats.

In yeast, where it is known as Gcn20, ABCF3 is a component of the general amino acid control

(GAAC) response to amino acid starvation, acting in a complex with Gcn1 [54]. Gcn20 binds to

Gcn1 via the latter's HEAT-containing N-terminal domain [46], an interaction that is conserved in

ABCF3 and Gcn1 of *Caenorhabditis elegans* [55]. As the HEAT domain is also found in eEF3, this

raises the possibility that Gcn20/ABCF3 and eEF3 interact in encoding organisms (fig. 3A).

Possible support for this comes from the observation that eEF3 overexpression impairs Gcn2

activation [56].

ABCF3 is widespread in eukaryotes, but absent in heterkont algae and archaeplastida. However,

these taxa encode ABCF4 and ABCF7 of unknown function, which have N-terminal domains

homologous to ABCF3. Thus, ABCF4 and ABCF7 may be the functional equivalents of ABCF3 in

these taxa, potentially interacting with HEAT domain-containing eEF3L in organisms that encode

the latter (fig. 3A). ABCF3 from four fungi (*Setosphaeria turcica,* NCBI protein accession number

XP_008030281.1; *Cochliobolus sativus,* XP_007703000.1; *Bipolaris oryzae,* XP_007692076.1; and

Pyrenophora teres, XP_003306113.1) are fused to a protein with sequence similarity to WHI2, an

activator of the yeast general stress response [57].

ABCF5 is a monophyletic group limited to fungi and green algae (*Volvox* and *Chlamydomonas*), with

a specific NTD and CTD (fig. 1 and 2A). ABCF5 is found in a variety of Ascomycete and

Basidiomycete fungi, including the yeast *Debaryomyces hansenii*, but is absent in yeasts *S. pombe*

and *S. cerevisiae*. ABCF4 is a polyphyletic group of various algal and amoeba protists that can not

be assigned to the ABCF5, or eEF3/eEF3L/New1 clades. In eukaryotic ABCF-specific phylogenetic

analysis, ABCF4/ABCF5/eEF3/eEF3L/New1 are separated from all other eukaryotic ABCFs with

moderate support (85% MLB fig. S3). ABCF6 represents a collection of algal and amoebal

sequences that associate with the ABCF1+ABCF2 clade with mixed support in phylogenetic

analysis (fig. 1).

Bacterial-like eukaryotic ABCFs

Five eukaryotic subfamilies are found in the bacteria-like subtree: algAF1, algUup, mEttA, algEttA

and cpYdiF (fig. 1). Given their affiliation with bacterial groups, they may have entered the cell with

an endosymbiotic ancestor. Indeed, chloroplast-targeting peptides are predicted at the N termini of

the majority of cpYdiF sequences, and mitochondrial localization peptides at most mEttA N

termini. The situation is less clear for the three remaining groups, with a mix of signal peptides, or

none at all being predicted across the group members (table S5).

Bacterial ABCFs comprise 30 subfamilies, most of which have unknown function

There are 30 groups of bacterial ABCFs, the most broadly distributed being YdiF (the subfamily is

given the name of the *Bacillus subtilis* protein as it is not present in *E. coli*) (table 1). This subfamily

is a paraphyletic grouping comprising ABCF sequences that can not confidently be classified into

any of the other subgroups (fig. 1). The next most broadly distributed group is Uup (*B. subtilis*

protein name YfmR), which itself is paraphyletic to EttA. *B. subtilis* does not encode EttA, but does

encode YbiT (*B. subtilis* name YkpA). It also encodes two ABCFs not present in *E. coli*: YfmM and

VmlR (also known as ExpZ). VmlR is in the ARE2 subfamily, and confers resistance to

virginiamycin M1 and lincomycin [19]. Insertional disruptants of all the chromosomal ABCF genes

in *B subtilis* strain 168 have been examined for resistance to a panel of nine MLS class antibiotics,

and only VmlR showed any hypersensitivity [19]. With the exception of EttA [16, 17], and the

seven antibiotic resistance ARE ABCFs (table 1), the biological roles of the other 22 bacterial

ABCFs are largely obscure.

<u>B. subtilis *ARE VmlR is a cytoplasmic protein that directly protects the ribosome from antibiotics*</u>

B. subtilis virginiamycin M and lincomycin resistance factor ABCF VmlR was originally annotated as

an ABC efflux transporter, i.e. a membrane protein [19, 58]. To probe VmlR's interaction with the

ribosome we took advantage of ATPase-deficient VmlR mutants generated by simultaneous

mutation of both glutamate residues for glutamine (EQ$_2$) [59] that lock ABC enzymes in an ATP-

bound active conformation [17, 60]. In the case of EttA, expression of the EQ$_2$ mutant results in a

dominant-negative phenotype as EttA incapable of ATP hydrolysis acts as a potent inhibitor of

protein synthesis and, consequently, bacterial growth [16]. We constructed C-terminally tagged

His$_6$-TEV-3xFLAG-tagged (HTF-tagged) wild type and EQ$_2$ (*vmlR*-HTF and *vmlR*EQ$_2$-HTF) under the

control of an IPTG-inducible P$_{hy-spank}$ promotor [61]. To probe the intracellular localization of VmlR,

we C-terminally tagged VmlR with the mNeonGreen fluorescent protein [62] under the control of

xylose inducible promoter protein P$_{xyl}$ [63]. We have validated the functionality of the fusion

constructs by lincomycin resistance assays using a Δ*vmlR* knock-out strain as a negative control

and a Δ*vmlR* knock-out strain expressing untagged VmlR under the control of an IPTG-inducible

P$_{hy-spank}$ promotor as a positive control (fig. S4A-C). While C-terminal tagging with either HTF or

mNeonGreen does not abolish VmlR's activity, the EQ$_2$ versions of the tagged proteins are unable

to protect from lincomycin (fig. S4B).

After establishing the functionality of the tagged VmlR constructs, we tested the effects of expression of either wild type or EQ$_2$ VmlR-HTF on *B. subtilis* growth in rich LB media (fig. 4A). Expression of the wild type protein in the Δ*vmlR* background has no detectable effect. In contrast, the EQ$_2$ version inhibits growth: while exponential growth is unaffected, the cells enter the stationary phase at lower cell densities, abruptly stopping growth instead of slowing down gradually (fig. 4A). Two factors are likely to cause the growth-phase specificity of the inhibitory effect. First, during the exponential growth cells efficiently dilute the toxic protein via cell division, and when the growth slows down, VmlR-EQ$_2$-HTF accumulates. Second, upon entering the early stationary phase, *B. subtilis* sequesters 70S ribosomes into inactive 100S dimers [64], and this decrease of active ribosome concentration could conceivably render the cells more vulnerable to the inhibitory effects of VmlR-EQ$_2$. We probed the interaction of wild type and EQ$_2$ VmlR-HTF with ribosomes using polysome analysis in sucrose gradients in combination with Western blotting (fig. 4B). While the wild type protein barely enters the gradient (most likely dissociating from the ribosomes during centrifugation), the EQ$_2$ version almost exclusively co-localises with the 70S peak fraction and is absent from the polysomal fractions (fig. 4B), suggesting co-sedimentation of a tight 70S:VmlR-EQ$_2$ complex.

Finally, having ascertained the functionality of VmlR-mNeonGreen (fig. S4C), we imaged *B. subtilis* cells expressing VmlR-mNeonGreen in the presence and absence of lincomycin (fig. 4C) and quantified the intensity of the fluorescent signal across the cell (fig. 4D). As a positive control for membrane localization we used WALP23-GFP [65] – an artificial model transmembrane helix WALP23 [66] fused with an N-terminal GFP label. We observe no evidence for association of VmlR with the membrane: the protein is clearly cytoplasmic, with a slight exclusion from the nucleoid in the presence of 5 μg/mL lincomycin. A likely explanation for this effect is the nucleoid compaction caused by inhibition of translation resulting in protein exclusion from the nucleoid-occupied space [67-69]. However, we cannot rule out that this general effect potentiated by specific interaction of VmlR with strongly nucleoid-excluded ribosomes.

<u>E. coli *ABCFs EttA, YbiT, YheS and Uup interact genetically and functionally with protein synthesis*</u>

<u>*and ribosome assembly*</u>

E. coli encodes four ABCFs: EttA, YbiT, YheS and Uup. An array of structural, biochemical and microbiological methods has been used to establish that EttA operates on the ribosome [16, 17]. Ribosomal association of the other *E. coli* ABCFs has not been shown. However, a recent PhD thesis by Dr. Katharyn L. Cochrane suggests that Uup genetically interacts with an enigmatic ribosome-associated factor, the translational GTPase BipA (TypA) [70]. The *E. coli bipA* knock-out strain is characterised by a decreased level of 50S subunits accompanied by an accumulation of pre-50S particles [71]. In the presence of its native substrate, GTP, BipA associates with mature 70S ribosomes [72], occupying the ribosomal A-site [73]. However, in the presence of stress alarmone (p)ppGpp – a molecular mediator of the stringent response [74] – BipA binds the 30S subunit [75].

We have set out to systematically probe the involvement of *E. coli* ABCFs in protein synthesis. We used two experimental systems. The first is geared towards low level constitutive expression of native, untagged wild type and EQ_2 proteins in a clinically relevant uropathogenic *E. coli* strain CFT073 [76]. For this, we cloned ABCF genes into a low copy pSC101 vector under control of a constitutive tet-promoter (P_{tet}) that in the original plasmid drives expression of the tetracycline efflux pump TcR [77]. Using the λRed-mediated gene disruption method [78] we generated a set of mutants of lacking each of the four ABCF genes, as well as a Δ*bipA* and Δ*bipA*Δ*uup* knock-out strains. The second system allows inducible high-level expression of tagged proteins in the avirulent BW25113 *E. coli* strain [79, 80]. We used wild type and EQ_2 mutants of EttA, YbiT, YheS and Uup with N-terminal FLAG-TEV-His$_6$ (FTH)-tags expressed from a low copy pBAD18 plasmid under an arabinose-inducible araBAD (P_{BAD}) promoter. The BW25113 strain can not metabolise arabinose (Δ*(araD-araB)567* genotype) [79], and therefore the inducer is not metabolised during the experiment.

First we tested the genetic interactions between *bipA* and all *E. coli* ABCFs in CFT073 background.

At 37°C the *bipA* CFT073 knock-out strain has no growth defect (fig. S5A), however at 18°C, the

Δ*bipA* strain displays a pronounced growth defect characteristic of strains defective in ribosome

assembly (fig. 5A) [81]. Ectopic expression of Uup efficiently suppresses the growth defect, while

deletion of *uup* in the *bipA* background exacerbates it (fig. 5A). Expression of EttA and Ybit have no

effect, but expression of YheS leads to a dramatic growth defect. Importantly, in the wild type

background, the expression of YheS has no effect on growth at 18°C (fig. S5B), indicating that the

genetic interaction between *bipA* and *yheS* is specific. As reported previously, disruption of *bipA*

leads to a dramatic ribosome assembly defect at low (18°C) temperature [71] (fig. 5B). The levels

of mature 70S ribosomes as well as 50S subunits are dramatically decreased, accompanied by an

accumulation of 50S assembly precursors (the peak marked with an asterisk) and free 30S

subunits. Ectopic expression of Uup partially suppresses these defects, and in the Δ*bipA*Δ*uup* strain

the defects are exacerbated. All of the effects described above are conditional on disruption of *bipA*

since neither disruption of individual ABCF genes nor simultaneous disruption of *uup* and *ettA* –

the only well-characterised ribosome-associated *E. coli* ABCF to date – causes cold-sensitivity (fig.

S5C) or affects polysome profiles (fig. S5D).

Next we set out to test the effects of EQ_2 versions of ABCFs on translation. We validated the

expression of the FTH-tagged ABCFs using Western blotting (fig. S6A). As was observed for

untagged Uup (fig. 5A), the expression of FTH-tagged Uup suppresses the cold sensitivity caused

by *bipA* deletion while YheS expression exacerbates the growth defect (fig. S6B-C). Expression of

the EQ_2 versions universally causes growth inhibition, both at 18°C in Δ*bipA* CFT073 (fig. S6C) and

at 37°C in the wild type BW25113 background (fig. 6). Overexpression of none of the wild type

ABCFs results in a growth defect (fig. 6A-D). Next we used a [35]S-methionine pulse-labeling assay as

a readout of translational inhibition. For all the ABCF-EQ_2s, the methionine incorporation

decreases, showing protein synthesis is clearly inhibited. The strongest effect is observed for EttA-

EQ_2 (fig. 6A) and YbiT-EQ_2 (fig. 6C), and the weakest is seen in for Uup-EQ_2 (fig. 6B).

 Phylogenetic analysis reveals putative AREs

Through phylogenetic analysis of predicted proteins and previously documented AREs with

sequences available in UniProt [42] and the Comprehensive Antibiotic Resistance Database (CARD)

[82], we have identified seven groups of AREs (fig. 1, table 1). Surprisingly, these can be quite

variable in their subdomain architecture (fig. 7A). Some AREs (ARE1-5) have experienced

extension of the Linker by on average around 30 amino acids compared to EttA, which is in line

with the observation that the extended Linker is in close contact with the bound antibiotic in the

case of ARE1 MsrE [33]. However, linker extension is not the rule for AREs; ARE7 (OptrA) has a

linker of comparable length to the *E. coli* ABCFs (fig. 7A). This supports the notion based on the

VmlR –ribosome co-structure that antibiotic protection by ABCFs can involve allosteric changes in

the ribosome as well as through direct interaction [34]. The Arm subdomain that in EttA interacts

with the L1 ribosomal protein and the L1 stalk rRNA (fig. 7A) varies in length among AREs, and the

CTD extension may or not be present (fig. 7B). Surprisingly, the Uup protein from cave bacterium

Paenibacillus sp. LC231, a sequence that is unremarkable among Uups, confers resistance to the

pleuromutilin antibiotic tiamulin [39] (fig. S1).

Actinobacteria are the source of many ribosome-targeting antibiotics [83], and they have evolved

measures to protect their own ribosomes, including ARE4 (OleB) and ARE5 (VarM). In addition to

these known AREs, Actinobacteria encode a number of other ABCFs specific to this phylum (AAF1-

6; table 1). It is possible that some – if not all – of these groups are in fact AREs. Other subfamilies

may also be unidentified AREs, but two particularly strong candidates are BAF2 and BAF3 that

have strong support for association with ARE5 (fig. 1) and are found in a wide range of bacteria

(table 1). PAF1 is also worthy of investigation; it is found in the genomes of several pathogens in

the proteobacterial genera *Vibrio*, *Enterobacter*, *Klebsiella*, *Serratia* and *Citrobacter*, but not

Escherichia (table S1). With the exception of antibiotic producers, known antibiotic resistance

ABCFs tend to have a variable presence within genera (fig. S7), probably because they are

frequently transferred by mobile elements such as plasmids and transposons. Therefore,

variability in the presence of a subfamily across species in a genus can be an indication that an

ABCF is an ARE. Taking into account phylogenetic relationships (fig. 1, fig. S1) and disjunction

within genera (fig. S7), we predict the following novel AREs: AAF1-5 (which tend to be found in

antibiotic producers), BAF1-3, FAF1-2, and PAF1.

ABCFs are polyproline-rich proteins

Curiously, we find that ABCFs from both bacteria and eukaryotes are often rich in polyproline

sequences, which are known to cause ribosome slow-down or stalling during translation [84]. This

stalling is alleviated by the elongation factor EF-P in bacteria [85, 86], and indeed EF-P is required

for full expression of EttA, which contains two XPPX motifs [87]. 56% of all the sequences in our

ABCF database contain at least two consecutive prolines, compared to an overall 37% of all the

proteins in the predicted proteomes considered here. There is a particularly proline-rich hotspot

in the C-terminal part of the linker (fig. 7B), which can be up to nine consecutive prolines long in

the case of Uup from *Novosphingobium aromaticivorans* (NCBI protein accession number

WP_028641352.1). Arms and CTD extensions are also polyproline hotspots; PPP is a common

motif in the Arm of EttA proteins, and polyprolines are frequent in YdiF, Uup and YheS CTD

extensions. Prolines are rigid amino acids, and conceivably their presence may support the tertiary

structures of ABCFs, particularly the orientation of the subdomain coiled coils [88].

Discussion

Towards a general model for non-eEF3 ABCF function

In the case of ABCFs that act on the assembled ribosome during translation, the E-site should be

vacant (i.e. not filled by an E-site tRNA) for the protein to bind. Specifically, this would be when the

E-site has not yet received a tRNA (during initiation, where ABCF50 (ABCF1) functions [13]) or

when an empty tRNA has dissociated and not been replaced (during slow or stalled translation

such as in the presence of an antibiotic, as in the case of AREs and EttA [16, 17, 34]). EttA has been

proposed to promote the first peptide bond after initiation through modulation of the PTC

conformation [16, 17]; similarly, allosteric effects acting on the PTC have been observed for the

ARE VmlR [34]. This structural modulation or stabilisation could conceivably be a general function

of ABCFs, with the specific ribosomal substrate differing depending on the stage of translation,

assembly, or cellular conditions. Differences in subdomains would determine both what is sensed

and the resulting signal. For instance the presence of the Arm would affect signal transmission

between the PTC and the L1 protein and/or the L1 stalk.

Evolution of AREs

In order to determine antibiotic resistance capabilities and track the transfer routes of resistance,

it is critical to be able to annotate antibiotic resistance genes in genomes. This requires

discrimination of antibiotic genes from homologous genes from which resistance functions have

evolved. At present this is not straightforward for ABCF AREs, as the distinction between potential

translation and antibiotic resistance factors is ambiguous. ABCF AREs have been compared to the

Tet family of antibiotic resistance proteins that evolved from trGTPase EF-G to remove tetracycline

from the ribosome [29]. However the distinction between Tet proteins and EF-G is much more

clear-cut, with Tet comprising a distinct lineage in the evolutionary history of trGTPases [35]. The

surprising lack of a clear sequence signature for antibiotic resistance in the ARE ABCFs suggests

that antibiotic resistance functions may evolve in multiple ways in ABCFs, and that ABCFs closely

related to AREs may have similar functions to the AREs, while not conferring resistance. For

example, there are multiple small molecules that bind the PTC and exit tunnel [89], and

conceivably ABCFs may be involved in sensing such cases, removing the small molecule, or

allowing translation of a subset of mRNAs to continue in its presence. Even macrolide antibiotics

that target the exit tunnel do not abrogate protein synthesis entirely, but rather reshape the

translational landscape [90, 91].

Conclusion

ABCFs are stepping into the limelight as important translation, ribosome assembly and antibiotic

resistance factors. We have found hydrolysis-incompetent EQ_2 mutants of all four *E. coli* ABCFs

inhibit protein synthesis, suggesting they all function on the ribosome. Overexpression of Uup

suppresses both the cold-sensitivity and the 50S ribosome assembly defect caused the loss of

translational GTPase BipA, suggesting that Uup is involved in the 50S ribosome subunit assembly,

either directly or indirectly, for example by fine-tuning expression of ribosomal proteins.

Additionally, we have added ARE2 VmlR to the repertoire of AREs confirmed to act on the

ribosome, joining the ranks of LsaA, VgaA and MsrE, all ARE1s. Considering the well-established

ribosome association of eukaryotic ABCFs, our results suggest that ribosome-binding is a general -

perhaps ancestral - feature of ABCFs. However, the ABCF family is diverse, and even within

subfamilies, there can be differences in subdomain architecture. We have identified clusters of

antibiotic resistance ARE ABCFs, and predicted likely new AREs. Strikingly, the AREs do not form a

clear monophyletic group, and either antibiotic resistance has evolved multiple times

independently from the ABCF diversity, or this is an innate ability of ABCFs, raising the possibility

of a general role of ABCFs in ribosome-binding small molecule sensing and signaling.

Methods

Sequence searching and classification

Predicted proteomes were downloaded from the NCBI genome FTP site (2nd December 2014). One

representative was downloaded per species of bacteria (i.e. not every strain). Seven additional

proteomes were downloaded from JGI (*Aplanochytrium kerguelense, Aurantiochytrium limacinum,*

Fragilariopsis cylindrus, Phytophthora capsici, Phytophthora cinnamomi, Pseudo-nitzschia

multiseries, and *Schizochytrium aggregatum*). Previously documented AREs were retrieved from

UniProt [42] and the Comprehensive Antibiotic Resistance Database (CARD) [82]. Taxonomy was

retrieved from NCBI, and curated manually where some ranks were not available.

An initial local BlastP search was carried out locally with BLAST+ v 2.2.3 [92] against a proteome

database limited by taxonomy to one representative per class or (order if there was no information

on class from the NCBI taxonomy database), using EttA as the query. Subsequent sequence

searching against the proteome collections used hmmsearch from HMMER 3.1b1, with HMMs
made from multiple sequence alignments of subfamilies, as identified below. The E value threshold
for hmmsearch was set to $1e^{-70}$, a value at which the subfamily models hit outside of the
eukaryotic-like or bacterial-like ABCF bipartitions, ensuring complete coverage while not picking
up sequences outside of the ABCF family.

Sequences were aligned with MAFFT v7.164b (default settings) and Maximum Likelihood
phylogenetic analyses were carried out with RAxML-HPC v.8 [93] on the CIPRES Science Gateway
v3 [94] using the LG model of substitution, after removing positions containing >50% gaps.
Additional phylogenetic analyses of representative sequences were carried out as described in the
section "phylogenetic analysis of representatives", below. Trees were visualised with FigTree v.
1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/) and visually inspected to identify putative
subfamilies that preferably satisfied the criteria of 1) containing mostly orthologues, and 2) had at
least moderate (>60% bootstrap support). These subfamilies were then isolated, aligned
separately and used to make HMM models. Models were refined with subsequent rounds of
searching and classification into subfamilies first by comparisons of the E values of HMM hits, then
by curating with phylogenetic analysis. After the final classification of all ABCF types in all
predicted protein sequences using HMMER, some manual correction was still required. For
example, cyanobacterial sequences always hit the chloroplast HMM with a more significant E value
than the bacterial model, and eEF3/New1/eEF3L sequences could not be reliably discriminated
between using E value comparisons. Therefore, the final classification is a manually curated
version of that generated from automatic predictions (table S2). All sequence handling was carried
out with bespoke Python scripts, and data was stored in a MySQL database (exported to Excel files
for the supplementary material tables). Identification of EFL and eEF1A in predicted proteomes
was carried out with HMMER, using the HMMs previously published for these trGTPases [35], The
E value cut-of was set to e-200, lenient enough to match both eEF1A or EFL, with assignment to
either protein subfamily made by E value comparisons as above.

Domain prediction

Domain HMMs were made from subalignments extracted from subfamily alignments. Partial and

poorly aligned sequences were excluded from the alignments. All significant domain hits (<E value

1e^{-3}) for each ABCF sequence were stored in the MySQL database. Sequence logos of domains were

created with Skylign [95]. Putative transit peptides for mitochondrial and plastid subcellular

localization were predicted with the TargetP web server hosted at the Technical University of

Denmark [96].

Phylogenetic analysis of representatives

For the representative tree of the ABCF family, taxa were selected from the ABCF database to

sample broadly across the tree of life, including eukaryotic protistan phyla, while also covering all

subfamilies of ABCFs. Sequences were aligned with MAFFT with the L-ins-i strategy [97] and

positions with >50% gaps, and several ambiguously aligned positions at the termini were

removed. The resulting 249 sequences, and 533 positions were subject to RAxML and IQ-TREE

Maximum Likelihood phylogenetic analysis, both run on the CIPRES Science Gateway v3 [94].

RAxML was run with the LG substitution matrix, as favoured by ProtTest 2.4 [98] and 100

bootstrap replicates. Bootstrapping with RaxML yields a value (maximum likelihood bootstrap

percentage, MLB) for how much of the input alignment supports a particular branch in the tree

topology, and therefore the reliability of that branch. These support values are indicated on

branches in the tree figures. In the case of IQ-TREE, the most appropriate model was selected by

the program during the run, which also favoured the LG substitution matrix. IQ-TREE was run with

its ultrafast bootstrapping approximation method to ascertain support values (UFB) for branches

out of 1000 replicates [99]. To test whether our trees made with ABC domains separately are

incompatible (as might indicate recombination), the RaxML and IQ-TREE analyses were repeated

with a dataset containing the ABC domains uncoupled from each other and aligned together.

Alignments were prepared as above, to make a data set of 204 alignment positions from 525 taxa

(Text S1).

Bayesian inference phylogenetic analysis was carried out with MrBayes v3.2.6, also on the CIPRES

gateway. The analysis was run for 1 million generations, after which the standard deviation of split

frequencies (SDSF) was 0.08. The mixed model setting was used for determining the amino acid

substitution matrix, which converged on WAG. RAxML analysis with the WAG model showed no

difference in topology for well-supported branches compared to the RAxML tree with the LG

model. Branch support values in this case are posterior probabilities, shown on the tree figure as

Bayesian Inference posterior probabilities (BIPP), on a scale of 0 to 1, with increasing probability.

For the rooted tree with ABCE as the outgroup, all ABCFs were selected from *Arabidopsis thaliana,*

Homo sapiens, Escherichia coli, Bacillus subtilis, Saccharomyces cerevisae, Schizosaccharomyces

pombe and *Methanococcus maripaludis*. Ambiguously aligned sites were identified and removed

manually. RAxML, IQ-TREE and MrBayes were carried out as above on the resulting 344 positions

from 35 sequences. The MrBayes analysis stopped automatically when the SDSF dropped to the

0.009 threshold, which was at 235,000 generations.

For the tree of eukaryotic and viral ABCFs rooted with YheS, sequences were extracted from the

ABCF database, and viral sequences were found in the NCBI protein database using BlastP. The

resulting 658 sequences were aligned with MAFFT and a RAxML analysis of 645 positions was

carried out as above. The alignments used to build the phylogenies presented in the main text as

supplementary information are available in text S1, along with trees that are used for ascertaining

branch support but not included as figures.

Structural analyses

Homology modeling was carried out using Swiss Model [100] with EttA (PDB ID 3J5S) as the

template structure. Because the linkers were lacking secondary structure, QUARK [101] was used

for *ab initio* structure modeling of these regions, and the resulting coils were aligned back to the

homology model using the structural alignment method of MacPyMOL [102]. The presence of

coiled coil regions was predicted with the COILS program hosted at the ExPASy Bioinformatics

Research Portal [103].

Construction of plasmids and bacterial strains

All bacterial strains and plasmids used in this study are described in **Supplementary Methods**

and listed in **Table 2**.

Table 2. Strains and plasmids used in the study.

Strain or Plasmid	Description	Reference
Strains: E.coli		
BW	BW25113 *E. coli*	[79]
CFT073	Uropathogenic *E. coli* O6:K2H1	[76]
CFTuup	Δ*uup* (locus tag c1085)	This study
CFTettA	Δ*yjjK* (locus tag c5478)	This study
CFTyheS	Δ*yheS* (locus tag c4127)	This study
CFTybiT	Δ*ybiT* (locus tag c0906)	This study
CFTbipA	Δ*yihK* (locus tag c4820)	This study
CFTbipA_pUup	Δ*yihK* with pSC-uup	This study
Strains: B. subtilis		
B. subtilis 168	*trpC2*	[104]
VHB5	*trpC2 ΔvmlR*	This study
VHB38	*trpC2 ΔvmlR amyE*::P$_{xyl}$-*vmlR-mNeoGreen* Spc^r	This study
VHB44	*trpC2 ΔvmlR thrC*::P$_{hy-spnak}$-*vmlR* Kan^r	This study
VHB45	*trpC2 ΔvmlR thrC*::P$_{hy-spnak}$-*vmlREQ2* Kan^r	This study
VHB91	*trpC2 ΔvmlR thrC*::P$_{hy-spnak}$-*vmlR-HTF* Kan^r	This study
VHB92	*trpC2 ΔvmlR thrC*::P$_{hy-spnak}$-*vmlREQ$_2$-HTF* Kan^r	This study
HS64	*trpC2 amyE*::P$_{xyl}$-*WALP23-gfp* Spc^r	[65]
Plasmids		
pKD4	λRed PCR template plasmid with Kan resistance cassette; Kan^r Amp^r	[78]
pKD13	λRed PCR template plasmid with Kan resistance cassette; Kan^r Amp^r	[78]
pKD46	λRed recombinase helper plasmid, temperature sensitive; Amp^r	[78]
pCP20	FLP recombinase encoding plasmid, temperature sensitive; Amp^r	[78]
pSC101	pSC101 empty vector; Kan^r	This study
pSC-ettA	Constitutive *ettA* overexpression plasmid; Kan^r	This study
pSC-uup	Constitutive *uup* overexpression plasmid; Kan^r	This study
pSC-yheS	Constitutive *yheS* overexpression plasmid; Kan^r	This study

pSC-ybiT	Constitutive *ybiT* overexpression plasmid; Kan^r	This study
pSC-bipA	Constitutive *bipA* overexpression plasmid; Kan^r	This study
pHT009	Integration plasmid ; Kan^r Amp^{r}Kan^r	This study
pSG1154	Integration plasmid; Spc^r Amp^r	[63]
pSHP2	Integration plasmid; Spc^r Amp^r	This study
VHp62	pAPNC with vmlR-HTF in SalI/BamHI sites; Spc^r Amp^r	Laboratory stock
VHp66	pAPNC with vmlREQ$_2$-HTF in SalI/BamHI sites; Spc^r Amp^r	Laboratory stock
pHT009-vmlR	pHT009 with vmlREQ2; Kan^r Amp^r	This study
pHT009-vmlR-HTF	pHT009 with vmlR-HTF in HindIII/SphI site; Kan^r Amp^r	This study
pHT009-vmlREQ2-HTF	pHT009 with vmlREQ2-HTF in HindIII/SphI site; Kan^r Amp^r	This study
pSHP2-vmlR	pSHP2 with vmlR in ApaI/EcoRI site ; Spc^r Amp^r	This study

Growth assays

Bacterial growth (OD_{600}) was monitored using a Bioscreen C (Oy Growth Curves Ab Ltd)

microplate reader in Honeycomb plates (150 μL culture per well) with continuous shaking (speed:

fast, amplitude: normal). All experiments with CFT073 were performed at 18°C unless stated

otherwise. Three biological replicates were averaged for each growth curve and the data presented

as geometric means ± standard deviation.

E. coli *transformed with pSC101-based expression plasmids:*

Overnight (16 h) cultures were pre-grown in LB medium supplemented with 50 μg/mL

kanamycin, diluted to OD_{600} of 0.03 in filtered LB and grown in Bioscreen C microplate reader as

described above.

E. coli *transformed with pBAD-based expression plasmids:*

Overnight (16 h) cultures were pre-grown in Neidhardt MOPS medium [105] supplemented with

0.1% of casamino acids, 0.4% glucose as a carbon source and 100 μg/mL carbenicillin, diluted to

OD_{600} of 0.03 in the same media but containing and 0.5% glycerol instead of 0.4% glucose as well

as supplemented with 0.5% arabinose and grown in Bioscreen C microplate reader as described

above.

Antibiotic resistance testing of tagged B. subtilis *VmlR*:

B. subtilis strains VHB38, VHB91 and VHB92 were pre-grown on LB plates overnight at 30℃. Fresh

individual colonies were used to inoculate filtered LB medium, either in the presence and absence

of 1 mM IPTG (for VHB91 and VHB92) or in the presence and absence of 0.3% xylose (for VHB38),

and OD_{600} adjusted to 0.01. The cultures were seeded on Honeycomb plates, and plates incubated

in a Bioscreen C at 37 °C with continuous shaking as described above for *E. coli* cultures. After 90

minute incubation ($OD_{600} \approx 0.1$) increasing concentrations of lincomycin (final concentration 0 - 5

μg/ml) were added and growth was monitored for additional 6 hours. Three biological replicates

were averaged for each growth curve and the data presented as geometric means ± standard

deviation.

Fluorescence microscopy

Fluorescence microscopy was carried out with cell grown to early-mid logarithmic growth phase

(OD_{600} of 0.2-0.5) in LB medium at 37 °C in the presence or absence of inducers. The used inducer

concentrations were 0.3% for VmlR-mNG and 1% for WALP23-GFP. If indicated, the cells were

incubated with 5μg/ml lincomycin upon shaking at 37 °C prior to the microscopy. The cells were

immobilised on microscopy slides covered with a thin film of 1.2% (w/v) agarose in H_2O as

described in detail elsewhere [106]. The microscopy was carried out with Nikon Eclipse Ti

equipped with Nikon Plan Apo 100x/1.40 Oil Ph3 objective, Sutter Instrument Company Lambda

LS xenon arc light source, and Photometrics Prime sCMOS camera. The images were captured

using Metamorph 7.7 (Molecular Devices) and analysed using Fiji [107].

Western blot analysis of FTH-tagged wt and EQ_2 E. coli ABCF proteins

Preparation of bacterial samples:

Bacteria were grown either at 37 °C (*E. coli* BW25113 derivatives) or 18°C (*E. coli ΔbipA* CFT073

derivatives) up to OD_{600} of 0.5 in 50 mL of Neidhardt MOPS minimal medium [105] supplemented

with 0.1% casamino acids (w/v), 0.5% glycerol (w/v) and 100 μg/mL carbenicillin and L-

arabinose was added to a final concentration of 0.2% (w/v). Cultures grown at 37°C were

harvested 10 minutes after induction by pouring them into precooled centrifuge bottles containing

100 g of crushed ice and centrifuged at 10,000 rpm for 10 minutes at 4 °C (Beckman JLA16.250

rotor). Cultures grown at 18°C were harvested 5 hours after induction by collecting into precooled

centrifuge bottles and pelleting at 10,000 rpm for 10 minutes at 4 °C (Beckman JA25.50 rotor).

Lysates were prepared the same as for polysome profiling of *E. coli* (see below).

Western blotting:

3 μg of total protein as determined by Bradford assay of each sample was resolved on 10% SDS-

PAGE gel and transferred to 0.2 μm nitrocellulose membrane (Trans-Blot® Turbo™ Transfer Pack,

Bio-Rad) using Turbo MIXED MW protocol in Trans-Blot® Turbo™ Transfer System (Bio-Rad). The

membrane was blocked in PBS-T (1x PBS 0.05% Tween-20) with 5% w/v nonfat dry milk at room

temperature for one hour. Antibody incubations were performed for one hour in 1% nonfat dry

milk in PBS-T with five 5-minute washes in fresh PBS-T between and after antibody incubations.

FTH-tagged ABCFs were detected using anti-Flag M2 primary (Sigma-Aldrich, F1804; 1:10,000

dilution) antibodies combined with anti-mouse-HRP secondary (Rockland; 610-103-040; 1:10,000

dilution) antibodies. AECL detection was performed on ImageQuant LAS 4000 (GE Healthcare)

imaging system using Pierce® ECL Western blotting substrate (Thermo Scientific).

Polysome profiling analysis of E. coli strains

Preparation of bacterial samples:

Overnight (16 h) cultures were pre-grown at 37°C in LB medium supplemented with 50 μg/mL

kanamycin in the case of strains transformed with pSC101-based expression plasmids. Overnight

cultures were diluted in filtered LB (33 mL cultures) and after 24 h growth at 18 °C harvested by

pouring into precooled centrifuge bottles and pelleting at 10,000 rpm for 10 minutes at 4 °C

(Beckman JA25.50 rotor). For the sake of convenience, cultures were diluted to different starting

densities (Table S3) to ensure that all of them reach $OD_{600} \approx 0.5$ simultaneously.

Preparation of clarified lysates:

Cell pellets were resuspended in 0.4 mL of Polymix buffer [108] (20 mM HEPES:KOH pH 7.5, 95

mM KCl, 5 mM NH$_4$Cl, 5 mM Mg(OAc)$_2$, 0.5 mM CaCl$_2$, 8 mM putrescine, 1 mM spermidine, 1 mM

DTT) and 200 μL of pre-chilled zirconium beads (0.1 mm) were added to each sample. Cellular

lysates were prepared by a FastPrep homogeniser (MP Biomedicals) (three 20 seconds pulses at

speed 6.0 mp/sec with chilling on ice for 1 minutes between the cycles) and clarified by

centrifugation at 21,000 g for 10 minutes at 4 °C. The supernatant was carefully collected avoiding

the lipid layer and cellular pellet, aliquoted, frozen in liquid nitrogen and stored at -80 °C until

further processing.

Sucrose gradient centrifugation:

After melting the frozen samples on ice, 2 A$_{260}$ units of each extract was loaded onto 5–25% (w/v)

sucrose density gradients in Polymix buffer, 5 mM Mg^{2+} [108]. Gradients were resolved at 35,000

rpm for 2.5 hours at 4 °C in SW41 rotor (Beckman) and analysed using Biocomp Gradient Station

(BioComp Instruments) with A$_{260}$ as a readout. The ribosome profiles presented were normalised

to the total area under the curve and are representative of at least three independent experiments

for each strain.

Polysome profiling and Western blot analysis of B. subtilis strains

Experiments were performed as described above for *E. coli* strains, with minor modifications.

Preparation of bacterial samples and preparation of clarified lysates:

VHB90 and VHB91 strains were pre-grown on LB plates overnight at 30 °C. Fresh individual

colonies were used to inoculate 200 mL LB cultures. The cultures were grown at 37 °C until OD$_{600}$

of 0.3 and IPTG was added to final concentration of 30 μM. After 30 min cells were collected by

centrifugation (8,000 rmp, 10 minutes), dissolved in 0.5 mL of Polymix buffer [108] (20 mM

HEPES:KOH pH 7.5, 95 mM KCl, 5 mM NH$_4$Cl, 10 mM Mg(OAc)$_2$, 0.5 mM CaCl$_2$, 8 mM putrescine, 1

mM spermidine, 1 mM DTT, 2 mM PMSF), lysed (FastPrep homogeniser (MP Biomedicals): four 20

seconds pulses at speed 6.0 mp/sec with chilling on ice for 1 minutes between the cycles), and

clarified by ultracentrifugation (14,800 rpm, 20 minutes).

Sucrose gradient centrifugation and Western blotting:

Clarified cell lysates were loaded onto 7–35% sucrose gradients in Polymix buffer [108] (20 mM

HEPES:KOH pH 7.5, 95 mM KCl, 5 mM NH_4Cl, 10 mM $Mg(OAc)_2$, 0.5 mM $CaCl_2$, 8 mM putrescine, 1

mM spermidine, 1 mM DTT) and subjected to centrifugation (35,000 rpm for 3 hours at 4 °C). C-

terminally HTF-tagged VmlR (wild type and EQ_2 mutant) and ribosomal protein L3 of the 50S

ribosomal subunit were detected using either anti-Flag M2 primary combined with anti-mouse-

HRP secondary antibodies or anti-L3 primary (a gift from Fujio Kawamura) combined with goat

anti-rabbit IgG-HRP secondary antibodies, respectively. All antibodies were used at 1:10,000

dilution.

L-[^{35}S]-methionine pulse-labelling

Preparation of bacterial samples:

Since glucose specifically inhibits the arabinose promoter, the cultures were grown in defined

Neidhardt MOPS medium [105] supplemented with 0.4% glycerol as a carbon source. One colony

of freshly transformed *E. coli* BW25113 cells expressing N-terminal FTH-tagged ABCFs (wild type

and EQ_2 mutants) from pBad vector was used to inoculate 10 mL of 1x MOPS media supplemented

with 0.4% glycerol 100 μg/mL carbenicillin, and the cultures were grown until early stationary

phase (about 24 hours). Stationary phase cells were diluted to OD_{600} of 0.04–0.07 in 25 mL of the

same media, grown at 37 °C with vigorous shaking (200 rpm) to OD_{600} of 0.15–0.2 and expression

of ABCFs was induced by addition of L-arabinose to the final concentration of 0.2%.

L-[^{35}S]-methionine pulse-labelling:

For radioactive pulse labelling, 1 μCi L-[^{35}S]-methionine (500μCi, PerkinElmer) aliquots were

prepared in 1.5 mL Eppendorf tubes. As a zero time point, 1 mL of cell culture was taken and mixed

with an aliquot of radioactive methionine just before inducing cells with L-arabinose.

Simultaneously a 1 mL aliquot was taken for an OD_{600} measurement. All consecutive samples were

processed similarly at designated time points after induction. ^{35}S-methionine incorporation was

stopped after 5 minutes by chloramphenicol added to the final concentration of 200 μg/mL.

Subsequent processing of samples differs in the case of scintillation counting and autoradiography.

Scintillation counting:

1 mL of culture was combined with 200 µL of 50% trichloroacetic acid (TCA), passed through a

GF/C filter (Whatman) prewashed with 5% TCA and unincorporated label was removed by

washing the filter with 5 mL of ice-cold 5% TCA followed by 5 mL of ice-cold 95% EtOH [109].

Filters were dried for at least 2 hours, and counted on a TRI-CARB 4910TR 110 V scintillation

counter (PerkinElmer) (5 mL of ScintiSafe 3 scintillation cocktail (FisherScientific) per sample,

pre-soaked with shaking for 15 minutes prior to counting).

Autoradiography:

1 mL cultures were pelleted by centrifugation, cell pellet washed with Phosphate Buffered Saline

(PBS) to remove unincorporated L-[^{35}S]-methionine and dissolved/lysed in 50 µL 1x SDS-loading

buffer. Samples were normalised by OD600 by addition of appropriate volume of 1x SDS-loading

buffer (50-80 µL according to OD600), and 10 µL of the sample was loaded onto 10% SDS-PAGE

and resolved electrophoretically (BioRad), gels were dried on Whatman paper, exposed on BAS

storage phosphor screen (GE Heathcare) overnight, and scanned by Typhoon imaging system (GE

Heathcare).

Acknowledgements

We are grateful to the anonymous reviewers in their suggestions of additional experiments and

analyses that strengthened the manuscript. We gratefully acknowledge Fujio Kawamura for

providing anti-L3 primary antibodies and Stijn Hendrik Peeters for constructing the pSHP2

plasmid. Thanks to Martin Carr for discussion and advice about eEF3 and EFL co-distribution.

Author contributions:

GCA conceived the study and carried out all the bioinformatic analyses except for sequence logo

construction and eEF1A/EFL/eEF3 co-distribution analysis, which was carried out by CKS. VH, VM,

MK, HS, HT, TT and MP designed experimental analyses. VM, MK, HT, MH, TS, MR, HS and JWG

carried out the experiments. VH, GCA, HS, TT, HT, VM and MK analyzed the data. GCA and VH

drafted the manuscript with input from VM, MK, HT, MH, CKS, JWG, TS, MP, TT and HS.

References

1. Skogerson, L. and E. Wakatama, *A ribosome-dependent GTPase from yeast distinct from elongation factor 2.* Proc Natl Acad Sci U S A, 1976. **73**(1): p. 73-6.

2. Andersen, C.B., et al., *Structure of eEF3 and the mechanism of transfer RNA release from the E-site.* Nature, 2006. **443**(7112): p. 663-8.

3. Kurata, S., et al., *Ribosome recycling step in yeast cytoplasmic protein synthesis is catalyzed by eEF3 and ATP.* Proc Natl Acad Sci U S A, 2010. **107**(24): p. 10854-9.

4. Triana-Alonso, F.J., K. Chakraburtty, and K.H. Nierhaus, *The elongation factor 3 unique in higher fungi and essential for protein biosynthesis is an E site factor.* J Biol Chem, 1995. **270**(35): p. 20473-8.

5. Pisarev, A.V., et al., *The role of ABCE1 in eukaryotic posttermination ribosomal recycling.* Mol Cell, 2010. **37**(2): p. 196-210.

6. Becker, T., et al., *Structural basis of highly conserved ribosome recycling in eukaryotes and archaea.* Nature, 2012. **482**(7386): p. 501-6.

7. Young, D.J., et al., *Rli1/ABCE1 Recycles Terminating Ribosomes and Controls Translation Reinitiation in 3'UTRs In Vivo.* Cell, 2015. **162**(4): p. 872-84.

8. Caetano-Anolles, D., et al., *Proteome evolution and the metabolic origins of translation and cellular life.* J Mol Evol, 2011. **72**(1): p. 14-33.

9. Davidson, A.L., et al., *Structure, function, and evolution of bacterial ATP-binding cassette systems.* Microbiol Mol Biol Rev, 2008. **72**(2): p. 317-64, table of contents.

10. Dean, M., A. Rzhetsky, and R. Allikmets, *The human ATP-binding cassette (ABC) transporter superfamily.* Genome Res, 2001. **11**(7): p. 1156-66.

11. Kerr, I.D., *Sequence analysis of twin ATP binding cassette proteins involved in translational control, antibiotic resistance, and ribonuclease L inhibition.* Biochem Biophys Res Commun, 2004. **315**(1): p. 166-73.

12. Vazquez de Aldana, C.R., M.J. Marton, and A.G. Hinnebusch, *GCN20, a novel ATP binding cassette protein, and GCN1 reside in a complex that mediates activation of the eIF-2 alpha kinase GCN2 in amino acid-starved cells.* EMBO J, 1995. **14**(13): p. 3184-99.

13. Paytubi, S., et al., *ABC50 promotes translation initiation in mammalian cells.* J Biol Chem, 2009. **284**(36): p. 24061-73.

14. Dong, J., et al., *The novel ATP-binding cassette protein ARB1 is a shuttling factor that stimulates 40S and 60S ribosome biogenesis.* Mol Cell Biol, 2005. **25**(22): p. 9859-73.

15. Li, Z., et al., *Rational extension of the ribosome biogenesis pathway using network-guided genetics.* PLoS Biol, 2009. **7**(10): p. e1000213.

16. Boël, G., et al., *The ABC-F protein EttA gates ribosome entry into the translation elongation*
cycle. Nat Struct Mol Biol, 2014. **21**(2): p. 143-51.

17. Chen, B., et al., *EttA regulates translation by binding the ribosomal E site and restricting*
ribosome-tRNA dynamics. Nat Struct Mol Biol, 2014. **21**(2): p. 152-9.

18. Reynolds, E.D. and J.H. Cove, *Resistance to telithromycin is conferred by msr(A), msrC and*
msr(D) in Staphylococcus aureus. J Antimicrob Chemother, 2005. **56**(6): p. 1179-80.

19. Ohki, R., et al., *Transcriptional termination control of a novel ABC transporter gene involved*
in antibiotic resistance in Bacillus subtilis. J Bacteriol, 2005. **187**(17): p. 5946-54.

20. Singh, K.V., G.M. Weinstock, and B.E. Murray, *An Enterococcus faecalis ABC homologue (Lsa)*
is required for the resistance of this species to clindamycin and quinupristin-dalfopristin.
Antimicrob Agents Chemother, 2002. **46**(6): p. 1845-50.

21. Novotna, G. and J. Janata, *A new evolutionary variant of the streptogramin A resistance*
protein, Vga(A)LC, from Staphylococcus haemolyticus with shifted substrate specificity
towards lincosamides. Antimicrob Agents Chemother, 2006. **50**(12): p. 4070-6.

22. Hot, C., N. Berthet, and O. Chesneau, *Characterization of sal(A), a novel gene responsible for*
lincosamide and streptogramin A resistance in Staphylococcus sciuri. Antimicrob Agents
Chemother, 2014. **58**(6): p. 3335-41.

23. Buche, A., C. Mendez, and J.A. Salas, *Interaction between ATP, oleandomycin and the OleB*
ATP-binding cassette transporter of Streptomyces antibioticus involved in oleandomycin
secretion. Biochem J, 1997. **321 (Pt 1)**: p. 139-44.

24. Ross, J.I., et al., *Inducible erythromycin resistance in staphylococci is encoded by a member of*
the ATP-binding transport super-gene family. Mol Microbiol, 1990. **4**(7): p. 1207-14.

25. Wang, Y., et al., *A novel gene, optrA, that confers transferable resistance to oxazolidinones*
and phenicols and its presence in Enterococcus faecalis and Enterococcus faecium of human
and animal origin. J Antimicrob Chemother, 2015. **70**(8): p. 2182-90.

26. Kitani, S., et al., *Characterization of varM Encoding Type II ABC Transporter in Streptomyces*
virginiae, a Virginiamycin M1 Producer. Actinomycetologica, 2010. **24**(2): p. 51-57.

27. Kerr, I.D., E.D. Reynolds, and J.H. Cove, *ABC proteins and antibiotic drug resistance: is it all*
about transport? Biochem Soc Trans, 2005. **33**(Pt 5): p. 1000-2.

28. Rice, L.B., *Progress and challenges in implementing the research on ESKAPE pathogens.* Infect
Control Hosp Epidemiol, 2010. **31 Suppl 1**: p. S7-10.

29. Wilson, D.N., *The ABC of Ribosome-Related Antibiotic Resistance.* MBio, 2016. **7**(3).

30. Sharkey, L.K.R. and A.J. O'Neill, *Antibiotic Resistance ABC-F Proteins: Bringing Target*
Protection into the Limelight. ACS Infect Dis, 2018. **4**(3): p. 239-246.

31. Sharkey, L.K., T.A. Edwards, and A.J. O'Neill, *ABC-F Proteins Mediate Antibiotic Resistance*
through Ribosomal Protection. MBio, 2016. **7**(2).

32.   Murina, V., et al., *Antibiotic resistance ABCF proteins reset the peptidyl transferase centre of*
the ribosome to counter translational arrest. Nucleic Acids Res, 2018. **46**(7): p. 3753-3763.

33.   Su, W., et al., *Ribosome protection by antibiotic resistance ATP-binding cassette protein.* Proc
Natl Acad Sci U S A, 2018. **115**(20): p. 5157-5162.

34.   Crowe-McAuliffe, C., et al., *Structural basis for antibiotic resistance mediated by the Bacillus*
subtilis ABCF ATPase VmlR. Proc Natl Acad Sci U S A, 2018.

35.   Atkinson, G.C., *The evolutionary and functional diversity of classical and lesser-known*
cytoplasmic and organellar translational GTPases across the tree of life. BMC Genomics,
2015. **16**: p. 78.

36.   Donhofer, A., et al., *Structural basis for TetM-mediated tetracycline resistance.* Proc Natl
Acad Sci U S A, 2012. **109**(42): p. 16900-5.

37.   Li, W., et al., *Mechanism of tetracycline resistance by ribosomal protection protein Tet(O).* Nat
Commun, 2013. **4**: p. 1477.

38.   Laurence, M., C. Hatzis, and D.E. Brash, *Common contaminants in next-generation sequencing*
that hinder discovery of low-abundance microbes. PLoS One, 2014. **9**(5): p. e97876.

39.   Pawlowski, A.C., et al., *A diverse intrinsic antibiotic resistome from a cave bacterium.* Nat
Commun, 2016. **7**: p. 13803.

40.   Schoner, B., et al., *Sequence similarity between macrolide-resistance determinants and ATP-*
binding transport proteins. Gene, 1992. **115**(1-2): p. 93-6.

41.   Peschke, U., et al., *Molecular characterization of the lincomycin-production gene cluster of*
Streptomyces lincolnensis 78-11. Mol Microbiol, 1995. **16**(6): p. 1137-56.

42.   The UniProt Consortium, *UniProt: the universal protein knowledgebase.* Nucleic Acids Res,
2017. **45**(D1): p. D158-D169.

43.   Carlier, L., et al., *The C-terminal domain of the Uup protein is a DNA-binding coiled coil motif.*
J Struct Biol, 2012. **180**(3): p. 577-84.

44.   Maurice, T.C., et al., *A highly conserved intraspecies homolog of the Saccharomyces cerevisiae*
elongation factor-3 encoded by the HEF3 gene. Yeast, 1998. **14**(12): p. 1105-13.

45.   Sarthy, A.V., et al., *Identification and kinetic analysis of a functional homolog of elongation*
factor 3, YEF3 in Saccharomyces cerevisiae. Yeast, 1998. **14**(3): p. 239-53.

46.   Marton, M.J., et al., *Evidence that GCN1 and GCN20, translational regulators of GCN4, function*
on elongating ribosomes in activation of eIF2alpha kinase GCN2. Mol Cell Biol, 1997. **17**(8):
p. 4474-89.

47.   Anand, M., et al., *Functional interactions between yeast translation eukaryotic elongation*
factor (eEF) 1A and eEF3. J Biol Chem, 2003. **278**(9): p. 6985-91.

48.   Anand, M., et al., *Domain and nucleotide dependence of the interaction between*
Saccharomyces cerevisiae translation elongation factors 3 and 1A. J Biol Chem, 2006.
281(43): p. 32318-26.

49.   Yamada, T., et al., *Expression of the gene encoding a translational elongation factor 3*
908   *homolog of Chlorella virus CVK2.* Virology, 1993. **197**(2): p. 742-50.

909   50.   Atkinson, G.C., et al., *An evolutionary ratchet leading to loss of elongation factors in*
910   *eukaryotes.* BMC Evol Biol, 2014. **14**(1): p. 35.

911   51.   Inoue, Y., et al., *Yeast prion protein New1 can break Sup35 amyloid fibrils into fragments in*
912   *an ATP-dependent manner.* Genes Cells, 2011. **16**(5): p. 545-56.

913   52.   Paytubi, S., et al., *The N-terminal region of ABC50 interacts with eukaryotic initiation factor*
914   *eIF2 and is a target for regulatory phosphorylation by CK2.* Biochem J, 2008. **409**(1): p. 223-
915   31.

916   53.   Kachroo, A.H., et al., *Evolution. Systematic humanization of yeast genes reveals conserved*
917   *functions and genetic modularity.* Science, 2015. **348**(6237): p. 921-5.

918   54.   Castilho, B.A., et al., *Keeping the eIF2 alpha kinase Gcn2 in check.* Biochim Biophys Acta,
919   2014. **1843**(9): p. 1948-68.

920   55.   Hirose, T. and H.R. Horvitz, *The translational regulators GCN-1 and ABCF-3 act together to*
921   *promote apoptosis in C. elegans.* PLoS Genet, 2014. **10**(8): p. e1004512.

922   56.   Visweswaraiah, J., et al., *Overexpression of eukaryotic translation Elongation Factor 3 (eEF3)*
923   *impairs GCN2 activation.* J Biol Chem, 2012.

924   57.   Kaida, D., et al., *Yeast Whi2 and Psr1-phosphatase form a complex and regulate STRE-*
925   *mediated gene expression.* Genes Cells, 2002. **7**(6): p. 543-52.

926   58.   Dar, D., et al., *Term-seq reveals abundant ribo-regulation of antibiotics resistance in bacteria.*
927   Science, 2016. **352**(6282): p. aad9822.

928   59.   Orelle, C., et al., *The conserved glutamate residue adjacent to the Walker-B motif is the*
929   *catalytic base for ATP hydrolysis in the ATP-binding cassette transporter BmrA.* J Biol Chem,
930   2003. **278**(47): p. 47002-8.

931   60.   Smith, P.C., et al., *ATP binding to the motor domain from an ABC transporter drives formation*
932   *of a nucleotide sandwich dimer.* Mol Cell, 2002. **10**(1): p. 139-49.

933   61.   Britton, R.A., et al., *Genome-wide analysis of the stationary-phase sigma factor (sigma-H)*
934   *regulon of Bacillus subtilis.* J Bacteriol, 2002. **184**(17): p. 4881-90.

935   62.   Shaner, N.C., et al., *A bright monomeric green fluorescent protein derived from*
936   *Branchiostoma lanceolatum.* Nat Methods, 2013. **10**(5): p. 407-9.

937   63.   Lewis, P.J. and A.L. Marston, *GFP vectors for controlled expression and dual labelling of*
938   *protein fusions in Bacillus subtilis.* Gene, 1999. **227**(1): p. 101-10.

939   64.   Akanuma, G., et al., *Ribosome dimerization is essential for the efficient regrowth of Bacillus*
940   *subtilis.* Microbiology, 2016. **162**(3): p. 448-458.

941   65.   Scheinpflug, K., et al., *Antimicrobial peptide cWFW kills by combining lipid phase separation*
942   *with autolysis.* Sci Rep, 2017. **7**: p. 44332.

66.   Schafer, L.V., et al., *Lipid packing drives the segregation of transmembrane helices into*
disordered lipid domains in model membranes. Proc Natl Acad Sci U S A, 2011. **108**(4): p.
1343-8.

67.   Jahn, N., S. Brantl, and H. Strahl, *Against the mainstream: the membrane-associated type I*
toxin BsrG from Bacillus subtilis interferes with cell envelope biosynthesis without increasing
membrane permeability. Mol Microbiol, 2015. **98**(4): p. 651-66.

68.   Lewis, P.J., S.D. Thaker, and J. Errington, *Compartmentalization of transcription and*
translation in Bacillus subtilis. EMBO J, 2000. **19**(4): p. 710-8.

69.   Sanamrad, A., et al., *Single-particle tracking reveals that free ribosomal subunits are not*
excluded from the Escherichia coli nucleoid. Proc Natl Acad Sci U S A, 2014. **111**(31): p.
11413-8.

70.   Cochrane, K.L., *Elucidating Ribosomes-Genetic Studies of the ATPase Uup and the Ribosomal*
Protein L1. 2015, University of Michigan. p. 147.

71.   Gibbs, M.R. and K. Fredrick, *Roles of elusive translational GTPases come to light and inform*
on the process of ribosome biogenesis in bacteria. Mol Microbiol, 2018. **107**(4): p. 445-454.

72.   deLivron, M.A. and V.L. Robinson, *Salmonella enterica serovar Typhimurium BipA exhibits*
two distinct ribosome binding modes. J Bacteriol, 2008. **190**(17): p. 5944-52.

73.   Kumar, V., et al., *Structure of BipA in GTP form bound to the ratcheted ribosome.* Proc Natl
Acad Sci U S A, 2015. **112**(35): p. 10944-9.

74.   Hauryliuk, V., et al., *Recent functional insights into the role of (p)ppGpp in bacterial*
physiology. Nat Rev Microbiol, 2015. **13**(5): p. 298-309.

75.   deLivron, M.A., et al., *A novel domain in translational GTPase BipA mediates interaction with*
the 70S ribosome and influences GTP hydrolysis. Biochemistry, 2009. **48**(44): p. 10533-41.

76.   Welch, R.A., et al., *Extensive mosaic structure revealed by the complete genome sequence of*
uropathogenic Escherichia coli. Proc Natl Acad Sci U S A, 2002. **99**(26): p. 17020-4.

77.   Cohen, S.N. and A.C. Chang, *Recircularization and autonomous replication of a sheared R-*
factor DNA segment in Escherichia coli transformants. Proc Natl Acad Sci U S A, 1973. **70**(5):
p. 1293-7.

78.   Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in Escherichia*
coli K-12 using PCR products. Proc Natl Acad Sci U S A, 2000. **97**(12): p. 6640-5.

79.   Grenier, F., et al., *Complete Genome Sequence of Escherichia coli BW25113.* Genome
Announc, 2014. **2**(5).

80.   Baba, T., et al., *Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants:*
the Keio collection. Mol Syst Biol, 2006. **2**: p. 2006 0008.

81.   Guthrie, C., H. Nashimoto, and M. Nomura, *Structure and function of E. coli ribosomes. 8.*
Cold-sensitive mutants defective in ribosome assembly. Proc Natl Acad Sci U S A, 1969. **63**(2):
p. 384-91.

82.   Jia, B., et al., *CARD 2017: expansion and model-centric curation of the comprehensive*
antibiotic resistance database. Nucleic Acids Res, 2017. **45**(D1): p. D566-D573.

83.   Mahajan, G.B. and L. Balachandran, *Antibacterial agents from actinomycetes - a review.*
Front Biosci (Elite Ed), 2012. **4**: p. 240-53.

84.   Lassak, J., D.N. Wilson, and K. Jung, *Stall no more at polyproline stretches with the translation*
elongation factors EF-P and IF-5A. Mol Microbiol, 2016. **99**(2): p. 219-35.

85.   Ude, S., et al., *Translation elongation factor EF-P alleviates ribosome stalling at polyproline*
stretches. Science, 2013. **339**(6115): p. 82-5.

86.   Peil, L., et al., *Distinct XPPX sequence motifs induce ribosome stalling, which is rescued by the*
translation elongation factor EF-P. Proc Natl Acad Sci U S A, 2013. **110**(38): p. 15265-70.

87.   Elgamal, S., et al., *EF-P dependent pauses integrate proximal and distal signals during*
translation. PLoS Genet, 2014. **10**(8): p. e1004553.

88.   Receveur, V., et al., *Dimension, shape, and conformational flexibility of a two domain fungal*
cellulase in solution probed by small angle X-ray scattering. J Biol Chem, 2002. **277**(43): p.
40887-92.

89.   Seip, B. and C.A. Innis, *How Widespread is Metabolite Sensing by Ribosome-Arresting Nascent*
Peptides? J Mol Biol, 2016. **428**(10 Pt B): p. 2217-27.

90.   Kannan, K., N. Vazquez-Laslop, and A.S. Mankin, *Selective protein synthesis by ribosomes*
with a drug-obstructed exit tunnel. Cell, 2012. **151**(3): p. 508-20.

91.   Almutairi, M.M., et al., *Co-produced natural ketolides methymycin and pikromycin inhibit*
bacterial growth by preventing synthesis of a limited number of proteins. Nucleic Acids Res,
2017. **45**(16): p. 9573-9582.

92.   Camacho, C., et al., *BLAST+: architecture and applications.* BMC Bioinformatics, 2009. **10**: p.
421.

93.   Stamatakis, A., *RAxML version 8: a tool for phylogenetic analysis and post-analysis of large*
phylogenies. Bioinformatics, 2014. **30**(9): p. 1312-3.

94.   Miller, M.A., W. Pfeiffer, and T. Schwartz. *Creating the CIPRES Science Gateway for Inference*
of Large Phylogenetic Trees. in *Gateway Computing Environments Workshop (GCE)*. 2010.
New Orleans, LA.

95.   Wheeler, T.J., J. Clements, and R.D. Finn, *Skylign: a tool for creating informative, interactive*
logos representing sequence alignments and profile hidden Markov models. BMC
Bioinformatics, 2014. **15**: p. 7.

96.   Emanuelsson, O., et al., *Locating proteins in the cell using TargetP, SignalP and related tools.*
Nat Protoc, 2007. **2**(4): p. 953-71.

97.   Katoh, K. and D.M. Standley, *MAFFT multiple sequence alignment software version 7:*
improvements in performance and usability. Mol Biol Evol, 2013. **30**(4): p. 772-80.

98.   Abascal, F., R. Zardoya, and D. Posada, *ProtTest: selection of best-fit models of protein*
evolution. Bioinformatics, 2005. **21**(9): p. 2104-5.

99.   Minh, B.Q., M.A. Nguyen, and A. von Haeseler, *Ultrafast approximation for phylogenetic*
bootstrap. Mol Biol Evol, 2013. **30**(5): p. 1188-95.

100.   Biasini, M., et al., *SWISS-MODEL: modelling protein tertiary and quaternary structure using*
evolutionary information. Nucleic Acids Res, 2014. **42**(Web Server issue): p. W252-8.

101.   Xu, D. and Y. Zhang, *Ab initio protein structure assembly using continuous structure*
fragments and optimized knowledge-based force field. Proteins, 2012. **80**(7): p. 1715-35.

102.   Schrodinger, L., *The PyMOL Molecular Graphics System, Version 1.8.* 2015.

103.   Lupas, A., M. Van Dyke, and J. Stock, *Predicting coiled coils from protein sequences.* Science,
1991. **252**(5009): p. 1162-4.

104.   Barbe, V., et al., *From a consortium sequence to a unified sequence: the Bacillus subtilis 168*
reference genome a decade later. Microbiology, 2009. **155**(Pt 6): p. 1758-75.

105.   Neidhardt, F.C., P.L. Bloch, and D.F. Smith, *Culture medium for enterobacteria.* J Bacteriol,
1974. **119**(3): p. 736-47.

106.   Te Winkel, J.D., et al., *Analysis of Antimicrobial-Triggered Membrane Depolarization Using*
Voltage Sensitive Dyes. Front Cell Dev Biol, 2016. **4**: p. 29.

107.   Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat Methods,
2012. **9**(7): p. 676-82.

108.   Antoun, A., et al., *Ribosome formation from subunits studied by stopped-flow and Rayleigh*
light scattering. Biol Proced Online, 2004. **6**: p. 35-54.

109.   Esposito, A.M. and T.G. Kinzy, *In vivo [35S]-methionine incorporation.* Methods Enzymol,
2014. **536**: p. 55-64.

1039

Supplementary file captions

S1 Table. Taxonomy of 4505 species, and their ABCF composition

All species considered in the analysis are listed, ordered by taxonomy. The number and identity of

ABCF subfamilies are recorded.

S2 Table. Classification of 16848 ABCF sequences into subfamilies, accompanied by domain

assignments

The unique sequence identifiers are included for retrieval of data from online repositories.

S3 Table. Domain coordinates

The domain coordinates for the representative sequences in fig. 3A are listed. In the second tab, all

the coordinates for each identified domain in all ABCFs are given. As these are from HMM hits, the

same domain can have more than one hit in each protein. For example, the ABC1 HMM always hits

the ABC2 domain, and vice versa. Duplicate domain hits were removed when generating fig 3A.

S4 Table. Presence and absence of EFL, eEF1A and eEF3 in eukaryotes

Where the distribution is unchanged within a specific taxonomic lineage, those rows are collapsed

down to one, and the highest common taxonomic rank is given. The full lineage data is available in

the second tab.

S5 Table. Transit peptide predictions

Predictions were made separately for plastid-containing and non-plastid containing eukaryotes.

The description of the output format is shown below the predictions.

Table S6. Primers used in the study

Table S7. The starting OD$_{600}$ of *E. coli* CFT073 and its derivatives.

1069 **S1 Figure. Ladderised version of the Figure 1 tree**

1070 All branch support values and taxon names including subfamily identity are shown. Branch

1071 colouring is as per Figure 1. Orange stars show Uup sequences that have truncated Arm

1072 subdomains.

1073

S2 Figure. Sequence logos of domains show amino acid biases

Sequence logos of each domain HMM. The height of stacked amino acids at each position show the

information content, in bits, with letters dividing the height according to their estimated

probability. Beneath the stacked amino acids there are three lines showing probabilities; line 1 is

occupancy, the probability of observing a letter –rather than a gap - at that position. Line 2 is the

probably of seeing an insertion at that position, and line 3 is the expected length of an insertion

following that position.

1081

1082 **S4 Figure. Functional testing of inducible *B. subtilis* VmlR-HTF and VmlR-NeonGreen fusion**

1083 **proteins.** All experiments were performed in filtered LB at 37℃ in the presence of increasing

1084 concentrations of lincomycin and presented as the geometric mean ± standard deviation (n = 3).

1085 HTF stands for C-terminal His_6-TEV- 3xFLAG tag.

1086

1087 **S5 Figure. Growth and polysome analysis of *E. coli* CFT073 wild type, Δ*bipA*, Δ*abcf* and**

1088 **Δ*bipA*Δ*uup* strains, and *E. coli* CFT073 wild type overexpressing ABCFs under the control of**

1089 **constitutive P$_{tet}$ promoter.** All experiments were performed in filtered LB at either 18℃ (A) or

1090 37℃ (B-D) and growth data are presented as geometric means ± standard deviation (n = 3).

1091

S6 Figure. Functional testing of FTH-tagged *E. coli* ABCF wild type and EQ$_2$ proteins.

Western blot (A) and growth assays (B and C) N-terminally FTH-tagged *E. coli* ABCF proteins

expressed under the control of arabinose-inducible P$_{BAD}$ promoter. Growth experiments were

performed in MOPS media at 18℃ and presented as geometric means ± standard deviation (n = 3).

Western blotting was performed either at 18°C or 37°C. FTH stands for N-terminal 3xFLAG-TEV-

His$_6$ tag.

1098

S7 Figure. AREs show variability in presence across species of the same genus, which can be

used to predict novel ARE ABCFs. Genera were selected that contained more than 20 species in

the ABCF database. For each subfamily in each of those genera, the percent of species in which that

subfamily is found, is plotted. Colder colours are those ABCFs that are more universal, as typical of

housekeeping genes. AREs on the other hand have a more patchy distribution, as shown by their

tendencies for warmer colours. Antibiotic producing genera are the exception, where AREs and

ARE-like subfamilies can be universal within a genus.

1106

1107   **S3 Figure. Maximum likelihood phylogeny of eukaryotic subgroup-type ABCFs from**

1108   **eukaryotes and viruses.**

1109   The tree is rooted with bacterial YheS sequences. Branch support is from 100 bootstrap replicates

1110   and branch length is proportional to the number of amino acid substitutions (see lower scale bar).

1111   Names are coloured by subfamily.

1112

1113   **S1 Text. Supplementary sequence alignments and phylogenetic trees.**

1114   The file contains sequence alignments used to generate Figures 1 (and S1), 2 and S3 in FASTA

1115   format, and Newick format phylogenetic trees that are not shown as figures. Each entity

1116   (alignment or tree) is separated by comments preceded by "#".

1117

1118   **S1 Materials and Methods.**

1119   **Supplementary methods describing the construction of plasmids and bacterial strains.**

1120

1121

Fig. 1 | The family tree of ABCFs has a bipartite structure corresponding to eukaryotic-like and bacterial (and organellar)-like sequences

The tree is a RaxML maximum likelihood phylogeny of representatives across the ABCF family with branch support values from 100 bootstrap replicates with RaxML (MLB), 1000 ultrafast bootstrap replicates with IQ-TREE (UFB) and Bayesian inference posterior probability (BIPP). The inset box shows the legend for subfamily and intersubfamily support; support values within subfamilies and that are less that 60% MLB are not shown. Species were chosen that sample broadly across the tree of ABCF-encoding life, sampling at least one representative from each subfamily. Green shading shows the eukaryotic type ABCFs; other subgroups are bacterial unless marked with a green shaded circle to indicate eukaryotic groups with potentially endosymbiotic origin. CpYdif contains both cyanobacterial and predicted chloroplast sequences. The full tree with taxon names and sequence IDs is shown in fig. S1. Branch lengths are proportional to amino acid substitutions as per the scale bar in the lower right. The asterisked branch is not supported by this data set, however it is supported at 85% MLB in phylogenetic analysis of the eukaryotic subgroup and its viral relatives, rooted with YheS (fig. S3). Branch lengths are proportional to amino acid substitutions as per the lower right scale bar.

Fig. 2 | Rooting with ABCE shows eukaryotic-like ABCFs nesting within bacterial-like ABCFs, with YheS as the sister group to the eukaryotic-like clade

Maximum likelihood phylogeny of representatives across the ABCF family, and ABCE sequences from the UniProt database. Branch support from 200 bootstrap replicates with RaxML (MBP), 1000 ultrafast bootstrap replicates with IQ-TREE and Bayesian inference posterior probability is indicated with the key in the inset box. Branch lengths are proportional to amino acid substitutions as per the inset scale bar.

Fig. 3 | Typical domain and subdomain architectures of ABCFs

1150   (A) Boxes show domains as predicted by HMMs. Full coordinates and sequence data for these

1151   examples are recorded in Table S3. Dotted lines shows possible interactions between the HEAT

1152   domain of eEF3/New1/eEF3L and the N-terminal domain of ABCF3, ABCF4 and ABCF7. (B)

1153   Predicted coiled coil regions of *E. coli* YheS along the protein length. Inset: cartoon representation

1154   of the coiled coil subdomains protruding from the core ABC domains.

1155

Fig. 4 | *B. subtilis* ARE VmlR is a cytoplasmic protein that directly protects the ribosome from

antibiotics

(A) Growth of wild type *B. subtilis* 168, isogenic Δ*vmlR* knockout as well as Δ*vmlR* knockout

expressing either wild type or EQ$_2$ version of VmlR under the control of IPTG-inducible P$_{hy\text{-}spank}$

promoter. Six biological replicates were averaged for each growth curve and the data presented as

geometric means ± standard deviation. (B) Polysome analysis and western blotting of Δ*vmlR B.*

subtilis expressing C-terminally HTF-tagged wild type and EQ$_2$ version of VmlR. (C) Phase contrast

and fluorescence images of uninhibited *B. subtilis* cells expressing VmlR-mNeonGreen (VmlR-mNG)

in the presence and absence of lincomycin (40 min incubation with 5 μg/ml), and a model

transmembrane protein WALP23-GFP are shown for comparison. (D) Fluorescence intensity

profiles were measured perpendicular to the cell length axis along a 325 nm wide and 5.8 μm long

line as indicated. Fluorescence intensity profiles of cells expressing WALP23-GFP [65], and cells

expressing VmlR-mNG in the presence and absence of lincomycin. The graph depicts the average

fluorescence intensity profiles and the corresponding standard deviations (n = 30).

1170

Fig. 5 | Overexpression of *E. coli* ABCF Uup suppresses cold sensitivity and ribosome

assembly defects caused by loss of translational GTPase BipA

Growth (A) and sucrose gradient polysome analysis (B) of CFT073 wild type, isogenic Δ*bipA* and

Δ*bipA*Δ*uup*, as well as CFT073 Δ*bipA* transformed with low-copy pSC vector expressing either BipA

or ABCFs EttA, Uup, UheS and YbiT under control of constitutive promoter P$_{tet}$. All experiments

were performed in filtered LB at 18°C and data are presented as geometric means ± standard

deviation (n = 3).

1178

Fig. 6 | Expression of *E. coli* ABCF-EQ$_2$ mutants inhibits growth and protein synthesis

Growth of wildtype *E. coli* BW2513 transformed with pBAD18 vector (grey trace) as well as *E. coli*

BW2513 expressing either wild type (black trace) or EQ$_2$ mutants (red trace) of EttA (A), Uup (B),

YbiT (C), and YheS (D) under the control of arabinose-inducible promoter P$_{BAD}$. Radiographs show

the effect of wild type and EQ$_2$ ABCF expression on protein synthesis, as probed by pulse labeling

with L-[^{35}S]-methionine. Expression was induced by the addition of L-arabinose to a final

concentration of 0.2% at time point 0, and efficiency of incorporation was quantified by

scintillation counting and visualised by autoradiography at 0 and 20-minute time points.

Scintillation counting data are presented as geometric means ± standard deviation (n = 3). All

experiments were performed at 37°C in Neidhardt MOPS medium [105] supplemented with 0.4%

glycerol as a carbon source. The inset cartoons are a representation of ABCF domains and sub-

domains, as per the legend in the lower box.

1191

Fig. 7 | AREs tend to have relatively long linker regions that potentially extend towards the

ribosome bound antibiotics

(A) The structure of EttA and its interacting ribosomal components from PDB 3J5S [17] is shown

alongside homology models of *Staphylococcus aureus* VgaA and *Entercoccus faecalis* LsaA, using

3J5S as the template, with *de novo* modeling of the linker regions. The dotted circle shows the

relative location of PTC-inhibiting antibiotics. Arm and linker regions are shaded in yellow and

turquoise respectively. (B) Extracts from the multiple sequence alignment of *E. coli* and *B. subtilis*

ABCFs, and representative AREs, containing the Arm (yellow shading) and Linker (turquoise

shading) subdomains. Alignment numbering is according to the EttA sequence. A boxed region

shows a region that is particularly rich in proline and polyproline in various ABCF family members.

1202

ARE4

ARE5

BAF2

AAF4

AAF5

AAF1

AAF3

BAF3

AAF2

PAF1

BAF1

ARE1

ARE3

ARE6

SAF1

ARE2

FAF1

YfmM

FAF2

ARE7

/ =YdiF

bdAF1

cpYdif

DAF1

DAF3

AAF6

DAF2

YbiT

algAF1

YheS

* = ABCF7

ABCF3

ABCF6

ABCF1 (ABC50)

ABCF5

ABCF4

/ =eEF3L

ABCF2 (Arb1)

New1

eEF3

Uup

algUup

EttA

mEttA

algEttA

★ Uup sequences with deleted arm

Group and inter-group branch support:

● 100% MLB 100%, UFB, 1.0 BIPP,
present in all uncoupled domain trees

● 85-99% MLB, >90% UFB, 1.0 BIPP

○ 70-84% MLB, 80-89% UFB, <1.0 BIPP

⊘ 60-69% MLB, 80-89% UFB, 1.0 BIPP

—— 0.1 substitutions

A

−IPTG |
- ΔvmlR + VmlR-EQ₂$^{Phy-spank}$ (red dotted)
- ΔvmlR + VmlR$^{Phy-spank}$ (black dotted)

+IPTG |
- ΔvmlR + VmlR$^{Phy-spank}$ (black solid)
- ΔvmlR + VmlR-EQ₂$^{Phy-spank}$ (red solid)

OD_{600}

Time, h

wt 168
ΔvmlR
37°C, LB

B

LB + 30 μM IPTG
37°C

70S

A_{260}, a.u.

ΔvmlR +
VmlR$^{Phy-spank}$
VmlR-EQ₂$^{Phy-spank}$

Polysomes

30S

50S

top
7%

bottom
35%

VmlR
L3
VmlR-EQ₂
L3

C

VmlR-mNG^{Pxyl} WALP23-GFP^{Pxyl}

37°C, LB + xylose

untreated   + Lnc 5 μg/mL

D

VmlR-mNG

WALP23-GFP

Fluorescence, a.u.

Length, μm

VmlR-mNG^{Pxyl} + Lnc   VmlR-mNG^{Pxyl} WALP23-GFP^{Pxyl}

A

EttA · VgaA model · LsaA model

(Labels within figure: Linker, P site tRNA, H86, L1, Arm, L1 stalk)

B

proline hotspot