Suboptimal eye movements for seeing fine details

Mehmet N. Ağaoğlu[*], Christy K. Sheehy[3], Pavan Tiruveedhula[1], Austin Roorda[1,2], Susana T.L. Chung[1,2]

[1]School of Optometry, University of California, Berkeley

[2]Vision Science Graduate Group, University of California, Berkeley

[3]Department of Neurology, University of California, San Francisco

*corresponding author, lead contact: mnagaoglu@gmail.com

Abstract

Human eyes are never stable, even during attempts of maintaining gaze on a visual target. Considering transient response characteristics of retinal ganglion cells, a certain amount of motion of the eyes is required to efficiently encode information and to prevent neural adaptation. However, excessive motion of the eyes leads to insufficient exposure to the stimuli which creates blur and reduces visual acuity. Normal miniature eye movements fall in between these extremes but it is unclear if they are optimally tuned for seeing fine spatial details. We used a state-of-the-art retinal imaging technique with eye tracking to address this question. We sought to determine the optimal gain (stimulus/eye motion ratio) that corresponds to maximum performance in an orientation discrimination task performed at the fovea. We found that miniature eye movements are tuned, but may not be optimal, for seeing fine spatial details.

We make large, rapid, voluntary eye movements – saccades, to redirect our gaze to accomplish
numerous visual tasks (e.g., searching for an object, reading a book, etc.). This is to form a fine-grained
representation of the external world by taking advantage of a part of the retina – the fovea, which has
the highest spatial resolution. However, our eyes are always in motion between epochs of saccades,
even when we try to maintain our gaze on an object. Miniature eye movements that we make during
fixation are often referred to as fixational eye movements (FEM). Different types of FEM have been
identified depending on their spatiotemporal characteristics[1,2]. Microsaccades are small jerky eye
movements, and share similar peak velocity-amplitude dynamics as larger saccades[3]. Drifts are
relatively slower and smoother but rather erratic eye movements that occur between (micro) saccades,
and have been usually modeled as various types of random walk or Brownian motion[4–6]. Lastly,
tremors are usually defined as very low-amplitude and high-frequency oscillatory movements that are
superimposed on drifts[7–9].

In addition to the non-uniform distribution of density and size of receptive fields of ganglion cells across
the retina[10–12], there is another unique property that differentiates our visual system from a
computer vision system – neural adaptation. For instance, retinal ganglion cells (RGC) are most
responsive to light transients and their responses decay with prolonged exposure[13,14]. Although such
a system is ideal for the detection of changes or movements that are crucial for survival in a natural
setting, it comes with a consequence. It has been known that in the absence of FEM, visual perception
fades away, and more so for small visual stimuli[15–17]. This suggests that retinal image motion is
essential for continuous and high-acuity vision. However, if FEM are too large or too fast, the light
intensity defining a visual stimulus will spread over a large population of cells each with an insufficient
exposure to the stimulus, resulting in nothing but smeared, ghost-like impressions. Therefore, there
must be an optimum movement between these two extremes whereby visual perception is maximized
such that the ability to perform a visual task is highest at that particular movement.

Previous research showed that naturally occurring or artificially induced irregular and continuous retinal
image drifts help in seeing fine spatial details[6,18]. Likewise, naturally occurring microsaccades or
sudden jumps of stimuli are known to counteract visual fading[19,20], and help redirect our gaze to
compensate for the non-homogeneous acuity within the fovea[21]. If there is a causal relationship
between FEM and visual perception, the latter should show a "tuning" function with different levels of
the former, analogous to orientation tuning of cells in the early visual areas where firing rate of a given
cell is continuously modulated by how close the orientation of a stimulus in its receptive field to its
"preferred orientation." In other words, direct manipulation of FEM, or the way retinal image moves as
a function of FEM should result in systematic changes in visual perception, where maximum
performance in a visual task would be obtained at the preferred or optimal FEM. From an evolutionary
point of view, FEM in normal vision can be thought as optimally tuned for seeing fine details at the
fovea. Here, we explicitly tested this hypothesis. Our results show that normal FEM are tuned, but not
quite optimal, for fine discrimination at the fovea. We also found that within the range of spatial
frequencies where the human visual system has highest contrast sensitivity, this relationship disappears
suggesting a higher tolerance for retinal image motion for coarse visual structures.

Results

Using a tracking scanning laser ophthalmoscope (TSLO)[22], we presented seven human subjects with a
high spatial frequency grating (12 cpd) for 900 ms while imaging their retina and tracking their eye
movements in real time (**Fig. 1a, b**). We systematically manipulated the way retinal image motion and
the actual FEM are related. The motion of the stimulus on the scanning raster was a function of the
estimated eye motion times a *gain* factor (**Fig. 1d**). A gain of 0 means that the stimulus position

remained fixed relative to the raster but slipped across the retina based on natural FEM. A gain of 1
means that the stimulus was stabilized on the retina, i.e., the retinal image motion due to FEM was
completely cancelled out. A gain of 0.5 refers to partial stabilization, i.e., the stimulus moved only half as
much as the eye motion. Assuming similar oculomotor behavior under different gains, a gain of -1
doubles the retinal slip of the stimulus compared to that under natural viewing (i.e., gain = 0), and a gain
of 2 results in the same retinal slip but in the opposite direction of what would occur under natural
viewing. Offline analyses of retinal videos for eye movement extraction revealed that eye tracking and
stimulus delivery were performed with near-perfect accuracy (~99%) for complete retinal stabilization
(gain = 1). For gain conditions other than 0 and 1, there was some trial-to-trial variability in accuracy of
stimulus delivery (**Fig. 1e, f**). Nevertheless, each gain condition resulted in a statistically distinct
distribution of effective gains centered at the desired gain (**Fig. 1e, f**). We measured subjects' ability to
discriminate the direction of the grating's orientation from vertical under different gain conditions. If
FEM are not tuned for fine discrimination at the fovea, then performance should not depend on gain
(the null hypothesis, **Fig. 1c**). On the other hand, if the retinal image motion due to FEM is tightly tuned
for fine discrimination, performance should manifest a non-monotonic relationship with gain, where a
particular value of gain results in the best (or worst) performance (**Fig. 1c**). The tuning hypothesis can
hold true in various ways. If FEM are optimal for fine discrimination at the fovea, then visual
performance should peak at the gain of 0. Alternatively, retinal image motion might be the primary
determinant of visual performance. If retinal image motion is always detrimental for seeing, visual
performance should be highest at the gain of 1. Retinal motion might also be beneficial regardless of the
underlying FEM. If that is the case, the lowest discrimination performance should occur when the
stimulus is fully stabilized on the retina.

Figure 1 *Manipulating the relationship between retinal image motion and eye motion with the TSLO. (a) An orientation discrimination task at the fovea. Subjects' view of a grating on the raster (left), and corresponding retinal image (right). Note that the stimulus is imprinted on the retinal image. (b) The luminance profile used to create grating patterns on the raster. The mean luminance of the grating was set to ~70% of the background and the contrast of grating was adjusted for each subject. (c) Predictions from the no tuning (null) and tuning hypotheses. The panels with blue and red outlines show various ways tuning can occur. (d) Sample eye motion and retinal image motion traces (black lines) and corresponding probability densities (red clouds) for different gains. The horizontal and vertical lines in the lower left corner of each panel represent 0.1°. Dimensions were adjusted for clarity. (e) Retinal ISOA as a function of eye ISOA across gains in Experiment 1. Different colors represent different gains, and subjects are coded by different symbols. Inset shows a close-up view of data for smallest retinal/eye motion. (f) The distribution of retinal/eye ISOA ratios for different gains, averaged across seven subjects. Vertical dotted lines show theoretical ISOA ratios, i.e., assuming that eye tracking, stimulus delivery, and offline eye movement extraction were perfect. Error bars represent ±SEM (n=7). Color conventions for gains are identical across all figures.*

We found that orientation discrimination performance is tuned to gain (**Fig. 2a, b**). Averaged across subjects, the peak performance occurred at a gain of 0.43 (95% confidence intervals: 0.12, 0.74), suggesting that partially reducing the effects of FEM is actually helpful in seeing fine spatial details. Results were similar when data from each subject were fitted separately (polynomial: t_6=3.165, p=0.019; Gaussian: t_6=2.600, p=0.041) (**Fig. 1b** and **Figure S1**). The exact choice of the tuning model (a quadratic

 polynomial or a Gaussian) did not matter (paired t-test: $t_{12}=3.165$, p=0.019). Bootstrapping tuning curve
fits to binary data (correct vs incorrect) also revealed no optimality in six of the seven subjects (**Figure S1**). These results suggest that FEM are tuned but not optimal for fine discrimination at the fovea, at least within the range of parameters investigated here.

Figure 2 *FEM are tuned, but not optimal, for fine discrimination at the fovea. **(a)** Proportion correct as a function of gain, averaged across subjects, in Experiment 1. A Gaussian tuning function was fit to all data (black curve) to estimate the optimum gain. Vertical white line represents optimal gain defined as the gain corresponding to the peak of the Gaussian. Shaded regions represent 95% confidence intervals of the optimum gain. **(b)** (Top) Average optimal gains based on individual tuning function fits along with individual optimal gains. A quadratic polynomial and a Gaussian tuning function resulted in statistically indistinguishable optimal gains. (Bottom) To compare the "no tuning" and "tuning" hypotheses in terms of how well they can explain our data, we computed Adjusted R^2 metric for the constant model and tuning models (a quadratic polynomial or a Gaussian), respectively. For all subjects, tuning models performed better. **(c)** (Top-right) the distribution of preferred retinal locus (PRL) across trials for one representative subject. Each symbol represents one trial. (Bottom-left) A close-up view of the central ~2.5° part of the retina. Note the systematic change in PRLs across gains. **(d, e)** Retinal image motion and eye motion ISOA as a function of gain. **(f, g)** Microsaccade rates and PRL eccentricity across gains. Optimal gain and confidence intervals in (a) are replotted in (d, e, f, and g). Error bars represent ±SEM (n=7).*

In order to check whether or not the tuning between performance and gain is limited only to fine
discrimination tasks, we repeated the experiment with a spatial frequency (3 cpd) at which the human
visual system has highest contrast sensitivity for static displays(23). The hypothesis was that the retinal
jitter due to FEM causes much less modulations in retinal ganglion cells with low spatial frequency
stimuli, therefore, gain manipulations should result in minimal or no change in performance. We found
no effect of gain on performance (**Fig. 3a**), despite the retinal image motion varied over two log units
across conditions (**Fig. 3d**). Different performance trends with gain in Experiments 1 and 2 cannot be
explained by differences in retinal motion, eye motion, microsaccade rate, or PRL eccentricity (**Fig. 2d-g**
vs **Fig. 3d-g**).

*Figure 3 FEM are not tuned for coarse discrimination at the fovea. (a) Proportion correct as a function of
gain in Experiment 2. (b) Average optimal gains based on individual tuning function fits along with
individual optimal gains. (c) The distribution of retinal/eye motion ISOA in Experiment 2. (d, e) Retinal
image motion and eye motion ISOA as a function of gain. (f, g) Microsaccade rates and PRL eccentricity
across gains. Optimal gain and confidence intervals in (a) are replotted in (d, e, f, and g). Error bars
represent ±SEM (n=7). Conventions are as in **Figure 2**.*

Next, we sought to determine what drives the strong dependency between performance and gain with
high spatial frequency gratings. If gain manipulation only modulates the retinal image motion, then the
answer would simply be retinal image motion, assuming no interference from extra-retinal mechanisms.
The approach taken in most retinal stabilization studies in the literature implicitly assumes that gain
manipulation only results in changes in retinal image motion. In other words, retinal image motion is
considered as the one and only mediator of performance. However, we found that gain modulates
multiple mediators. We computed two-dimensional probability density of stimulus locations on the

retina and eye positions on the raster (e.g., **Fig. 2d**), and quantified, on a trial-by-trial basis, the extent of
retinal image motion and eye motion by the isoline area (ISOA) containing roughly 68% of the
retinal/eye motion traces (**Fig. 2d, e** and **Figure S1**). As expected, the minimum retinal ISOA occurred
when the stimulus was stabilized on the retina (i.e., gain = 1) but the pattern of changes in retinal ISOA
as a function of gain revealed an asymmetric "V" shape around the gain of 1 (**Fig. 2d**). This asymmetry
can be explained by differences in oculomotor behavior of subjects across different gains. More
specifically, consistent with previous literature(24), subjects made smooth pursuit-like eye movements
for gains of 1 and larger, which resulted in larger eye ISOAs (**Fig. 2e**). This change in behavior occurred as
soon as the retinal slip is no longer in a direction that is consistent with eye motion, in line with recent
perceptual observations(25). In addition, subjects made slightly more microsaccades for negative gains
where retinal image motion is amplified (**Fig. 2f** and **Figure S1**). In addition, although each trial started
with a fixation cross at the center of the raster, the preferred retinal locus (PRL) during grating
presentation, defined here as the retinal location corresponding to peak probability density of retinal
stimulus locations, also changed with gain (**Fig. 2c, f, g** and **Figure S1**). To determine what really drives
the relationship between gain and performance, one must take these mediators into account. In a
regression-based mediation analysis following the most commonly used four-step approach(26), we
found that (i) gain has a significant effect on performance, (ii) gain significantly modulated all four
mediators (retinal ISOA, eye ISOA, microsaccade rate, and PRL eccentricity), (iii) all mediators
individually, with the exception of microsaccade rate, are significant predictors of performance, (iv) gain
remains a significant predictor of performance even when the effects of all significant mediators are
taken into account (**Fig. 4**). In order to determine whether or not mediators can account for the data as
well as gain by itself, we performed a series of linear-mixed effects regression analyses (**Figure S4**). In
terms of explained variance and log-likelihood, where number of factors is not penalized, several purely
mediator-based models could surpass the models based on gain only, suggesting that mediators
identified here might fully account for how gain modulates performance. However, as the Bayes
Information Criterion (BIC) differences show, none of the mediator-based models could outperform the
simple model that is based only on gain. Through additional regression analyses and model comparisons
using BIC, we confirmed that performance cannot be fully accounted by mediators alone (**Figure S4**).

Figure 4 *Teasing apart contributions of different mediators.* ***(a)*** *The first step in mediation analysis is to*
establish a significant relationship between gain (G) and proportion correct (PC). Since the tuning
hypothesis predicts a quadratic relationship between G and PC, we included the G^2 in our regression
analyses. ***(b)*** *Second, whether or not gain is a significant predictor of each covarying factor (Ret: retinal*

*ISOA, Eye: eye ISOA, PRL: PRL eccentricity, MR: microsaccade rate) is established. **(c)** The third step tests*
*separately for a significant effect each mediator on performance. **(d)** Finally, gain and mediators with a*
significant correlation on performance are used to explain performance. Red and blue colors represent
statistically significant negative and positive effects whereas gray lines represent insignificant
*relationships. The final model in **(d)** shows that even when all significant mediators are taken into*
*account, gain still has a significant effect on performance. **(e)** When all mediators are included,*
*regardless of the outcome of **(c)**, gain remains to be a significant factor. Thickness of each line represent*
the absolute value of the standardized effect size.

Discussion

"Tuning" refers to a relationship between an independent variable and an outcome measure, where a
certain level of the former is more preferable than others. Optimality in this context refers to achieving
the best possible outcome in the face of several antagonist factors. Throughout the vast literature on
FEM and visual perception, the word "optimal" has been used quite liberally in regard to spatiotemporal
properties of FEM(1,27–32) although there has never been an explicit test for addressing it. Here, we
tested whether visual performance in a fine orientation discrimination task would show tuning as a
function of the relationship between the retinal image motion and actual eye movements. We found
strong tuning for a fine-detail discrimination task (Experiment 1) but not for a coarse discrimination task
(Experiment 2). The absence of tuning in Experiment 2, despite up to a two log-unit change in retinal
motion across conditions, suggests a very high tolerance for motion. Surprisingly, the optimal gain in
Experiment 1 was obtained at a gain value between 0 and 1, suggesting that partially compensating for
FEM can be beneficial.

Our results might seem inconsistent with previous reports where complete retinal stabilization resulted
in impaired discrimination performance(6,18). A simple interpolation between the two extremes
suggests a monotonic impairment in visual performance with better compensation for FEM. This
apparent inconsistency may not be real. First, it is technically possible to get impaired performance with
complete stabilization *and* a nonzero optimal gain at the same time (which was the case for five out of
seven subjects, **Figure S1**). Second, none of the existing studies explored the range of gains used here
for discrimination tasks at the fovea. In addition, in previous studies, fading that resulted from retinal
stabilization was quantified by threshold elevations, but the degree to which fading occurs depends on
many variables such as stimulus duration, size, contrast, eccentricity, equipment used, etc.(16,33–36).
Early studies on retinal stabilization used small stimuli extending only a few arcmin, and was closely
surrounded by other visual cues coming from the apparatus(8,16,17). More recent studies used foveally
presented gratings extending several degrees of visual angle far from display boundaries(6,21), or
parafoveally presented diffraction-limited stimuli covering only a few cones within a visible raster
covering 1-1.3 deg(18). The paradigm used here was somewhere in between; we presented through
natural optics of the eye a grating that covers the fovea and is situated within a visible raster covering 10
deg. Therefore, it is possible to make qualitative comparisons across aforementioned studies, however,
it is not feasible to extrapolate previous studies to the conditions investigated here.

Our results are highly consistent with recent theoretical work that has been successfully used to account
for performance impairment due to retinal stabilization(5,37,38). According to this framework, there are
two distinct mechanisms that work in tandem, one for estimating FEM from RGC responses across the
retina which negates the need for an extra-retinal mechanism to properly decode spatial information,
and another one for making an *optimal* inference about the spatial layout of the stimuli. The presence of
a global motion compensation mechanism for FEM was demonstrated by a striking visual illusion(39).
Surprisingly, when receptive field size and density across the retina and the statistics of FEM under

normal viewing conditions are factored in, this model predicted that normal human FEM are not optimal
for high acuity tasks(5,37). This theory also predicts that larger stimulus sizes and peripheral cues would
improve discrimination at the fovea since estimating FEM would be easier and more accurate in these
conditions. It is possible that the absence of optimality might have arisen since the scanning raster was
always visible in the present work. Although several lines of evidence against this prediction have been
presented(40), they turned out to be lacking technical precision and proper controls to directly test this
prediction(41).

We have identified several mediator factors that could explain a significant portion of the variability in
the data. Note that the presence of these mediators is not due to the equipment used or stimulus
parameters, but reflects the inevitable consequence of foveal presentation of the stimuli. None of these
mediators have been reported quantitatively or used to account for data in the previous literature about
the roles of FEM. Parafoveal (or peripheral) presentation of stabilized stimuli may not activate all of the
aforementioned mediators (e.g., eye ISOA), however, non-foveal presentation of stimuli would defeat
the purpose of this study since one cannot make strong inferences about foveal viewing with
peripherally presented stimuli. Alternatively, an experiment where stimulus moves in an incongruent
manner to avoid chasing can be performed, however, it is unclear whether or not small amplitudes of
stimulus motion would still lead to pursuit-like eye movements. The way we chose to address what
factors underlie the tuning between performance and gain reported here is to perform a mediation
analysis(42). This analysis showed that even when retinal motion, eye motion, PRL eccentricity, and
microsaccade rate were factored in, gain still had a significant direct effect on performance. This finding
suggests that (i) there are additional mediators not considered here, or (ii) "post-retinal" factors such as
changes in attentional engagement in the task depending on gain value might be at play.

In order to assign extra-retinal factors a role for perception during FEM, one needs to factor out all
possible retinal factors such as retinal ISOA, velocity, acceleration, PRL eccentricity, initial retinal
position of the stimuli, etc. Obviously, these factors are not independent from each other, limiting the
use of mediation analysis described here. Admittedly, the optimal gain might also be affected by these
mediators. A way to compensate for their effects for the purpose of estimating optimal gain might be
normalizing performance by each mediator and then testing for tuning. However, this exacerbates the
problem since (i) whether a covarying factor is a positive mediator (reducing the effect) or a negative
one (increasing the effect) is not known a priori, (ii) the relative contribution of each mediator is
different but normalization assumes equal contribution, and (iii) each mediator has a different scale of
change across conditions, which could result in numerical instabilities and prevent accurate
determination of the optimal gain. Point (iii) can be addressed by log-transforming some mediators (e.g.,
retinal ISOA) and/or standardizing them, and point (i) can be addressed by using the outcome of a
mediation analysis to guide the normalization process, but point (ii) cannot be readily addressed. On the
other hand, since visual performance comes about via mediators, there may not be a need for
normalizing performance before computing optimal gains. From this perspective, they are not just
artifacts to be removed, but the actual underlying factors of visual function. The logic is that whatever
the exact value of optimal gain is, visual performance results from an interplay between various
mediators, and it may not be possible to uniformly sample the multidimensional space defined by
multiple mediators. A case in point, it seems that foveal presentation of a stimulus almost always leads
to smooth pursuit-like oculomotor behaviors when the retinal projection of it is stabilized(24).

Human retina is non-homogeneous, even within the fovea. Therefore, making microsaccades to redirect
gaze to enjoy the highest acuity part of the retina is a reasonable strategy(21,28). Microsaccades are not
always initiated voluntarily, however, and recent studies claimed that they often occur after a period of
low retinal slip and are executed to avoid fading(43). Their occurrence seems to be coupled to heartbeat

as well(44). Based on the finding that microsaccades cause widespread activity across the visual system
and help temporally synchronize neural activity, some researchers supported the view that
microsaccades, among other FEM, contribute most to visual function(20,45,46). Although the rate of
microsaccades in our experiments was very low, we still performed a series of analyses on
microsaccades made by all observers to test these hypotheses (**Figure S2**). The low rate of
microsaccades itself, especially when retinal image motion was minimized, is an evidence against a
primary role for microsaccades for visual processing. In response to partial or complete retinal
stabilization for instance, subjects made larger drifts rather than larger or more frequent microsaccades.
Moreover, we found evidence for both low retinal slip and gaze redirection, although the evidence for
the latter was stronger. More specifically, we found that retinal image velocity was slightly reduced
immediately before microsaccade onset, and most microsaccades were made to bring the retinal
projection of the stimuli closer to the PRL.

There are several other facts to be considered when functional roles of drifts and microsaccades are to
be determined. First, RGCs are most responsive to light transients, and the time constant of their
responses can vary from 30 to 100 ms(47). Second, although the initial burst activity of RGCs in response
to a light transient is highly precise, prolonged presentation breaks this temporal synchrony, and the
tonic neural activity demonstrates quite a bit of variability(48,49). Encoding spatial information using a
rate code with a few spikes necessitates the accumulation of information over time to improve the
signal to noise ratio. The presence of FEM makes encoding of spatial information via rate coding even
less reliable by further increasing variability in spiking activity. Third, FEM create retinal motion signals
that are well beyond motion detection thresholds but not perceived.

From an evolutionary standpoint, it is unclear which of the facts listed so far was the root cause for the
others. For instance, whether RGCs prefer light transients and do not respond as strongly after
prolonged presentation due to FEM, or FEM exist due to the temporal characteristics of RGC responses
is a hard problem to address. In addition, a recent modeling work demonstrated potential alternatives
to spatial encoding via rate coding, where FEM do not pose problems to be solved by the visual system,
but instead they are part of the solution to efficient information encoding(32). This also renders the
mystery of how the visual system differentiates motion due to FEM from those of external objects a
non-issue since if we actually see via FEM, why correct for them? In fact, drifts transform the spectral
content of retinal stimulation into spatiotemporal frequencies to which the early visual system is most
sensitive(31), but it is unclear whether this is an epiphenomenon or a result targeted by an active and/or
adaptive process. However, the current implementation of this model relies on weak assumptions, one
of which is that drifts are cyclic (sinusoidal) motions (to drive phase-locking mechanism) within time
courses that reflect average fixation duration (~300 ms). Except in very few instances, we did not
observe such patterns (**Figure S3**).

Some visual/cortical impairments (e.g., amblyopia, central vision loss) result in "abnormal" FEM(50,51).
In a computer vision system with limited spatial resolution or blurry optics, it is theoretically possible to
achieve "super-resolution" or de-blurring by moving a sensor array. Therefore, we think that to classify
FEM as abnormal, one needs to consider several factors such as the amount of blur, receptive field sizes,
and contrast sensitivity at the PRL. Otherwise, a genuine strategy of a perfectly normal oculomotor
system might be misinterpreted as an artifact. In the case of central vision loss, the use of peripheral PRL
leads to changes in all these factors, and it is quite possible that apparently abnormal FEM in these
patients might be a way to compensate for these changes. In fact, recent studies on the effects of retinal
image motion in peripheral vision reported improvements in reading and discrimination performance
with increased motion(52,53).

The statistics of FEM may change when a subject's head is restrained compared to head-free
viewing(54). The amplitudes of FEM increase under free viewing. Measurements using a Dual-Purkinje
tracker showed that drifts from the two eyes show minimal correlation under head-fixed conditions, and
they become mostly conjugate under head-free conditions. However, retinal imaging via a binocular
TSLO revealed almost complete conjugacy under head-fixed conditions(55). Nonetheless, the conditions
reported here may demonstrate a special case of oculomotor control, which need not be optimized
since outside the laboratory, we always view the environment with freely moving body and head. In
addition, since the stimulus presentation was monocular in the present study, it remains to be seen
whether similar tuning functions would be obtained with binocular presentation. It may be that
binocular viewing increases the tolerance of the visual system to retinal image motion due to FEM even
for high spatial frequencies due to redundancy from the second eye. Finally, the different patterns of
results in the two experiments reported here also suggest that the relationship between FEM and
frequency might be a continuum, from no tuning to optimal tuning with increasing spatiotemporal
frequency. Future endeavors along these lines will require denser sampling of the frequency space as
well as accurate eye tracking combined with fast stimulus delivery to both eyes.

Methods

Participants

Seven human subjects (including the first author, S1) with normal or corrected-to-normal vision (20/20
or better in each eye) participated in the study. All seven subjects took part in Experiment 1. Three of
the seven subjects participated in Experiment 2. All subjects, except the first author, were naïve as to
the purpose and the details of the experiments. All subjects gave written informed consent prior to the
experiments. All experimental procedures were approved by the Institutional Review Board at the
University of California, Berkeley.

Apparatus

For stimulus delivery and eye tracking, we used a custom-built tracking scanning laser ophthalmoscope
(TSLO)(22). The TSLO has a diffraction-limited optical design, provides high-fidelity imaging of the retina,
and more importantly, offers online tracking of eye movements. For the experiments presented here,
we used a 10 x 10 deg^2 (512 x 512 pixels^2) field of view (FOV), which yielded a pixel size of 1.17 arcmin. A
large FOV enabled us to capture videos with rich retinal structure which, in turn, allowed accurate
image-based eye tracking and stimulus delivery. The horizontal scanner operates at 16 kHz whereas the
vertical scanner operates at 1/512 of this rate to record full frames at 30 frames per sec. An 840 nm
super luminescent diode with a 50 nm bandwidth was used to scan the retina. Visual stimuli were
delivered by manipulating the laser beam via an acousto-optic modulator with the output controlled by
a 14-bit digital to analog converter. Therefore, the stimuli had a negative contrast on the dim red raster
created by the scanner (i.e., appeared black on a red background). Details of online eye movement
tracking have been reported elsewhere (56–58). Briefly, each frame was broken into 32 horizontal strips
(16 x 512 pixel^2) and each strip was cross-correlated with a reference frame acquired earlier. The
horizontal and vertical shifts required to match a strip to the reference frame represent a measure of
the relative motion of the eye. This method results in an eye movement sampling rate of 960 Hz. These
computations occur in near real-time (with 2.5±0.5 ms delay) and allows accurate stimulus delivery at
specific retinal locations.

Stimuli and Procedures

The task was to report the orientation of a sinusoidal grating from vertical (2AFC, clockwise or
counterclockwise). The amount of tilt was ±45° from vertical and the spatial frequency of the grating

was 12 cpd in Experiment 1 (**Fig. 1a**) and 3 cpd in Experiment 2. Prior to each experiment, the contrast of
the grating was adjusted for each subject to yield ~80% correct discrimination performance under the
natural viewing condition (i.e., gain = 0). The size of the grating was 3 x 0.75 deg^2. Horizontal edges were
smoothed using a cosine profile. Mean luminance (as measured indirectly from laser power) of the
grating was kept at ~70% of the luminance of the scanning raster for all subjects, regardless of the
contrast of the grating (**Fig. 1b**). All experiments were performed in a dark room, and subjects were dark
adapted for about half an hour prior to any data collection. For some subjects (three out of seven), pupil
of the imaged eye was dilated to maintain good retinal image quality throughout the session.

Each trial started with a fixation cross (0.2 deg) presented at the center of the raster. The experimenter
manually acquired a reference frame for online tracking at the start of each trial. Following a random
delay (up to 1 sec), the stimulus was presented for 900 ms (flickered at 30 Hz). Subjects responded via a
gamepad, had unlimited time to respond, and could have a break at any point during a block of trials.
The set of gains used were -1, -0.5, 0, 0.5, 1, 1.5, and 2. Different gains were interleaved within a block
of trials. All observers completed at least 100 trials per gain. Right after the completion of the main
experiment and before data analyses, subject S4 was tested for a second run in Experiment 1 since her
retinal image quality and head stability during the experiment was poor, causing retinal stabilization to
fail in many of her trials. In the second run, we used finer steps of gain (from -0.25 to 1.25 in steps of
0.125) and subject S4 again ran at least 100 trials per gain. However, all results presented in the main
text include only the data from the first run for S4. The results from both runs for S4 are shown in **Figure
S3**.

Retinal video analysis

Retinal videos were analyzed offline for five main reasons. First, we sought to determine how well and
where the stimulus was delivered on a trial-by-trial basis. Second, online eye tracking was performed by
using raw retinal images which were corrupted by high frequency noise and low frequency luminance
gradients. In order to get more accurate eye motion estimates relatively less dependent on changes in
overall brightness and uniformity of retinal images across trials, one needs to perform several pre-
processing steps on retinal images. To this end, we performed the following image processing steps
before computing eye motion. Trimming, detection and removal of frames during which subjects
blinked, extracting stimulus position and removal of the stimulus (replaced by random noise patterns
whose statistics –mean and standard deviation, matched to the rest of the frame), gamma correction,
bandpass filtering (for removal of high frequency noise and low frequency brightness gradients), and
making a reference frame. Third, during online eye tracking, if the peak of normalized cross-correlation
between a strip and the reference frame was below 0.3, possibly due to (i) bad image quality, (ii)
excessive distortion of image features due to a rapid eye movement, (iii) insufficient amount of overlap
between the strip and the reference frame due to large eye motion, or (iv) blinks, the stimulus was not
delivered. By offline processing of retinal videos, we also sought to inspect each and every frame of
retinal videos and discard the trials if the stimuli was delivered inaccurately or was not delivered at all in
more than two frames per trial (note that with this criterion, trials where subjects blinked were also
discarded). This procedure resulted in removal of 28.8% (1305/4525) and 8.7% (206/2353) of all trials in
Experiment 1 and 2, respectively. Fourth, since the reference frames used for online tracking were
basically snapshots of the retina taken manually by the experimenter, and since the eyes are almost
never stationary, the reference frames themselves might have some distortions due to these motions.
By offline processing, we created a relatively motion-free reference frame for each and every trial
separately in an iterative process. This process started by selecting one of the frames in a retinal video
as the reference frame, and computing eye motion. After each iteration, a new reference frame was
built by using computed eye motion and individual strips. We performed three iterations for each video,

and the reference frames made in the last iteration were used for the final computation of eye motion.
The strip height and sampling rate used for the final strip analysis were 25 pixels and 540 Hz,
respectively. Fifth, during offline analysis, we could interpolate the cross-correlation maps around where
the peak occurs to achieve subpixel resolution (one tenth of a pixel, 0.12 arcmin) in computing eye
motion.

Post-processing

Following strip analysis of individual videos, the computed eye motion traces were subjected to several
post-processing steps. First, eye motion traces were "re-referenced" to a larger reference frame created
by retinal videos recorded in a separate session where subjects were asked to fixate at different position
on the scanning raster. This essentially allowed us to capture images from different part of the retina
and tile them on a larger ("global") reference frame. Re-referencing was needed since each and every
video had a slightly different ("local") reference frame (since reference frames were created for each
video separately), and hence, the absolute values of the eye motion would differ across trials. This step
is required also for computing the absolute retinal position of the stimulus across all trials for a given
subject. Re-referencing was performed by adding a constant shift to the previously computed eye
motion traces, where the amount of shift was defined as the position of the local reference frame on
the global one. After re-referencing, eye motions were then converted to visual degrees, low-pass
filtered (passband and stopband frequencies of 80 and 120 Hz, respectively, with 65 dB attenuation in
the stopband), and median filtered (with a window of 9 samples, ~17 ms) to reduce frame-rate artifacts
(30 Hz noise and its harmonics). Filtered traces were used to compute retinal motion of the stimulus
(defined as the difference between stimulus motion and eye motion). We quantified the amount of eye
and retinal motion on a trial-by-trial basis as the 68% isoline area(59) (referred to as ISOA in the main
text), which corresponds to the area of the 0.68 cumulative probability isoline (**Fig. 1d**). This also roughly
corresponds to the area covered by 68% percent of the motion samples. In the case of retinal motion,
this metric quantifies the retinal area traversed by the stimulus in a given trial, and for eye motion, it
represents the area of the raster (in world centered coordinates) over which the eye moved. The
distribution of the ratio of retinal ISOA and eye ISOA reveals how well the stimulus was delivered in
different conditions (see Fig. X). The theoretical ratio for a given gain is defined as

$$\frac{ISOA_{ret}}{ISOA_{eye}} = sgn(1-g)(1-g)^2,$$

where g represents gain, and $sgn(.)$ represents the signum function which was introduced to
differentiate between gains that result in the same retinal motion magnitude but in opposite directions
(**Fig. 1f**, and **Fig. 3c**).

We also computed the preferred retinal locus (PRL) of the stimulus for each trial. PRL was defined as the
retinal location corresponding to the highest probability density of stimulus presence. The probability
densities were computed by the "kernel density estimation via diffusion" method(60) with a slight
modification. More specifically, the kernel bandwidth was set to one-sixth of the standard deviation of
the eye (or retinal) motion (as in Kwon et al.(61)). The median PRL in the trials where the gain was 0 (i.e.,
natural viewing) was taken as the location of the fovea and trial-to-trial PRL eccentricity was calculated
with respect to this quantity. Finally, we identified microsaccades by using a median-based velocity
threshold(62). Eye motion traces from all trials were visually inspected to ensure that microsaccade
detection was performed correctly.

The PRL estimated when gain is 0 reflects the true PRL. Since PRL is mostly determined by the position of
the stimulus at the beginning of a trial (e.g., when gain is 1, the stimulus will stay at the start position),

the estimated PRLs in other gain conditions do not necessarily reflect the preferences of the subjects.
However, they demonstrate the idiosyncratic eye movements which govern the starting position of the
stimulus. Nevertheless, to keep a consistent nomenclature, we used the term PRL across all conditions.

Statistics

In order to test the tuning and no tuning hypotheses, we fit performance with a flat line and a quadratic
polynomial (as well as a Gaussian, although polynomial and Gaussian fits produced almost identical
results), and compared the adjusted R^2 values as a metric of goodness of fit.

Due to foveal presentation of the stimuli, different gains led to different idiosyncratic oculomotor
behaviors which could not be controlled during the experiments. We quantified several covarying
factors such as retinal ISOA, eye ISOA, PRL eccentricity, and microsaccade rate. The exact choice of
covarying factors was driven by the need to account for main retinal and eye movement-related metrics.
It is possible to estimate retinal image velocity, acceleration, or components of retinal image motion
parallel or perpendicular to the orientation of the gratings. It is also possible to quantify these metrics in
multiple ways, such as by their mean, standard deviation, minimum, maximum, or any combination of
these together. However, since most of these metrics are strongly related to each other, adding
different variants of them does not add much explanatory power. In addition, eye position traces
extracted from retinal videos tend to have frame-rate artifacts, i.e., more power than normal at
temporal frequencies around the frame rate of the videos. The frame-rate artifact gets amplified for
velocity and acceleration due to differentiation, and more importantly, the severity of the effect
interacts with different gain conditions (due to changes in oculomotor behavior). Position estimates are
not influenced as much and indirectly captures the effect of its derivatives.

In order to determine how much of the effect of gain on performance is mediated by the
aforementioned covarying factors, we performed a linear-mixed effects regression-based mediation
analysis[42]. We followed the commonly-used four-step approach suggested by Baron and Kenny[26].
The aim of this analysis was to determine whether or not gain had a significant direct effect on
discrimination performance, even after taking into account the effects of mediators (i.e., significant
covarying factors). Mediation analysis can be done in many ways depending on the causal relationship
between the independent variable and mediators. When there are multiple mediators, say n, there are
$2^{(n!)}$ possible ways of decomposing total effect size. In our case, n=4, this yields 16,777,216
possibilities[63]. Since finding the best way of organizing mediators to account for data is beyond the
scope of the present work, we chose a simple case where all mediators were treated as independent
factors that are directly modulated by only the independent variable, the gain, and they did not have
interactions among each other nor with the gain. However, they were allowed to covary with subjects,
and each had a fixed slope and a random intercept to account for individual differences in mediator
values.

Author contributions

498 MNA and STLC conceived the idea and designed the experiments. MNA and STLC performed the
499 experiments. MNA analyzed the data and wrote the manuscript. CKS, PT, and AR provided technical
500 support for the TSLO system (hardware, electronics, and software), and CKS, PT, AR, and STLC reviewed
501 the manuscript.

Competing financial interests

503 MNA and STLC have no financial interest. AR, CKS, and PT hold a patent for the design of the TSLO, and
504 have financial interest in C.Light Technologies.

Acknowledgements

We would like to thank Jonathan Patrick, Arun Kumar Krishnan and Haluk Ogmen for his comments on the manuscript, and Harold Bedell for many discussions of our results. This study was supported by grants R01-EY012810 (STLC), R01-EY017707 (STLC), R01-EY023591 (AR), P30-EY003176 (core grant) from the National Institutes of Health, and UCSF-CTSI grant TL1-TR001871 (CKS).

References

1. Martinez-Conde S, Macknik SL, Hubel DH. The role of fixational eye movements in visual perception. Nat Rev Neurosci. 2004;5(3):229–40.

2. Rucci M, Poletti M. Control and Functions of Fixational Eye Movements. Annu Rev Vis Sci. 2015;1(1):annurev-vision-082114-035742.

3. Otero-Millan J, Troncoso XG, Macknik SL, Serrano-Pedraza I, Martinez-Conde S. Saccades and microsaccades during visual fixation, exploration, and search: Foundations for a common saccadic generator. J Vis. 2008 Dec 1;8(14):21–21.

4. Engbert R, Mergenthaler K, Sinn P, Pikovsky A. An integrated model of fixational eye movements and microsaccades. Proc Natl Acad Sci. 2011;108(39):E765–70.

5. Burak Y, Rokni U, Meister M, Sompolinsky H. Bayesian model of dynamic image stabilization in the visual system. Proc Natl Acad Sci. 2010;107(45):19525–30.

6. Rucci M, Iovin R, Poletti M, Santini F. Miniature eye movements enhance fine spatial detail. Nature. 2007;447(7146):852–5.

7. Ratliff F, Riggs L. Involuntary motions of the eye during monocular fixation. J Exp Psychol. 1950;40(6):687–701.

8. Ditchburn R, Ginsborg BL. Involuntary Eye Movements During Fixation. J Physiol. 1953;119(1940):1–17.

9. Ko H kyoung, Snodderly DM, Poletti M. Eye movements between saccades: Measuring ocular drift and tremor. Vision Res. 2016;122:93–104.

10. Curcio C, Sloan K, Kalina R, Hendrickson A. Human photoreceptor topography. J Comp Neurol. 1990;4(292):497–523.

11. Watson AB. A formula for human retinal ganglion cell receptive field density as a function of visual field location. J Vis. 2016;14(2014):1–17.

12. Dacey DM, Petersen MR. Dendritic field size and morphology of midget and parasol ganglion cells of the human retina. Proc Natl Acad Sci U S A. 1992;89(20):9666–70.

13. Kaplan E, Benardete E. The dynamics of primate M retinal ganglion cells. Vis Neurosci. 1999;16:355–68.

14. Benardete EA, Kaplan E. The receptive field of the primate P retinal ganglion cell, I: Linear dynamics. Vis Neurosci. 1997;14(1997):169–85.

15. Ditchburn RW, Ginsborg BL. Vision with a stabilized retinal image. Nature. 1952;170:36–7.

16. Riggs LA, Ratliff F, Cornsweet JC, Cornsweet TN. The disappearance of steadily fixated visual test

objects. J Opt Soc Am. 1953;43(6):495–501.

17.   Yarbus AL. Perception of Objects Stationary Relative to the Retina. In: Eye Movements and Vision.
Boston, MA: Springer US; 1967. p. 59–101.

18.   Ratnam K, Harmening WM, Roorda A. Benefits of retinal image motion at the limits of spatial
vision. J Vis. 2017;17((1):30):1–11.

19.   Costela FM, McCamy MB, Macknik SL, Otero-Millan J, Martinez-Conde S. Microsaccades restore
the visibility of minute foveal targets. PeerJ. 2013;1:e119.

20.   Martinez-Conde S, Otero-Millan J, Macknik SL. The impact of microsaccades on vision: towards a
unified theory of saccadic function. Nat Rev Neurosci. 2013;14(2):83–96.

21.   Poletti M, Listorti C, Rucci M. Microscopic eye movements compensate for nonhomogeneous
vision within the fovea. Curr Biol. 2013;23(17):1691–5.

22.   Sheehy CK, Yang Q, Arathorn DW, Tiruveedhula P, de Boer JF, Roorda A. High-speed, image-based
eye tracking with a scanning laser ophthalmoscope. Biomed Opt Express. 2012 Oct 1;3(10):2611.

23.   Kelly DH. Visual Contrast Sensitivity. Opt Acta Int J Opt. 1977;24(2):107–29.

24.   Poletti M, Listorti C, Rucci M. Stability of the visual world during eye drift. J Neurosci. 2010 Aug
18;30(33):11143–50.

25.   Arathorn D, Stevenson S, Yang Q, Tiruveedhula P, Roorda A. How the unstable eye sees a stable
and moving world. J Vis. 2013;13:1–19.

26.   Baron RM, Kenny DA. The moderator–mediator variable distinction in social psychological
research: Conceptual, strategic, and statistical considerations. J Pers Soc Psychol.
1986;51(6):1173–82.

27.   Skavenski AA, Hansen RM, Steinman RM, Winterson BJ. Quality of retinal image stabilization
during small natural and artificial body rotations in man. Vision Res. 1979 Jan 1;19(6):675–83.

28.   Cornsweet TN. Determination of the Stimuli for Involuntary Drifts and Saccadic Eye Movements. J
Opt Soc Am. 1956;46(11):987–8.

29.   Ditchburn R, Fender DH, Mayne S. Vision with Controlled Movements of the Retinal Image. J
Physiol. 1959;145:98–107.

30.   Gerrits HJM, Vendrik AJH. Artificial movements of a stabilized image. Vision Res. 1970 Dec
1;10(12):1443–56.

31.   Kuang X, Poletti M, Victor JD, Rucci M. Temporal encoding of spatial information during active
visual fixation. Curr Biol. 2012;22(6):510–4.

32.   Ahissar E, Arieli A. Seeing via Miniature Eye Movements: A Dynamic Hypothesis for Vision. Front
Comput Neurosci. 2012 Nov 8;6:89.

33.   Coppola D, Purves D. The extraordinarily rapid disappearance of entopic images. Proc Natl Acad
Sci. 1996;93(15):8001–4.

34.   Kelly DH. Motion and vision. I. Stabilized images of stationary gratings. J Opt Soc Am.
1979;69(9):1266–74.

35.   Kelly DH. Motion and vision. II. Stabilized spatio-temporal threshold surface. J Opt Soc Am.
1979;69(10):1340–9.

36.   Riggs LA, Tulunay SÜ. Visual Effects of Varying the Extent of Compensation for Eye Movements*.
J Opt Soc Am. 1959 Aug 1;49(8):741.

37.   Pitkow X, Sompolinsky H, Meister M. A neural computation for visual acuity in the presence of
eye movements. PLoS Biol. 2007;5(12):2898–911.

38.   Anderson AG, Olshausen BA, Ratnam K, Roorda A. A neural model of high-acuity vision in the
presence of fixational eye movements. In: 2016 50th Asilomar Conference on Signals, Systems
and Computers. IEEE; 2016. p. 588–92.

39.   Murakami I, Cavanagh P. A jitter after-effect reveals motion-based stabilization of vision. Nature.
1998 Oct 22;395(6704):798–801.

40.   Wehrhahn C. Psychophysical and physiological evidence contradicts a model of dynamic image
stabilization. Proc Natl Acad Sci U S A. 2011 Mar 8;108(10):E35; author reply E36.

41.   Burak Y, Rokni U, Meister M, Sompolinsky H. Reply to Wehrhahn: Experimental requirements for
testing the role of peripheral cues in dynamic image stabilization. Proc Natl Acad Sci U S A. 2011
Mar 8;108(10):E36.

42.   MacKinnon DP, Fairchild AJ, Fritz MS. Mediation Analysis. Annu Rev Psychol. 2007;58(1):593–614.

43.   Engbert R, Mergenthaler K. Microsaccades are triggered by low retinal image slip. Proc Natl Acad
Sci U S A. 2006;103(18):7192–7.

44.   Ohl S, Wohltat C, Kliegl R, Pollatos O, Engbert R. Microsaccades Are Coupled to Heartbeat. J
Neurosci. 2016;36(4):1237–41.

45.   Masquelier T, Portelli G, Kornprobst P. Microsaccades enable efficient synchrony-based coding in
the retina: a simulation study. Sci Rep. 2016;6(October 2015):24086.

46.   McCamy MB, Otero-Millan J, Macknik SL, Yang Y, Troncoso XG, Baer SM, et al. Microsaccadic
Efficacy and Contribution to Foveal and Peripheral Vision. J Neurosci. 2012;32(27):9194–204.

47.   Brien BJO, Isayama T, Richardson R, Berson DM, O'Brien BJ. Intrinsic physiological properties of
cat retinal ganglion cells. J Physiol. 2002;538(Pt 3):787–802.

48.   Berry MJ, Warland DK, Meister M. The structure and precision of retinal spike trains. Proc Natl
Acad Sci. 1997;94(10):5411–6.

49.   Reich DS, Victor JD, Knight BW, Ozaki T, Kaplan E. Response variability and timing precision of
neuronal spike trains in vivo. J Neurophysiol. 1997;77(5):2836–41.

50.   Kumar G, Chung STL. Characteristics of fixational eye movements in people with macular disease.
Invest Ophthalmol Vis Sci. 2014 Aug;55(8):5125–33.

51.   Chung STL, Kumar G, Li RW, Levi DM. Characteristics of fixational eye movements in amblyopia:
Limitations on fixation stability and acuity? Vision Res. 2015 Sep 1;114:87–99.

52.   Patrick JA, Roach NW, McGraw P V. Motion-based super-resolution in the peripheral visual field. J
Vis. 2017 Sep 18;17(9):15.

53. Watson LM, Strang NC, Scobie F, Love GD, Seidel D, Manahilov V. Image Jitter Enhances Visual
Performance when Spatial Resolution Is Impaired. Investig Opthalmology Vis Sci. 2012 Sep
6;53(10):6004.

54. Poletti M, Aytekin M, Rucci M. Head-Eye Coordination at a Microscopic Scale. Curr Biol.
2015;25(24):3253–9.

55. Stevenson SB, Sheehy CK, Roorda A. Binocular eye tracking with the Tracking Scanning Laser
Ophthalmoscope. Vision Res. 2016;118:98–104.

56. Arathorn DW, Yang Q, Vogel CR, Zhang Y, Tiruveedhula P, Roorda A. Retinally stabilized cone-
targeted stimulus delivery. Opt Express. 2007 Oct 17;15(21):13731.

57. Yang Q, Arathorn DW, Tiruveedhula P, Vogel CR, Roorda A. Design of an integrated hardware
interface for AOSLO image capture and cone-targeted stimulus delivery. Opt Express. 2010 Aug
16;18(17):17841.

58. Mulligan JB. Recovery of motion parameters from distortions in scanned images. Proc Image
Regist Work. 1997;281–92.

59. Castet E, Crossland M. Quantifying Eye Stability During a Fixation Task: A Review of Definitions
and Methods. Seeing Perceiving. 2012;25(5):449–69.

60. Botev ZI, Grotowski JF, Kroese DP. Kernel density estimation via diffusion. Ann Stat. 2010 Oct
11;38(5):2916–57.

61. Kwon M, Nandy AS, Tjan BS. Rapid and persistent adaptability of human oculomotor control in
response to simulated central vision loss. Curr Biol. 2013;23(17):1663–9.

62. Engbert R, Kliegl R. Microsaccades uncover the orientation of covert attention. Vision Res.
2003;43(9):1035–45.

63. Daniel RM, De Stavola BL, Cousens SN, Vansteelandt S. Causal mediation analysis with multiple
mediators. Biometrics. 2015;71(1):1–14.

Supplemental Figures

Figure S1 Individual results from Experiment 1. (a) Proportion correct and all mediators quantified in the present study, binned based on gain. (b) Bootstrapping tuning curve fits (top) and optimal gains (bottom) for each subject by using binary data (correct vs incorrect). White lines represent the fits corresponding to median parameters. Shaded regions (top) represent 2.5-97.5% percentiles of the bootstrapped distributions of fitted curves. For each panel, bootstrapping was done by resampling the individual trial data with replacement 1000 times. Vertical dashed lines (bottom) represents the median optimal gains. The data from second run of S4 were combined with the first run and analyzed together, and are shown here in the rightmost panels.

Figure S2 Analyses of microsaccades. **(a)** Amplitude and direction distribution of microsaccades, combined across seven subjects, in Experiment 1. **(b)** The retinal position of the stimuli at the start (black squares) and end (red circles) of microsaccades. Clearly, the primary role of microsaccades was redirecting gaze to compensate for non-homogeneous vision. **(c)** Retinal image velocity just before microsaccades. The blue and red lines represent downward and upward microsaccades (within ±45° from vertical was considered as upward). **(d)** Retinal position of the stimuli across microsaccades. In **(c)** and **(d)**, the panels on the left and right represent data from horizontal and vertical component of the eye movements, respectively.

Figure S3 Ocular drifts under normal viewing conditions. To qualitatively test the assumption that drifts
are cyclic motions within the time scale of typical fixation (~300 ms), we randomly sampled 20 eye
position traces from three subjects in the zero-gain condition, and computed the power spectra of both
the horizontal and vertical components in **(a)** a 300 ms and **(b)** ~900 ms time windows. If indeed, drifts
show three to five cycles per ~300 ms, this should be visible as clear peaks in the power spectra, and
disappear when power spectra are computed over a longer time scale. However, except for a few cases,
we did not encounter such motions. For some subjects, much lower-frequency fluctuations were visible,
but these fluctuations are too slow to be of any use for fast and efficient temporal encoding. For this
particular figure, eye position traces were further filtered with a low-pass filter with a cut-off frequency
of 25 Hz. Power spectra are shown with a linear frequency axis with limits from 3 to 30 Hz.

Figure S4 Contributions of gain and mediators in explaining variance. **(a)** Change in (left) explained variance, (middle) log-likelihood, and (right) BIC with addition of mediators. Note that the sign of ΔBIC is flipped so that red color represents superiority of a model on the vertical axis with respect to another one on the horizontal axis. G: Gain, R: retinal ISOA, E: eye ISOA, P: PRL eccentricity, M: microsaccade rate. **(b)** Can mediators fully account for the effects of gain? Here, we explicitly tested whether having gain in addition to mediators improve statistical models substantially. The right diagonal in each panel represents the exact contribution of the gain term. The red squares represent the final model in the mediation analysis (G+G^2+R+E+P) (**Fig. 4d**). In general, adding gain was helpful only when there are three or less mediators in the regression model.