Species comparison of liver proteomes reveals enhanced lipid metabolism, reduced mitochondrial respiration and enhanced expression of detoxifying enzymes in the long-lived naked mole-rat

Authors:

Ivonne Heinze[1*], Martin Bens[1*], Enrico Calzia[2*], Susanne Holtze[3], Oleksandr Dakhovnik[1], Arne Sahm[1], Joanna M. Kirkpatrick[1], Karol Szafranski[1], Natalie Romanov[4], Stephan Singer[5], Maria Ermolaeva[1], Matthias Platzer[1#], Thomas Hildebrandt[3#] and Alessandro Ori[1#]

[1] Leibniz Institute on Aging – Fritz Lipmann Institute (FLI), Jena, Germany
[2] Institute of Anesthesiological Pathophysiology and Process Engineering, University of Ulm, Germany
[3] Leibniz Institute for Zoo & Wildlife Research (IZW), Berlin, Germany
[4] Structural and Computational Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany
[5] Institute of Pathology, University Hospital Heidelberg, Heidelberg, Germany

[*] co-first authors
[#] co-senior authors

Correspondence should be addressed to: matthias.platzer@leibniz-fli.de or HILDEBRAND@izw-berlin.de or alessandro.ori@leibniz-fli.de

Abstract:

Mammals display wide range of variation in their lifespan. Lifespan is generally correlated to body size, but outliers such as human and the naked mole-rat (NMR, *Heterocephalus glaber*) exist. Investigating the molecular networks that distinguish long- from short-lived species has proven useful to identify determinants of longevity. Here, we compared the liver of long-lived NMRs and the phylogenetically closely related, shorter-lived, guinea pigs (GP, *Cavia porcellus*) using an integrated transcriptomic and proteomic approach. We found that NMRs express substantially higher levels (up to 30 fold) of a restricted number of longevity-associated proteins that confer enhanced buffering against oxidative stress. Moreover, NMR livers display a unique expression pattern of mitochondrial proteins that result in distinct metabolic features of their mitochondria. For instance, we observed a generally reduced respiration rate associated with lower protein levels of respiratory chain components, particularly complex I, and increased capacity to utilize fatty acids. Interestingly, the same molecular networks are affected during aging in both NMR and humans, supporting a direct link to the extraordinary longevity of both species. Finally, we used our analysis to identify novel longevity pathways, and validated one of them experimentally in the phylogenetically distantly related nematode *C. elegans.*

49 **Introduction:**
50
Among mammals lifespan generally correlates with other life-history parameters
such as gestation period and body mass [1]. In this perspective, a subterranean
rodent, the naked mole-rat (*Heterocephalus glaber,* NMR), and humans represent
two species outliers by having an exceptionally long lifespan relatively to their
body mass. NMRs are eusocial animals that live in colonies where only a
subgroup of animals is devoted to reproduction (usually a queen and one male
called pasha) [2]. NMRs exhibit other exceptional traits including lifelong
fertility, resistance to infection, high regenerative capacity, resistance to cancer
and diabetes, reviewed in [3,4]. For these reasons, NMRs have drawn attention of
multiple studies aimed at identifying the molecular mechanisms behind their
extreme longevity and resistance to age-related diseases. Comparative genome
analysis has revealed positively selected genes in NMR [5,6], and RNA-seq
analysis revealed minimal changes in gene expression during aging [5,7],
supporting the view of enhanced maintenance of homeostasis in NMRs at the
molecular level. NMRs possess enhanced protein stability and increased
proteasomal activity [8,9], negligible levels of cellular senescence [10], over-
activation of pathways that contribute to stress resistance (e.g., the nuclear
factor erythroid2-related factor 2 (NFE2L2, previously NRF2/Nrf2) and p53)
[3,11], atypical expression of extracellular matrix components, such as high
molecular mass hyaluronan, that confer resistance against cancer development
[12]. Intriguingly, NMRs have higher steady-state levels of oxidative damage
compared to, e.g., mouse [8,13], and possess mitochondria with unusual
morphology in the heart and skeletal muscle [14]. However, NMRs appear to be
protected from the age-dependent increase in oxidative damage that manifest in
other species [8], presumably due to enhanced detoxifying systems [15].
76
There is a wealth of evidence linking energy metabolism to the aging process and
organism lifespan. Dietary interventions and nutrient sensing pathways have
been shown to play a central role in modulating aging in different organisms
[16–18]. Studies of genes under positive selection pressure across related
species that show different lifespan have highlighted genes involved in
mitochondrial homeostasis balance [19], likely influencing both energy
metabolism and hormetic responses affecting lifespan [20]. Age-dependent
changes of mitochondrial ultrastructure and activity have been described both in
flies and mice [21].
86
There is emerging evidence that metabolic changes in the NMR contribute to
adaptation to its ecosystem [22], allow to act as a "superorganism" with its
eusocial life style [4], and might be related to its extreme longevity. However, the
relationship between NMR metabolism and longevity has not yet been
investigated. Given its central role in organism metabolism, we set out to
investigate the liver of NMRs in order to identify novel molecular signatures of
longevity. Since ecological adaptations are more likely to affect gene expression
(Fraser 2013), and mechanisms of aging act both at the transcript and protein
level [23], we performed a cross-species comparison between NMR and the
shorter-lived guinea pig (*Cavia porcellus,* GP) using an integrated proteomic and
transcriptomic approach. In order to investigate cross-species differences in the

context of aging, we additionally analyzed livers from young and old NMRs, and
related the identified changes to human aging by studying the liver proteome of
12 individuals aged between 31 and 88 years. Finally, we validated one of the
newly identified longevity pathways to be a mediator of lifespan in the nematode
C. elegans.
103

Results:

Cross-species comparison of liver proteomes reveals elevated expression of
detoxifying enzymes in naked mole rats

We first compared the liver proteomes of 4 young adult male NMRs (2.7-3.8 year
old (yo)) and GPs (0.7-1 yo) using mass spectrometry (Table S1). For each
animal, we obtained a quantitative proteome profile by liquid chromatography
tandem mass spectrometry, and estimated absolute protein abundances using
the iBAQ method [24]. In order to directly compare the two species, we mapped
both NMR and GP proteins to the respective human orthologs, and used these as
the reference for comparison (see Material and Methods). This allowed us to
perform a quantitative comparison of the two species using estimated absolute
abundances for 3248 protein groups quantified by at least two unique
proteotypic peptides in both species (Figure 1A and Table S2). In order to
validate our approach, we obtained RNA-seq data from the same samples and
determined transcript-level fold changes between the two species. For this, only
those reads that exclusively mapped to conserved regions were used, a method
used for transcriptomic cross-species comparisons [7] (Table S2). Protein and
transcript fold changes displayed a significant positive correlation (Pearson
R=0.52, p<2.2e-16, Figure 1B), which is in line with comparisons performed
within the same species [25]. These data indicate that our strategy can reveal
meaningful differences in protein abundance between species and that many of
these changes are driven by changes in transcript levels.

We next asked whether our comparison would reveal known longevity-
associated proteins. We found strikingly higher levels of peroxiredoxin 1
(PRDX1) and thioredoxin reductase 1 (TXNRD1) in NMRs as compared to GPs
(Figure 1C and D). *PRDX1* and *TXNRD1* are both target genes of the transcription
factor NFE2L2, they play a crucial role in maintaining cell redox homeostasis,
and both have been shown to influence lifespan in multiple species by buffering
ROS and promoting proteostasis [26–28]. Interestingly, both PRDX1 and
TXNRD1 are cytosolic enzymes, and their mitochondrial counterparts (PRDX5,
TXNRD2) are instead expressed at similar levels in both species (Table S2).

We next wondered whether we could identify similar differences in other organs.
For this purpose, we compared RNA-seq data from heart, skin, kidney and
cerebellum across the two species and found that increased transcript levels of
PRDX1 occur exclusively in the liver (Figure 1E and F). This suggests that
increased level PRDX1 and TXNRD1 might be linked to a specific metabolic
activity of the NMR liver.

Figure 1. Cross-species comparison of the liver proteome integrated with RNA-seq from the same animals. (A) Liver proteomes from 4 adult naked mole rats (NMRs) and 4 guinea pigs (GPs) were compared using quantitative mass spectrometry. Hierarchical clustering based on the correlation between proteome profiles based on 3248 protein groups quantified across the two species. **(B)** Correlation between fold changes estimated at the protein level by quantitative mass spectrometry and at the transcript level by RNA-seq. For RNA-seq analysis only reads mapping to conserved regions between the two species were considered. **(C and D)** PRDX1 and TXNRD1 as examples of longevity-associated proteins that show drastically increased abundance in NMR vs. GP. Each bar represents the abundance estimated for one animal. **(E and F)** Comparison of transcript levels of *PRDX1* and *TXNRD1* across multiple tissues. For both genes, transcript levels are increased in NMR vs. GP in the liver (q<2.2x10^{-300} for *PRDX1*; q=1.5x10^{-45} for *TXNRD1*), while they show similar abundances in the other tissues examined. Related to Tables S1 and S2.

165 *The liver of naked mole rats has a unique metabolism characterized by reduced*
166 *mitochondrial respiration and enhanced lipid metabolism*
167
Intrigued by this finding, we used gene set enrichment analysis to investigate
differences in pathways and molecular networks between the long-lived and
shorter-lived species. Our analysis returned pathways linked to energy
metabolism (Figure 2A and Table S3). In particular, we found pathways related
to lipid metabolism to be up regulated, and gene sets related to oxidative
phosphorylation to be down regulated in NMR (q<0.05). Among up-regulated
proteins involved in lipid metabolism, we found enzymes responsible for fatty
acid beta-oxidation (e.g., ACOX2 and ACOX3), and lipid (e.g., ACACA and ACSL5),
cholesterol (e.g., MVD and DHCR24) and bile acids biosynthesis (e.g., AMACR)
compared to GP (Figure 2A and B and Table S2). Many of these are direct target
of the nuclear receptor peroxisome proliferator-activated receptor alpha
(PPARα), a master regulator of energy metabolism linked to aging [29] (Figure
2B). The majority of these enzymes showed higher abundance in the liver of
NMRs both at transcript and protein level, irrespectively of their sub-cellular
localization (Figure 2B).
183
In order to exclude that such differences would arise from a different organelle
composition of the NMR liver cells, we analyzed the distribution of fold changes
for proteins belonging to different cell compartments and found slight, but
significantly lower general levels of mitochondrial proteins (mean $\log_2$ fold
change=-0.17, p=0.01, Welch two sample t-test), and higher levels of lysosomal
proteins (mean $\log_2$ fold change=0.2, p=0.005, Welch two sample t-test) in NMR
compared to GP.
191
Regarding oxidative phosphorylation, NMRs showed reduced abundance of a
subset of mitochondrial respiratory chain components (Figure 2C). Interestingly,
these differences manifested almost exclusively at the protein level, and affected
to different extents the respiratory chain complexes, with components of
complex I being the most strongly reduced (mean $\log_2$ fold change=-1.53,
p=7.2x10^{-5}, Welch two sample t-test, Figure 2D). The opposite changes of
mitochondrial enzymes involved in lipid metabolism and respiratory chain
components indicate that NMR livers possess a distinct composition of their
mitochondrial proteome.
201
In order to investigate whether proteomic differences result in altered
mitochondrial activity of NMR liver parenchyma, we performed *ex vivo*
measurements of cellular respiration in liver extracts of NMR by means of high-
resolution respirometry (see Material and Methods). We measured
mitochondrial respiration from an independent group of adolescent (1 yo, n=4)
and mature adult (10 yo, n=4) male NMRs and compared them to mature adult
male mice (6-8 mo, n=4). We first compared the contribution of fatty acids to
mitochondrial activity using octanoylcarnitine and malate as substrates. After
this, maximum coupled respiration-state (OXPHOS-state) was established by
further addition of glutamate and succinate. Finally, we completed the test
sequence by adding carbonyl cyanide p-(trifluoromethoxy)-phenylhydrazone
(FCCP) in order to achieve the maximum uncoupled respiration-state. We

observed (Figure 2E) that NMRs and mice showed similar rates of mitochondrial
respiration supported by fatty acids. In contrast, NMR livers show reduced
maximum mitochondrial activity compared to mouse liver. In line with the
reduced abundance of respiratory chain components, mitochondrial oxygen
consumption normalized per wet tissue volume was significantly reduced in
both maximum OXPHOS (35% (p=0.011) and 37% (p=0.014) in 1 and 10 yo
NMRs, respectively; one-way ANOVA followed by Bonferroni's t-test), and
maximum uncoupled states (51% (p<0.001) and 54% (p<0.001)). This implies a
~2-fold higher fatty acid supported mitochondrial respiration in NMRs
compared to mice (Figure 2F). Taken together, our data indicate a marked
rearrangement of energy metabolism in the liver of NMR characterized by an
enhanced lipid metabolism and globally reduced level of mitochondrial
respiration.
227

Figure 2. Distinctive features of NMR liver metabolism. (A) Gene set enrichment analysis was performed on proteomic data. Gene sets are plotted according to the $\log_{10}$ value of the calculated enrichment score. Positive and negative values are used for gene sets showing higher and lower abundance in NMR, respectively. Because of the redundancy among the significantly affected gene sets (q<0.05), we highlighted representative categories of gene sets. The complete list of enriched gene sets is available in Table S3. **(B)** Protein (P) and transcript (T) fold changes for genes involved in lipid metabolism ("REACTOME METABOLISM OF LIPIDS AND LIPOPROTEINS", combined RNA-seq and proteome q<0.001), and **(C)** oxidative phosphorylation that are significantly affected in NMR vs. GP (selection criteria as in B). **(D)** Fold changes comparison for genes of the different complexes of the respiratory chain. Light blue and orange boxes indicate protein and transcript fold changes, respectively. **(E)** Mitochondrial oxygen flux supported by octanoylcarnitine and malate, compared to maximum coupled (OXPHOS) and maximum uncoupled respiration in liver of NMR (1 and 10 yo) and 6-8 mo male mice. **(F)** Ratio of fatty acid supported respiration to maximum OXPHOS and uncoupled respiration in liver of NMR and mouse. In E and F, reported values are averages obtained from n=4 animals per experimental group ± standard deviation. *=p<0.05; **=p<0.001; n.s=not significant. Related to Table S3.

250 *Cross-species- and aging-related changes correlate*
251

Given the existing evidences linking both lipid metabolism and oxidative
phosphorylation to lifespan [30,31], we investigated how these pathways are
affected during aging in NMRs. We therefore compared the liver proteome of
young (3-4 yo) and old (>20 yo) NMRs using quantitative mass spectrometry
(Figure 3A, Table S1 and S4). Gene Ontology analysis, based on 5237 protein
groups quantified by at least 2 unique peptides, revealed changes that are typical
of the aging process (Figure 3B). These include increased inflammation and
immune response-related proteins [23], and accumulation of extracellular
matrix proteins [32]. Interestingly, we found a statistically significant overlap
between proteins differentially abundant in NMR vs. GP, and proteins whose
abundance is affected by NMR aging (99 proteins, $p=2\times10^{-4}$, Fisher's exact test,
Figure 3C). Additionally, we observed a negative correlation between protein
fold changes across species and NMR aging (Pearson R=-0.38, p=0.0001; Figure
3D), resulting in a significant difference between cumulative distributions of
NMR vs. GP fold changes for proteins up- or down-regulated in NMR aging
(p=0.002, Kolmogorov-Smirnov-test; Figure 3E). The directionalities of the
differences indicate that proteins with decreasing expression during NMR aging
tend to have a higher level in young NMR compared to GP, whereas, proteins
with increasing expression during NMR aging tend to start from a lower level in
young NMR than in GP. Statistical significance for overlap (p=0.008, Fisher's
exact test), anti-correlation of fold changes (Pearson R=-0.18, $p=1.8\times10^{-8}$), and
difference between cumulative distributions ($p=3.1\times10^{-12}$, Kolmogorov-Smirnov-
test) can also be observed from RNA-seq data obtained from the same animals
(Figure S1). In particular, we found among the 30 liver proteins up in NMR vs. GP
and down during NMR aging 13 linked to lipid or fatty acid and 5 to xenobiotic
metabolism (Figure 3F). Our data suggest that this group of proteins might be
involved in sustaining the longevity of NMRs, and their decline during aging
might contribute to a phenotypically/functionally not yet reported reduction of
liver function.
281

Figure 3. The impact of aging on the NMR liver proteome (A) Livers from 3 young (2.7-3.8 yo) and 3 old (>20 yo) NMRs were compared by Tandem Mass Tags (TMT) based quantitative mass spectrometry. Hierarchical clustering based on the correlation between proteome profiles based on 5237 protein groups cross-quantified between the two age groups (Table S4). **(B)** Gene set enrichment analysis. Gene sets are plotted according to the $\log_{10}$ value of the calculated enrichment score. Positive and negative values are used for gene sets showing higher and lower abundance in old NMRs, respectively. Selected significantly affected gene sets (q<0.05) are highlighted. The complete list of enriched gene sets is available in Table S5. **(C)** Overlap between proteins differentially expressed in NMR vs. GP and affected by aging in NMR. **(D)** Comparison between cross-species and aging-related fold changes for the 99 proteins significantly different in both comparisons. **(E)** Cumulative distributions of significant NMR vs. GP fold differences for the 99 proteins also significantly up- (red) or down- (blue) regulated in aging. **(F)** The 62 proteins with significant but opposite fold changes in both comparisons (Table S6). Proteins involved in lipid/fatty acid metabolism and xenobiotic metabolism are highlighted in green and orange, respectively. Related to Figure S1 and Tables S4, S5 and S6.

305
306
Figure S1. Correlations between liver transcript level differences of NMR
vs. GP and during NMR aging. (A) Significant overlap between differentially
expressed genes (DEGs) in NMR vs. GP and in aging of NMR. **(B)** Comparison
between cross-species and aging-related fold changes for the 875 DEGs
significant in both comparison (q<0.01) shows significant negative correlation.
(C) Cumulative distributions of NMR vs. GP fold changes for the 875 DEGs also
significantly up- (red) or down- (blue) regulated in NMR aging. The x-axis was
restricted to ±8 for display purpose. Related to Figure 3 and Table S6.
315

316   *Aging affects similar pathways in both NMR and human liver*
317
In order to generalize our findings to other species and in particular to human
aging, we analyzed the proteome of donor liver samples from 12 individuals
aged between 31 and 88 years (Table S1). In this case, we used formalin fixed
and paraffin embedded (FFPE) samples and quantitatively compared the
proteomes using mass spectrometry. Principal Component Analysis (PCA) based
on 3211 quantified protein groups revealed separation of the proteome profiles
based on the age of the donor (Figure 4A). Guided by the PCA analysis, we split
the individuals into two groups defined as young (below 50 years of age) and old
(above 66 years), and analyzed differential protein expression for 3064 protein
groups quantified across all samples (Table S7). Multiple lines of evidence
indicate that aging affects similar pathways in both human and NMR liver. First,
as in NMR, GO enrichment analysis revealed an age-dependent decline of
proteins involved in lipid metabolism and detoxification of xenobiotics, and an
increase of proteins related to immune response (Figure 4B and Table S7), and
inflammation markers such as RELA/p65 (Figure 4C). Second, enzymes involved
in lipid synthesis such as ACACA and DHCR24, which were found to be expressed
at higher level in NMR vs. GP and to decline during NMR aging, showed a
negative correlation with the age of the donor (Figure 4D and 4E). Enzymes
involved in fatty acid beta-oxidation, including ACAA2 and HADHA, also showed
a trend of lower abundance in livers from older individuals (Figure S2A). Third,
these changes in metabolic enzymes underline a more general reorganization of
the liver proteome that is characterized by a significant reduction of both
mitochondrial and peroxisomal proteins during aging in both NMR and human
(Figure 4F). Fourth, multiple proteins involved in different steps of the
xenobiotic metabolism showed similar trends. Cytochrome P450s, a subset of
Glutathione S-transferases (GSTs), and UDP-glucuronosyltransferases (UGTs)
were in most cases expressed at higher levels in NMR vs. GP and showed an age-
dependent decline both in NMR and in human (Figures 4G and S2B). Taken
together these data indicate that conserved pathways are affected in NMR and
humans during aging.

348
349
Figure 4. The impact of aging on the human liver proteome and comparison
to NMR aging. (A) FFPE liver samples from 12 human donors aged 31-88 years
old were analysed by Data Independent Acquisition (DIA) quantitative mass
spectrometry. Principal Component Analysis (PCA) of the proteome profiles
based on 3211 protein groups quantified. **(B)** Gene Ontology (GO) enrichment
analysis based on differential protein expression between young (<47 yo, n=6)
and old (>66 yo, n=5) donors. One donor aged 58 yo was excluded from
differential expression analysis. GO categories are plotted according to the $\log_{10}$
value of the calculated enrichment p-value. Positive and negative values are used
for gene sets showing higher and lower abundance in old individuals,
respectively. Selected significantly affected GO terms (p<0.001) are highlighted.
The complete list of enriched GO terms is available in Table S7. **(C)** The
inflammation marker RELA shows a steady increase of abundance with age. **(D**
and E) Selected examples of enzymes involved in lipid metabolism being up
regulated in NMR vs. GP and decreasing during aging both in NMRs and humans.
(F) Mitochondrial and peroxisomal proteins decrease with age in both NMR and
human liver. Distributions of fold-changes were calculated separately for proteins
assigned to different cellular compartments. Mitochondrial (n=953, p=1.2x10^{-37} Welch
two sample t-test for NMR; n=716, p=1.4x10^{-21} Welch two sample t-test for
human) and peroxisomal proteins (n=95, p=4.9x10^{-6} Welch two sample t-test for
NMR; n= 74, p=8.9x10^{-3} Welch two sample t-test for human) showed a significant

lower abundance in old samples as compared to the rest of the quantified
proteins. **(G)** Major categories of detoxifying enzymes differentially expressed in
NMR vs. GP and affected by aging in both NMR and humans (only significantly
affected cases are shown for each group; cut-offs: NMR vs. GP and NMR aging,
combined q<0.05; human proteome aging q<0.1). Related to Figure S2 and Table
S7.
377

378
379

380
381
Figure S2. Examples of enzymes involved in fatty acid beta-oxidation and
xenobiotic metabolism that decrease during aging in human liver. (A)
Additional examples of enzymes involved in lipid metabolism decreasing during
aging in human liver. **(B)** Detoxifying enzymes decreasing during aging in both
NMR and humans. Only significantly affected genes are shown; cut offs: NMR
aging, combined q<0.05; human proteome aging q<0.1; n.s. = no significant cases
detected. Related to Figure 4.
389

390 *The detoxifying enzyme SULT1C3 mediates lifespan extension in C. elegans*
391
In order to demonstrate that proteins expressed at higher level in NMR are
mechanistically linked to longevity, we focused on proteins involved in
xenobiotic metabolism, and in particular the phase II enzyme SULT1C3 and its
paralog SULT1C2. These enzymes are expressed at prominent level in NMR and
they strongly decline during aging (Figure 5A and Figure S3). The *C. elegans*
ortholog of SULTIC2/3 is SSU-1 (coded by *ssu-1*)*,* and it was previously shown to
be expressed at high levels in long-lived and stress-resistant dauer larvae
(surviving up to 4 months compared to 2-3 weeks lifespan of a reproducing
adult). High expression levels of *ssu-1* are thus associated with extended
longevity also in worms [33]. The feeding of wild type worms with RNAi against
ssu-1 from L4 larval stage on resulted in significant reduction of their life span
(median life span reduction of 13%) suggesting that *ssu-1* is required for normal
lifespan of nematodes (Figure 5B). The remarkable stress resistance of dauer
larvae is largely due to enhanced expression of pro-survival and detoxification
genes mediated by DAF-16/FOXO transcription factor [34,35]. To test whether
the FOXO pathway regulates the expression of ssu-1 during adulthood and aging,
we performed an epistasis experiment treating *daf-16* deficient nematodes with
ssu-1 RNAi. The lifespan reducing effect of the *ssu-1* RNAi is also observed in *daf-*
16 mutants (median lifespan reduction of 12%) (Figure 5C), suggesting that the
function of *ssu-1* in normal aging is regulated by factors other than DAF-
16/FOXO.
413

415
416
Figure 5. Knock-down of SSU-1 (*ssu-1)* reduces lifespan in wild type and
***daf-16(mu86)* mutant *C. elegans*. (A)** The orthologous mammalian phase II
conjugation enzyme SULT1C3 is significantly more abundant in NMR compared
to GP, and it declines with aging. **(B)** Wildtype (N2, Bristol strain) and **(C)** *daf-*
16(mu86) mutant worms were treated with *ssu-1* and control empty vector (e.v.)
RNAi from the L4 (pre-adult) developmental stage. Survival was scored daily; the
significance of lifespan shortening was determined by Mantel-Cox Log rank test
(p<0.0001 in both cases). Related to Figure S3.
425

427
428
429 **Figure S3. SULT1C2 decreases with aging in NMR.** Related to Figure 5.
430
431

432   **Discussion:**
433
Cross-species comparisons based on transcriptional profiling have highlighted
pathways that correlate with lifespan [1]. In this work, we decided to
concentrate on the NMR, as an outlier of exceptional longevity, and directly
relate the identified proteome changes to the ones observed in humans, another
of those outliers. In our NMR vs. GP comparative approach, we generally
observed a good correlation between transcriptome and proteome differences
confirming that a great fraction of adaptation to local ecosystems occurs via
changes in gene expression that translate into abundance changes of proteins.
However, we have identified changes, particularly among complexes of the
mitochondrial respiratory chain, which manifested exclusively at the proteome
level. Importantly some of these changes are inline with measureable difference
in mitochondrial activity in NMR.
446
Our cross-species analysis revealed that the liver of NMRs possesses three major
characteristics compared to GP: (i) lower rate of mitochondrial respiration, due
to reduced level of complex I; (ii) higher reliance on fatty acids for energy
production, deriving from increased abundance of enzymes responsible for lipid
turnover; and (iii) increased expression of detoxifying enzymes. Although we
cannot exclude that some of the observed changes derive from differences in diet
between the two compared species, the fact that mitochondrial and oxidative
phosphorylation genes were identified to be differentially expressed also in
NMRs vs. wild mice [7] supports the peculiarity of NMR liver metabolism among
rodents. Importantly, we have also shown that the aging process itself negatively
affects the abundance of proteins involved in lipid metabolism and detoxification
processes both in NMR and humans. These pathways are similarly affected with
aging also in mice [36].
460
Two major questions arise from our work: how NMRs have evolved their
particular liver metabolism, and how does this contribute to the extreme
longevity of these animals? Multiple studies have previously linked the
composition of the mitochondrial respiratory chain to lifespan extension in
multiple species [31]: altered composition of the respiratory chain has been
show to induce a hormetic response that can extend lifespan in *C.elegans* [37];
mild inhibition of complex I leads to increased lifespan in the short-lived fish *N.*
furzeri [38]; low abundance of the matrix arm of complex I predicts longevity in
mice [39]; fibroblasts isolated from long lived individuals including centenarians
show altered mitochondrial activity with lower complex I driven ATP synthesis
[40].
472
Similarly, lipid homeostasis and signaling has been linked to health and longevity
[30], and changes in lipid metabolism have been shown to mediate the positive
effects of anti-aging dietary interventions. Dietary interventions such as calorie
restriction (CR) or fasting influence lipid metabolism [41,42]. Transcriptomic
and metabolomic measurements showed that CR promotes fatty acid fueling of
mitochondria in liver of mice and it is accompanied by changes in body fat
composition [43]. Both fatty acid oxidation and lipid metabolism pathways as
well as xenobiotic metabolism are induced by CR in mouse liver via epigenetic

reprogramming [44]. Changes in lipid metabolism are mechanistically linked to
activation of stress response pathways that mediate enhanced proteostasis [45],
and fatty acid oxidation and functional peroxisomes are required to maintain
mitochondria network homeostasis and promote longevity in *C. elegans* [46]. In
humans, shifts in body composition accompany aging including a decrease of
lean mass and accumulation of body fat. Fatty acid oxidation by respiring tissue
decreases with age in humans [47,48], however there are discordant reports
[49]. A decrease in fatty acid oxidation during aging can lead to adipose tissue
accumulation and thus contribute to increased systemic inflammation, a major
risk factor for aging-associated disease such as type II diabetes [50]. Our data
show that in both NMR and human liver there is a progressive decline of
enzymes responsible for fatty acid turnover. These alterations might contribute
to changes in energy metabolism that favor the accumulation of adipose tissue
and increased inflammation at older age.
495
From a mechanistic point of view, it is conceivable that adaptation to the
particular ecosystem of NMRs has selected for characteristics of energy
metabolism that in turn enabled extreme longevity via activation of stress
pathways (Figure 6). Among these, the NFE2L2 pathway, which control the
expression of many of the detoxifying enzymes that we found increased in NMR
vs. GP, was shown to have enhanced activity in NMR [11]. The activity of the
same pathways tend to decline during aging, as shown here by the decline of
their target genes in both NMR and humans, and in different systems [51,52]. It
is therefore tempting to speculate that their higher basal activity in the NMR
might contribute to its enhanced stress resistance, and ultimately delay the aging
process. In line with this hypothesis, genes encoding for both, respiratory
electron transport chain and response to oxidative stress have been shown to be
under positive selection in the NMR [6]. Additionally, similar molecular networks
(lipid metabolism and oxidative stress pathways) are involved in the social
status transition from worker to breeder in NMR [53], suggesting common
evolutionary constrains and molecular mechanism underlying longevity and
eusociality, i.e. reproductive animals' lifelong fecundity coupled with
extraordinary life- and healthspan in the NMR [54] and even extended lifespan in
other mole-rats [55,56] and longevity. Further work is required to elucidate in
detail which aspects of liver metabolism are sufficient to promote lifespan, and
what is the molecular basis mediating positive systemic effects that support
organism health by delaying aging in NMR.
518

520
521
522 **Figure 6.   Proposed   model   linking   metabolic   changes   to   stress**
523 **response/detoxifying pathways in NMR and during aging.**
524

Materials and methods:

Samples

Young NMR and GP liver tissue samples were obtained at the Leibniz Institute for Zoo and Wildlife Research, IZW (Berlin, Germany). Old NMR liver tissue samples were obtained from the Stockholm Zoo. Sampling and animal procedures were approved by the local ethics committee of the "Landesamt für Gesundheit und Soziales", Berlin, Germany (reference numbers: #ZH 156, G02217/12, and T 0073/15) and were compliant with the national and institutional animal care guidelines. NMRs were kept inside artificial burrow systems, consisting of acrylic glass containers interconnected with tubes. Except during cleaning and management procedures, animals were kept in complete darkness and supplied daily *ad libitum* with fresh vegetables. Temperature and humidity were kept stable at 27.0±2.0°C and 85.0±5.0%, respectively. GPs (breed: Dunkin Hartley HsdDhl:DH, Harlan Laboratories, AN Venray, Netherlands) were kept pairwise in standardized GP cages with a 12h light-dark cycle. They were fed commercial pellets and fresh vegetables daily; hay and vitamin C enriched water were provided *ad libitum*. Temperature and humidity ranged between 18.0±2.0°C and 45.0±5.0%, respectively. For tissue collection, animals were euthanized by surgical decapitation under general anaesthesia (Isofluran CP, CP-Pharma, Burgdorf, Germany). Tissue samples were fresh frozen and stored in liquid nitrogen before transcriptome or proteome analysis.

Human liver tissue samples were provided by the tissue bank of the National Center for Tumor Diseases (NCT, Heidelberg, Germany) in accordance with the regulations of the tissue bank and the approval of the ethics committee of Heidelberg University. These samples were obtained in a transplant setting to check the quality of the donor liver by histology before implantation. The age of the donors ranges from 31-88. The tissue samples were formalin fixed, paraffin-embedded (FFPE) and slides were stained with hematoxylin and eosin (H&E). These full-section H&E slides were re-evaluated by a trained pathologist (SS) confirming that each of the samples used for proteomic analyses did not show any overt pathomorphological changes (e.g. necrosis or significant inflammatory or fatty changes).

RNA sequencing analysis.

Reference transcripts

Reference transcripts for NMR are based on recently published de novo transcriptome assembly [57]. Reference transcripts for GP were obtained by *de novo* transcriptome assembly of ten different tissue samples as described in [53], using the human transcriptome as a reference for gene symbol assignment. Both transcript sets were mapped to the corresponding genomes (NMR UCSC hetgla2, GP UCSC cavpor3) in two steps: BLAT was used to identify the locus and then SPLIGN (v1.39.8) was applied to splice align the transcript sequence within BLAT locus.

Transcript quantification
For NMR age comparison RNA-seq data were aligned to the reference genome
utilizing STAR (v2.4.1d) with a maximum mismatch of 6% and a minimum
aligned length of 90%. Reads mapped to multiple loci were discarded. Gene
expression was quantified using HTSEQ (v0.6.1p1) based on the aligned
reference transcripts. For cross-species comparison (NMR vs. GP) orthologous
transcribed regions were determined using PosiGene [58] with parameter
'prank=0 max_anchor_gaps_hard=100 rs=NMR'. RNA-seq data were aligned to
corresponding orthologous transcribed regions in NMR and GP reference
transcripts using bowtie2 (2.2.9) with the parameter '--very-sensitive-local'.
DESeq2 (v1.6.3) was used to identify DEGs after correction of p-values using
Benjamini Hochberg (FDR, denoted as 'q') for NMR aging and cross-species
comparison.
587
Data availability
RNA-seq data of GP used in *de novo* assembly were deposited at Sequence Read
Archive (SRP104222). RNA-seq data for NMR age and cross-species comparisons
were deposited in Gene Expression Omnibus (GSE98744, GSE98744).   The
corresponding gene annotation for NMR and GP are available as a gff3-file
(ftp://genome.leibniz-fli.de/pub/nmr2017/).
594
595   *Sample preparation for mass spectrometry*
596
NMR and GP fresh frozen liver samples for label free cross species comparison
Frozen tissue samples of NMR and GP livers (between 20 and 40 mg, Table S1)
were collected into Precellys Lysing kit tubes (Keramik-kit 1.4/2.8 mm, 2 mL
(CKM)) containing 200 µL of protein solubilization buffer (80 µM Tris pH 8.0, 80
µM DTT and 4% SDS) and processed directly. Samples were homogenized in the
Precellys 24 homogenizer (Bertin Instruments, France) at 5000 rpm for 30
seconds at 4 °C. Samples were then spun down and the supernatant transferred
to a 1.5 mL Eppendorf tube. Samples were sonicated using a Bioruptor Plus
(Diagenode) for 7.5 minutes (5 cycles: 1 min on, 30 sec off, 20 °C) using the high
setting, and then boiled for 10 min at 95°C. A second round of sonication (as
before) followed the boiling. Samples were spun down at 20800x *g* for 5 minutes
and the lysate supernatant transferred to fresh tubes. Protein concentration was
determined by BCA assay (Pierce) using standard protocol and adjusted to 10
µg/µL using solubilization buffer. 5 µL of tissue lysate, corresponding to 50 µg
protein, was taken for preparation for MS. Cysteine residues were alkylated by
adding 1 µL of, 200 mM iodoacetamide to a final concentration of 15 mM
(incubated for 30 min at room temperature in the dark). Reaction was quenched
by addition of 1 µL of 200 mM DTT. Sample clean-up proceeded following a
modified SP3 protocol. Sera-Mag Speed Beads (#45152105050250 and
#65152105050250, Thermo Scientific) were mixed 1:1, rinsed with water and
stored as a 40 µg/µL stock solution in 4°C, as described in [59]. 4 µL of beads
stock was added to the reaction tube and mixed by pipetting then 11 µL
acetonitrile containing 5% (v/v) formic acid was added. Samples were incubated
for 8 minutes at room temperature to allow protein bindings to the beads. Next,
tubes were placed on the magnetic rack. Supernatant was removed and
discarded. Beads were washed twice with 180 µL of 70% (v/v) ethanol and once

with 180 µL of 100% acetonitrile. After removal of acetonitrile beads were air-
dried for 60 sec and then resuspended in 7 µL of digestion buffer (6 µL 4 M urea
in 100 mM ammonium bicarbonate and 1 µL of 1 µg/µL of LysC (Wako)).
Samples were sonicated for 5 min in water bath, incubated for 5 min at 37°C and
then mixed by pipetting. Digestion was allowed to proceed for 4 h at 37°C. After
the first step of digestion, beads were resuspended by pipetting, urea was diluted
to the final concentration of 1.5 M and 1 µL of 1 µg/µL of sequencing grade
trypsin (Promega) was added to samples. Digestion was performed overnight at
37°C. After digestion, beads were resuspended by pipetting. 100% acetonitrile
was added to the final concentration of 95% (v/v) and samples were incubated
for 8 min at room temperature. Tubes were placed on the magnetic rack and
washed twice with 100% acetonitrile. Supernatant was removed and beads air-
dried and reconstituted in 20 µL of 2% DMSO followed by 5 min of sonication in
the water bath. Samples were resuspended by pipetting and placed on the
magnetic rack. Supernatant containing peptides was transferred to a fresh tube
and acidified with 2 µL of 1% (v/v) formic acid prior to pre-fractionation by high
pH reverse phase chromatography. Offline high pH reverse phase fractionation
was performed using an Agilent 1260 Infinity HPLC System equipped with a
binary pump, degasser, variable wavelength UV detector (set to 220 and 254
nm), peltier-cooled autosampler (set at 10°C) and a fraction collector. The
column was a Waters XBridge C18 column (3.5 µm, 100 x 1.0 mm, Waters) with a
Gemini C18, 4 x 2.0 mm SecurityGuard (Phenomenex) cartridge as a guard
column. The solvent system consisted of 20 mM ammonium formate (pH 10.0) as
mobile phase (A) and 100% acetonitrile as mobile phase (B). The separation was
accomplished at a mobile phase flow rate of 0.1 mL/min using a linear gradient
from 100% A to 35 % B in 61 min. Thirty-four fractions were collected along
with the LC separation, which were subsequently pooled into 10 fractions.
Pooled fractions were dried in a Speed-Vac and then stored at -80°C until LC-
MS/MS analysis.
652
<u>NMR frozen liver samples for TMT-based comparison of young and old samples</u>
For each experimental animal (Table S1), 100 µg protein lysate from the bead-
beaten stock of tissue described above were taken up to a final volume of 50 µL
with 100 mM HEPES buffer, pH 8.5. 5 µL of 2% SDS was added, prior to
biorupting (5 cycles: 1 min on, 30 sec off, 20 °C) at the highest settings. Samples
were spun down at 20800x g for 1 minute and the lysate supernatant transferred
to fresh tubes. Reduction was performed with 2.9 µL DTT (200 mM) for 15
minutes at 45 °C before alkylation with 200 mM IAA (5 µL, 30 minutes, room
temperature, in the dark). Proteins were then precipitated with 4 volumes ice
cold acetone to 1 volume sample and left overnight at -20 °C. The samples were
then centrifuged at 20800x g for 30 minutes, 4 °C. After removal of the
supernatant, the precipitates were washed twice with 500 µL 80% (v/v) acetone
(ice cold). After each wash step, the samples were vortexed, then centrifuged
again for 2 minutes at 4°C. The pellets were then allowed to air-dry before being
dissolved in digestion buffer (50 µL, 3M urea in 0,1M HEPES, pH 8; 1 µg LysC)
and incubated for 4 h at 37 °C with shaking at 600 rpm. Then the samples were
diluted 1:1 with milliQ water (to reach 1.5M urea) and were incubated with 1 µg
trypsin for 16 h at 37 °C. The digests were then acidified with 10% trifluoroacetic
acid and then desalted with Waters Oasis® HLB µElution Plate 30µm in the

presence of a slow vacuum. In this process, the columns were conditioned with
3x100 µL solvent B (80% (v/v) acetonitrile; 0.05% (v/v) formic acid) and
equilibrated with 3x 100 µL solvent A (0.05% (v/v) formic acid in milliQ water).
The samples were loaded, washed 3 times with 100 µL solvent A, and then eluted
into PCR tubes with 50 µL solvent B. The eluates were dried down with the speed
vacuum centrifuge and dissolved in 200 mM HEPES buffer, pH 8.5 for TMT
labeling. 25 µg peptides were taken for each labeling reaction at 1 µg/µL
concentration. TMT-6plex reagents (Thermo Scientific) were reconstituted in 41
µL 100% anhydrous DMSO. TMT labeling was performed by addition of 2.5 µL of
the TMT reagent. After 30 minutes of incubation at room temperature, with
shaking at 600 rpm in a thermomixer (Eppendorf) a second portion of TMT
reagent (2.5 µL) was added and incubated for another 30 minutes. The reaction
was quenched with 1 µL of 20 mM lysine in 100 mM ammonium bicarbonate.
After checking labeling efficiency, samples were pooled (48 µg total), cleaned
once again with Oasis and subjected to high pH fractionation prior to MS
analysis. Offline high pH reverse phase fractionation was performed as described
above with the following modifications for TMT labeled samples: (i) the
separation was accomplished at a mobile phase flow rate of 0.1 mL/min using a
non-linear gradient from 95% A to 40 % B in 91 min; (ii) 48 fractions were
collected along with the LC separation that were subsequently pooled into 16
fractions.
693
<u>Human FFPE liver samples</u>
The specimens were cut on a microtome into 5 µm thick sections and mounted
on glass slides. Slides were deparaffinized in xylene for 2x 5 minutes, rehydrated
in 100% ethanol for 2x 5 minutes, and then washed in 96% (v/v), 70% (v/v),
50% (v/v) ethanol and milliQ water for 1x 5 minutes each. Region of interest
were gently scraped using a scalpel and transferred to a PCR tube containing 100
µL of protein solubilization buffer (80 µM Tris pH 8.0, 80 µM DTT and 4% SDS)
and processed directly. Samples were sonicated using a Bioruptor Plus
(Diagenode) for 25.2 min (15 cycles: 1 min on, 30 sec off) at 20°C using the high
setting, and then boiled for 1h at 99°C. Sonication followed by boiling was
performed twice. Cysteine residues were alkylated by adding 200 mM
iodoacetamide to a final concentration of 15 mM (incubated for 30 min at room
temperature in the dark). Reaction was quenched by addition of 10 µL of 200
mM DTT. Protein were then acetone precipitated, digested and desalted as
described above for NMR samples (aging comparison), with the exceptions that
0.5 µg of both LysC and trypsin were used instead of 1 µg to accommodate for the
lower amount of protein extract employed, and no TMT labeling was performed.
711
712   *Mass spectrometry data acquisition*
713
<u>Label free analysis of NMR and GP liver samples</u>
For label free experiments, each fraction from the 4 GP and 4 NMR samples,
separated by high pH, were resuspended in 10 µL reconstitution buffer (5%
(v/v) acetonitrile, 0.1% (v/v) TFA in water) and 8 µL were injected. Peptides
were separated using the nanoAcquity UPLC system (Waters) fitted with a
trapping (nanoAcquity Symmetry C18, 5 µm, 180 µm x 20 mm) and an analytical
column (nanoAcquity BEH C18, 2.5 µm, 75 µm x 250 mm). The outlet of the

analytical column was coupled directly to an Orbitrap Fusion Lumos (Thermo
Fisher Scientific) using the Proxeon nanospray source. Solvent A was water,
0.1% (v/v) formic acid and solvent B was acetonitrile, 0.1% (v/v) formic acid.
The samples were loaded with a constant flow of solvent A at 5 μL/min, onto the
trapping column. Trapping time was 6 min. Peptides were eluted via the
analytical column at a constant flow of 0.3 μL/min, at 40°C. During the elution
step, the percentage of solvent B increased in a linear fashion from 5% to 7% in
10 minutes, then from 7% B to 30% B in a further 105 min and to 45% B by 130
min. The peptides were introduced into the mass spectrometer via a Pico-Tip
Emitter 360 μm OD x 20 μm ID; 10 μm tip (New Objective) and a spray voltage of
2.2kV was applied. The capillary temperature was set at 300°C. Full scan MS
spectra with mass range 375-1500 *m/z* were acquired in profile mode in the
Orbitrap with resolution of 120000 using the quad isolation. A first batch of
samples (NMR: F1-6074, M1-1449; GP: #18, #19) was acquired with the
following settings. The RF on the ion funnel was set to 60%. The filling time was
set at maximum of 100 ms with an AGC target of 4×10^5 ions and 1 microscan.
The peptide monoisotopic precursor selection was enabled along with relaxed
restrictions if too few precursors were found. The most intense ions (instrument
operated for a 3 second cycle time) from the full scan MS were selected for MS2,
using quadrupole isolation and a window of 1.6 Da. An intensity threshold of 5
$x10^3$ ions was applied. HCD was performed with collision energy of 35%. A
maximum fill time of 30 ms with an AGC target of 1×10^4 for each precursor ion
was set. MS2 data were acquired in centroid in the ion trap, in Rapid scan mode,
with fixed first mass of 120 *m/z*. The dynamic exclusion list was with a maximum
retention period of 60 sec and relative mass window of 10 ppm. In order to
improve the mass accuracy, internal lock mass correction using a background ion
(*m/z* 445.12003) was applied. For data acquisition and processing of the raw
data Xcalibur 4.0 (Thermo Scientific) and Tune version 2.0 were employed. As a
consequence of method optimization, the following parameters were changed
for a second batch of samples (NMR: #0713, #4925; GP: #23, #28): RF on the ion
funnel was set to 40%, AGC target to 2×10^5, quadrupole isolation window to 1.4
Da, HCD collision energy to 30%, fill time to 300 ms, AGC target to 2×10^3, and
the instrument was set to inject ions for all available parallelizable time. Since
the two batches of samples were block randomized (i.e. both contained the same
number of NMR and GP samples), the usage of two different methods did not
influence the outcome of our comparison, as shown by the expected clustering of
the samples according to the species of origin (Figure 1B).
758
<u>TMT analysis of NMR young and old samples</u>
For TMT-6plex experiments, fractions were resuspended in 10 μL reconstitution
buffer (5% (v/v) acetonitrile, 0.1% (v/v) TFA in water) and 3.5 μL were injected.
Peptides were analyzed using the same LC-MS/MS setup described above with
the following modifications. Peptides were eluted using a linear gradient from
5% to 7% in 10 minutes, then from 7% B to 30% B in a further 105 min and to
45% B by 130 min. Full scan MS spectra with mass range 375-1500 *m/z* were
acquired in profile mode in the Orbitrap with resolution of 60000 using the quad
isolation. The RF on the ion funnel was set to 40%. The filling time was set at
maximum of 100 ms with an AGC target of 4×10^5 ions and 1 microscan. The
peptide monoisotopic precursor selection was enabled along with relaxed

restrictions if too few precursors were found. The most intense ions (instrument
operated for a 3 second cycle time) from the full scan MS were selected for MS2,
using quadrupole isolation and a window of 1 Da. HCD was performed with
collision energy of 35%. A maximum fill time of 50 ms for each precursor ion was
set. MS2 data were acquired with fixed first mass of 120 *m/z*. The dynamic
exclusion list was with a maximum retention period of 60 sec and relative mass
window of 10 ppm. For the MS3, the precursor selection window was set to the
range 400-2000 m/z, with an exclude width of 18 *m/z* (high) and 5 *m/z* (low).
The most intense fragments from the MS2 experiment were co-isolated (using
Synchronus Precursor Selection = 8) and fragmented using HCD (65%). MS3
spectra were acquired in the Orbitrap over the mass range 100-1000 *m/z* and
resolution set to 30000. The maximum injection time was set to 105 ms and the
instrument was set not to inject ions for all available parallelizable time.
783
<u>Data Independent Acquisition (DIA) for human FFPE samples</u>
Peptides were spiked with retention time HRM kit (Biognosys AG), and analyzed
using the same LC-MS/MS setup described above with the following
modifications. Approx. 1 µg for Data Dependent Acquisition (DDA) and 3 µg for
DIA analysis were loaded. Peptides were eluted via a non-linear gradient from 0
% to 40 % in 120 minutes. Total runtime was 145 minutes, including clean-up
and column re-equilibration. The RF lens was set to 30%. For spectral library
generation, a pooled sample was generated by mixing equal portion of each
sample, injected 12 times, and measured in DDA mode. The conditions for DDA
data acquisition were as follows: Full scan MS spectra with mass range 350-
1650 *m/z* were acquired in profile mode in the Orbitrap with resolution of
60000. The filling time was set at maximum of 50 ms with limitation of 2×10^5
ions. The "Top Speed" method was employed to take the maximum number of
precursor ions (with an intensity threshold of 5×10^4) from the full scan MS for
fragmentation (using HCD collision energy, 30%) and quadrupole isolation (1.4
Da window) and measurement in the Orbitrap (resolution 15000, fixed first
mass 120 *m/z*), with a cycle time of 3 seconds. The MIPS (monoisotopic
precursor selection) peptide algorithm was employed but with relaxed
restrictions when too few precursors meeting the criteria were found. The
fragmentation was performed after accumulation of 2×10^5ions or after filling
time of 22 ms for each precursor ion (whichever occurred first). MS/MS data
were acquired in centroid mode. Only multiply charged (2^+ - 7^+) precursor ions
were selected for MS/MS. Dynamic exclusion was employed with maximum
retention period of 15s and relative mass window of 10 ppm. Isotopes were
excluded. For data acquisition and processing Tune version 2.1 was employed.
For the DIA data acquisition the same gradient conditions were applied to the LC
as for the DDA and the MS conditions were varied as follows: Full scan MS
spectra with mass range 350-1650 *m/z* were acquired in profile mode in the
Orbitrap with resolution of 120000. The filling time was set at maximum of 20
ms with limitation of 5×10^5 ions. DIA scans were acquired with 34 mass
window segments of differing widths across the MS1 mass range with a cycle
time of 3 seconds. HCD fragmentation (30% collision energy) was applied and
MS/MS spectra were acquired in the Orbitrap with a resolution of 30000 over
the mass range 200-2000 *m/z* after accumulation of 2×10^5 ions or after filling

time of 70 ms (whichever occurred first). Ions were injected for all available
parallelizable time. Data were acquired in profile mode.
820
821   *Mass spectrometry data analysis*
822
<u>Label free cross-species comparison of NMR and GP liver samples</u>
The Andromeda search engine [60], part of MaxQuant (version 1.5.3.28) [61] was
used to search the data. The data for GP and NMR were searched separately
against translated species-specific reference transcripts (see RNA sequencing
analysis). Database with a list of common contaminants were appended in both
cases. The data were searched with the following modifications:
Carbamidomethyl (C) (Fixed), and Oxidation (M) and Acetyl (Protein N-term)
(Variable). The mass error tolerance for the full scan MS spectra was set at 20
ppm and for the MS/MS spectra at 0.5 Da. A maximum of 2 missed cleavages
were allowed. Peptide and protein level 1% FDR were applied using a target-
decoy strategy [62]. iBAQ (label free quantification) values from the MaxQuant
output were used to perform cross-species differential protein expression
analysis using scripts written in R (v3.4.1). After removal of reverse and
contaminant hits, only protein groups quantified by at least two unique peptides
were retained. Common human gene symbols were used to combine iBAQ values
for NMR and GP samples. Only protein groups quantified in at least two animals
per group were retained when comparing protein abundances between NMR
and GP. To reduce technical variation, data were $\log_2$ transformed and quantile-
normalized using the preprocessCore library. Protein differential expression was
evaluated using the limma package [63]. Differences in protein abundances were
statistically determined using the Student's t test moderated by the empirical
Bayes method. P values were adjusted for multiple testing using the Benjamini-
Hochberg method [64] (Table S2).
846
<u>TMT-based analysis of young and old NMR livers</u>
TMT-6plex data were processed using Proteome Discoverer v2.0 (Thermo Fisher
Scientific). Data were searched against the NMR fasta database using Mascot
v2.5.1 (Matrix Science) with the following settings: Enzyme was set to trypsin,
with up to 1 missed cleavage. MS1 mass tolerance was set to 10 ppm and MS2 to
0.5 Da. Carbamidomethyl cysteine was set as a fixed modification and oxidation
of Methionine as variable.   Other modifications included the TMT-6plex
modification from the quan method used. The quan method was set for reporter
ions quantification with HCD and MS3 (mass tolerance, 20 ppm). The false
discovery rate for peptide-spectrum matches (PSMs) was set to 0.01 using
Percolator [65]. Reporter ion intensity values for the filtered PSMs were
exported and processed using in-house written R scripts to remove common
contaminants and decoy hits Additionally only PSMs having reporter ion
intensities above 1×10^3 in all the relevant TMT channels were retained for
quantitative analysis. Only protein groups quantified by at least two unique
peptides were analyzed for differential expression between young and old NMR.
Data were analysed using the MSnbase package [66]. Reporter ion intensities
were $\log_2$-transformed and normalized using the vsn package [67]. Peptide-level
data were summarized into their respective protein groups by taking the median

value. Differential protein expression was assessed using the limma package, as
described above (Table S4).
868
<u>DIA analysis of FFPE human samples</u>
For library creation, the DDA data was searched using the Andromeda search
engine built in MaxQuant (version 1.5.3.28). The data were searched against a
human database (Swiss-Prot entries of the Uniprot KB database release 2016_01,
20198 entries) with a list of common contaminants appended, as well as the
HRM peptide sequences. The data were searched with the following
modifications: Carbamidomethyl (C) (Fixed) and Oxidation (M)/ Acetyl (Protein
N-term) (Variable). The mass error tolerance for the full scan MS and MS/MS
spectra was set at 20 ppm. A maximum of 1 missed cleavage was allowed. The
identifications were filtered to satisfy FDR of 1 % on peptide and protein level. A
spectral library was created from the MaxQuant output of the DDA runs
combined using Spectronaut (version 10, Biognosys AG). This library contained
58296 precursors, corresponding to 4624 protein groups using Spectronaut
protein inference. DIA data were then uploaded and searched against this
spectral library. Precursor matching, protein inference and quantification was
performed in Spectronaut using default settings [68]. Differential protein
expression was evaluated using a pairwise t-test performed at the precursor
level followed by multiple testing correction according to [69]. The data
(candidate table, Table S7) was exported from Spectronaut and used for further
data analyses (see below).
889
890 *Data analysis*
891
For integrated analysis of RNA-seq and proteomic data, transcripts and protein
groups were matched using the corresponding gene symbol, P values were
combined using the Fisher method, followed by correction for multiple testing
using the Benjamini-Hochberg method [64]. Gene set enrichments (Figures 2A
and 3B) were performed with the R package *gage* [70] using gene set definitions
from the Molecular Signatures Database (MSigDB, C2 v5.1) [71]. Gene Ontology
enrichment analysis (Figure 4B) was performed the list of quantified proteins
that were ranked according to the level of differential expression (fold change)
using GOrilla (Eden et al., 2009) followed by GO term redundancy reduction
performed by REViGO [72] using default settings.
902
903 *Measurements of mitochondrial activity*
904
Mitochondrial respiration was measured in homogenized liver tissue samples of
NMRs and mice by means of high-resolution respirometry using the Oroboros®
Oxygraph-2K (Oroboros Instruments, Innsbruck, Austria). This device allows for
simultaneous recording the O_2 concentration in two parallel chambers calibrated
for 2 ml of respiration medium containing 110 mM D-Sucrose (Sigma 84097), 60
mM K-lactobionate (Aldrich 153516), 0.5 mM ethylene glycol tetra acetic acid
(Sigma E4378), 1 g/L bovine serum albumin free from essentially fatty acids
(Sigma A 6003), 3 mM $MgCl_2$ (Scharlau MA0036), 20 mM taurine (Sigma T0625),
10 mM KH_2PO_4 (Merck 104873), 20mM HEPES (Sigma H7523), adjusted to pH
7.1 with KOH and equilibrated with 21% O_2 at 37°C. Mitochondrial respiration

was quantified in terms of oxygen flux (JO_2) calculated as the rate of change of
the O_2 concentration in the chambers normalized for wet tissue volume.
The liver tissue homogenates were generated from 40-50 mg of wet tissue
samples suspended in 2 ml of ice-cold respiration medium. Aliquots of the
homogenates were added to each oxygraph chamber in order to obtain a final
amount of 4 mg of NMR liver tissue or 2 mg of mouse liver tissue per chamber.
The different amount of tissue was chosen in order to obtain similar absolute
$JO2$-values, i.e. $JO2$-values not normalized per wet weight, in both species. Every
sample was measured in duplicates; the mean values from both chambers were
used for statistical analysis.
The titration sequence used for the experiments was as follows: 2 mM malate + 1
mM octanoylcarnitine, 5 mM ADP, 10 μM cytochrome c, 10 mM glutamate, 10
mM succinate, 2.5 μM oligomycin, 1 μM carbonyl cyanide p-(trifluoromethoxy)-
phenylhydrazone (FCCP), 0.5 μM rotenone, and 5 μM antimycin A. The JO_2-values
after addition of octanoyl carnitine, malate, ADP, and cytochrome c allow
quantifying fatty acid oxidation. The addition of cytochrome c after ADP is
required to test the integrity of the outer mitochondrial membrane. If
homogenization steps damaged the mitochondrial membrane, addition of
cytochrome c induces an increase of the respiratory values. The maximum
oxidative capacity of the mitochondrial respiratory chain in the coupled state
(maximum OxPhos) was then determined after the subsequent addition of
glutamate and succinate. Further injections of the ATP synthase inhibitor
oligomycin and of the uncoupler FCCP allowed obtaining the maximum
respiratory activity in the uncoupled state. In the next two steps, complex I and
complex III were sequentially inhibited by administration of rotenone and
antimycin A respectively. Finally sequentially injecting 2 mM ascorbate and 0.5
mM of the complex IV substrate tetramethyl phenylene diamine (TMPD) in the
parallel chambers allowed for selectively quantifying the activity of the
cytochrome-c-oxidase (COX). Part of the JO_2 induced by the injection of TMPD is
caused by auto-oxidation of this compound. Therefore, inhibiting the COX by 40
μM sodium sulfide allowed to quantify and thus to subtract this auto-oxidation
related part from the total JO_2 value under TMPD.
947
948 *C.elegans lifespan measurements*
949
HT115 bacteria containing specific RNAi constructs were grown on lysogeny
broth agar plates supplemented with ampicillin and tetracycline. Plates were
kept at 4°C. Overnight cultures were grown in lysogeny broth media containing
ampicillin. RNAi expression was induced by adding 1 mM
isopropylthiogalactoside (IPTG) and incubating the cultures at 37°C for
20 minutes before seeding bacteria on NGM agar supplemented with ampicillin
and 3 mM IPTG. Synchronized L4 larvae were placed on 60 mm dishes containing
RNAi expressing bacteria at a density of 70 worms per plate. Worms were
transferred to new plates on a daily basis until adulthood day 6 (AD6) and later
transferred to new plates every 3-4 days. The number of dead animals was
scored daily. The analysis of the lifespan data including statistics was performed
using GraphPad Prism software.
962

Heinze, Bens, Calzia, et al.

963   **Acknowledgements:**
964
965   We thank Stefan Pietsch for support with bioinformatics analysis. We gratefully
966   acknowledge support from the FLI proteomics, sequencing, and bioinformatics
967   core facilities. The FLI is a member of the Leibniz Association and is financially
968   supported by the Federal Government of Germany and the State of Thuringia. We
969   thank Veronika Geißler and other members of NCT tissue bank for their support.
970

971   **Author contributions:**
972
973   IH, EC, SH, MV performed experiments and analysed data; MB, AS, KS, NR
974   analysed data; MP, TH, AO, SS, JMK, KS, EC, ME designed experiments; MP, TH
975   and AO oversaw the project and wrote the manuscript.
976

References:

1. Fushan AA, Turanov AA, Lee S-G, Kim EB, Lobanov A V, Yim SH, et al. Gene expression defines natural changes in mammalian lifespan. Aging Cell. 2015;14: 352–65. doi:10.1111/acel.12283

2. Sahm A, Bens M, Szafranski K, Holtze S, Groth M, Goerlach M, et al. Long-lived rodents reveal signatures of positive selection in genes associated with lifespan and eusociality. doi.org. Cold Spring Harbor Laboratory; 2017; 191999. doi:10.1101/191999

3. Lewis KN, Mele J, Hornsby PJ, Buffenstein R. Stress resistance in the naked mole-rat: the bare essentials - a mini-review. Gerontology. Karger Publishers; 2012;58: 453–62. doi:10.1159/000335966

4. Skulachev VP, Holtze S, Vyssokikh MY, Bakeeva LE, Skulachev M V., Markov A V., et al. Neoteny, Prolongation of Youth: From Naked Mole Rats to "Naked Apes" (Humans). Physiol Rev. 2017;97: 699–720. doi:10.1152/physrev.00040.2015

5. Kim EB, Fang X, Fushan AA, Huang Z, Lobanov A V., Han L, et al. Genome sequencing reveals insights into physiology and longevity of the naked mole rat. Nature. 2011;479: 223–227. doi:10.1038/nature10533

6. Fang X, Seim I, Huang Z, Gerashchenko M V., Xiong Z, Turanov AA, et al. Adaptations to a Subterranean Environment and Longevity Revealed by the Analysis of Mole Rat Genomes. Cell Rep. 2014;8: 1354–1364. doi:10.1016/j.celrep.2014.07.030

7. Yu C, Li Y, Holmes A, Szafranski K, Faulkes CG, Coen CW, et al. RNA sequencing reveals differential expression of mitochondrial and oxidation reduction genes in the long-lived naked mole-rat when compared to mice. PLoS One. Public Library of Science; 2011;6: e26729. doi:10.1371/journal.pone.0026729

8. Pérez VI, Buffenstein R, Masamsetti V, Leonard S, Salmon AB, Mele J, et al. Protein stability and resistance to oxidative stress are determinants of longevity in the longest-living rodent, the naked mole-rat. Proc Natl Acad Sci U S A. 2009;106: 3059–64. doi:10.1073/pnas.0809620106

9. Rodriguez KA, Edrey YH, Osmulski P, Gaczynska M, Buffenstein R. Altered composition of liver proteasome assemblies contributes to enhanced proteasome activity in the exceptionally long-lived naked mole-rat. PLoS One. Public Library of Science; 2012;7: e35890. doi:10.1371/journal.pone.0035890

10. Buffenstein R. Negligible senescence in the longest living rodent, the naked mole-rat: Insights from a successfully aging species [Internet]. Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology. 2008. pp. 439–445. doi:10.1007/s00360-007-0237-5

11. Lewis KN, Wason E, Edrey YH, Kristan DM, Nevo E, Buffenstein R. Regulation of Nrf2 signaling and longevity in naturally long-lived rodents. Proc Natl Acad Sci U S A. National Academy of Sciences; 2015;112: 3722–7. doi:10.1073/pnas.1417566112

12. Tian X, Azpurua J, Hine C, Vaidya A, Myakishev-Rempel M, Ablaeva J, et al. High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. Nature. 2013;499: 346–349. doi:10.1038/nature12234

13. Andziak B, O'Connor TP, Qi W, DeWaal EM, Pierce A, Chaudhuri AR, et al. High oxidative damage levels in the longest-living rodent, the naked mole-rat. Aging Cell. 2006;5: 463–471. doi:10.1111/j.1474-9726.2006.00237.x

14. Holtze S, Eldarov CM, Vays VB, Vangeli IM, Vysokikh MY, Bakeeva LE, et al. Study of age-dependent structural and functional changes of mitochondria in skeletal muscles and heart of naked mole rats (Heterocephalus glaber). Biochem.

1028   2016;81: 1429–1437. doi:10.1134/S000629791612004X

1029 15. Lewis KN, Andziak B, Yang T, Buffenstein R. The naked mole-rat response to
1030   oxidative stress: just deal with it. Antioxid Redox Signal. Mary Ann Liebert, Inc.;
1031   2013;19: 1388–99. doi:10.1089/ars.2012.4911

1032 16. Finkel T. The metabolic regulation of aging. Nat Med. 2015;21: 1416–1423.
1033   doi:10.1038/nm.3998

1034 17. Fontana L, Partridge L. Promoting Health and Longevity through Diet: From
1035   Model Organisms to Humans. Cell. 2015;161: 106–118.
1036   doi:10.1016/j.cell.2015.02.020

1037 18. Davinelli S, Willcox DC, Scapagnini G. Extending healthy ageing: nutrient sensitive
1038   pathway and centenarian population. Immun Ageing. 2012;9: 9.
1039   doi:10.1186/1742-4933-9-9

1040 19. Sahm A, Bens M, Platzer M, Cellerino A. Parallel evolution of genes controlling
1041   mitonuclear balance in short-lived annual fishes. Aging Cell. 11 Feb 2017.
1042   doi:10.1111/acel.12577

1043 20. Houtkooper RH, Mouchiroud L, Ryu D, Moullan N, Katsyuba E, Knott G, et al.
1044   Mitonuclear protein imbalance as a conserved longevity mechanism. Nature.
1045   Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights
1046   Reserved.; 2013;497: 451–7. doi:10.1038/nature12188

1047 21. Brandt T, Mourier A, Tain LS, Partridge L, Larsson N-G, Kühlbrandt W. Changes of
1048   mitochondrial ultrastructure and function during ageing in mice and Drosophila.
1049   Elife. 2017;6. doi:10.7554/eLife.24662

1050 22. Park TJ, Reznick J, Peterson BL, Blass G, Omerbašić D, Bennett NC, et al. Fructose-
1051   driven glycolysis supports anoxia resistance in the naked mole-rat. Science (80-).
1052   2017;356: 307–311. doi:10.1126/science.aab3896

1053 23. Ori A, Toyama BH, Harris MS, Bock T, Iskar M, Bork P, et al. Integrated
1054   Transcriptome and Proteome Analyses Reveal Organ-Specific Proteome
1055   Deterioration in Old Rats. Cell Syst. Elsevier; 2015;1: 224–237.
1056   doi:10.1016/j.cels.2015.08.012

1057 24. Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, et al. Global
1058   quantification of mammalian gene expression control. Nature. Nature Publishing
1059   Group, a division of Macmillan Publishers Limited. All Rights Reserved.;
1060   2011;473: 337–42. doi:10.1038/nature10098

1061 25. Vogel C, Marcotte EM. Insights into the regulation of protein abundance from
1062   proteomic and transcriptomic analyses. Nat Rev Genet. 2012;13: 227–32.
1063   doi:10.1038/nrg3185

1064 26. Nyström T, Yang J, Molin M. Peroxiredoxins, gerontogenes linking aging to
1065   genome instability and cancer. Genes Dev. 2012;26: 2001–8.
1066   doi:10.1101/gad.200006.112

1067 27. Hanzén S, Vielfort K, Yang J, Roger F, Andersson V, Zamarbide-Forés S, et al.
1068   Lifespan Control by Redox-Dependent Recruitment of Chaperones to Misfolded
1069   Proteins. Cell. Elsevier; 2016;166: 140–151. doi:10.1016/j.cell.2016.05.006

1070 28. Biteau B, Karpac J, Supoyo S, DeGennaro M, Lehmann R, Jasper H. Lifespan
1071   extension by preserving proliferative homeostasis in Drosophila. Kim SK, editor.
1072   PLoS Genet. Public Library of Science; 2010;6: 1–15.
1073   doi:10.1371/journal.pgen.1001159

1074 29. Erol A. The Functions of PPARs in Aging and Longevity. PPAR Res. Hindawi;
1075   2007;2007: 39654. doi:10.1155/2007/39654

1076 30. Bustos V, Partridge L. Good Ol' Fat: Links between Lipid Signaling and Longevity.
1077   Trends Biochem Sci. 2017; doi:10.1016/j.tibs.2017.07.001

31. Cellerino A, Ori A. What have we learned on aging from omics studies? Semin Cell
Dev Biol. 2017; doi:10.1016/j.semcdb.2017.06.012

32. Delire B, Lebrun V, Selvais C, Henriet P, Bertrand A, Horsmans Y, et al. Aging
enhances liver fibrotic response in mice through hampering extracellular matrix
remodeling. Aging (Albany NY). Impact Journals, LLC; 2016;9: 98–113.
doi:10.18632/aging.101124

33. Hattori K, Inoue M, Inoue T, Arai H, Tamura H. A novel sulfotransferase
abundantly expressed in the dauer larvae of Caenorhabditis elegans. J Biochem.
2006;139: 355–62. doi:10.1093/jb/mvj041

34. Fielenbach N, Antebi A. C. elegans dauer formation and the molecular basis of
plasticity. Genes Dev. 2008;22: 2149–65. doi:10.1101/gad.1701508

35. Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, et al.
Genes that act downstream of DAF-16 to influence the lifespan of Caenorhabditis
elegans. Nature. 2003;424: 277–83. doi:10.1038/nature01789

36. White RR, Milholland B, MacRae SL, Lin M, Zheng D, Vijg J. Comprehensive
transcriptional landscape of aging mouse liver. BMC Genomics. BioMed Central;
2015;16: 899. doi:10.1186/s12864-015-2061-8

37. Durieux J, Wolff S, Dillin A. The cell-non-autonomous nature of electron transport
chain-mediated longevity. Cell. 2011;144: 79–91. doi:10.1016/j.cell.2010.12.016

38. Baumgart M, Priebe S, Groth M, Hartmann N, Menzel U, Pandolfini L, et al.
Longitudinal RNA-seq analysis of vertebrate aging identifies mitochondrial
complex i as a small-molecule-sensitive modifier of lifespan. Cell Syst. 2016;2:
122–132. doi:10.1016/j.cels.2016.01.014

39. Miwa S, Jow H, Baty K, Johnson A, Czapiewski R, Saretzki G, et al. Low abundance
of the matrix arm of complex I in mitochondria predicts longevity in mice. Nat
Commun. Nature Publishing Group; 2014;5: 3837. doi:10.1038/ncomms4837

40. Sgarbi G, Matarrese P, Pinti M, Lanzarini C, Ascione B, Gibellini L, et al.
Mitochondria hyperfusion and elevated autophagic activity are key mechanisms
for cellular bioenergetic preservation in centenarians. Aging (Albany NY). Impact
Journals, LLC; 2014;6: 296–310. doi:10.18632/aging.100654

41. Sengupta S, Peterson TR, Laplante M, Oh S, Sabatini DM. mTORC1 controls
fasting-induced ketogenesis and its modulation by ageing. Nature. Nature
Research; 2010;468: 1100–1104. doi:10.1038/nature09584

42. Luis NM, Wang L, Ortega M, Deng H, Katewa SD, Li PW-L, et al. Intestinal IRE1 Is
Required for Increased Triglyceride Metabolism and Longer Lifespan under
Dietary Restriction. Cell Rep. Elsevier; 2016;17: 1207–1216.
doi:10.1016/j.celrep.2016.10.003

43. Mitchell SJ, Madrigal-Matute J, Scheibye-Knudsen M, Fang E, Aon M, González-
Reyes JA, et al. Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in
Mice. Cell Metab. Elsevier; 2016;23: 1093–1112. doi:10.1016/j.cmet.2016.05.027

44. Hahn O, Grönke S, Stubbs TM, Ficz G, Hendrich O, Krueger F, et al. Dietary
restriction protects from age-associated DNA methylation and induces epigenetic
reprogramming of lipid metabolism. Genome Biol. 2017;18: 1194.
doi:10.1186/s13059-017-1187-1

45. Kim HE, Grant AR, Simic MS, Kohnz RA, Nomura DK, Durieux J, et al. Lipid
Biosynthesis Coordinates a Mitochondrial-to-Cytosolic Stress Response. Cell.
2016;166: 1539–1552.e16. doi:10.1016/j.cell.2016.08.027

46. Weir HJ, Yao P, Huynh FK, Escoubas CC, Goncalves RL, Burkewitz K, et al. Dietary
Restriction and AMPK Increase Lifespan via Mitochondrial Network and
Peroxisome Remodeling. Cell Metab. 2017; doi:10.1016/j.cmet.2017.09.024

47.   Toth M, Tchernof A. Lipid metabolism in the elderly. Eur J Clin Nutr. Nature
1129   Publishing Group; 2000;54: S121–S125. doi:10.1038/sj.ejcn.1601033

1130   48.   Solomon TPJ, Marchetti CM, Krishnan RK, Gonzalez F, Kirwan JP. Effects of aging
1131   on basal fat oxidation in obese humans. Metabolism. 2008;57: 1141–1147.
1132   doi:10.1016/j.metabol.2008.03.021

1133   49.   St-Onge M-P, Gallagher D. Body composition changes with aging: the cause or the
1134   result of alterations in metabolic rate and macronutrient oxidation? Nutrition.
1135   NIH Public Access; 2010;26: 152–5. doi:10.1016/j.nut.2009.07.004

1136   50.   Franceschi C, Campisi J, LR M, J C, JL K, HY C. Chronic Inflammation
1137   (Inflammaging) and Its Potential Contribution to Age-Associated Diseases.
1138   Journals Gerontol Ser A Biol Sci Med Sci. Oxford University Press; 2014;69: S4–S9.
1139   doi:10.1093/gerona/glu057

1140   51.   Done AJ, Gage MJ, Nieto NC, Traustadóttir T. Exercise-induced Nrf2-signaling is
1141   impaired in aging. Free Radic Biol Med. 2016;96: 130–138.
1142   doi:10.1016/j.freeradbiomed.2016.04.024

1143   52.   Safdar A, deBeer J, Tarnopolsky MA. Dysfunctional Nrf2–Keap1 redox signaling in
1144   skeletal muscle of the sedentary old. Free Radic Biol Med. 2010;49: 1487–1493.
1145   doi:10.1016/j.freeradbiomed.2010.08.010

1146   53.   Bens M, Szafranski K, Holtze S, Sahm A, Groth M, Kestler HA, et al. Naked mole-rat
1147   transcriptome signatures of socially-suppressed sexual maturation and links of
1148   reproduction to aging. bioRxiv. Cold Spring Harbor Laboratory; 2017; 221333.
1149   doi:10.1101/221333

1150   54.   Buffenstein R, Jarvis JUM. The Naked Mole Rat--A New Record for the Oldest
1151   Living Rodent. Sci Aging Knowl Environ. 2002;2002: 7pe–7.
1152   doi:10.1126/sageke.2002.21.pe7

1153   55.   Dammann P, Šumbera R, Maßmann C, Scherag A, Burda H. Extended Longevity of
1154   Reproductives Appears to be Common in Fukomys Mole-Rats (Rodentia,
1155   Bathyergidae). de Polavieja GG, editor. PLoS One. 2011;6: e18757.
1156   doi:10.1371/journal.pone.0018757

1157   56.   Dammann P, Burda H. Sexual activity and reproduction delay ageing in a
1158   mammal. Curr Biol. 2006;16: R117–R118. doi:10.1016/j.cub.2006.02.012

1159   57.   Bens M, Sahm A, Groth M, Jahn N, Morhart M, Holtze S, et al. FRAMA: from RNA-
1160   seq data to annotated mRNA assemblies. BMC Genomics. 2016;17: 54.
1161   doi:10.1186/s12864-015-2349-8

1162   58.   Sahm A, Bens M, Platzer M, Szafranski K. PosiGene: automated and easy-to-use
1163   pipeline for genome-wide detection of positively selected genes. Nucleic Acids
1164   Res. 2017;45: e100–e100. doi:10.1093/nar/gkx179

1165   59.   Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsveld J.
1166   Ultrasensitive proteome analysis using paramagnetic bead technology. Mol Syst
1167   Biol. 2014;10: 757. Available:
1168   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4299378&tool=pm
1169   centrez&rendertype=abstract

1170   60.   Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen J V., Mann M. Andromeda: A
1171   Peptide Search Engine Integrated into the MaxQuant Environment. J Proteome
1172   Res. 2011;10: 1794–1805. doi:10.1021/pr101065j

1173   61.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
1174   p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat
1175   Biotechnol. 2008;26: 1367–72. doi:10.1038/nbt.1511

1176   62.   Elias JE, Gygi SP. Target-decoy search strategy for increased confidence in large-
1177   scale protein identifications by mass spectrometry. Nat Methods. 2007;4: 207–

1178   214. doi:10.1038/nmeth1019

1179   63.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
1180   differential expression analyses for RNA-sequencing and microarray studies.
1181   Nucleic Acids Res. 2015;43: e47–e47. doi:10.1093/nar/gkv007

1182   64.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
1183   Powerful Approach to Multiple Testing [Internet]. Journal of the Royal Statistical
1184   Society. Series B (Methodological). WileyRoyal Statistical Society; 1995. pp. 289–
1185   300. doi:10.2307/2346101

1186   65.   Brosch M, Yu L, Hubbard T, Choudhary J. Accurate and sensitive peptide
1187   identification with Mascot Percolator. J Proteome Res. 2009;8: 3176–81.
1188   doi:10.1021/pr800982s

1189   66.   Gatto L, Lilley KS. MSnbase-an R/Bioconductor package for isobaric tagged mass
1190   spectrometry data visualization, processing and quantitation. Bioinformatics.
1191   2012;28: 288–289. doi:10.1093/bioinformatics/btr645

1192   67.   Huber W, von Heydebreck A, Sültmann H, Poustka A, Vingron M. Variance
1193   stabilization applied to microarray data calibration and to the quantification of
1194   differential expression. Bioinformatics. 2002;18 Suppl 1: S96-104.

1195   68.   Bruderer R, Bernhardt OM, Gandhi T, Miladinović SM, Cheng L-Y, Messner S, et al.
1196   Extending the limits of quantitative proteome profiling with data-independent
1197   acquisition and application to acetaminophen-treated three-dimensional liver
1198   microtissues. Mol Cell Proteomics. 2015;14: 1400–10.
1199   doi:10.1074/mcp.M114.044305

1200   69.   Storey JD. A direct approach to false discovery rates. J R Stat Soc Ser B (Statistical
1201   Methodol. Blackwell Publishers; 2002;64: 479–498. doi:10.1111/1467-
1202   9868.00346

1203   70.   Luo W, Friedman MS, Shedden K, Hankenson KD, Woolf PJ. GAGE: generally
1204   applicable gene set enrichment for pathway analysis. BMC Bioinformatics.
1205   2009;10: 161. doi:10.1186/1471-2105-10-161

1206   71.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.
1207   Gene set enrichment analysis: a knowledge-based approach for interpreting
1208   genome-wide expression profiles. Proc Natl Acad Sci U S A. National Academy of
1209   Sciences; 2005;102: 15545–50. doi:10.1073/pnas.0506580102

1210   72.   Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO summarizes and visualizes long
1211   lists of gene ontology terms. PLoS One. Public Library of Science; 2011;6: e21800.
1212   doi:10.1371/journal.pone.0021800

1213