Analysis of Spounaviruses as a Case Study for the Overdue Reclassification of Tailed Bacteriophages

Jakub Barylski[1], François Enault[2], Bas E. Dutilh[3,4], Margo B.P. Schuller[3], Robert A.

Edwards[5], Annika Gillis[6], Jochen Klumpp[7], Petar Knezevic[8], Mart Krupovic[9], Jens H.

Kuhn[10], Rob Lavigne[11], Hanna M. Oksanen[12], Matthew B. Sullivan[13], Johannes Wittmann[14],

Igor Tolstoy[15], J. Rodney Brister[15], Andrew M. Kropinski[16], Evelien M. Adriaenssens[17*]

This paper is dedicated to Hans-Wolfgang Ackermann, a pioneer of prokaryotic virus

electron microscopy and taxonomy, who died on February 12[th], 2017, at the age of 80. He

was involved in the early stages of this study, and his input is dearly missed.

[1]Adam Mickiewicz University, Institute of Experimental Biology, Department of Molecular

Virology, Poznań, Poland

[2]Université Clermont Auvergne, CNRS, LMGE, F-63000, Clermont-Ferrand, France

[3]Theoretical Biology and Bioinformatics, Department of Biology, Utrecht University,

Utrecht, The Netherlands

[4]Theoretical Biology and Bioinformatics, Department of Biology, Utrecht University,

Utrecht, The Netherlands

[5]*Departments of Biology and Computer Science, San Diego State University, San Diego, CA,*

USA

[6]*Laboratory of Food and Environmental Microbiology, Université Catholique de Louvain,*

Louvain-la-Neuve, Belgium

[7]*Institute of Food, Nutrition and Health, ETH Zurich, Switzerland*

[8]*Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad,*

Serbia

[9]*Unit of Molecular Biology of the Gene in Extremophiles, Department of Microbiology,*

Institut Pasteur, Paris, France

[10]*Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious*

Diseases, National Institutes of Health, Fort Detrick, Frederick, USA

[11]*Laboratory of Gene Technology, KU Leuven, Belgium*

[12]*Department of Biosciences, University of Helsinki, Helsinki, Finland; and Institute of*

Biotechnology, University of Helsinki, Helsinki, Finland

[13]*Departments of Microbiology and Civil, Environmental, and Geodetic Engineering, The*

Ohio State University, Columbus, OH, USA

[14]*Leibniz-Institut DSMZ—Deutsche Sammlung von Mikroorganismen und Zellkulturen*

GmbH, Braunschweig, Germany

[15]*National Center for Biotechnology Information, National Library of Medicine, National*

Institutes of Health, Bethesda, MD, USA

[16]*Departments of Food Science, Molecular and Cellular Biology; and Pathobiology,*

University of Guelph, Guelph, Ontario, Canada

[17]*Institute of Integrative Biology, University of Liverpool, Biosciences Building, Crown*

Street, Liverpool L69 7ZB, United Kingdom; evelien.adriaenssens@gmail.com,

evelien.adriaenssens@liv.ac.uk; corresponding author*

ABSTRACT

It is almost a cliché that tailed bacteriophages of the order *Caudovirales* are the most

abundant and diverse viruses in the world. Yet, their taxonomy still consists of a single order

with just three families: *Myoviridae*, *Siphoviridae*, and *Podoviridae*. Thousands of newly

discovered phage genomes have recently challenged this morphology-based classification,

revealing that tailed bacteriophages are genomically even more diverse than once thought.

Here, we evaluate a range of methods for bacteriophage taxonomy by using a particularly

challenging group as an example, the Bacillus phage SPO1-related viruses of the myovirid

subfamily *Spounavirinae*. Exhaustive phylogenetic and phylogenomic analyses indicate that

the spounavirins are consistent with the taxonomic rank of family and should be divided into

at least five subfamilies. This work is a case study for virus genomic taxonomy and the first

step in an impending massive reorganization of the tailed bacteriophage taxonomy.

KEYWORDS

Virus taxonomy; bacteriophage taxonomy; *Caudovirales*; spounaviruses; phylogenetics;

phylogenomics

By the end of 2017, 3,033 complete genomes of tailed phages were available in the

National Center for Biotechnology Information (NCBI) RefSeq database and a further 18,753

partial genomes were found in International Nucleotide Sequence Database Collaboration

databases (Karsch-Mizrachi et al. 2012; O'Leary et al. 2016). The classification of this massive

group is the formal responsibility of the Bacterial and Archaeal Viruses Subcommittee of the

International Committee on the Taxonomy of Viruses (ICTV). In recent years, we (the

Subcommittee) focused on classifying newly described phages into species and genera, within

established viral families (Lavigne et al. 2008, 2009, Adriaenssens et al. 2015, 2017; Krupovic

et al. 2016). However, once our attention shifted towards higher order relationships, we found

that the ranks available in virus taxonomy (*species*, *genus*, *subfamily*, *family*, and *order*) were

no longer sufficient for the description of phage diversity. The limitation is particularly acute

in the case of the order *Caudovirales*—the most abundant and diverse group of viruses (Paez-

Espino et al. 2016; Roux et al. 2016; Nishimura et al. 2017). Indeed, the diversity of

caudovirads surpasses that of any other virus taxon. A recent analysis of the dsDNA virosphere

using a bipartite network approach, whereby viral genomes are connected via shared gene

families, demonstrated that the global network of dsDNA viruses consists of at least 19

modules, 11 of which correspond to caudovirads (Iranzo et al. 2016). Each of eight remaining

modules encompasses one or more families of eukaryotic or archaeal viruses. Consequently,

each of the 11 caudovirad modules could be considered a separate family. Despite this

remarkable diversity, all caudovirads are currently classified into three families - *Myoviridae,*

Podoviridae, and *Siphoviridae*. These families were historically established on morphological

features alone, forming an artificial classification ceiling.

In this study, the Subcommittee explored the diversity of the order *Caudovirales* on the

example of the *Spounavirinae* subfamily, a large group of myoviruses that forms one of the

above-mentioned caudovirad modules (Iranzo et al. 2016; Bolduc et al. 2017). The subfamily

was proposed in 2009 by Lavigne et al. to harbor Bacillus phage SPO1, Staphylococcus phage

Twort, Staphylococcus phage K, Staphylococcus phage G1, Listeria phage P100, and Listeria

phage A511 (Lavigne et al. 2009). The unifying characteristics of members of this subfamily

are: the host belongs to the bacterial phylum *Firmicutes*; strictly virulent lifestyle; myovirion

morphology; terminally redundant, non-permuted dsDNA genome 127–157 kb in length; and

"considerable amino acid homology" (Klumpp et al. 2010). The strictly virulent lifestyle of

these viruses has been somewhat disputed (Schuch and Fischetti 2009; Yuan et al. 2015) but

still remains a rule of thumb for the taxon inclusion. Since the inception of the subfamily, the

number of its members has grown significantly, and its taxonomic structure was contested

several times (Klumpp et al. 2010; Barylski et al. 2014; Iranzo et al. 2016; Krupovic et al.

2016; Adriaenssens et al. 2017; Bolduc et al. 2017). At present, the *Spounavirinae* subfamily

includes six genera (*Kayvirus, P100virus, Silviavirus, Spo1virus, Tsarbombavirus and*

Twortvirus) and three unassigned species (*Enterococcus virus phiEC24C, Lactobacillus virus*

Lb338-1 and Lactobacillus virus LP65).

Here, we reevaluated the current classification of spounavirins and related viruses and

outlined a better fitting scheme, in the process also reaffirming the need for major changes in

phage taxonomy that will better accommodate the observed genomic diversity.

MATERIALS & METHODS

Creation of the Dataset

Genome sequences of known spounavirins and spouna-like viruses were retrieved from

GenBank or (preferably) RefSeq databases based on literature data, ICTV and taxonomic

classifications provided by the NCBI. Records representing genomes of candidate

spounavirins were retrieved by searching the same databases with the tBLASTx algorithm

using as a queries terminase and major capsid proteins of Bacillus phage SPO1,

Staphylococcus phage Twort, Bacillus phage Bastille, Listeria phage A511, Enterococcus

phage φEF24C, and Lactobacillus phage LP65 [type isolates of the original subfamily,

(Altschul et al. 1990; Brister et al. 2015)]. Sequences were manually curated and pre-

clustered using Cluster Analysis of Sequences (CLANS; E-value cut-off 1e-10) to confirm

their spounaviral affiliation (Frickey and Lupas 2004). This search yielded a set of 93

complete virus genomes, which were used in the following analyses (Supplementary Table

1).

The genomes were re-annotated using PROKKA with the settings --kingdom Viruses, -

-E-value 1e-6 (Seemann 2014). All original genome sequences are available from NCBI

(accession number information listed in Supplementary Table 1) and the reannotated

genomes from Github (github.com/evelienadri/herelleviridae).

Genome-based Analyses

Gegenees (Ågren et al. 2012) was used to analyze genome similarities (fragment length

200 bp; step length 100 bp). Pairwise identities between all genomes under study were

determined using BLASTn and tBLASTx algorithms with default parameters (Camacho et al.

2009). Symmetrical identity scores (% SI) were calculated for each pairwise comparison

using the formula:

$\% \text{ SI} = 2.0 \text{ x } \frac{HL \text{ x } HI}{QL + SL}$

in which the HL is defined as the hit length of the BLAST hit, HI is defined as the

percentage hit identity, QL is defined as the query length, and SL is defined as the subject

length.

Symmetrical identity scores were converted into distances using the formula:

$\text{Distance} = \sqrt[2]{1.0 - \%SI \div 100}$

The resulting distance matrix was hierarchically clustered (complete linkage) using the

hclust function of R (Development Core Team 2008). Trees were visualized using Itol (Letunic

and Bork 2007).

Additionally, pairwise comparisons of the nucleotide sequences using VICTOR, a

Genome-BLAST Distance Phylogeny (GBDP) method, were conducted under settings

recommended for prokaryotic viruses (Meier-Kolthoff et al. 2014; Meier-Kolthoff and Göker

2017). The resulting intergenomic distances (including 100 replicates each) were used to infer

a balanced minimum evolution tree with branch support via FASTME including subtree

pruning and regrafting post-processing (Lefort et al. 2015) for each of the formulas D0, D4,

and D6, respectively. Trees were visualized with FigTree (Rambaud 2007). Taxon

demarcations at the species, genus and family rank were estimated with the OPTSIL program

(Göker et al. 2009), the recommended clustering thresholds (Meier-Kolthoff and Göker 2017),

and an F value (fraction of links required for cluster fusion) of 0.5 (Meier-Kolthoff et al. 2014).

Proteome-based Analyses

The Phage Proteomic Tree was constructed as described previously (Rohwer and

Edwards 2002) and detailed at

https://github.com/linsalrob/PhageProteomicTree/tree/master/spounavirus. Briefly, the

protein sequences were extracted and clustered using BLASTp. These clusters were refined

by Smith-Waterman alignment using CLUSTALW version 2 (Larkin et al. 2007).

Alignments were scored using open-source PROTDIST from the phylogeny inference

package (PHYLIP) (Felsenstein 1989). Alignment scores were averaged and weighted as

described previously (Rohwer and Edwards 2002) resulting in the final tree.

Orthologous protein clusters (OPCs) were constructed using GET_HOMOLOGUES

software, which utilizes several independent clustering methods (Contreras-Moreira and

Vinuesa 2013). To capture as many evolutionary relationships as possible, a greedy

COGtriangles algorithm was applied with a 50% sequence identity threshold, 50% coverage

threshold, and an E-value cut-off equal to 1e-10 (Kristensen et al. 2010). The results were

converted into an orthologue matrix with the "compare_clusters" script (part of the

GET_HOMOLOGUES suite) (Felsenstein 1989).

The OPCs defined above were used to compute the genomic fluidity for each pair of

genomes. For two genomes i and j:

Fluidity(i,j) $= \frac{Ui+Uj}{Mi+Mj}$

with Ui being the number of genes of i not found in j and Mi being the number of

genes in i (Kislyuk et al. 2011). The resulting distance matrix was hierarchically clustered

(complete linkage) using the hclust function of R (Development Core Team 2008). Trees

were visualized using Itol (Letunic and Bork 2007).

Multiple alignments were generated for each OPC using Clustal Omega (Sievers et al.

2011). For each cluster, the amino acid identity between all protein pairs inside a cluster was

determined using multiple alignment. For all genome pairs, the AAI (Konstantinidis and

Tiedje 2005) was then computed and transformed into distance using the formula:

Distance $= \frac{100-AAI}{100}$

The resulting distance matrix was clustered and visualized as described above.

OPCs and multiple alignments for each cluster were used to determine a distance

similar to the distance used to generate the Phage Proteomic Tree. To estimate protein

distances, in this case, the distance (dist) function of the seqinR package (Charif et al.

2005)was preferred to PROTDIST of the PHYLIP package (Felsenstein 1989) as the

resulting distances are between 0 and 1. Proteomic distances were then computed using the

same formula as for the Phage Proteomic Tree. The results were clustered and visualized as

described above.

The Dice score is based on reciprocal BLAST searches between all pairs of genomes A

and B (Mizuno et al. 2013). The total summed bit-scores of all tBLASTx hits with ≥30%

identity, alignment length ≥30 amino acids, and E-value ≤0.01 was converted to a distance

DAB as follows:

$$DAB = 1 - \frac{SAB + SBA}{SAA + SBB}$$

in which SAB and SBA represent the summed bit-scores between tBLASTx searches of

A versus B, and B versusA, respectively, while SAA and SBB represent the summed tBLASTx

bit-scores of the self-queries of A and B, respectively. The resulting distance matrix was

clustered with BionJ (Gascuel 1997).

To investigate a genomic synteny-based classification signal, we developed a gene order-

based metric built on dynamic programming, the Gene Order Alignment Tool (GOAT, Schuller

et al.: Python scripts are available on request, manuscript in preparation). GOAT first identified

protein-coding genes in the 93 spounavirin and spouna-like virus genomes using Prodigal

V2.6.3 in anonymous mode (Hyatt et al. 2010), and assigned them to the latest pVOGs

(Grazziotin et al. 2017)). pVOG alignments (9,518) were downloaded (http://dmk-

brain.ecn.uiowa.edu/pVOGs/) and converted to profiles of hidden Markov models (HMM)

using HMMbuild (HMMer 3.1b2, (Finn et al. 2011)). Proteins were assigned to pVOGs using

HMMsearch (E-value <10-2) and used to generate a synteny profile of every genome. GOAT

accounted for gene replacements and distant homology by using an all-vs-all similarity matrix

between pVOG pairs based on HMM-HMM similarity (HH-suite 2.0.16) (Söding et al. 2005)).

Distant HHsearch similarity scores between protein families were calculated as the average of

reciprocal hits and used as substitution scores in the gene order alignment. The GOAT

algorithm identified the optimal gene order alignment score between two virus genomes by

implementing semi-global dynamic programming alignment based only on the order of pVOGs

identified on every virus genome. To account for virus genomes being cut at arbitrary positions

during sequence assembly, GOAT transmutes the gene order at all possible positions and in

both sense and antisense directions in search of the optimal alignment score. The optimal

GOAT alignment score GAB between every pair of virus genomes A and B, was converted to

a distance DAB as follows:

$$DAB = 1 - \frac{GAB + GBA}{GAA + GBB}$$

in which GAB and GBA represent the optimal GOAT score between A and B, and B and

A, respectively, while GAA and GBB represent the GOAT scores of the self-alignments of A

and B, respectively. This pairwise distance matrix was clustered with BionJ (Gascuel 1997).

Prokka re-annotated genomes were used to create pan-, core-, and accessory genomes of

all selected spounavirins and spouna-like viruses (Seemann 2014). The annotations were

analyzed using Roary (Page et al. 2015) with a 50% length BLASTp identity threshold for

homologous genes. Roary functions as follows: CD-HIT (Fu et al. 2012) was used to pre-

cluster protein sequences and perform an all-vs-all comparison of protein sequences with

BLASTp to identify orthologs and paralogs within the genomes. Markov cluster algorithm

(MCL) (Enright et al. 2002) was then used to cluster the genomes based on the presence and

absence of the accessory genes. The gene presence-absence output table from Roary was then

imported into R and pairwise shared gene contents were calculated for each combination of

genomes   using   a   custom   R-script   (available   from

github.com/evelienadri/herelleviridae/tree/master). The resulting tree file was visualized using

FigTree v1.4.3 (Rambaud 2007).

Single Protein Phylogenies

Based on the OPC and pVOG analyses, we chose nine well-annotated protein clusters

present in all 93 spounavirins and spouna-like viruses. Selected clusters included: DNA

helicases, major capsid proteins, tail sheath proteins, two different groups of baseplate

proteins, and four clusters with no known function. The members of these clusters were

aligned using Clustal Omega with default parameters (Sievers et al. 2011). Resulting

alignments were analyzed with ProtTest 3.4 (Darriba et al. 2011) to determine a suitable

protein evolution model (only variations of models compatible with downstream software

like JTT (Jones et al. 1992) and WAG (Whelan and Goldman 2001) were considered).

Estimated models were used to generate phylograms with FastTree 2.1.7 (Price et al. 2010).

The program implements the approximately maximum-likelihood method with Shimodaira-

Hasegawa tests to generate the tree and calculate support of the splits. This approach is much

faster than "traditional" maximum-likelihood methods with negligible accuracy loss (Price et

al. 2010; Darriba et al. 2011; Liu et al. 2011).

RESULTS

General Overview

To determine the phylogenetic relationship between 93 known and alleged spounavirins,

we used genomic, proteomic and marker gene-based comparative strategies. Regardless of the

adopted phylogenetic approach applied, five separate, clear-cut clusters were identified. We

believe that these clusters have a common origin and ought to come together under one

umbrella taxon. We suggest to name this taxon "*Herelleviridae,*" in honor of the 100th

anniversary of the discovery of prokaryotic viruses by Félix d'Hérelle (Table 1, Figs. 1-3 and

Supplementary Table 1). The first cluster (here suggested to retain the name *Spounavirinae*)

groups *Bacillus*-infecting viruses that are similar to Bacillus phage SPO1. The second cluster

includes *Bacillus*-infecting viruses that resemble phage Bastille instead (named

"*Bastillevirinae*" after the type species (Barylski et al. 2014)). The third cluster

("*Brockvirinae*," named in honor of Thomas D. Brock, a microbiologist known for discovery

of hyperthermophiles who worked on *Streptococcus* phages early in his career) comprises

currently unclassified viruses of enterococci that are similar to Enterococcus phage φEF24C.

The fourth cluster ("*Twortvirinae*," named in honor of Frederick William Twort, the

bacteriologist who discovered prokaryotic viruses in 1915) gathers staphylococci-infected

viruses that are similar to Staphylococcus phage Twort. The remaining cluster

("*Jasinskavirinae*," named in honor of Stanisława Jasińska-Lewandowska who was one of the

first to study *Listeria* and its viruses) consists of viruses infecting *Listeria* that are similar to

Listeria phage P100. The classification in five clusters left three viruses unassigned at this rank:

Lactobacillus phage Lb338, Lactobacillus phage LP65, and Brochothrix phage A9.

These robust clusters can be further subdivided into smaller clades that correspond well

with the currently accepted genera. The evidence supporting this suggested taxonomic re-

classification is presented in the following sections.

Genome-based Analyses

BLASTn analysis revealed that the genomes of several viruses were similar enough to

consider them strains of the same species (they shared >95% nucleotide identity,

Supplementary Fig. 1). The Staphylococcus viruses fell into four distinct, yet closely related

groups corresponding to the established genera Twortvirus, Sep1virus, Silviavirus, and

Kayvirus (Supplementary Fig. 1). With the exception of Enterococcus phage EFDG1, all

Enterococcus viruses clustered as a clade representing a new genus (here suggested to be

named "Kochikohdavirus" after the place of origin of the type virus of the clade,

Enterococcus phage φEF24C (Uchiyama et al. 2008a, 2008b)). The Bacillus viruses clustered

into the established genera Spo1virus, Cp51virus, Bastillevirus, Agatevirus, B4virus,

Bc431virus, Nit1virus, Tsarbombavirus, and Wphvirus, with three species remaining

unassigned at the genus rank (Table 1). These results were also confirmed with the Virus

Classification and Tree Building Online Resource (VICTOR), a genome-BLAST distance

phylogeny (GBDP) method (Supplementary Fig. 2) (Meier-Kolthoff and Göker 2017) and the

Dice score (Supplementary Fig. 3), a tBLASTx-based measure that compares whole genome

sequences at the amino acid level (Mizuno et al. 2013).

291   The patterns coalesced at a higher taxonomic level when the genomes were analyzed

292 using tBLASTx (Supplementary Fig. 4). The *Enterococcus* viruses clustered into a single group

293 sharing 41% genome identity, whereas the *Bacillus* viruses fell into two major groups, a group

294 combining the genera *Spo1virus* and *Cp51virus*, and the remainder. All *Staphylococcus* viruses

295 clustered above ≈36% genome identity, whereas *Listeria* viruses grouped with more than 79%

296 genome identity. Overall, all these genomes were related at the level of at least 15% genome

297 identity. *Lactobacillus* and *Brochothrix* viruses remained genomic orphans, peripherally

298 related to the remainder of the viruses in this assemblage.

300 Table 1. Suggested new classification of the 93 spounavirins and spouna-like viruses in the new family "Herelleviridae."[a]

Order	Family	Subfamily	Genus	Species
Caudovirales	*"Herelleviridae"*	*"Bastillevirinae"*	*Agatevirus*	*Bacillus virus Agate, Bacillus virus Bobb, Bacillus virus Bp8pC* (Bp8p-T)
			B4virus	*Bacillus virus AvesoBmore, Bacillus virus B4* (B5S), *Bacillus virus Bigbertha, Bacillus virus Riley, Bacillus virus Spock, Bacillus virus Troll*
			Bastillevirus	*Bacillus virus Bastille, Bacillus virus CAM003, "Bacillus virus Evoli", "Bacillus virus HoodyT"*
			Bc431virus	*Bacillus virus Bc431, Bacillus virus Bcp1, Bacillus virus BCP82, Bacillus virus JBP901*
			Nit1virus	*Bacillus virus Grass, Bacillus virus NIT1, Bacillus virus SPG24*
			Tsarbombavirus	*Bacillus virus BCP78* (BCU4), *Bacillus virus TsarBomba*
			Wphvirus	*Bacillus virus BPS13, Bacillus virus Hakun), Bacillus virus Megatron* (Eyuki), *Bacillus virus WPh, "Bacillus virus BPS10C"*
		"*Brockvirinae*"	"*Kochikohdavirus*"	"*Enterococcus virus ECP3*", "*Enterococcus virus EF24C*" (phiEFC24C-P2), "*Enterococcus virus EFLK1*"
			Unassigned	"*Enterococccus virus EFDG1*"
		"*Jasinskavirinae*"	*P100virus*	*Listeria virus A511, Listeria virus P100* (List-36, LMSP-25, AvB_LmoM_AG20, LP-125, LP-064, LP-083-2, LP-124, LP-125, LP-048, LMTA-34, LMTA-94, LMTA-148, LMTA-57, WIL-1)
		Spounavirinae	*Cp51virus*	*Bacillus virus CP51, Bacillus virus JL, Bacillus virus Shanette*
			Spo1virus	*Bacillus virus Camphawk, Bacillus virus SPO1*
			Unassigned	*"Bacillus virus Mater", "Bacillus virus Moonbeam", "Bacillus virus SIOphi"*
		"*Twortvirinae*"	*Kayvirus*	*Staphylococcus virus G1, Staphylococcus virus G15, Staphylococcus virus JD7, Staphylococcus virus K, Staphylococcus virus MCE2014, Staphylococcus virus P108, Staphylococcus virus Rodi, Staphylococcus virus S253, Staphylococcus virus S25-4, Staphylococcus virus SA12, "Staphylococcus virus Sb1"* (676Z, A3R, A5W, Fi200W, IME-SA1, IME-SA118, IME-SA119, IME-SA2, ISP, MSA6, P4W, SA5, Staph1N, Team1)
			Silviavirus	*Staphylococcus virus Remus* (Romulus), *Staphylococcus virus SA11*
			Sep1virus	*Staphylococcus virus IPLAC1C, Staphylococcus virus SEP1*
			Twortvirus	*Staphylococcus virus Twort*
		Unassigned	Unassigned	*"Lactobacillus virus Lb338", "Lactobacillus virus LP65", "Brochothrix virus A9"*

301 [a] The species listed here are representing the 93 genome dataset on which all analyses have been performed. Species names ratified in 2017 and

302 later are not included. Phage isolates at the subspecies or strain level are indicated between brackets. Non-ICTV-ratified taxa are indicated

303 between quotation marks.

Figure 1. Genome-based clustering trees of 93 spounavirin and spouna-like viruses. A)

Clustering was performed using nucleotide similarities (BLASTn) or B) translated nucleotide

similarities (tBLASTx,). Genomes were compared in a pairwise fashion using Gegenees,

transformed into a distance matrix, clustered using R, and visualized as trees using Interactive

tree of life (Itol). The trees were rooted at Brochotrix phage A9. Genera and suggested

subfamilies are delineated with colored squares and colored circles, respectively.

Proteome-based Analyses

The virus proteomic tree showed five robust groupings corresponding with the

suggested subfamilies (Fig. 2). Viruses that infect *Bacillus* fell into two groups as described

before, represented by the revised *Spounavirinae* subfamily and the suggested new subfamily

"*Bastillevirinae*." Similarly, the *Listeria* and *Staphylococcus* viruses formed their own

clusters, "*Jasinskavirinae*" and "*Twortvirinae*", respectively. This clustering suggests that the

major *Bacillus, Listeria,* and *Staphylococcus* virus groups are represented, but that further

representatives are required from the under-sampled groups. The suggested "*Brockvirinae*"

subfamily is under-sampled, and the grouping observed in the tree was not as well-supported

as the other clusters.

Among 1,296 singleton proteins and 2,070 protein clusters defined using the

orthologous protein clusters (OPC) approach, we identified 12 clusters common for all viruses

(Table 2, Supplementary Table 2). Classification of the viral proteins using prokaryotic virus

orthologous groups (pVOGs) showed that 38 pVOGs were shared between all 93 virus

genomes (Table 2, Supplementary Table 3). This finding was in stark contrast with the results

from core genome analysis using Roary, which revealed only one core gene (the tail tube

protein gene). Upon closer inspection of the gene annotations, we found that these analyses

might have been confounded by the presence of introns and inteins in many of the core genes

(Supplementary Figs. 5–6). Indeed, many genes of spounavirins and related viruses are invaded

by mobile introns or inteins (Goodrich-Blair et al. 1990; Lavigne and Vandersteegen 2013).

These gaps in coding sequences challenge gene prediction tools and introduce additional bias

in similarity-based cluster algorithms.

Figure 2. Predicted proteome-based clustering trees of 93 spounavirin and spouna-like

viruses. A) Clustering was performed using the Phage Proteomics Tree approach and B)

proteomic distance. Distances were calculated pairwise between all sets of predicted

proteomes, clustered with R, and visualized using Itol. The trees were rooted at Brochotrix

phage A9. Genera and suggested subfamilies are delineated with colored squares and colored

circles, respectively.

The pairwise comparison of the predicted proteome content of the viruses revealed a

very low overall similarity at the protein level (Supplementary Fig. 7). Most viruses shared less

than 10% of their proteins. However, at the suggested new subfamily rank, we observed

obvious virus groups sharing their proteomes. The *Enterococcus* viruses ("*Brockvirinae*")

shared over 35% of their protein content. The members of the *Bacillus* virus genera *Spo1virus*

and *Cp51virus* of the subfamily *Spounavirinae* (*sensu stricto*) had approximately 20% of their

proteins in common, whereas the *Bacillus* virus genera *Bastillevirus*, *B4virus*, *Bc431virus*,

Agatevirus, *Nit1virus*, *Tsarbombavirus*, and *Wphvirus* ("*Bastillevirinae*") and the

Staphylococcus virus genera *Kayvirus*, *Silviavirus*, and *Twortvirus* ("*Twortvirinae*") shared

over 25% and over 30% of their predicted proteomes, respectively.

Genomic fluidity is a measure of the dissimilarity of genomes evaluated at the gene

level (Kislyuk et al. 2011). Accordingly, the genomic fluidity results followed those obtained

using proteome content analysis (Supplementary Fig. 8). Despite a high genomic fluidity for

most of these viruses, the newly suggested subfamilies and genera were all supported.

The topology of the dendrogram obtained using the average amino acid identity (AAI)

approach also supported the suggested new taxonomic scheme (Supplementary Fig. 8). The

AAI was greater than 35% within each subfamily and greater than 67% within each genus. The

AAI of all viruses analyzed in this study was not lower than 22%. The members of the genus

Wphvirus had the lowest AAI (76%) and the lowest AAI for a pair of proteomes (67% between

Bacillus phage W.Ph. and Bacillus phage Eyuki) but surprisingly they had a mid-range

genomic fluidity (0.15), suggesting that the protein sequences of wphviruses might have

evolved rapidly.

Table 2. Core genes with putative annotated functions identified in all 93 spounavirin

and spouna-like virus genomes.

Putative function of the core gene identified[a]	pVOG[b] / OPC[b] ID	Identification method
DnaB-like helicase	VOG0025, OPC6121	OPC, pVOG
Baseplate J-like protein	VOG4691, VOG4644, OPC6132	OPC, pVOG
Tail sheath protein	VOG0067, OPC6142	OPC, pVOG
Terminase large subunit	VOG0051, OPC6160	pVOG
Major capsid protein	VOG0061, OPC6148	OPC, pVOG
Prohead protease	VOG4568, OPC6150	pVOG
Portal protein	VOG4556, OPC6151	OPC, pVOG
DNA primase	VOG4551	pVOG
DNA polymerase I	VOG0668, OPC6097	OPC, pVOG
RNA polymerase	VOG0118	pVOG
Recombination exonuclease	VOG4575	pVOG
Recombination endonuclease	VOG0083	pVOG
Tail tape measure protein	VOG0069	pVOG
Tail tube protein	VOG0068, OPC6141	OPC, pVOG, Roary

[a] The full lists of orthologous proteins and pVOGs are available in Supplementary

Tables 2 and 3, respectively.

[b] pVOG, prokaryotic virus orthologous group; OPC, orthologous protein clusters.

The pangenome of the spounavirins and spouna-like viruses (4,182 genes) as calculated

using Roary (Page et al. 2015) was further analyzed by clustering the genomes based on the

presence or absence of the accessory genes (Supplementary Fig. 9). The obtained tree

supported the current division of the viruses into approved genera and the suggested new

subfamilies.

Many virus genomes are thought to be highly modular, with recombination and

horizontal gene transfer potentially resulting in "mosaic genomes" (Juhala et al. 2000;

Krupovic et al. 2011). By clustering the spounavirin and spouna-like virus genomes based

solely on the gene order of their genomes, we investigated whether gene synteny was preserved

(Supplementary Fig. 10). The results revealed that genomic rearrangements leave a measurable

evolutionary signal in all lineages, since the genomic architecture analysis clustered all viruses

according the suggested taxa. The potential exception was Bacillus phage Moonbeam

(Cadungog et al. 2015). However, we did not observe the high modularity that may be expected

with rampant mosaicism. The lack of considerable mosaicism supports the recent findings by

Bolduc et al. that, at most, about 10% of reference virus genomes have a high degree of

mosaicism (Bolduc et al. 2017). Thus, while the gene order in viruses belonging to the newly

suggested family "*Herelleviridae*" is not necessarily strictly conserved, we observed a clear

evolutionary pattern that is consistent with the sequence-based approaches tested in this study.

Single Protein Phylogenies

The phylogenetic trees based on comparisons of the major capsid, tail sheath, and DnaB-

like helicase proteins are presented in Figure 3. All nine phylograms based on OPC are included

in the trees in Supplementary Figure 11. For nearly all single marker trees, the topologies

supported the suggested taxonomic scheme. Generally, each taxon is represented as a separate

branch on the dendrogram. Notable exceptions could be found in two trees based on

hypothetical proteins (OPCs 10357 and 10386). The first protein places the revised subfamily

Spounavirinae as a subclade of "*Bastillevirinae*" and the second protein shuffled viruses from

the genera *Silviavirus* and *Kayvirus*. This result may indicate that some degree of horizontal

gene transfer occurs between groups, which share common hosts.

Figure 3. Phylogenetic trees based on comparisons of major protein clusters of amino

acid sequences of spounavirin and spouna-like viruses. Amino acid sequences from A) the

major capsid protein, B) tail sheath protein, and C) helicase were aligned with Clustal Omega,

and trees were generated using FastTree maximum likelihood with Shimidaira-Hasegawa tests.

The scale bar represents the number of substitutions per site. The trees were rooted at

Brochotrix phage A9. Genera and suggested subfamilies are delineated with colored squares

and colored circles, respectively.

DISCUSSION

Taxonomic methods must constantly develop to keep up with the ever-increasing pace

of virus discovery. In the rapidly expanding field of phage studies, this requirement proved to

be problematic, and although there are more than 3000 publicly available caudovirad genomes,

only 873 have been officially classified by the ICTV (Davison 2017). The remaining genomes

are provisionally stashed within "unclassified" bins attached to the order or associated families

(Adams et al. 2017; Simmonds et al. 2017). We believe that this work is an important step

toward solving the problem of these "phage orphans". This study represents the first example

of a true taxonomic assessment from an "ensemble of methods". We are encouraged that the

combination of genome sequence analyses, virus proteomic trees, core protein clusters, gene

order genomic synteny (GOAT), and single gene phylogenies yields consistent and

complementary results. Convergence of the results reasserts the usefulness of genome-based

classification at a higher taxonomic rank and the ability of these methods to accommodate viral

diversity.

All evidence considered, we suggest that the spounavirins should be removed from the

family *Myoviridae* and given a family rank. Hence, we propose establishing a new family

"*Herelleviridae*", in the order *Caudovirales* next to a smaller *Myoviridae* family. The new

family would contain five subfamilies: *Spounavirinae* (sensu stricto), "*Bastillevirinae*",

"*Twortvirinae*", "*Jasinkavirinae*", and "*Brockvirinae*", each comprising the genera listed in

Table 1 (with additional information in S1 Table). The suggested classification corresponds

well with the taxonomy of the hosts and leaves only 3% of viruses within the new family

unassigned at the genus and subfamily rank. These unassigned viruses may represent clades at

the genus and subfamily rank that are still under-sampled.

We believe that detachment of spounavirins from their original taxon will soon be

followed by abolishment of the *Podoviridae*, *Myoviridae* and *Siphoviridae* families, in

combination with the addition of new taxon ranks (e.g., class) required to accommodate the

observed diversity of tailed phages. Substitution of the current families with a set of new

"phylogenomic" ones will more faithfully reflect the genetic relationships of these viruses. This

change does not remove the historically established virus morphotypes observed among

caudovirads: myovirids forming particles with contractile tails, siphovirids forming particles

with long non-contractile tails, and podovirids forming particles with short non-contractile

ones. By disconnecting morphotype and family classification, taxonomically related clades can

be grouped across different morphotypes. Such an approach would solve the problems of the

muviruses that are suggested to be classified in the family "*Saltoviridae*" (Hulo et al. 2015)

and potentially the broad set of Escherichia phage lambda-related viruses that are currently

distributed among the families *Siphoviridae* and *Podoviridae* (Grose and Casjens 2014).

REFERENCES

Adams M.J., Lefkowitz E.J., King A.M.Q., Harrach B., Harrison R.L., Knowles N.J.,

Kropinski A.M., Krupovic M., Kuhn J.H., Mushegian A.R., Nibert M.L., Sabanadzovic

S., Sanfaçon H., Siddell S.G., Simmonds P., Varsani A., Zerbini F.M., Orton R.J., Smith

D.B., Gorbalenya A.E., Davison A.J. 2017. 50 years of the International Committee on

Taxonomy of Viruses: progress and prospects. Arch. Virol. 162:1441–1446.

Adriaenssens E.M., Edwards R., Nash J.H.E., Mahadevan P., Seto D., Ackermann H.-W.,

Lavigne R., Kropinski A.M. 2015. Integration of genomic and proteomic analyses in the

classification of the *Siphoviridae* family. Virology. 477:144–154.

Adriaenssens E.M., Krupovic M., Knezevic P., Ackermann H.-W., Barylski J., Brister J.R.,

Clokie M.R.C., Duffy S., Dutilh B.E., Edwards R.A., Enault F., Jang H. Bin, Klumpp J.,

Kropinski A.M., Lavigne R., Poranen M.M., Prangishvili D., Rumnieks J., Sullivan

M.B., Wittmann J., Oksanen H.M., Gillis A., Kuhn J.H. 2017. Taxonomy of prokaryotic

viruses: 2016 update from the ICTV bacterial and archaeal viruses subcommittee. Arch.

Virol. 162:1153–1157.

Ågren J., Sundström A., Håfström T., Segerman B. 2012. Gegenees: Fragmented alignment

of multiple genomes for determining phylogenomic distances and genetic signatures

unique for specified target groups. PLoS One. 7:e39107.

Altschul S.F., Gish W., Miller W., Myers E.W., Lipman D.J. 1990. Basic local alignment

search tool. J. Mol. Biol. 215:403–410.

Barylski J., Nowicki G., Goździcka-Józefiak A. 2014. The discovery of phiAGATE, a novel

phage infecting *Bacillus pumilus*, leads to new insights into the phylogeny of the

subfamily *Spounavirinae*. PLoS One. 9:e86632.

Bolduc B., Jang H. Bin, Doulcier G., You Z.-Q., Roux S., Sullivan M.B. 2017. vConTACT:

an iVirus tool to classify double-stranded DNA viruses that infect Archaea and Bacteria.

PeerJ. 5:e3243.

Brister J.R., Ako-adjei D., Bao Y., Blinkova O. 2015. NCBI Viral Genomes Resource.

Nucleic Acids Res. 43:D571–D577.

Cadungog J.N., Khatemi B.E., Hernandez A.C., Kuty Everett G.F. 2015. Complete genome

sequence of *Bacillus megaterium* myophage Moonbeam. Genome Announc. 3:e01428-

14.

Camacho C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., Madden T.L.

2009. BLAST+: architecture and applications. BMC Bioinformatics. 10:421.

Charif D., Thioulouse J., Lobry J.R., Perrière G. 2005. Online synonymous codon usage

analyses with the ade4 and seqinR packages. Bioinformatics. 21:545–547.

Contreras-Moreira B., Vinuesa P. 2013. GET_HOMOLOGUES, a versatile software package

for scalable and robust microbial pangenome analysis. Appl. Environ. Microbiol.

79:7696–7701.

Darriba D., Taboada G.L., Doallo R., Posada D. 2011. ProtTest 3: Fast selection of best-fit

models of protein evolution. Bioinformatics. 27:1164–1165.

Davison A.J. 2017. Journal of General Virology – Introduction to "ICTV Virus Taxonomy

Profiles." J. Gen. Virol. 98:1–1.

Development Core Team R. 2008. R: A language and environment for statistical computing.

Available from http://www.r-project.org.

Enright A.J., Van Dongen S., Ouzounis C.A. 2002. An efficient algorithm for large-scale

detection of protein families. Nucleic Acids Res. 30:1575–1584.

Felsenstein J. 1989. PHYLIP - Phylogeny inference package - v3.2. Cladistics. 5:164–166.

Finn R.D., Clements J., Eddy S.R. 2011. HMMER web server: Interactive sequence

similarity searching. Nucleic Acids Res. 39:29–37.

Frickey T., Lupas A. 2004. CLANS: A Java application for visualizing protein families based

on pairwise similarity. Bioinformatics. 20:3702–3704.

Fu L., Niu B., Zhu Z., Wu S., Li W. 2012. CD-HIT: Accelerated for clustering the next-

generation sequencing data. Bioinformatics. 28:3150–3152.

Gascuel O. 1997. BIONJ: an improved version of the NJ algorithm based on a simple model

of sequence data. Mol. Biol. Evol. 14:685–695.

Göker M., García-Blázquez G., Voglmayr H., Tellería M.T., Martín M.P. 2009. Molecular

taxonomy of phytopathogenic fungi: A case study in Peronospora. PLoS One. 4:8–10.

Goodrich-Blair H., Scarlato V., Gott J.M., Xu M.-Q., Shub D.A. 1990. A self-splicing group

I intron in the DNA polymerase gene of *Bacillus subtilis* bacteriophage SPO1. Cell.

63:417–424.

Grazziotin A.L., Koonin E. V, Kristensen D.M. 2017. Prokaryotic Virus Orthologous Groups

(pVOGs): a resource for comparative genomics and protein family annotation. Nucleic

Acids Res. 45:D491–D498.

Grose J.H., Casjens S.R. 2014. Understanding the enormous diversity of bacteriophages: The

tailed phages that infect the bacterial family *Enterobacteriaceae*. Virology. 468–

470:421–443.

Hulo C., Masson P., Le Mercier P., Toussaint A. 2015. A structured annotation frame for the

transposable phages: A new proposed family "Saltoviridae" within the *Caudovirales*.

Virology. 477:155–163.

Hyatt D., Chen G.L., LoCascio P.F., Land M.L., Larimer F.W., Hauser L.J. 2010. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics. 11.

Iranzo J., Krupovic M., Koonin E. V. 2016. The double-stranded DNA virosphere as a

modular hierarchical network of gene sharing. MBio. 7:e00978-16.

Jones D.T., Taylor W.R., Thornton J.M. 1992. The rapid generation of mutation data

matrices from protein sequences. Bioinformatics. 8:275–282.

Juhala R.J., Ford M.E., Duda R.L., Youlton A., Hatfull G.F., Hendrix R.W. 2000. Genomic

sequences of bacteriophages HK97 and HK022: pervasive genetic mosaicism in the

lambdoid bacteriophages. J. Mol. Biol. 299:27–51.

Karsch-Mizrachi I., Nakamura Y., Cochrane G. 2012. The International Nucleotide Sequence

Database Collaboration. Nucleic Acids Res. 40:D33–D37.

Kislyuk A.O., Haegeman B., Bergman N.H., Weitz J.S. 2011. Genomic fluidity: An

integrative view of gene diversity within microbial populations. BMC Genomics. 12:32.

Klumpp J., Lavigne R., Loessner M.J., Ackermann H.-W. 2010. The SPO1-related

bacteriophages. Arch. Virol. 155:1547–61.

Konstantinidis K.T., Tiedje J.M. 2005. Towards a genome-based taxonomy for prokaryotes.

J. Bacteriol. 187:6258–6264.

Kristensen D.M., Kannan L., Coleman M.K., Wolf Y.I., Sorokin A., Koonin E. V,

Mushegian A. 2010. A low-polynomial algorithm for assembling clusters of orthologous

groups from intergenomic symmetric best matches. Bioinformatics. 26:1481–7.

Krupovic M., Dutilh B.E., Adriaenssens E.M., Wittmann J., Vogensen F.K., Sullivan M.B.,

Rumnieks J., Prangishvili D., Lavigne R., Kropinski A.M., Klumpp J., Gillis A., Enault

F., Edwards R.A., Duffy S., Clokie M.R.C., Barylski J., Ackermann H.-W., Kuhn J.H.

2016. Taxonomy of prokaryotic viruses: update from the ICTV bacterial and archaeal

viruses subcommittee. Arch. Virol. 161:1095–1099.

Krupovic M., Prangishvili D., Hendrix R.W., Bamford D.H. 2011. Genomics of bacterial and

archaeal viruses: Dynamics within the prokaryotic virosphere. Microbiol. Mol. Biol.

Rev. 75:610–635.

Larkin M.A., Blackshields G., Brown N.P., Chenna R., McGettigan P.A., McWilliam H.,

Valentin F., Wallace I.M., Wilm A., Lopez R., Thompson J.D., Gibson T.J., Higgins

D.G. 2007. Clustal W and Clustal X version 2.0. Bioinformatics. 23:2947–2948.

Lavigne R., Darius P., Summer E.J., Seto D., Mahadevan P., Nilsson A.S., Ackermann H.W.,

Kropinski A.M. 2009. Classification of *Myoviridae* bacteriophages using protein

sequence similarity. BMC Microbiol. 9:224.

Lavigne R., Seto D., Mahadevan P., Ackermann H.-W., Kropinski A.M. 2008. Unifying

classical and molecular taxonomic classification: analysis of the *Podoviridae* using

BLASTP-based tools. Res. Microbiol. 159:406–414.

Lavigne R., Vandersteegen K. 2013. Group I introns in *Staphylococcus* bacteriophages.

Future Virol. 8:997–1005.

Lefort V., Desper R., Gascuel O. 2015. FastME 2.0: A comprehensive, accurate, and fast

distance-based phylogeny inference program. Mol. Biol. Evol. 32:2798–2800.

Letunic I., Bork P. 2007. Interactive Tree Of Life (iTOL): An online tool for phylogenetic

tree display and annotation. Bioinformatics. 23:127–128.

Liu K., Linder C.R., Warnow T. 2011. RAxML and FastTree: Comparing two methods for

large-scale maximum likelihood phylogeny estimation. PLoS One. 6:e27731.

Meier-Kolthoff J.J.P., Hahnke R.L., Petersen J., Scheuner C., Michael V., Fiebig A., Rohde

C., Rohde M., Fartmann B., Goodwin L.A., Chertkov O., Reddy T.B.K., Pati A.,

Ivanova N.N., Markowitz V.V., Kyrpides N.C.N., Woyke T., Göker M., Klenk H.-P.H.-

P.H., Pagani I., Liolios K., Jansson J., Chen I., Smirnova T., Nosrat B., Markowitz V.V.,

Kyrpides N.C.N., Lapage S., Sneath P., Lessel E., Skerman V., Seeliger H., Clark W.,

Blattner F., Plunkett G., Bloch C., Perna N., Burland V., Riley M., Vides J.C., Glasner

J., Rode C., Mayhew G., Gregor J., Davis N., Kirkpatrick H., Goeden M., Rose D., Mau

B., Shao Y., Wu D., Hugenholtz P., Mavromatis K., Pukall R., Dalin E., Ivanova N.N.,

Kunin V., Goodwin L.A., Wu M., Tindall B., Hooper S., Pati A., Lykidis A., Spring S.,

Anderson I., D P., Escherich T., Skerman V., McGowan V., Sneath P., Kauffmann F.,

Ørskov F., Ørskov I., Filannino P., Azzi L., Cavoski I., Vincentini O., Rizzello C.,

Gobbetti M., Cagno R. Di, Schumann P., Pukall R., Farnleitner A., Kreuzinger N.,

Kavka G., Grillenberger S., Rath J., Mach R., Tee T., Chowdhury A., Maranas C.,

Shanks J., Wen M., Bond-Watts B., Chang M., Rosano G., Ceccarelli E., Donovan C.,

Bramkamp M., Kuzminov A., Kang Z., Zhang C., Zhang J., Jin P., Zhang J., Du G.,

Chen J., Whitfield C., Roberts I., Cooper K., Mandrell R., Louie J., Korlach J., Clark T.,

Parker C., Huynh S., Chain P., Ahmed S., Carter M., Allocati N., Masulli M., Alexeyev

M., Ilio C. Di, Kaper J., Nataro J., Mobley H., Auch A., Jan M. Von, Klenk H.-P.H.-

P.H., Göker M., Meier-Kolthoff J.J.P., Auch A., Klenk H.-P.H.-P.H., Göker M., Meier-

Kolthoff J.J.P., Klenk H.-P.H.-P.H., Göker M., Göker M., Cleland D., Saunders E.,

Lapidus A., Nolan M., Lucas S., Hammon N., Deshpande S., Cheng J.-F., Tapia R., Han

C., Goodwin L.A., Pitluck S., Liolios K., Pagani I., Ivanova N.N., Mavromatis K., Pati

A., Chen A., Palaniappan K., Land M., Hauser L., Chang Y.Y.-J., Jeffries C., Detter J.,

Beck B., Woyke T., Bristow J., Eisen J., Markowitz V.V., Welch R., Scheutz F.,

Strockbine N., Koser S., Topley W., Wilson G., Huys G., Cnockaert M., Janda J.,

Swings J., Field D., Garrity G., Gray T., Morrison N., Selengut J., Sterk P., Tatusova T.,

Thomson N., Allen M., Angiuoli S., Ashburner M., Axelrod N., Baldauf S., Ballard S.,

Boore J., Cochrane G., Cole J., Dawyndt P., Vos P. De, Pamphilis C. de, Edwards R.,

Faruque N., Feldman R., Gilbert J., Gilna P., Glockner F., Goldstein P., Guralnick R.,

Haft D., Hancock D., Field D., Amaral-Zettler L., Cochrane G., Cole J., Dawyndt P.,

Garrity G., Gilbert J., Glöckner F., Hirschman L., Karsch-Mizrachi I., Klenk H.-P.H.-

P.H., Knight R., Kottmann R., Kyrpides N.C.N., Meyer F., Gil I.S., Sansone S.-A.,

Schriml L., Sterk P., Tatusova T., Ussery D., White O., Wooley J., Woese C., Kandler

O., Weelis M., Garrity G., Bell J., Lilburn T., Garrity G., Bell J., Lilburn T., Williams

K., Kelly D., Brenner D., Castellani A., Chalmers A., Ashburner M., Ball C., Blake J.,

Botstein D., Butler H., Cherry J., Davis A., Dolinski K., Dwight S., Eppig J., Harris M.,

Hill D., Issel-Tarver L., Kasarskis A., Lewis S., Matese J., Richardson J., Ringwald M.,

Rubin G., Sherlock G., Consortium G., Vaas L., Sikorski J., Michael V., Göker M.,

Klenk H.-P.H.-P.H., Vaas L., Sikorski J., Hofner B., Fiebig A., Buddruhs N., Klenk H.-

P.H.-P.H., Göker M., Chang Y.Y.-J., Feingold D., Boer H., Maaheimo H., Koivula A.,

Penttila M., Richard P., Xiao Z., Xu P., Göker M., Klenk H.-P.H.-P.H., Mavromatis K.,

Land M., Brettin T., Quest D., Copeland A., Clum A., Goodwin L.A., Woyke T.,

Lapidus A., Klenk H.-P.H.-P.H., Cottingham R., Kyrpides N.C.N., Markowitz V.V., I-

M A.C., Palaniappan K., Chu K., Szeto E., Grechkin Y., Ratner A., Jacob B., Huang J.,

Williams P., Huntemann M., Anderson I., Mavromatis K., Ivanova N.N., Kyrpides

N.C.N., Gemeinholzer B., Dröge G., Zetzsche H., Haszprunar G., Klenk H.-P.H.-P.H.,

Güntsch A., Berendsohn W., Wägele J., Zerbino D., Birney E., Gordon D., Abajian C.,

Green P., Hyatt D., Chen G., LoCascio P., Land M., Larimer F., Hauser L., Mavromatis

K., Ivanova N.N., Chen I., Szeto E., Markowitz V.V., Kyrpides N.C.N., Finn D.,

Clements J., Eddy S., Lowe T., Eddy S., Nawrocki E., Kolbe D., Eddy S., Markowitz

V.V., Ivanova N.N., Chen I., Chu K., Kyrpides N.C.N., Bland C., Ramsey T., Sabree F.,

Lowe M., Brown K., Kyrpides N.C.N., Hugenholtz P., Edgar R., Wayne L., Brenner D.,

Colwell R., Grimont P., Kandler O., Krichevsky M., Moore L., Moore W., Murray R.,

Stackebrandt E., Starr M., Truper H., Tindall B., Rosselló-Móra R., Busse H., Ludwig

W., Kämpfer P., Kaas R., Friis C., Ussery D., Aarestrup F., Clermont O., Bonacorsi S.,

Bingen E., Clermont O., Gordon D., Brisse S., Walk S., Denamur E., Clermont O.,

Christenson J., Denamur E., Gordon D., Sahl J., Morris C., Rasko D., Patil K., McHardy

A., Thorne J., Kishino H., Meier-Kolthoff J.J.P., Auch A., Klenk H.-P.H.-P.H., Göker

M., Letunic I., Bork P., Desper R., Gascuel O., Lukjancenko O., Wassenaar T., Ussery

D., Touchon M., Hoede C., Tenaillon O., Barbe V., Baeriswyl S., Bidet P., Bingen E.,

Bonacorsi S., Bouchier C., Bouvet O., Calteau A., Chiapello H., Clermont O., Cruveiller

S., Danchin A., Diard M., Dossat C., Karoui M. El, Frapy E., Garry L., Ghigo J., Gilles

A., Johnson J., Bouguenec C. Le, Lescat M., Mangenot S., Martinez-Jehanne V., Matic

I., Nassif X., Oztas S., Zuo G., Xu Z., Hao B., Abt B., Han C., Scheuner C., Lu M.,

Lapidus A., Nolan M., Lucas S., Hammon N., Deshpande S., Cheng J.-F., Tapia R.,

Goodwin L.A., Pitluck S., Mavromatis K., Mikhailova N., Huntemann M., Pati A., Chen

A., Palaniappan K., Land M., Hauser L., Brambilla E.-M., Rohde M., Spring S., Gronow

S., Göker M., Woyke T., Bristow J., Eisen J., Markowitz V.V., Abt B., Göker M.,

Scheuner C., Han C., Lu M., Misra M., Lapidus A., Nolan M., Lucas S., Hammon N.,

Deshpande S., Chang J.-F., Tapia R., Goodwin L.A., Pitluck S., Liolios K., Pagani I.,

Ivanova N.N., Mavromatis K., Mikhailova N., Huntemann M., Pati A., Chen A.,

Palaniappan K., Land M., Hauser L., Brambilla E.-M., Rohde M., Spring S., Gronow S.,

Anderson I., Scheuner C., Göker M., Mavromatis K., Hooper S., Porat I., Klenk H.-

P.H.-P.H., Ivanova N.N., Kyrpides N.C.N., Frank O., Pradella S., Rohde M., Scheuner

C., Klenk H.-P.H.-P.H., Göker M., Petersen J., Göker M., Scheuner C., Klenk H.-P.H.-

P.H., Stielow J., Menzel W., Spring S., Scheuner C., Lapidus A., Lucas S., Rio T. Del,

Tice H., Copeland A., Cheng J.-F., Chen F., Nolan M., Saunders E., Pitluck S., Liolios

K., Ivanova N.N., Mavromatis K., Lykidis A., Pati A., Chen A., Palaniappan K., Land

M., Hauser L., Chang Y.Y.-J., Jeffries C., Goodwin L.A., Detter J., Brettin T., Rohde

M., Göker M., Woyke T., Bristow J., Stackebrandt E., Scheuner C., Göker M.,

Schumann P., Verbarg S., Göker M., Scheuner S., Schumann P., Stackebrandt E.,

Altschul S., Madden T., Schaffer A., Zhang J., Zhang Z., Miller W., Lipman D., Li L.,

Stoeckert C., Roos D., Edgar R., Thompson J., Thierry J.-C., Poch O., Castresana J.,

Meusemann K., Reumont B. von, Simon S., Roeding F., Strauss S., Kuck P.,

Ebersberger I., Walzl M., Pass G., Breuers S., Achter V., Haeseler A. von, Burmester T.,

Hadrys H., Wagele J., Misof B., Felsenstein J., Fitch W., Goloboff P., Stamatakis A.,

Pattengale N., Alipour M., Bininda-Emonds O., Moret B., Stamatakis A., Swofford D.,

Klenk H.-P.H.-P.H., Göker M., Enright A., Dongen S. van, Ouzounis C., Albuquerque

L., Rainey F., Nobre M.F., Costa M. da, Fricke W., McDermott P., Mammel M., Zhao

S., Johnson T., Rasko D., Fedorka-Cray P., Pedroso A., Whichard J., Leclerc J., White

D., Cebula T., Ravel J., Brinkkötter A., Klöss H., Alpert C., Lengeler J., Göker M.,

Garcáa-Blázquez G., Voglmayr H., Telleráa M., Martán M., Staley J., Krieg N.,

Tindall B., Kampfer P., Euzeby J., Oren A., Lan R., Reeves P. 2014. Complete genome

sequence of DSM 30083T, the type strain (U5/41T) of *Escherichia coli*, and a proposal

for delineating subspecies in microbial taxonomy. Stand. Genomic Sci. 9:2.

Meier-Kolthoff J.P., Göker M. 2017. VICTOR: genome-based phylogeny and classification

of prokaryotic viruses. Bioinformatics. 33:3396–3404.

Mizuno C.M., Rodriguez-Valera F., Kimes N.E., Ghai R. 2013. Expanding the marine

virosphere using metagenomics. PLoS Genet. 9:e1003987.

Nishimura Y., Watai H., Honda T., Mihara T., Omae K., Roux S., Blanc-Mathieu R.,

Yamamoto K., Hingamp P., Sako Y., Sullivan M.B., Goto S., Ogata H., Yoshida T.

2017. Environmental viral genomes shed new light on virus-host interactions in the

ocean. mSphere. 2:e00359-16.

O'Leary N.A., Wright M.W., Brister J.R., Ciufo S., Haddad D., McVeigh R., Rajput B.,

Robbertse B., Smith-White B., Ako-Adjei D., Astashyn A., Badretdin A., Bao Y.,

Blinkova O., Brover V., Chetvernin V., Choi J., Cox E., Ermolaeva O., Farrell C.M.,

Goldfarb T., Gupta T., Haft D., Hatcher E., Hlavina W., Joardar V.S., Kodali V.K., Li

W., Maglott D., Masterson P., McGarvey K.M., Murphy M.R., O'Neill K., Pujar S.,

Rangwala S.H., Rausch D., Riddick L.D., Schoch C., Shkeda A., Storz S.S., Sun H.,

Thibaud-Nissen F., Tolstoy I., Tully R.E., Vatsan A.R., Wallin C., Webb D., Wu W.,

Landrum M.J., Kimchi A., Tatusova T., DiCuccio M., Kitts P., Murphy T.D., Pruitt

K.D. 2016. Reference sequence (RefSeq) database at NCBI: current status, taxonomic

expansion, and functional annotation. Nucleic Acids Res. 44:D733–D745.

Paez-Espino D., Eloe-Fadrosh E.A., Pavlopoulos G.A., Thomas A.D., Huntemann M.,

Mikhailova N., Rubin E., Ivanova N.N., Kyrpides N.C. 2016. Uncovering Earth's

virome. Nature. 536:425–430.

Page A.J., Cummins C.A., Hunt M., Wong V.K., Reuter S., Holden M.T.G., Fookes M.,

Falush D., Keane J.A., Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome

analysis. Bioinformatics. 31:3691–3693.

Price M.N., Dehal P.S., Arkin A.P. 2010. FastTree 2 - Approximately maximum-likelihood

trees for large alignments. PLoS One. 5:e9490.

Rambaud. 2007. FigTree. Available from http://tree.bio.ed.ac.uk/software/figtree/.

Rohwer F., Edwards R. 2002. The Phage Proteomic Tree: a genome-based taxonomy for

phage. J. Bacteriol. 184:4529–4535.

Roux S., Brum J.R., Dutilh B.E., Sunagawa S., Duhaime M.B., Loy A., Poulos B.T.,

Solonenko N., Lara E., Poulain J., Pesant S., Kandels-Lewis S., Dimier C., Picheral M.,

Searson S., Cruaud C., Alberti A., Duarte C.M., Gasol J.M., Vaqué D., Bork P., Acinas

S.G., Wincker P., Sullivan M.B. 2016. Ecogenomics and potential biogeochemical

impacts of globally abundant ocean viruses. Nature. 537:689–693.

Schuch R., Fischetti V.A. 2009. The secret life of the anthrax agent Bacillus anthracis:

Bacteriophage-mediated ecological adaptations. PLoS One. 4:e6532.

Seemann T. 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics. 30:2068–

2069.

Sievers F., Wilm A., Dineen D., Gibson T.J., Karplus K., Li W., Lopez R., McWilliam H.,

Remmert M., Söding J., Thompson J.D., Higgins D.G. 2011. Fast, scalable generation of

high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst.

Biol. 7:539.

Simmonds P., Adams M.J., Benkő M., Breitbart M., Brister J.R., Carstens E.B., Davison

A.J., Delwart E., Gorbalenya A.E., Harrach B., Hull R., King A.M., Koonin E. V,

Krupovic M., Kuhn J.H., Lefkowitz E.J., Nibert M.L., Orton R., Roossinck M.J.,

Sabanadzovic S., Sullivan M.B., Suttle C.A., Tesh R.B., van der Vlugt R.A., Varsani A.,

Zerbini F.M. 2017. Consensus statement: Virus taxonomy in the age of metagenomics.

Nat. Rev. Microbiol. 15:161–168.

Söding J., Biegert A., Lupas A.N. 2005. The HHpred interactive server for protein homology

detection and structure prediction. Nucleic Acids Res. 33:W244-2488.

Uchiyama J., Rashel M., Maeda Y., Takemura I., Sugihara S., Akechi K., Muraoka A.,

Wakiguchi H., Matsuzaki S. 2008a. Isolation and characterization of a novel

Enterococcus faecalis bacteriophage phiEF24C as a therapeutic candidate. FEMS

Microbiol. Lett. 278:200–206.

Uchiyama J., Rashel M., Takemura I., Wakiguchi H., Matsuzaki S. 2008b. In silico and in

vivo evaluation of bacteriophage phiEF24C, a candidate for treatment of *Enterococcus*

faecalis infections. Appl. Environ. Microbiol. 74:4149–4163.

Whelan S., Goldman N. 2001. A general empirical model of protein evolution derived from

multiple protein families using a maximum-likelihood approach. Mol. Biol. Evol.

18:691–699.

Yuan Y., Peng Q., Wu D., Kou Z., Wu Y., Liu P., Gao M. 2015. Effects of actin-like proteins

encoded by two *Bacillus pumilus* phages on unstable lysogeny, revealed by genomic

analysis. Appl. Environ. Microbiol. 81:339–350.

FUNDING

This work was supported by the National Science Centre, Poland (grant number

2016/23/D/NZ2/00435) to JB; the Netherlands Organization for Scientific Research (NWO)

(Vidi grant number 864.14.004) to MBPS and BED; the US National Science Foundation

(grant numbers DUE-132809 and MCB-1330800) to RAE; the University of Helsinki

funding for Instruct research infrastructure (Virus and Macromolecular Complex Production,

ICVIR) to HMO; the Chargé de Recherches fellowship from the National Fund for Scientific

Research, FNRS, Belgium to AG; the EUed Horizon 2020 Framework Programme for

Research and Innovation, 'Virus-X' (grant number 685778) to FE; the Gordon and Betty

Moore Foundation Investigator Award (grant number GBMF#3790) to MBS; the Battelle

Memorial Institute's prime contract with the US National Institute of Allergy and Infectious

Diseases (NIAID) (Contract number HHSN272200700016I) to JHK. The content of this

publication does not necessarily reflect the views or policies of the US Department of Health

and Human Services or of the institutions and companies affiliated with the authors.

ACKNOWLDEGMENTS

We thank Laura Bollinger, Integrated Research Facility at Fort Detrick, for technical

writing services. E.M.A. would like to thank P.H. Nel for assistance with R scripting.