Genomic, proteomic, and phylogenetic analysis of spounaviruses

indicates paraphyly of the order *Caudovirales*

Short title: Taxonomy of SPO1-like phages

Jakub Barylski[1], François Enault[2], Bas E. Dutilh[3,4], Margo B.P. Schuller[3], Robert A. Edwards[5],

Annika Gillis[6], Jochen Klumpp[7], Petar Knezevic[8], Mart Krupovic[9], Jens H. Kuhn[10], Rob

Lavigne[11], Hanna M. Oksanen[12], Matthew B. Sullivan[13], Johannes Wittmann[14], Igor Tolstoy[15],

J. Rodney Brister[15], Andrew M. Kropinski[16], Evelien M. Adriaenssens[17*]

[1]Adam Mickiewicz University, Institute of Experimental Biology, Department of Molecular

Virology, Poznań, Poland, [2]Université Clermont Auvergne, CNRS, LMGE, F-63000, Clermont-

Ferrand, France, [3]Theoretical Biology and Bioinformatics, Department of Biology, Utrecht

University, Utrecht, The Netherlands, [4]Theoretical Biology and Bioinformatics, Department

of Biology, Utrecht University, Utrecht, The Netherlands, [5]Departments of Biology and

Computer Science, San Diego State University, San Diego, CA, USA, [6]Laboratory of Food and

Environmental Microbiology, Université Catholique de Louvain, Louvain-la-Neuve, Belgium,

[7]Institute of Food, Nutrition and Health, ETH Zurich, Switzerland, [8]Department of Biology

and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad, Serbia, [9]Unit of Molecular

Biology of the Gene in Extremophiles, Department of Microbiology, Institut Pasteur, Paris,

France, [10]Integrated Research Facility at Fort Detrick, National Institute of Allergy and

Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, USA, [11]Laboratory

of Gene Technology, KU Leuven, Belgium, [12]Department of Biosciences, University of

Helsinki, Helsinki, Finland; and Institute of Biotechnology, University of Helsinki, Helsinki,

Finland, [13]Departments of Microbiology and Civil, Environmental and Geodetic Engineering,

The Ohio State University, Columbus, OH, USA, [14]Leibniz-Institut DSMZ—Deutsche

Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany,

[15]National Center for Biotechnology Information, National Library of Medicine, National

Institutes of Health, Bethesda, MD, USA, [16]Departments of Food Science, Molecular and

Cellular Biology; and Pathobiology, University of Guelph, Guelph, Ontario, Canada,

[17]Institute of Integrative Biology, University of Liverpool, Biosciences Building, Crown Street,

Liverpool L69 7ZB, United Kingdom; orcid.org/0000-0003-4826-5406

[*]evelien.adriaenssens@gmail.com (EMA)

Abstract

Since the mid-20th century, prokaryotic double-stranded DNA viruses producing tailed
particles ("tailed phages") were grouped according to virion tail morphology. In the early
1980s, these viruses were classified into the families *Myoviridae*, *Siphoviridae*, and
Podoviridae, later included in the order *Caudovirales*. However, recent massive sequencing
of prokaryotic virus genomes revealed that caudovirads are extremely diverse. The official
taxonomic framework does not adequately reflect caudovirad evolutionary relationships.
Here, we reevaluate the classification of caudovirads using a particularly challenging group
of viruses with large dsDNA genomes: SPO1-like viruses associated with the myovirid
subfamily *Spounavirinae*. Our extensive genomic, proteomic, and phylogenetic analyses
reveal that some of the currently established caudovirad taxa, especially at the family and
subfamily rank, can no longer be supported. Spounavirins alone need to be elevated to
family rank and divided into at least five major clades, a first step in an impending massive
reorganization of caudovirad taxonomy.

Introduction

Prokaryotic virus taxonomy is the formal responsibility of the Bacterial and Archaeal Viruses

Subcommittee of the International Committee on Taxonomy of Viruses (ICTV). In recent

years, the Subcommittee has focused on classifying newly described double-stranded DNA

viruses producing tailed particles ("tailed phages") into species and genera included in

existing families in the order *Caudovirales* [1–5]. At the species rank, a similarity threshold of

95% nucleotide sequence identity is used for classification, i.e. viruses are classified into the

same species if they shared ≥95% identity over the entire length of their genomes.

Examination of wild populations of caudovirads infecting cyanobacteria demonstrated that

such a 95% threshold robustly captures formally delineable species using population genetic

metrics [6]. Extrapolation of such thresholds to viruses of the global surface oceans has

resulted in population-scale ecological understanding of thousands of new virus "species"

[7], which, however, do not have official status. At the genus rank, cohesive groups have

been defined for viruses sharing a significant genome similarity(≥60% nucleotide identity),

gene synteny, and a core gene set. This framework has helped to relatively rapidly establish

official low-rank taxonomic positions for newly isolated and sequenced viruses [1,2,5,8,9].

The currently available ranks in virus taxonomy (in ascending order *species*, *genus*,

subfamily, *family*, and *order*) limit the description of the full diversity of prokaryotic viruses.

This limitation is particularly acute in the case of the order *Caudovirales*, which represents

the most abundant and diverse group of viruses in the environment [10–12]. Indeed, the

diversity of caudovirads greatly surpasses that of other bacterial, archaeal and eukaryotic

double-stranded (ds)DNA viruses. A recent analysis of the dsDNA virosphere using a

bipartite network approach, whereby viral genomes are connected via shared gene families,

demonstrated that the global network of dsDNA viruses consists of at least 19 modules, 11

of which correspond to caudovirads [13]. The eight remaining modules encompass one or

more families of eukaryotic or archaeal viruses. Consequently, each of the caudovirad

modules could be considered to represent a separate family. Despite this remarkable

sequence diversity, all caudovirads are currently classified into three families—*Myoviridae,*

Podoviridae, and *Siphoviridae*. These families were historically established on the

morphological, not genomic, features of their members, forming an artificial classification

ceiling. In this study, the Subcommittee explored the diversity of the order *Caudovirales*

based on the example of a large group of Bacillus phage SPO1-related viruses (*Myoviridae*:

Spounavirinae), which forms a discrete caudovirad module [13,14].

The type virus of this group, Bacillus phage SPO1, was isolated in 1964 from soil in

Osaka, Japan, using *Bacillus subtilis* as a host [15]. Bacillus phage SPO1 has been extensively

studied ever since. Morphologically, the particle possesses an icosahedral head 87 nm in

diameter and a 140-nm long (18.6-nm wide) contractile tail of "stacked-disk" appearance

that is terminated by a complex baseplate structure [16,17]. The packaged genome is a

145.7-kb long, terminally redundant (13.1-kb) DNA molecule with thymine completely

replaced by 5-hydroxymethyluracil (HMU) [18–20]. This genome encodes at least 204

proteins and five tRNAs [21].

In 1995, Bacillus phage SPO1 was assigned to the species *Bacillus phage SP8*

(renamed *Bacillus phage SPO1* in 1996 and *Bacillus virus SPO1* in 2015), which in 1996 was

included into a monospecific myovirid genus currently called *Spo1virus* [1,22,23]. The

subfamily "*Spounavirinae*" was proposed in 2009 by Lavigne et al. to harbor Bacillus phage

SPO1, Staphylococcus phage Twort, Staphylococcus phage K, Staphylococcus phage G1,

Listeria phage P100, and Listeria phage A511 [4]. This subfamily became official in 2012 [24].

The unifying characteristics of members of this subfamily as described by Klumpp et al. are

that "(a) the[ir] host organisms are bacteria of the phylum Firmicutes; (b) [they] are strictly

virulent myovirids; (c) all… feature common morphological properties; (d) [their genomes]

consist of a terminally redundant, non-permuted dsDNA molecule of 127–157 kb in size; and

(e) [they] share considerable amino acid homology" [20]. The inclusion requirement of a

strictly lytic lifestyle became controversial when it was observed that a few related viruses

(Bacillus phages Bcp1, Bp8p-T, and Bp8p-C) can persist in host cultures without causing the

immediate lysis [25,26]. It remains unknown whether this persistence is due to virion

entrapment inside bacillus spores or some other kind of semi-stable virus-host relationship.

109   By 2015, the subfamily *Spounavirinae* had been expanded to include five genera

(*Kayvirus*, *P100virus*, *Silviavirus*, *Spo1virus*, and *Twortvirus*) and three unassigned or

"floating" species (*Enterococcus virus phiEC24C*, *Lactobacillus virus Lb338-1*, and

Lactobacillus virus LP65). It was already clear that *Bacillus virus SPO1* represented one of

two major lineages within the subfamily. Consequently, spounavirins were divided in the

Bacillus phage SPO1-like viruses, which have modified DNA (HMU) and which produce

particles possessing generally shorter tails; and, the Staphylococcus phage Twort-like

viruses, which feature non-modified DNA and produce particles with longer tails. Both

Bacillus phage SPO1-like and Staphylococcus phage Twort-like virus particles feature double-

ringed baseplates and visible capsomeres as a morphologic hallmark [20,27]. A third group,

represented by Bacillus phages sharing limited similarity with Bacillus phage SPO1 but

related to phage Bastille had already been proposed at that time but not yet officially

recognized [28]. In 2016, additional genera of Bacillus phage SPO1-like viruses (henceforth

"spouna-like viruses") were established, but not all of these were immediately included in

the subfamily *Spounavirinae* [1,2].

In this study, we reevaluated the classification of spounavirins and spouna-like

viruses. To this end, a well-defined set of 93 viruses was analyzed using complementary DNA

and protein sequence analysis tools and phylogenetic methods. Our results indicate that the

subfamily *Spounavirinae* fails to adequately reflect the diversity of its current members, and

we therefore outline a better fitting classification scheme.

Results

General overview

To determine the phylogenetic relationship between 93 known and alleged spounavirins,

we employed genomic, proteomic and marker gene-based comparative strategies.

Regardless of the adopted phylogenetic approach applied, five separate, clear-cut clusters

were identified. We believe that they clearly have a common origin and ought to come

together under one caudoviral umbrella taxon. We propose to name this taxon

"*Herelleviridae*," in honor of the 100th anniversary of the discovery of prokaryotic viruses by

Félix d'Hérelle (Table 1, Figs 1-3, S1 Table). The first cluster (here suggested to retain the

name *Spounavirinae*) groups *Bacillus*-infecting viruses that are similar to Bacillus phage

SPO1. The second cluster ("*Bastillevirinae*," named after the type species *Bacillus virus*

Bastille [28]) includes *Bacillus*-infecting viruses that have only limited similarity to Bacillus

phage SPO1 and resemble Bacillus phage Bastille instead. The third cluster ("*Brockvirinae*,"

named in honor of Thomas D. Brock [1926–], an American microbiologist and educator

known for his discovery of hyperthermophiles, who worked on Streptococcus phages early

in his career) comprises currently unclassified viruses of enterococci that are similar to

Enterococcus phage φEF24C. The fourth cluster ("*Twortvirinae*," named in honor of

Frederick William Twort (1877–1950), the English bacteriologist who discovered prokaryotic

viruses in 1915) gathers staphylococci-infected viruses that are similar to Staphylococcus

phage Twort, whereas the remaining cluster ("*Jasinskavirinae*," named in honor of

Stanislawa Jasińska-Lewandowska (1921–1998), Polish scientist who was one of the first to

study *Listeria* and their viruses) consists of viruses infecting *Listeria* that are similar to

Listeria phage P100, the type isolate of the *P100virus* genus. The classification in five

clusters left three viruses unassigned at this rank: Lactobacillus phage Lb338, Lactobacillus

phage LP65, and Brochothrix phage A9.

These robust clusters can be further subdivided into smaller clades that correspond

well with the currently accepted genera. The evidence supporting this suggested taxonomic

re-classification is presented in the following sections.

Table 1. Suggested new classification of the 93 spounavirins and spouna-like viruses in the new caudoviral family "*Herelleviridae*".

Order	Family	Subfamily	Genus	Species[a]	Viruses
Caudovirales	*"Herelleviridae"*	**"***Bastillevirinae***"**	*Agatevirus*	*Bacillus virus Agate, Bacillus virus Bobb, Bacillus virus Bp8pC*	Bp8p-T
			B4virus	*Bacillus virus AvesoBmore, Bacillus virus B4, Bacillus virus Bigbertha, Bacillus virus Riley, Bacillus virus Spock, Bacillus virus Troll*	B5S
			Bastillevirus	*Bacillus virus Bastille, Bacillus virus CAM003, "Bacillus virus Evoli", "Bacillus virus HoodyT"*	
			Bc431virus	*Bacillus virus Bc431, Bacillus virus Bcp1, Bacillus virus BCP82, Bacillus virus JBP901*	
			Nit1virus	*Bacillus virus Grass, Bacillus virus NIT1, Bacillus virus SPG24*	
			Tsarbombavirus	*Bacillus virus BCP78, Bacillus virus*	BCU4

				TsarBomba	
			Wphvirus	*Bacillus virus BPS13, Bacillus virus Hakuna, Bacillus virus Megatron, Bacillus virus WPh, "Bacillus virus BPS10C"*	Eyuki
		"*Brockvirinae*"	"*Kochikohdavirus*"	"*Enterococcus virus ECP3*", "*Enterococcus virus EF24C*", "*Enterococcus virus EFLK1*"	phiEFC24C-P2
			Unassigned	"*Enterococccus virus EFDG1*"	
		"*Jasinskavirinae*"	P100virus	*Listeria virus A511, Listeria virus P100*	List-36, LMSP-25, AvB_LmoM_AG20, LP-125, LP-064, LP-083-2, LP-124, LP-125, LP-048, LMTA-34, LMTA-94, LMTA-148, LMTA-57, WIL-1
		Spounavirinae	*Cp51virus*	*Bacillus virus CP51, Bacillus virus JL, Bacillus virus Shanette*	
			Spo1virus	*Bacillus virus Camphawk, Bacillus*	

				virus SPO1	
			Unassigned	*"Bacillus virus Mater", "Bacillus virus Moonbeam", "Bacillus virus SlOphi"*	
		"*Twortvirinae*"	*Kayvirus*	*Staphylococcus virus G1, Staphylococcus virus G15, Staphylococcus virus JD7, Staphylococcus virus K, Staphylococcus virus MCE2014, Staphylococcus virus P108, Staphylococcus virus Rodi, Staphylococcus virus S253, Staphylococcus virus S25-4, Staphylococcus virus SA12, "Staphylococcus virus Sb1"*	676Z, A3R, A5W, Fi200W, IME-SA1, IME-SA118, IME-SA119, IME-SA2, ISP, MSA6, P4W, SA5, Staph1N, Team1
			Silviavirus	*Staphylococcus virus Remus, Staphylococcus virus SA11*	Romulus
			Sep1virus	*Staphylococcus virus IPLAC1C,*	

				Staphylococcus virus SEP1	
			Twortvirus	*Staphylococcus virus Twort*	
		Unassigned	Unassigned	*"Lactobacillus virus Lb338",*	
				"Lactobacillus virus LP65",	
				"Brochothrix virus A9"	

[a] The species listed here are representing the 93 genome dataset on which all analyses have been performed. Species ratified in 2017 and later

have not been included here.

Genome-based analyses

BLASTn analysis revealed that the genomes of several viruses were similar enough to

consider them strains of the same species (they shared >95% nucleotide identity, S1 Fig).

The *Staphylococcus* viruses fell into four distinct, yet closely related groups corresponding to

the established genera *Twortvirus, Sep1virus, Silviavirus*, and *Kayvirus* (S1 Fig). With the

exception of Enterococcus phage EFDG1, all *Enterococcus* viruses clustered as a clade

representing a new genus (here suggested to be named "*Kochikohdavirus*" after the place of

origin of the type virus of the clade, Enterococcus phage φEF24C; [29,30]). The *Bacillus*

viruses clustered into the established genera *Spo1virus, Cp51virus, Bastillevirus, Agatevirus*,

B4virus, Bc431virus, Nit1virus, Tsarbombavirus, and *Wphvirus*, with three species remaining

unassigned at the genus rank (Table 1). These results were also confirmed with VICTOR, a

genome-BLAST distance phylogeny (GBDP) method (S2 Fig) and the Dice score (S3 Fig) [31],

a tBLASTx-based measure that compares whole genome sequences at the amino acid level.

Fig 1: Genome-based clustering trees of 93 spounavirin and spouna-like viruses.
Clustering was performed using nucleotide similarities (BLASTn, A) or translated nucleotide
similarities (tBLASTx, B). Genomes were compared in a pairwise fashion using Gegenees,
transformed into a distance matrix, clustered using R and visualized as trees using Itol. The
trees were rooted at Brochotrix phage A9. Genera are delineated with colored squares and
suggested subfamilies with colored circles.

The patterns coalesced at a higher taxonomic level when the genomes were

analyzed using tBLASTx (S4 Fig). The *Enterococcus* viruses clustered into a single group

sharing 41% genome identity, whereas the *Bacillus* viruses fell into two major groups, a

group combining the genera *Spo1virus* and *Cp51virus*, and the remainder. All *Staphylococcus*

viruses clustered above ≈36% genome identity, whereas *Listeria* viruses grouped with more

than 79% genome identity. Overall, all these genomes were related at the level of at least

15% genome identity. *Lactobacillus* and *Brochothrix* viruses remained genomic orphans,

peripherally related to the remainder of the viruses in this assemblage.

Predicted proteome-based analyses

The virus proteomic tree shows four robust groupings mainly determined by the hosts that

the viruses infect, corresponding largely with the suggested subfamilies (Fig 2). Viruses that

infect *Bacillus* fell into two groups as described before, represented by the revised

Spounavirinae subfamily and the suggested new subfamily "*Bastillevirinae*." Similarly, the

Listeria and *Staphylococcus* viruses formed their own clusters, "*Jasinskavirinae*" and

"*Twortvirinae*", respectively. This clustering suggests that the major *Bacillus, Listeria,* and

Staphylococcus virus groups are represented, but that further representatives are required

from the under-sampled groups. The suggested "*Brockvirinae*" subfamily is under-sampled,

and the grouping observed in the tree is not as well-supported as the other clusters.

Fig 2: Predicted proteome-based clustering trees of 93 spounavirin and spouna-like viruses. Clustering was performed using the Phage Proteomics Tree approach (A) and proteomic distance (B). Distances were calculated pairwise between all sets of predicted proteomes, clustered with R and visualized using Itol. The trees were rooted at Brochotrix phage A9. Genera are delineated with colored squares and suggested subfamilies with colored circles.

Among 1,296 singleton proteins and 2,070 protein clusters defined using the orthologous protein clusters (OPC) approach, we identified 12 clusters common for all viruses (Table 2, S2 Table). Classification of the viral proteins using prokaryotic virus orthologous groups (pVOGs) showed that 38 pVOGs were shared between all 93 virus genomes (Table 2, S3 Table). This finding was in stark contrast with the results from core

genome analysis using Roary, which revealed only one core gene (the tail tube protein

gene). Upon closer inspection of the gene annotations, we found that these analyses might

have been confounded by the presence of introns and inteins in many of the core genes

(S5–S6 Figs). Indeed, many genes of spounavirins and related viruses are invaded by mobile

introns or inteins [32,33]. These gaps in coding sequences challenge gene prediction tools

and introduce additional bias in similarity-based cluster algorithms.

Table 2: Core genes with putative annotated functions identified in all 93 spounavirin and

spouna-like virus genomes.

Putative function of the core gene identified[a]	pVOG[b] / OPC[b] ID	Identification method
DnaB-like helicase	VOG0025, OPC6121	OPC, pVOG
Baseplate J-like protein	VOG4691, VOG4644, OPC6132	OPC, pVOG
Tail sheath protein	VOG0067, OPC6142	OPC, pVOG
Terminase large subunit	VOG0051, OPC6160	pVOG
Major capsid protein	VOG0061, OPC6148	OPC, pVOG
Prohead protease	VOG4568, OPC6150	pVOG
Portal protein	VOG4556, OPC6151	OPC, pVOG
DNA primase	VOG4551	pVOG
DNA polymerase I	VOG0668, OPC6097	OPC, pVOG
RNA polymerase	VOG0118	pVOG
Recombination exonuclease	VOG4575	pVOG

Recombination endonuclease	VOG0083	pVOG
Tail tape measure protein	VOG0069	pVOG
Tail tube protein	VOG0068, OPC6141	OPC, pVOG, Roary

[a] The full lists of orthologous proteins and pVOGs are available in S2 Table and S3 Table,

respectively.

[b] pVOG, prokaryotic virus orthologous group; OPC, orthologous protein clusters.

The pairwise comparison of the predicted proteome content of the viruses revealed

a very low overall relatedness at the protein level (S7 Fig S7). The majority of viruses shared

less than 10% of their proteins. However, at the suggested new subfamily rank, we observed

obvious virus groups sharing their proteomes. The *Enterococcus* viruses ("*Brockvirinae*")

shared over 35% of their protein content. The members of the *Bacillus* virus genera

Spo1virus and *Cp51virus* of the subfamily *Spounavirinae* (sensu stricto) had approximately

20% of their proteins in common, whereas the *Bacillus* virus genera *Bastillevirus*, *B4virus*,

Bc431virus, *Agatevirus*, *Nit1virus*, *Tsarbombavirus*, and *Wphvirus* ("*Bastillevirinae*") and the

Staphylococcus virus genera *Kayvirus*, *Silviavirus*, and *Twortvirus* ("*Twortvirinae*") shared

over 25% and over 30% of their predicted proteomes, respectively.

Genomic fluidity is a measure of the dissimilarity of genomes evaluated at the gene

level [34]. Accordingly, the genomic fluidity results followed those obtained using proteome

content analysis (S8 Fig). Despite a high genomic fluidity for most of these viruses, the newly

suggested subfamilies and genera were all supported.

The topology of the dendrogram obtained using the average amino acid identity

(AAI) approach also supported the suggested new taxonomic scheme (S8 Fig). The AAI was

greater than 35% within each subfamily and greater than 67% within each genus. The AAI of

all viruses analyzed in this study was not lower than 22%. The members of the genus

Wphvirus had the lowest AAI (76%) and the lowest AAI for a pair of proteomes (67%

between Bacillus phage W.Ph. and Bacillus phage Eyuki) but surprisingly they had a mid-

range genomic fluidity (0.15), suggesting that the protein sequences of wphviruses might

have evolved rapidly.

249   The pangenome of the spounavirins and spouna-like viruses as calculated using

250   Roary [35], consisting of 4,182 genes, was further analyzed by clustering the genomes based

251   on the presence or absence of the accessory genes (S9 Fig). The obtained tree supported the

252   current division of the viruses into approved genera and the suggested new subfamilies.

253   Many virus genomes are thought to be highly modular, with recombination and

horizontal gene transfer potentially resulting in "mosaic genomes" [36,37]. By clustering the

spounavirin and spouna-like virus genomes based solely on the gene order of their

genomes, we investigated whether the gene synteny was preserved (S10 Fig). The results

revealed that genomic rearrangements leave a measurable evolutionary signal in all

lineages, since the genomic architecture analysis clustered all viruses according the

suggested taxa with the potential exception of Bacillus phage Moonbeam [38]. However, we

did not observe the high modularity that may be expected with rampant mosaicism. The

lack of rampant mosaicism supports the recent findings by Bolduc et al. that at most about

10% of reference virus genomes have a high degree of mosaicism [14]. Thus, while the gene

order in viruses belonging to the newly suggested family "*Herelleviridae*" is not necessarily

strictly conserved, we observed a clear evolutionary pattern that is consistent with the

sequence-based approaches tested in this study.

Single protein phylogenies

The phylogenetic trees based on comparisons of the major capsid, tail sheath, and DnaB-like

helicase proteins are presented in Fig 3. All nine phylograms based on conserved proteins

(OPC) are depicted in the trees in S11 Fig. For nearly all single marker trees, the topologies

supported the suggested taxonomic scheme. Generally, each taxon is represented as a

separate branch on the dendrogram. Notable exceptions could be found in two trees based

on hypothetical proteins (OPCs 10357 and 10386). The first (10357) places revised the

subfamily *Spounavirinae* as a subclade of "*Bastillevirinae*" and the second (10386) shuffled

silviaviruses and kayviruses. This result may indicate that some degree of horizontal gene

transfer occurs between groups, which share common hosts.

Fig 3: Single gene phylogenies of the major capsid protein (MCP, A), tail sheath protein
(TSP, B) and helicase (C) amino acid sequences of 93 spounavirin and spouna-like viruses.
Amino acid sequences were aligned with Clustal Omega and trees were generated using

281 FastTree maximum likelihood with Shimidaira-Hasegawa tests. The scale bar represents the
282 number of substitutions per site. The trees were rooted at Brochotrix phage A9. Genera are
283 delineated with colored squares and suggested subfamilies with colored circles.
284

Discussion

Using the conventional definition of "tailed phage" families, *Myoviridae*, *Podoviridae*, and

Siphoviridae (order *Caudovirales*), researchers effectively classified caudovirads for decades

[39,40]. However, the classification of these viruses, defined by a traditional morphology-

based approach, has been contested with the advent of high-throughput sequencing. The

steadily increasing number of available genomes and debates on the impact of horizontal

gene transfer, which marked the late 1990s and early 2000s, resulted in a decade-long

moratorium on the introduction of any new taxonomy for prokaryotic viruses [41]. The

increasing discrepancy between the official taxonomic framework and the emerging high-

volume genomics and metagenomics research left ≈90% of prokaryotic viruses known from

genome sequences unclassified beyond the family rank, i.e. they were classified as orphan

species in a family. Consequently, prokaryotic virus diversity was vastly underappreciated,

and virus genome curation remained in disarray. Recently, rather than implementing a

'repeal and replace' strategy for prokaryotic virus taxonomy, the Committee has introduced

a holistic system, involving virus particle morphology, overall DNA and protein sequence

identities, and phylogeny—an approach used for classification of all other viruses [1,2].

Successful modern taxonomic approaches must scale up to accommodate the

increasing pace of prokaryotic virus discovery and be effective across the 4,470 putative

complete prokaryotic virus sequences currently deposited in GenBank and other databases

[42]. These scaling requirements have remained problematic, and although there are more

than 3,400 publicly available caudovirad genomes, only 873 have been officially classified by

the ICTV by now [43]. The remaining genomes are provisionally stashed within

"unclassified" bins attached to the order *Caudovirales* or its member families, either

because they still need to be classified in a species/genus/subfamily, or because they may

be unidentified isolates of already classified viruses.

The growth in the number of prokaryotic virus genome sequences now supports the

application of a range of genomic analyses for robust taxonomic classification [44,45].

Meanwhile, phylogenomic approaches are yielding to network-based approaches to better

reflect the evolution of viral genomes [9,14]. These network-based approaches help to

organize the viral sequence space into statistically-resolvable "viral clusters." These clusters

are approximately equivalent to ICTV-recognized genera and provide a taxonomic

description that better reflects evolutionary relationships. Given the high computational

demand of network-based approaches, however, the development of centralized resources

and authority-monitored cyberinfrastructure, such as the iVirus platform on Cyverse [46] or

the Joint Genome Institute Integrated Microbial Genomes Virus resource IMG/VR [47], will

have to assist the prokaryotic virus community with large dataset computation and

classification.

Based on the results of this study, we suggest that the group of spounavirin and

spouna-like viruses be removed from the family *Myoviridae* and be given a family rank.

Hence, we propose establishing the suggested family "*Herelleviridae*", in the order

Caudovirales next to a smaller *Myoviridae* and the established *Podoviridae* and *Siphoviridae*

families. The new family would contain five subfamilies: *Spounavirinae* (sensu stricto),

"*Bastillevirinae*", "*Twortvirinae*", "*Jasinkavirinae*", and "*Brockvirinae*", each comprising the

ICTV-established genera listed in Table 1 (with additional information in S1 Table).

This study represents the first example of a true taxonomic assessment from an

'ensemble of methods'. The *de facto* taxon splitting suggested here results from the

observed diversity of prokaryotic viruses. We are encouraged that the combination of

genome BLAST analyses, virus proteomic trees, core protein clusters, genomic synteny

(GOAT), and single gene phylogenies yields consistent and complementary results, showing

the robustness of the suggested taxa. In addition, the suggested genera correspond well

with the taxonomy of the hosts (*Bacillus, Listeria, Staphylococcus*, and *Enterococcus*)

indicating broader microbiological consistency. Moreover, only approximately 3% of viruses

are left as unassigned at the genus and subfamily rank at this time within this group. These

unassigned viruses may represent clades at the genus and subfamily rank that are still

under-sampled.

This work demonstrates the usefulness of genome-based classification at a higher

taxonomic rank and its ability to accommodate the complex viral diversity. Substitution of

the families *Podoviridae*, *Myoviridae*, and *Siphoviridae* with a set of new families which

more faithfully reflect the true genetic relationships of the viruses would clarify the

taxonomic situation. However, this change does not remove the historically established

virus morphotypes observed in the nature among caudovirads: myovirids forming particles

with contractile tails, siphovirids forming particles with long non-contractile tails, and

podovirids forming particles with short non-contractile ones. By disconnecting morphotype

and family classification of caudovirads, taxonomically related clades can be grouped across

morphotypes. This grouping includes the muviruses suggested to be classified in the family

"*Saltoviridae*" [48] and potentially the broad set of Escherichia phage lambda-related

viruses that are currently distributed among the families *Siphoviridae* and *Podoviridae* [49].

We believe that abolishment of the *Podoviridae*, *Myoviridae* and *Siphoviridae*

families will soon be followed by the "upgrade" of existing viral taxonomy with additional

taxon ranks required to accommodate the observed diversity in an orderly manner.

Materials and methods

Creation of the dataset

Genome sequences of known spounavirins and spouna-like viruses were retrieved from
GenBank or (preferably) RefSeq databases in based on literature data, ICTV and taxonomic
classifications provided by the National Center for Biotechnology Information (NCBI).
Records representing genomes of candidate spounavirins and spouna-like viruses were
retrieved by searching the same databases with the tBLASTx algorithm using terminase and
major capsid proteins of several type virus isolates as a query (i.e., Bacillus phage SPO1,
Staphylococcus phage Twort, Bacillus phage Bastille, Listeria phage A511, Enterococcus
phage φEF24C, and Lactobacillus phage LP65) [50,51]. Sequences were manually curated
and pre-clustered using CLANS (E-value cut-off 1e-10) to confirm their spounaviral affiliation
[52]. This search yielded a set of 93 complete virus genomes, which were used in the
following analyses (S1 Table).

The coding sequences in the genomes were re-annotated using PROKKA with the
settings --kingdom Viruses, --E-value 1e-6 [53]. All genome sequences are available from
NCBI (accession number information listed in S1 Table) or from Github
(github.com/evelienadri/herelleviridae).

Genome-based analyses

Gegenees [54] was used in BLASTn and tBLASTx modes (fragment length 200 bp; step length

100 bp) to analyze virus genome nucleotide similarities. Pairwise identities between all

genomes under study were determined using BLASTn and tBLASTx algorithms with default

parameters [55]. Symmetrical identity scores (% SI) were calculated for each pairwise

comparison using the formula

% SI = 2.0 x $\frac{HL \; x \; HI}{QL + SL}$ (1)

in which the HL is defined as the hit length of the BLAST hit, HI is defined as the percentage

hit identity, QL is defined as the query length, and SL is defined as the subject length.

Symmetrical identity scores were converted into distances using the formula

Distance = $\sqrt[2]{1.0 - \%SI \div 100}$ (2)

The resulting distance matrix was hierarchically clustered (complete linkage) using the

hclust function of R [56]. Trees were visualized using Itol [57].

All pairwise comparisons of the nucleotide sequences using VICTOR, a Genome-

BLAST Distance Phylogeny (GBDP) method, were conducted under settings recommended

for prokaryotic viruses [58,59]. The resulting intergenomic distances (including 100

replicates each) were used to infer a balanced minimum evolution tree with branch support

via FASTME including subtree pruning and regrafting (SPR) post-processing [60] for each of

the formulas D0, D4, and D6, respectively. Trees were visualized with FigTree [61]. Taxon

demarcations at the species, genus and family rank were estimated with the OPTSIL

program [62], the recommended clustering thresholds [59], and an F value (fraction of links

required for cluster fusion) of 0.5 [58].

Proteomic tree

The Phage Proteomic Tree was constructed as described previously [63] and detailed at

https://github.com/linsalrob/PhageProteomicTree/tree/master/spounavirus. Briefly, the

protein sequences were extracted and clustered using BLASTp. These clusters were refined

by Smith-Waterman alignment using CLUSTALW version 2 [64]. Alignments were scored

using PROTDIST from the PHYLIP package [65]. Alignment scores were averaged and

weighted as described previously [63] resulting in the final tree.

Core protein clusters

Orthologous proteins were clustered using GET_HOMOLOGUES software, which utilizes

several independent clustering methods [66]. To capture as many evolutionary relationships

as possible, a greedy COGtriangles algorithm was applied with a 50% sequence identity

threshold, 50% coverage threshold, and an E-value cut-off equal to 1e-10 [67]. The results

were converted into an orthologue matrix with the "compare_clusters" script (part of the

GET_HOMOLOGUES suite) [65].

The orthologous protein clusters (OPCs) defined above were used to compute the

genomic fluidity for each pair of genomes. For two genomes i and j:

$$\text{Fluidity}(i,j) = \frac{Ui + Uj}{Mi + Mj} \tag{3}$$

 with Ui being the number of genes of i not found in j and Mi being the number of genes in i

[34]. The resulting distance matrix was hierarchically clustered (complete linkage) using the

hclust function of R [56]. Trees were visualized using Itol [57].

Multiple alignments were generated for each OPC using Clustal Omega [68]. For each

cluster, the amino acid identity between all protein pairs inside a cluster were determined

using multiple alignment. For all genome pairs, the AAI [69] was then computed and

transformed into distance using the formula:

Distance = $\frac{100 - AAI}{100}$ (4)

The resulting distance matrix was clustered and visualized as described above.

OPCs and multiple alignments for each cluster were used to determine a distance

similar to the distance used to generate the Phage Proteomic Tree. To estimate protein

distances, in this case, the dist function of the seqinR package [70]was preferred to

PROTDIST of the PHYLIP package [65] as the resulting distances are between 0 and 1.

Proteomic distances were then computed using the same formula as for the Phage

Proteomic Tree. The results were clustered and visualized as described above.

The Dice score is based on reciprocal BLAST searches between all pairs of genomes A

and B [31]. The total summed bitscoresof all tBLASTx hits with ≥30% identity, alignment

length ≥30 amino acids, and E-value ≤0.01 was converted to a distance DAB as follows:

DAB = $1 - \frac{S_{AB} + S_{BA}}{S_{AA} + S_{BB}}$ (5)

In which SAB SAB and SBA represent the summed bitscores between tBLASTx searches of A

versus B, and B versusu A, respectively, while SAA and SBB represent the summed tBLASTx

bitscores of the self-queries of A and B, respectively. The resulting distance matrix was

clustered with BionJ [71].

To investigate a genomic synteny-based classification signal, we developed a

geneorder-based metric built on dynamic programming, the Gene Order Alignment Tool

(GOAT, Schuller et al.: Python scripts are available on request, manuscript in preparation).

GOAT first identified protein-coding genes in the 93 spounavirin and spouna-like virus

genomes using Prodigal V2.6.3 in anonymous mode [72], and assigned them to the latest

pVOGs [73]). pVOG alignments (9,518) were downloaded (http://dmk-

brain.ecn.uiowa.edu/pVOGs/) and converted to profiles of hidden Markov models (HMM)

using HMMbuild (HMMer 3.1b2, [74]). Proteins were assigned to pVOGs using HMMsearch

(E-value <10-2) and used to generate a synteny profile of every genome. GOAT accounted

for gene replacements and distant homology by using an all-vs-all similarity matrix between

pVOG pairs based on HMM-HMM similarity (HH-suite 2.0.16) [75]). Distant HHsearch

similarity scores between protein families were calculated as the average of reciprocal hits

and used as substitution scores in the gene order alignment. The GOAT algorithm identified

the optimal gene order alignment score between two virus genomes by implementing semi-

global dynamic programming alignment based only on the order of pVOGs identified on

every virus genome. To account for virus genomes being cut at arbitrary positions during

sequence assembly, GOAT transmutes the gene order at all possible positions and in both

sense and antisense directions in search of the optimal alignment score. The optimal GOAT

alignment score GAB between every pair of virus genomes A and B, was converted to a

distance DAB as follows:

$\quad$ DAB $= 1 - \dfrac{G_{AB} + G_{BA}}{G_{AA} + G_{BB}}$ $\qquad\qquad$ (6)

in which GAB and GBA represent the optimal GOAT score between A and B, and B and A,

respectively, while GAA and GBB represent the GOAT scores of the self-alignments of A and

B, respectively. This pairwise distance matrix was clustered with BionJ [71].

Prokka re-annotated genomes were used to create pan-, core-, and accessory

genomes of all selected spounavirins and spouna-like viruses [53]. The annotations were

analyzed using Roary [35] with a 50% length BLASTp identity threshold for homologous

genes. Roary functions as follows: CD-HIT [76] was used to pre-cluster protein sequences

and perform an all-vs-all comparison of protein sequences with BLASTp to identify orthologs

and paralogs within the genomes. MCL [77] was then used to cluster the genomes based on

the presence and absence of the accessory genes. The resulting tree file was visualized using

FigTree v1.4.3 [61]. The tree was rooted in Brochothrix phage A9. The gene presence-

absence output table from Roary was then imported into R and using a custom R-script

(available from github.com/evelienadri/herelleviridae/tree/master). Pairwise shared gene

contents were calculated for each combination of genomes.

Single gene phylogenies

Based on the OPC and pVOG analyses, we chose nine well-annotated protein clusters

present in all 93 spounavirins and spouna-like viruses. Selected clusters included: DNA

helicases, major capsid proteins, tail sheath proteins, two different groups of baseplate

proteins, and four clusters with no known function. The members of these clusters were

aligned using Clustal Omega with default parameters [47]. Resulting alignments were

analyzed with ProtTest 3.4 [59] to determine a suitable protein evolution model (only

variations of models compatible with downstream software like JTT and WAG were

considered). Estimated models were used to generate phylograms with FastTree 2.1.7 [60].

The program implements the approximately maximum-likelihood method with Shimodaira-

Hasegawa tests to generate the tree and calculate support of the splits. This approach is

much faster than "traditional" maximum-likelihood methods with negligible accuracy loss

[59–61].

Figure legends

Fig 1: Genome-based clustering trees of 93 spounavirin and spouna-like viruses. Clustering was performed using nucleotide similarities (BLASTn, A) or translated nucleotide similarities (tBLASTx, B). Genomes were compared in a pairwise fashion using Gegenees, transformed into a distance matrix, clustered using R and visualized as trees using Itol. The trees were rooted at Brochotrix phage A9. Genera are delineated with colored squares and suggested subfamilies with colored circles.

Fig 2: Predicted proteome-based clustering trees of 93 spounavirin and spouna-like viruses. Clustering was performed using the Phage Proteomics Tree approach (A) and proteomic distance (B). Distances were calculated pairwise between all sets of predicted proteomes, clustered with R and visualized using Itol. The trees were rooted at Brochotrix phage A9. Genera are delineated with colored squares and suggested subfamilies with colored circles.

Fig 3: Single gene phylogenies of the major capsid protein (MCP, A), tail sheath protein (TSP, B) and helicase (C) amino acid sequences of 93 spounavirin and spouna-like viruses. Amino acid sequences were aligned with Clustal Omega and trees were generated using FastTree maximum likelihood with Shimidaira-Hasegawa tests. The scale bar represents the number of substitutions per site. The trees were rooted at Brochotrix phage A9. Genera are delineated with colored squares and suggested subfamilies with colored circles.

Supporting information

S1 Fig. Heatmap of the blastn-based nucleotide similarities between pairs of genomes as calculated with Gegenees at default parameters.

S2 Fig. Genome-blast Distance Phylogeny as calculated using VICTOR.

S3 Fig. Heatmap of the DICE coefficient calculated between each pair of genomes.

S4 Fig. Heatmap of the tblastx-based nucleotide similarities between pairs of genomes as calculated with Gegenees at default parameters.

S5 Fig. Heatmap of the pairwise comparison of all genomes visualized as percentage of shared orthologous proteins (OPCs) as calculated on original GenBank files.

S6 Fig. Heatmap of the pairwise comparison of all genomes visualized as percentage of shared orthologous proteins (OPCs) as calculated on reannotated genomes.

S7 Fig. Heatmap of the pairwise comparison of all genomes visualized as percentage of shared proteins as calculated with Roary on reannotated genome files.

S8 Fig. Clustering trees of genomic fluidity and amino acid identity calculated pairwise between all genomes using orthologous protein clusters.

S9 Fig. Accessory genome clustering tree, calculated based on the presence and absence of accessory genes in each genome.

S10 Fig. Heatmap and clustering tree calculated by the Gene Order Alignment Tool and visualized as a distance matrix between all genome pairs.

S11 Fig. Maximum Likelihood trees of single gene phylogenies using protein clusters present in all 93 genomes.

S1 Table. Overview of the 93 phage genomes used in this study.

S2 Table. Complete list of all orthologous proteins identified in the set of 93 spounavirin and spouna-like virus genomes.

540   **S3 Table.** Complete list of prokaryotic virus orthologous groups identified in the set of 93

541   spounavirin and spouna-like virus genomes.

Acknowledgments

We thank Laura Bollinger, Integrated Research Facility at Fort Detrick, for technical writing

services. EMA would like to thank PH Nel for assistance with R scripting. JB was supported

by the National Science Centre (Poland SONATA 12 grant number 2016/23/D/NZ2/00435).

MBPS and BED were supported by the Netherlands Organization for Scientific Research

(NWO) Vidi grant 864.14.004. RAE was supported by grants DUE-132809 and MCB-1330800

from the US National Science Foundation. HMO was supported by University of Helsinki

funding for Instruct research infrastructure (Virus and Macromolecular Complex Production,

ICVIR). AG was supported by a *Chargé de Recherches* fellowship from the National Fund for

Scientific Research (FNRS, Belgium). FE was supported by the EUed Horizon 2020

Framework Programme for Research and Innovation ('Virus-X', project no. 685778). MBS

would like to acknowledge the Gordon and Betty Moore Foundation Investigator Award

(GBMF#3790) for funding.

Competing interests

558   All authors, except for MBPS, are members of the Bacterial and Archaeal Viruses

559   Subcommittee of the International Committee on the Taxonomy of Viruses (ICTV). The

560   authors declare no conflicts of interest.

References

References

1.   Krupovic M, Dutilh BE, Adriaenssens EM, Wittmann J, Vogensen FK, Sullivan MB, et al.

Taxonomy of prokaryotic viruses: update from the ICTV bacterial and archaeal viruses

subcommittee. Arch Virol. Springer Vienna; 2016;161: 1095–1099.

doi:10.1007/s00705-015-2728-0

2.   Adriaenssens EM, Krupovic M, Knezevic P, Ackermann H-W, Barylski J, Brister JR, et al.

Taxonomy of prokaryotic viruses: 2016 update from the ICTV bacterial and archaeal

viruses subcommittee. Arch Virol. Springer Vienna; 2017;162: 1153–1157.

doi:10.1007/s00705-016-3173-4

3.   Lavigne R, Seto D, Mahadevan P, Ackermann H-W, Kropinski AM. Unifying classical

and molecular taxonomic classification: analysis of the Podoviridae using BLASTP-

based tools. Res Microbiol. 2008;159: 406–414. doi:10.1016/j.resmic.2008.03.005

4.   Lavigne R, Darius P, Summer EJ, Seto D, Mahadevan P, Nilsson AS, et al. Classification

of Myoviridae bacteriophages using protein sequence similarity. BMC Microbiol.

2009;9: 224. doi:10.1186/1471-2180-9-224

5.   Adriaenssens EM, Edwards R, Nash JHE, Mahadevan P, Seto D, Ackermann H-W, et al.

Integration of genomic and proteomic analyses in the classification of the

Siphoviridae family. Virology. 2015;477: 144–154. doi:10.1016/j.virol.2014.10.016

6.   Gregory AC, Solonenko SA, Ignacio-Espinoza JC, LaButti K, Copeland A, Sudek S, et al.

Genomic differentiation among wild cyanophages despite widespread horizontal

gene transfer. BMC Genomics. BMC Genomics; 2016;17: 930. doi:10.1186/s12864-

016-3286-x

7.   Brum JR, Ignacio-Espinoza JC, Roux S, Doulcier G, Acinas SG, Alberti A, et al. Patterns

and ecological drivers of ocean viral communities. Science (80-). 2015;348: 1261498–

1261498. doi:10.1126/science.1261498

8.   Holmfeldt K, Solonenko N, Shah M, Corrier K, Riemann L, Verberkmoes NC, et al.

Twelve previously unknown phage genera are ubiquitous in global oceans. Proc Natl

Acad Sci U S A. 2013;110: 12798–12803. doi:10.1073/pnas.1305956110

9.   Lima-Mendez G, Van Helden J, Toussaint A, Leplae R. Reticulate representation of

evolutionary and functional relationships between phage genomes. Mol Biol Evol.

2008;25: 762–777. doi:10.1093/molbev/msn023

10.   Nishimura Y, Watai H, Honda T, Mihara T, Omae K, Roux S, et al. Environmental viral

genomes shed new light on virus-host interactions in the ocean. mSphere. 2017;2:

e00359-16. doi:10.1128/mSphere.00359-16

11.   Paez-Espino D, Eloe-Fadrosh EA, Pavlopoulos GA, Thomas AD, Huntemann M,

Mikhailova N, et al. Uncovering Earth's virome. Nature. 2016;536: 425–430.

doi:10.1038/nature19094

12.   Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, et al. Ecogenomics and

potential biogeochemical impacts of globally abundant ocean viruses. Nature.

2016;537: 689–693. doi:10.1038/nature19366

13.   Iranzo J, Krupovic M, Koonin E V. The double-stranded DNA virosphere as a modular

hierarchical network of gene sharing. MBio. 2016;7: e00978-16.

doi:10.1128/mBio.00978-16

14.   Bolduc B, Jang H Bin, Doulcier G, You Z-Q, Roux S, Sullivan MB. vConTACT: an iVirus

tool to classify double-stranded DNA viruses that infect Archaea and Bacteria. PeerJ.

2017;5: e3243. doi:10.7717/peerj.3243

15.   Okubo S, Strauss B, Stodolsky M. The possible role of recombination in the infection

of competent Bacillus subtilis by bacteriophage deoxyribonucleic acid. Virology.

1964;24: 552–562. doi:10.1016/0042-6822(64)90207-7

16. Parker ML, Eiserling FA. Bacteriophage SPO1 structure and morphogenesis I. Tail

structure and length regulation. J Virol. 1983;46: 239–249.

17. Duda RL, Hendrix RW, Huang WM, Conway JF. Shared architecture of bacteriophage

SPO1 and herpesvirus capsids. Curr Biol. 2006;16: R11–R13.

doi:10.1016/j.cub.2006.02.014

18. Hemphill HE, Whiteley HR. Bacteriophages of Bacillus subtilis. Bacteriol Rev. 1975;39:

257–315.

19. Okubo S, Yanagida T, Fujita DJ, Olsson-Wilhelm BM. The genetics of bacteriophage

SPO1. Biken J. 1972;15: 81–97.

20. Klumpp J, Lavigne R, Loessner MJ, Ackermann H-W. The SPO1-related bacteriophages.

Arch Virol. 2010;155: 1547–61. doi:10.1007/s00705-010-0783-0

21. Stewart CR, Casjens SR, Cresawn SG, Houtz JM, Smith AL, Ford ME, et al. The genome

of Bacillus subtilis bacteriophage SPO1. J Mol Biol. 2009;388: 48–70.

doi:10.1016/j.jmb.2009.03.009

22. Murphy FA, Fauquet CM, Bishop DHL, Ghabrial SA, Jarvis AW, Martelli GP, et al. Sixth

report of the International Committee on Taxonomy of Viruses. Arch Virol Suppl.

1995;10: 590.

23. Ackermann H-W, Elzanowski A, Fobo G, Stewart G. Relationships of tailed phages: a

survey of protein sequence identity. Arch Virol. 1995;140: 1871–1884.

doi:10.1007/BF01384350

24. Adams MJ, Carstens EB. Ratification vote on taxonomic proposals to the International

Committee on Taxonomy of Viruses (2012). Arch Virol. 2012;157: 1411–1422.

doi:10.1007/s00705-012-1299-6

25.   Schuch R, Fischetti VA. The secret life of the anthrax agent Bacillus anthracis:

636   Bacteriophage-mediated ecological adaptations. PLoS One. 2009;4: e6532.

637   doi:10.1371/journal.pone.0006532

638   26.   Yuan Y, Peng Q, Wu D, Kou Z, Wu Y, Liu P, et al. Effects of actin-like proteins encoded

639   by two Bacillus pumilus phages on unstable lysogeny, revealed by genomic analysis.

640   Appl Environ Microbiol. 2015;81: 339–350. doi:10.1128/AEM.02889-14

641   27.   Klumpp J, Dorscht J, Lurz R, Bielmann R, Wieland M, Zimmer M, et al. The terminally

642   redundant, nonpermuted genome of Listeria bacteriophage A511: a model for the

643   SPO1-like myoviruses of gram-positive bacteria. J Bacteriol. 2008;190: 5753–65.

644   doi:10.1128/JB.00461-08

645   28.   Barylski J, Nowicki G, Goździcka-Józefiak A. The discovery of phiAGATE, a novel phage

646   infecting Bacillus pumilus, leads to new insights into the phylogeny of the subfamily

647   Spounavirinae. Dąbrowska K, editor. PLoS One. 2014;9: e86632.

648   doi:10.1371/journal.pone.0086632

649   29.   Uchiyama J, Rashel M, Maeda Y, Takemura I, Sugihara S, Akechi K, et al. Isolation and

650   characterization of a novel Enterococcus faecalis bacteriophage phiEF24C as a

651   therapeutic candidate. FEMS Microbiol Lett. 2008;278: 200–206. doi:10.1111/j.1574-

652   6968.2007.00996.x

653   30.   Uchiyama J, Rashel M, Takemura I, Wakiguchi H, Matsuzaki S. In silico and in vivo

654   evaluation of bacteriophage phiEF24C, a candidate for treatment of Enterococcus

655   faecalis infections. Appl Environ Microbiol. 2008;74: 4149–4163.

656   doi:10.1128/AEM.02371-07

657   31.   Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. Expanding the marine virosphere

658   using metagenomics. Rocha EPC, editor. PLoS Genet. 2013;9: e1003987.

doi:10.1371/journal.pgen.1003987

32.   Goodrich-Blair H, Scarlato V, Gott JM, Xu M-Q, Shub DA. A self-splicing group I intron

in the DNA polymerase gene of bacillus subtilis bacteriophage SPO1. Cell. 1990;63:

417–424. doi:10.1016/0092-8674(90)90174-D

33.   Lavigne R, Vandersteegen K. Group I introns in Staphylococcus bacteriophages.

Future Virol. 2013;8: 997–1005. doi:10.2217/fvl.13.84

34.   Kislyuk AO, Haegeman B, Bergman NH, Weitz JS. Genomic fluidity: An integrative view

of gene diversity within microbial populations. BMC Genomics. 2011;12: 32.

doi:10.1186/1471-2164-12-32

35.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid

large-scale prokaryote pan genome analysis. Bioinformatics. 2015;31: 3691–3693.

doi:10.1093/bioinformatics/btv421

36.   Juhala RJ, Ford ME, Duda RL, Youlton A, Hatfull GF, Hendrix RW. Genomic sequences

of bacteriophages HK97 and HK022: pervasive genetic mosaicism in the lambdoid

bacteriophages. J Mol Biol. 2000;299: 27–51. doi:10.1006/jmbi.2000.3729

37.   Krupovic M, Prangishvili D, Hendrix RW, Bamford DH. Genomics of bacterial and

archaeal viruses: Dynamics within the prokaryotic virosphere. Microbiol Mol Biol Rev.

2011;75: 610–635. doi:10.1128/MMBR.00011-11

38.   Cadungog JN, Khatemi BE, Hernandez AC, Kuty Everett GF. Complete genome

sequence of Bacillus megaterium myophage Moonbeam. Genome Announc. 2015;3:

e01428-14. doi:10.1128/genomeA.01428-14

39.   Ackermann H-W, DuBow MS. Viruses of prokaryotes. Boca Raton, FL, USA: CRC Press;

1987.

40.   Ackermann H-W. Classification of bacteriophages. In: Calendar R, editor. The

Bacteriophages. 2nd ed. New York, NY, USA: Oxford University Press; 2006. pp. 8–17.

41.   Tolstoy I, Kropinski AM, Brister JR. Bacteriophage taxonomy: an evolving discipline. In:

Azeredo J, Sillankorva S, editors. Methods in Molecular Biology: Bacteriophage

Therapy. Springer Nature; doi:Forthcoming

42.   Karsch-Mizrachi I, Nakamura Y, Cochrane G. The International Nucleotide Sequence

Database Collaboration. Nucleic Acids Res. 2012;40: D33–D37.

doi:10.1093/nar/gkr1006

43.   Davison AJ. Journal of General Virology – Introduction to "ICTV Virus Taxonomy

Profiles." J Gen Virol. 2017;98: 1–1. doi:10.1099/jgv.0.000686

44.   Simmonds P, Adams MJ, Benkő M, Breitbart M, Brister JR, Carstens EB, et al.

Consensus statement: Virus taxonomy in the age of metagenomics. Nat Rev

Microbiol. 2017;15: 161–168. doi:10.1038/nrmicro.2016.177

45.   Adams MJ, Lefkowitz EJ, King AMQ, Harrach B, Harrison RL, Knowles NJ, et al. 50 years

of the International Committee on Taxonomy of Viruses: progress and prospects.

Arch Virol. 2017;162: 1441–1446. doi:10.1007/s00705-016-3215-y

46.   Bolduc B, Youens-Clark K, Roux S, Hurwitz BL, Sullivan MB. iVirus: facilitating new

insights in viral ecology with software and community data sets imbedded in a

cyberinfrastructure. ISME J. 2017;11: 7–14. doi:10.1038/ismej.2016.89

47.   Paez-Espino D, Chen IMA, Palaniappan K, Ratner A, Chu K, Szeto E, et al. IMG/VR: A

database of cultured and uncultured DNA viruses and retroviruses. Nucleic Acids Res.

2017;45: D457–D465. doi:10.1093/nar/gkw1030

48.   Hulo C, Masson P, Le Mercier P, Toussaint A. A structured annotation frame for the

transposable phages: A new proposed family "Saltoviridae" within the Caudovirales.

Virology. 2015;477: 155–163. doi:10.1016/j.virol.2014.10.009

49.   Grose JH, Casjens SR. Understanding the enormous diversity of bacteriophages: The

708   tailed phages that infect the bacterial family Enterobacteriaceae. Virology. 2014;468–

709   470: 421–443. doi:10.1016/j.virol.2014.08.024

710   50.   Brister JR, Ako-adjei D, Bao Y, Blinkova O. NCBI Viral Genomes Resource. Nucleic Acids

711   Res. 2015;43: D571–D577. doi:10.1093/nar/gku1207

712   51.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.

713   J Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

714   52.   Frickey T, Lupas A. CLANS: A Java application for visualizing protein families based on

715   pairwise similarity. Bioinformatics. 2004;20: 3702–3704.

716   doi:10.1093/bioinformatics/bth444

717   53.   Seemann T. Prokka: Rapid prokaryotic genome annotation. Bioinformatics. 2014;30:

718   2068–2069. doi:10.1093/bioinformatics/btu153

719   54.   Ågren J, Sundström A, Håfström T, Segerman B. Gegenees: Fragmented alignment of

720   multiple genomes for determining phylogenomic distances and genetic signatures

721   unique for specified target groups. PLoS One. 2012;7: e39107.

722   doi:10.1371/journal.pone.0039107

723   55.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

724   architecture and applications. BMC Bioinformatics. 2009;10: 421. doi:10.1186/1471-

725   2105-10-421

726   56.   Development Core Team R. R: A language and environment for statistical computing

727   [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2008. Available:

728   http://www.r-project.org

729   57.   Letunic I, Bork P. Interactive Tree Of Life (iTOL): An online tool for phylogenetic tree

730   display and annotation. Bioinformatics. 2007;23: 127–128.

doi:10.1093/bioinformatics/btl529

58.   Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, et al.

Complete genome sequence of DSM 30083T, the type strain (U5/41T) of Escherichia

coli, and a proposal for delineating subspecies in microbial taxonomy. Stand Genomic

Sci. 2014;9: 2. doi:10.1186/1944-3277-9-2

59.   Meier-Kolthoff JP, Göker M. VICTOR: genome-based phylogeny and classification of

prokaryotic viruses. Bioinformatics. 2017;33: 3396–3404.

doi:10.1093/bioinformatics/btx440

60.   Lefort V, Desper R, Gascuel O. FastME 2.0: A comprehensive, accurate, and fast

distance-based phylogeny inference program. Mol Biol Evol. 2015;32: 2798–2800.

doi:10.1093/molbev/msv150

61.   Rambaud. FigTree [Internet]. 2007 [cited 1 Aug 2015]. Available:

http://tree.bio.ed.ac.uk/software/figtree/

62.   Göker M, García-Blázquez G, Voglmayr H, Tellería MT, Martín MP. Molecular

taxonomy of phytopathogenic fungi: A case study in Peronospora. PLoS One. 2009;4:

8–10. doi:10.1371/journal.pone.0006319

63.   Rohwer F, Edwards R. The Phage Proteomic Tree: a genome-based taxonomy for

phage. J Bacteriol. Am Soc Microbiol; 2002;184: 4529–4535.

doi:10.1128/JB.184.16.4529

64.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.

Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23: 2947–2948.

doi:10.1093/bioinformatics/btm404

65.   Felsenstein J. PHYLIP - Phylogeny inference package - v3.2. Cladistics. 1989;5: 164–

166. doi:10.1111/j.1096-0031.1989.tb00562.x

66. Contreras-Moreira B, Vinuesa P. GET_HOMOLOGUES, a versatile software package for

scalable and robust microbial pangenome analysis. Appl Environ Microbiol. 2013;79:

7696–7701. doi:10.1128/AEM.02411-13

67. Kristensen DM, Kannan L, Coleman MK, Wolf YI, Sorokin A, Koonin E V, et al. A low-

polynomial algorithm for assembling clusters of orthologous groups from

intergenomic symmetric best matches. Bioinformatics. 2010;26: 1481–7.

doi:10.1093/bioinformatics/btq229

68. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation

of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst

Biol. 2011;7: 539. doi:10.1038/msb.2011.75

69. Konstantinidis KT, Tiedje JM. Towards a genome-based taxonomy for prokaryotes. J

Bacteriol. 2005;187: 6258–6264. doi:10.1128/JB.187.18.6258-6264.2005

70. Charif D, Thioulouse J, Lobry JR, Perrière G. Online synonymous codon usage analyses

with the ade4 and seqinR packages. Bioinformatics. 2005;21: 545–547.

doi:10.1093/bioinformatics/bti037

71. Gascuel O. BIONJ: an improved version of the NJ algorithm based on a simple model

of sequence data. Mol Biol Evol. 1997;14: 685–695.

doi:10.1017/CBO9781107415324.004

72. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. BMC Bioinformatics.

2010;11. doi:10.1186/1471-2105-11-119

73. Grazziotin AL, Koonin E V, Kristensen DM. Prokaryotic Virus Orthologous Groups

(pVOGs): a resource for comparative genomics and protein family annotation. Nucleic

Acids Res. 2017;45: D491–D498. doi:10.1093/nar/gkw975

74.   Finn RD, Clements J, Eddy SR. HMMER web server: Interactive sequence similarity

780   searching. Nucleic Acids Res. 2011;39: 29–37. doi:10.1093/nar/gkr367

781   75.   Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology

782   detection and structure prediction. Nucleic Acids Res. 2005;33: W244-2488.

783   doi:10.1093/nar/gki408

784   76.   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: Accelerated for clustering the next-generation

785   sequencing data. Bioinformatics. 2012;28: 3150–3152.

786   doi:10.1093/bioinformatics/bts565

787   77.   Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale

788   detection of protein families. Nucleic Acids Res. 2002;30: 1575–1584. doi:doi:

789   10.1093/nar/30.7.1575