TITLE: Age-accelerated cognitive decline in asymptomatic adults with CSF β-amyloid

AUTHORS: Lindsay R. Clark, PhD, Sara E. Berman, BS, Derek Norton, MS, Rebecca L. Koscik, PhD, Erin Jonaitis, PhD, Kaj Blennow, MD, PhD, Barbara B. Bendlin, PhD, Sanjay Asthana, MD, Sterling C. Johnson, PhD, Henrik Zetterberg, MD, PhD, Cynthia M. Carlsson, MD, MS

AFFILIATIONS AND CONTACT INFORMATION:
Lindsay R. Clark, PhD[a,b,c], lrclark@medicine.wisc.edu
Sara E. Berman, BS[b,d], seberman@medicine.wisc.edu
Derek Norton, MS[b,e], dlnorton2@wisc.edu
Rebecca L. Koscik, PhD[c], rekoscik@wisc.edu
Erin Jonaitis, PhD[c], jonaitis@wisc.edu
Kaj Blennow, MD, PhD[f,g], Kaj.Blennow@neuro.gu.se
Barbara B. Bendlin, PhD[b,c], bbb@medicine.wisc.edu
Sanjay Asthana, MD[a,b], sa@medicine.wisc.edu
Sterling C. Johnson, PhD[a,b,c], scj@medicine.wisc.edu
Henrik Zetterberg, MD, PhD[f,g,h], henrik.zetterberg@clinchem.gu.se
Cynthia M. Carlsson, MD, MS[a,b,c], cmc@medicine.wisc.edu

[a]Geriatric Research Education and Clinical Center, William. S. Middleton Memorial Veterans Hospital, Madison WI 53705, USA.
[b]Alzheimer's Disease Research Center, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA.
[c]Wisconsin Alzheimer's Institute, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA.
[d]Medical Scientist and Neuroscience Training Programs, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA.
[e]Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA.
[f]Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden.
[g]Department of Psychiatry and Neurochemistry, Institute of Neuroscience & Physiology, The Sahlgrenska Academy at the University of Gothenburg, Molndal, Sweden
[h]Department of Molecular Neuroscience, University College London, Institute of Neurology, Queen Square, London, UK

Corresponding author: Lindsay R. Clark, Ph.D., William S. Middleton Memorial VA Hospital, 2500 Overlook Terrace (11G), GRECC, Madison, WI, 53705, USA. Phone: 608-263-4405; Fax: 608-265-3091; Email: lrclark@medicine.wisc.edu.

Title characters (with spaces): 75   Abstract word count: 250
Manuscript word count: 2991   References: 24
Tables and Figures: 5
Supplementary Material: 482 words, 3 references, 5 tables
Search Terms: Alzheimer's disease, cognitive aging, cognitive neuropsychology in dementia, neuropsychological assessment, cerebrospinal fluid

AUTHOR CONTRIBUTIONS

Lindsay Clark: Drafting/revising the manuscript, study concept or design, analysis or interpretation of data, statistical analysis.
Sara Berman: Drafting/revising the manuscript, study concept or design, analysis or interpretation of data, statistical analysis.
Derek Norton: Analysis or interpretation of data, statistical analysis.
Rebecca Koscik: Drafting/revising the manuscript, Analysis or interpretation of data, statistical analysis.
Erin Jonaitis: Drafting/revising the manuscript, Analysis or interpretation of data, statistical analysis.
Kaj Blennow: Analysis or interpretation of data, Contribution of vital reagents/tools/patents.
Barbara Bendlin: Analysis or interpretation of data, obtaining funding.
Sanjay Asthana: Study concept or design, study supervision or coordination, obtaining funding. Dr. Asthana reports no disclosures.
Sterling Johnson: Drafting/revising the manuscript, study concept or design, analysis or interpretation of data, study supervision or coordination, obtaining funding.
Henrik Zetterberg: Analysis or interpretation of data, contribution of vital reagents/tools/patents.
Cynthia Carlsson: Study concept or design, analysis or interpretation of data, acquisition of data, study supervision or coordination.

AUTHOR DISCLOSURES

Dr. Clark reports no disclosures.
Ms. Berman reports no disclosures.
Mr. Norton reports no disclosures.
Dr. Koscik reports no disclosures.
Dr. Jonaitis reports no disclosures.
Dr. Blennow holds the Torsten Söderberg professorship in Medicine.
Dr. Bendlin reports no disclosures.
Dr. Johnson reports no disclosures.
Dr. Zetterberg is a Wallenberg Academy Fellow.
Dr. Carlsson is a site principal investigator for a project jointly funded by the NIH and Lilly (A4 study).

STUDY FUNDING

The project described was supported by the Clinical Translational Science Award (CTSA) program through the NIH National Center for Advancing Translational Sciences (NCATS), and grant UL1TR00427. Funding support was also provided by NIH grants ADRC P50 AG033514 (SA), R01 AG027161 (SCJ), R01 AG021155 (SCJ), R01AG037639 (BBB), F30 AG054115 (SEB), T32 GM007507 (SEB) and T32 GM008692 (SEB).

ABSTRACT

Objective: Compare cognitive and hippocampal volume (HCV) trajectories in asymptomatic middle-aged and older adults with positive cerebrospinal fluid (CSF) markers of β-amyloid (Aβ) or tau to adults without an AD-associated biomarker profile.

Method: 392 adults enrolled in a longitudinal cohort study (Wisconsin Registry for Alzheimer's Prevention or Wisconsin Alzheimer's Disease Research Center) completed a lumbar puncture and at least two biennial or annual neuropsychological evaluations. Cutoffs for $A\beta_{42}$, total tau, and phosphorylated tau were developed via receiver operating characteristic curve analyses on a sample of 78 participants (38 dementia, 40 controls). These cutoffs were applied to a separate sample of 314 cognitively healthy adults (mean age at CSF collection = 61.5) and mixed-effects regression analyses tested linear and quadratic interactions of biomarker group x age at each visit on cognitive and HCV outcomes.

Results: 215 participants (69%) were biomarker negative (preclinical AD Stage 0), 46 (15%) were Aβ+ only (preclinical AD Stage 1), 25 (8%) were Aβ+ and tau+ (preclinical AD Stage 2), and 28 (9%) were tau+ only. Both Stage 1 and Stage 2 groups exhibited greater rates of linear decline on story memory and processing speed measures, and non-linear decline on list-learning and set-shifting measures compared to Stage 0. The tau+ only group did not significantly differ from Stage 0 in rates of cognitive decline.

Conclusion: In an asymptomatic at-risk cohort, elevated CSF Aβ (with or without elevated tau) was associated with greater rates of cognitive decline, with the specific pattern of decline varying across cognitive measures.

INTRODUCTION

Although most studies of preclinical Alzheimer's disease (AD) focus on older adults, recent studies report that middle-aged adults with CSF biomarkers of both beta-amyloid (Aβ) and tau exhibit more rapid decline on cognitive and clinical measures than those with only one abnormal biomarker[1, 2]. These studies support guidelines defining preclinical AD as the presence of Aβ and neurodegeneration, while designating the presence of only one feature as "asymptomatic at-risk for AD"[3]. However, prior studies examined change on cognitive composite scores or global screening measures and it remains unclear whether the presence of either Aβ or tau in isolation is associated with decline within specific cognitive domains, such as memory. Additionally, although cutoff values defining normal or abnormal levels of Aβ and tau are useful clinically, examining relationships between biomarkers and clinical symptoms along a continuum may provide additional information.

Our analysis was designed to replicate and build upon prior work by: 1) identifying Aβ and tau positivity in a longitudinal cohort sample of cognitively healthy middle-aged and older adults, 2) comparing biomarker groups on longitudinal neuropsychological performance across multiple measures and 3) investigating relationships between continuous variables of Aβ, tau, and cognitive performance. We hypothesized that adults with both Aβ and tau positivity would exhibit greater rates of cognitive decline compared to biomarker negatives. Based on prior work showing associations between Aβ and cognitive decline[4, 5], we further hypothesized that those with Aβ+ would exhibit

greater decline on memory measures, whereas tau+ adults would not differ from biomarker negatives.

METHODS

Participants

Participants included 392 middle-aged or older community-dwelling adults enrolled in longitudinal cohort studies of Wisconsin Registry for Alzheimer's Prevention (WRAP) [6] (*n*=141) or the Wisconsin Alzheimer's Disease Research Center (WADRC) clinical core (*n*=251). These cohorts include cognitively healthy and impaired participants, are enriched for at-risk adults with family history of AD, and conduct study evaluations on an annual or biennial basis. Cognitive status was determined by consensus conference panel based on National Institute on Aging-Alzheimer's Association (NIA-AA) criteria[7, 8]. The current study included participants with dementia in the development of CSF cutoff values, but included cognitively healthy middle-aged and older adults in all remaining analyses. Exclusion criteria consisted of only one study visit completed, relevant CSF or diagnosis data unavailable, diagnosis of mild cognitive impairment (MCI) or Impaired-not MCI at baseline or LP visit, or diagnosis of dementia that reverted to MCI at subsequent visits. Participants with incomplete neuropsychological data were included if data for at least two visits were available. Participants from the WRAP cohort were younger at baseline than those from the WADRC, but samples were similar in sex distribution, education, and *APOE* genotype (see Supplemental Table 1 for characteristics of cognitively healthy participants across cohorts).

Standard Protocol Approvals, Registrations, and Patient Consents

The inclusion of human subjects in this study was approved by the University of Wisconsin-Madison Institutional Review Board and all participants provided informed consent.

Procedures

CSF was collected in the morning after a minimum 12-hour fast. A Sprotte spinal needle was inserted into the L3-L4 or L4-L5 vertebral interspace and 22 mL of CSF was removed via gentle extraction into polypropylene syringes. Within 30 minutes of collection, the CSF was combined, gently mixed, centrifuged to remove red blood cells or other debris, aliquoted into 0.5-mL polypropylene tubes, and stored at -80°C. Samples were sent in batches at two time points for analysis at the Clinical Neurochemistry Laboratory at the Sahlgrenska Academy of the University of Gothenburg, Sweden. All samples were analyzed according to protocols approved by the Swedish Board of Accreditation and Conformity Assessment using one batch of reagents (intraassay coefficients of variation < 10%) for each batch. Board-certified laboratory technicians blinded to clinical diagnosis performed all analyses on one occasion for each of the two batches. CSF samples were assayed for total tau (t-tau), phosphorylated tau (p-tau$_{181}$), amyloid beta 1–42 (Aβ42), and amyloid beta 1–40 (Aβ40) using commercially available enzyme-linked immunosorbent assay methods (INNOTEST assays, Fujirebio, Ghent, Belgium; Triplex assays, MSD Human Aβ peptide ultra-sensitive kit, Meso Scale Discovery, Gaithersburg, MD). Additional details on batch-to-batch conversions is provided in the Supplemental Material.

A comprehensive neuropsychological assessment was completed at each visit. Measures of memory and executive functioning were included in analyses due to evidence that these demonstrate decline in preclinical AD[9, 10]. Overlapping measures within these domains across cohorts assessed verbal memory (Rey Auditory Verbal Learning Test [RAVLT] Total Trials 1-5 and Delayed Recall [11], Wechsler Memory Scale-Revised Logical Memory Story A [LM] Immediate and Delayed Recall) [12] and executive functioning (Trailmaking Test Part B [TMT-B] [13], Animal Fluency, Wechsler Adult Intelligence Scale-Revised Digit Symbol [14]). A subset of 205 participants completed at least two MRI scans and were included in secondary analyses of hippocampal volume (HCV) change (see Supplemental Material for MRI details).

Statistical analyses

Statistical analyses were conducted in R version 3.3.1 [15]. Cutoff values for CSF assays were developed using receiver operating characteristic (ROC) curve analysis in the pROC package (version 1.8) [16] in a sample of 38 participants with clinical diagnosis of dementia due to AD based on NIA-AA criteria[8] without reference to CSF biomarkers and 40 late middle-age (ages 48-64) stable cognitively healthy adults at lower risk for AD (*APOE* ε4 non-carrier, no family history of AD). Youden's J (sensitivity + specificity - 1), which maximizes both the sensitivity and specificity of a diagnostic test, was used.

To reduce potential risk of researcher assessment bias, a non-overlapping sample of 314 cognitively healthy participants (mean LP age 61.5) were included in subsequent

analyses. We compared biomarker groups on demographic and AD risk characteristics using chi-square and analysis of variance (ANOVA) tests. We compared mean neuropsychological performance and HCV among biomarker groups at the visit closest to the LP using analysis of covariance (ANCOVA) models with age at LP (mean = 61.5), sex (reference group = female), and years of education (mean = 16.3) as covariates. Comparisons of HCV also included TIV (mean = 1464.8 mm^3) as a covariate.

To test if longitudinal change on the seven neuropsychological measures and HCV varied across biomarker groups, linear mixed-effects models were conducted using the lme4 package version 1.1-12[17]. Fixed effects included sex, years of education, practice effects (number of exposures to test [18]), biomarker group (4 levels), age (at each visit), and the interaction of age x biomarker group. To allow for acceleration of cognitive decline with increasing age, two quadratic terms, age^2 and age^2 x biomarker group, were included in all models and removed if non-significant. To minimize collinearity in the linear and quadratic age terms, the age variable was centered on the sample mean. All models included random effects of intercept and slope nested within subject. The overall significance of the interaction term was assessed by likelihood ratio tests comparing the primary model and a model that did not include the interaction term. *P-*values for fixed effect coefficients were calculated using asymptotic properties of the estimates[19]. Statistical significance was defined as *p* < .05.

To investigate the relationship between cognitive or HCV change and continuous Aβ42 or tau values, we conducted two identical models to those above (excluding biomarker

group terms). The first included predictors of Aβ42 (centered), p-tau (centered), age x Aβ42, age x p-tau, Aβ42 x p-tau, and age x Aβ42 x p-tau. The second model included effects of p-tau/Aβ42 and age x p-tau/Aβ42. Since t-tau was highly correlated with p-tau (r = .85, p < .001), we only included p-tau in these models.

RESULTS

Biomarker cutoffs

Table 1 details sample characteristics. All biomarker cutoffs had a minimum sensitivity and specificity of 70% and 90%, respectively (Supplemental Table 2). The ratios of tau to Aβ42 exhibited sensitivities and specificities ≥ 90% and greater area under the curve (AUC) values than Aβ42 (p < .05), Aβ42/Aβ40 (p < .05), and p-tau (p < .01).

Characteristics of biomarker groups

Of the 314 cognitively healthy participants, 53 (17%) had a positive tau biomarker (either p-tau ≥ 59.5 (n=40; 13%) or t-tau ≥ 461.26 (n=42; 13%)) and 76 (24%) had a positive amyloid biomarker (either Aβ42(ln) ≤ 6.156 [back-transformed value = 471.54] (n=44; 14%) or Aβ42/Aβ40 ≤ 0.09 (n=67; 21%)). Each participant was classified into one of four groups: Stage 0 = biomarker negative, Stage 1 = Aβ+ and tau-, Stage 2 = Aβ+ and tau+, or Tau+ = tau+ and Aβ-.

The majority of participants were negative for both Aβ and tau (Stage 0 = 68.5%). 14.6% were positive for Aβ only (Stage 1), 8% were positive for tau only, and 8.9% were positive for both Aβ and tau (Stage 2). Stages 0 and 1 did not differ on mean t-tau (p =

.10) or p-tau (p = .41). Stage 0 had lower Aβ42 than the tau+ group (p < .001), but did not differ on the Aβ42/Aβ40 ratio (p = .97). The Stage 2 group was the oldest and the Stage 0 group was the youngest (p < .001). The Stage 1 and 2 groups included greater proportions of *APOE* ε4 carriers (63-64%) compared with the Stage 0 or tau+ groups (28-38%). There were no differences between biomarker groups in sex, years of education, family history of AD, or source cohort (Table 2).

Cognitive trajectories across biomarker groups

At the visit closest to the LP, there were no significant differences in cognitive performance or HCV across biomarker groups (see Table 2), with the exception of processing speed (Digit Symbol).

Longitudinal neuropsychological performance for each biomarker group is displayed in Figure 1. Results from likelihood ratio tests ($\chi^2(3)$) indicated that age^2 x biomarker group accounted for a significant amount of variation in change on RAVLT Delay (χ^2 = 9.74, p = .02) and similar but non-significant variation in change on RAVLT Total (χ^2 = 7.11, p = .07) and TMT-B (χ^2 = 6.89, p = .08). Compared to the Stage 0 group, both Stage 1 and 2 groups showed more rapid, non-linear decline with age on the RAVLT Delay (*p's* < .05), whereas the Stage 2 group only showed more rapid, non-linear decline on the RAVLT Total (p = .02). Compared to the Stage 0 group, the Stage 1 group showed more rapid, non-linear change with age on TMT-B (p = .02). In contrast, the tau+ group did not significantly differ from the Stage 0 group. Age^2 x biomarker group was non-

significant for the remaining outcomes ($p's > .43$). Model parameters are displayed in Table 3.

For the remaining outcomes (in which the quadratic term was not associated with cognitive performance), results from likelihood ratio tests ($\chi^2(3)$) indicated that the interaction between age x biomarker group accounted for a significant amount of variation in change on LM Immediate ($\chi^2 = 11.74$, $p < .01$), LM Delay ($\chi^2 = 12.77$, $p < .01$), and Digit Symbol ($\chi^2 = 13.21$, $p < .01$). For all three outcomes, Stage 1 and Stage 2 exhibited greater age-related decline than Stage 0 ($p < .05$). In contrast, the tau+ group did not differ from the Stage 0 group in rates of cognitive change. Age-related change in HCV did not differ by biomarker group. Sensitivity analyses conducted on WRAP and ADRC cohorts separately revealed similar directions of effects, but slight heterogeneity in magnitude of beta-weights possibly due to baseline age differences across cohorts (see Supplemental Material).

Cognitive trajectories and continuous CSF values
There were no significant interactions between Aβ42 x p-tau x age^2; this term was removed from subsequent analyses. The three-way interaction between Aβ42 x p-tau x age was statistically significant for LM Delay ($B = .01$, $p = .03$), in which the relationship between p-tau and longitudinal story memory performance was dependent on Aβ42. Similar to results above, two-way interactions between age^2 x Aβ42 were significant for RAVLT Delay (age^2: $B = -.004$, $p < .01$; age^2 x Aβ42: $B = 0.01$, $p=.03$), RAVLT Total (age^2: $B = -.01$, $p = .05$; age^2 x Aβ42: $B = 0.03$, $p < .01$), and TMT-B (age^2: $B = .0002$, p

< .001; age^2 x Aβ42: B = -.0004, p = .03), indicating that lower CSF Aβ42 (higher brain amyloid) was associated with greater non-linear decline. For outcomes for which age^2 x Aβ42 was non-significant, greater amyloid burden was associated with greater linear decline (significant age x Aβ42 interaction) for LM Immediate (B = .22, p = .001), LM Delay (B = 0.22, p < .01), Digit Symbol (B = .48, p < .01), and Animal Fluency (B = .34, p < .01). In contrast, there were no interactions between age (linear or quadratic) x p-tau (Figure 2).

Age^2 x p-tau/Aβ42 was significant for TMT-B (B = .004, p < .01) and marginal for RAVLT Delay (B = -0.1, p = .08). Age x p-tau/Aβ42 was significant for all other outcomes with the exception of HCV, indicating that elevated AD biomarkers were associated with greater decline on RAVLT Total (B = -3.6, p < .01), LM Immediate (B = -2.3, p < .001), LM Delay (B = -2.3, p < .001), Digit Symbol (B = -3.0, p = .02), and Animal Fluency (B = -2.6, p < .01).

DISCUSSION

In this sample of over 300 non-demented middle-aged and older adults enriched for AD risk due to family history of AD, approximately one-third were positive for CSF biomarkers of AD (Aβ or tau). Those with Aβ positivity (with or without tau positivity), about 24% of the sample, exhibited significantly greater decline on neuropsychological measures than biomarker negative adults, whereas those with only tau positivity did not differ from biomarker negatives.

Growing evidence suggests that elevated Aβ is associated with cognitive decline in asymptomatic adults[4, 5, 20], but the contribution of tau to these findings is unclear. Studies incorporating both Aβ and tau observed Aβ and tau positive adults showed greater decline on a cognitive composite score[1] or global cognitive screening measure[21]. Our study extends these findings by demonstrating that adults with elevated Aβ only also show greater decline on specific measures of memory and executive functioning, and that this decline is more rapid at older ages. Although composites representing multiple domains reduce multiple comparisons, they may also mask signal from individual cognitive domains (e.g., memory). Future studies investigating additional factors that may differ between Aβ+ and Aβ- asymptomatic adults, such as vascular risk factors, are needed to determine if other factors cause or exacerbate decline in Aβ+ asymptomatic adults.

Novel results from this study demonstrate that cognitive change associated with Aβ varies by age and cognitive measure. While performance on story memory and processing speed declined linearly with age and Aβ burden, non-linear decline on list-learning and set-shifting tasks suggest faster rates of decline with advancing age in the presence of amyloid burden. These results have potentially important implications for choosing appropriate outcome measures in clinical trials. For example, if a trial is enrolling older adults, it may be more optimal to choose a list-learning memory measure outcome since it would be expected to decline more rapidly in older adults with AD pathology. However, processing speed measures may be most sensitive to detection of early decline associated with Aβ, as this was the only measure on which the Aβ+

groups performed more poorly at the LP visit and has been shown to be predictive of

progression to cognitive impairment in a similar cohort [22].

In the context of the recently proposed amyloid/tau/neurodegeneration (A/T/N)

biomarker classification system[23], our findings suggest that those characterized as

A+/T- are at similar risk of decline to those characterized as A+/T+. However, we did not

fully examine neurodegeneration as markers of both total and phosphorylated tau were

incorporated into the tau positivity classification. In this sample, we did not observe

cross-sectional or longitudinal differences among biomarker groups in hippocampal

volume, unlike a prior study showing that individuals characterized as Stage 2 had

accelerated hippocampal atrophy[24]. It is possible these differences are due to the

younger age of our cohort, which may not be expected to show structural brain changes

at this stage. As hippocampal volume may be less specific to AD-related changes (e.g.,

could reflect hippocampal sclerosis), future studies will incorporate additional structural

imaging markers that may be more sensitive to early neurodegeneration and more

specific to AD (e.g., cortical thickness patterns) than hippocampal volume alone.

Furthermore, results from the majority of models including continuous CSF markers

were similar to those using a group variable based on cutoffs (e.g., lower CSF Aβ42

was associated with worsening performance, whereas elevated tau was not),

suggesting that dichotomizing continuous biomarker variables did not result in

significant loss of information and that Aβ exerts an independent effect from tau on

cognition. Although A/T/N is a biomarker classification system, measuring cognitive

trajectories is essential for validating these criteria and providing prognostic information needed as biomarkers become more readily available in clinical settings.

An important limitation of the current study was inclusion of only CSF biomarkers. Although this approach is consistent with prior studies, future analyses will incorporate both CSF and neuroimaging biomarkers to provide greater reliability in classification of preclinical AD. Our sample contained a smaller proportion of adults with markers of only tau+ (8%) compared to other studies (11-23%), perhaps due to the younger mean age of our cohort, the method by which we defined the cutoffs, or the relatively small sample from which the cutoffs were derived. These results are based on longitudinal cohorts that include a majority of Caucasian, highly educated adults from the Midwest region of the United States and may be less generalizable to other populations. Despite these limitations, our results suggest that in an asymptomatic cohort at increased risk for AD, elevated Aβ is associated with significantly greater rates of decline, with the specific patterns of decline varying by age and cognitive measure examined.

ACKNOWLEDGMENTS

We gratefully acknowledge the assistance of researchers and staff at the Wisconsin Registry for Alzheimer's Prevention and Wisconsin Alzheimer's Disease Research Center for assistance in recruitment and data collection. Most importantly, we thank the dedicated WRAP and WADRC participants for their continued support and participation in this research.

TABLES

Table 1. Sample characteristics

Variable	ROC sample	Cognitively Healthy sample
N (Total = 392)	78[a]	314
Age at visit 1 (mean, range)	64.5 (47-92)	58.8 (37-85)
Age at lumbar puncture (LP) visit (mean, range)	65.4 (48-93)	61.5 (43-86)
Months between visit 1 and LP (mean, range)	10.9 (0-91)	33.5 (0-134)
Female (*n*, %)	42 (54%)	218 (69%)
Education (mean, SD, range)	15.3 (2.6; 8-20)	16.3 (2.5; 8-25)
APOE ε4+ (*n*, %)	27 (35%)	135 (43%)
Years in study (mean, SD, range)	2.8 (2.6; 0-11)	5.8 (3.5; 1-13)
Natural-log beta-amyloid ($A\beta_{42}$) (mean, SD, range)	6.3 (0.4; 5.3-7.2)	6.5 (0.3; 5.6-7.5)
Total tau (t-tau) (mean, SD, range)	528.1 (366.0; 67.2-1633.0)	324.7 (153.3; 67.2-1085.0)
Phosphorylated tau (p-tau) (mean, SD, range)	58.2 (29.6; 17.1-152.0)	43.5 (15.5; 12-114)
$A\beta_{42}/A\beta_{40}$ (mean, SD, range)	0.08 (0.03; 0.04-0.13)	0.1 (0.02; 0.04-0.2)
t-tau/$A\beta_{42}$ (mean, SD, range)	1.2 (1.1; 0.1-4.5)	0.5 (0.4; 0.1-3.5)
p-tau/$A\beta_{42}$ (mean, SD, range)	0.1 (0.1; 0.03-0.5)	0.1 (0.04; 0.02-0.3)
Diabetes	6 (8%)	16 (5%)
Hypertension	28 (36%)	61 (19%)
Hypercholesterolemia	34 (44%)	122 (39%)
History of stroke or TIA	3 (4%)	0 (0%)
Prescribed cognitive-enhancing medication[b]	37 (47%)	0 (0%)

ROC=Receiver Operator Characteristic Curve; AD=Alzheimer's disease; APOE=Apolipoprotein
[a]*n*=40 cognitively healthy controls (51%) and *n*=38 participants with clinical diagnoses of dementia due to Alzheimer's disease (49%); [b]Donepezil, Memantine, Galantamine, or Rivastigmine

Table 2. Biomarker group characteristics (*n*=314) (mean [SD] or n (%))

	Biomarker Negative (*n*=215)	Amyloid + Tau – (*n*=46)	Tau + Amyloid – (*n*=25)	Amyloid + Tau + (*n*=28)	*p*-value	*Effect Size (f² or φc)*
NIA-AA, 2011	Stage 0	Stage 1		Stage 2		
IWG-AA, 2016	N/A	Asymptomatic at-risk	Asymptomatic at-risk	Preclinical AD		
Age (study visit 1)	57.7 (8.2)	59.3 (6.9)	59.6 (10.7)	65.9 (9.0)	<.001	.08
Age (lumbar puncture)	60.2 (7.6)	62.5 (6.9)	62.3 (9.5)	68.7 (7.0)	<.001	.10
Education (years)	16.2 (2.5)	16.6 (2.8)	16.2 (2.4)	16.5 (2.7)	.84	.003
Sex (Women)	149 (69%)	33 (72%)	17 (68%)	19 (68%)	.98	.02
APOE ε4 carriers	81 (38%)	29 (63%)	7 (28%)	18 (64%)	.001	.24
AD family history +	182 (85%)	35 (76%)	17 (68%)	23 (82%)	.15	.13
Total Tau	271.6 (82.6)	295.0 (111.1)	499.6 (164.4)	625.3 (167.1)	<.001	1.2
Phosphorylated Tau	38.4 (10.2)	39.4 (12.3)	65.2 (7.2)	69.8 (16.6)	<.001	.99
Aβ42 (ln)	6.6 (0.2)	6.2 (0.2)	6.9 (0.2)	6.2 (0.2)	<.001	1.2
Aβ42/Aβ40	.11 (.01)	.08 (.01)	.11 (.01)	.06 (.01)	<.001	1.8
Characteristics at biomarker visit (estimated marginal means and standard errors)						
RAVLT Total Trials 1-5	52.5 (0.6)	52.9 (1.1)	52.0 (1.5)	51.9 (1.5)	.94	.001
RAVLT Delayed Recall	10.8 (0.2)	10.1 (0.4)	10.4 (0.5)	10.4 (0.5)	.37	.01
WMS-R LM Immediate	15.0 (0.3)	15.0 (0.5)	15.2 (0.7)	15.0 (0.7)	.99	.001
WMS LM Delay	13.7 (0.3)	13.9 (0.5)	14.0 (0.7)	14.4 (0.7)	.78	.004
Trailmaking Test Part B	59.0 (1.8)	63.5 (3.4)	61.1 (4.6)	58.7 (4.5)	.66	.01
Digit Symbol	59.2 (0.7)	55.0 (1.4)	57.4 (1.9)	56.9 (1.8)	.03	.03
Animal Fluency	24.2 (0.5)	23.9 (0.8)	23.3 (1.2)	22.4 (1.2)	.51	.01
Hippocampal volume (n=202)	7858.9 (90.2)	7793.1 (148.3)	8069.7 (210.8)	7674.4 (184.8)	.53	.01

APOE=Apolipoprotein; AD=Alzheimer's disease; Aβ = Beta-amyloid; ln=natural log; RAVLT=Rey Auditory Verbal Learning Test; WMS-R LM=Wechsler Memory Scale-Revised Logical Memory Story A subtest; f^2 = Cohen's f2; $φ_c$ = Cramer's V

Table 3. Parameter estimates from linear mixed-effects models

Fixed Effects	RAVLT TOTAL TRIALS 1-5	RAVLT DELAYED RECALL	LM IMMEDIATE RECALL	LM DELAYED RECALL	DIGIT SYMBOL	TRAILS B (lg10)
	B (SE)	B (SE)	B (SE)	B (SE)	B (SE)	B (SE)
Intercept	45.0 (2.5)***	8.8 (0.9)***	9.4 (1.0)***	8.3 (1.1)***	50.2 (3.2)***	2.8 (0.0)***
Biomarker Group						
Aβ-/Tau-	--	--	--	--	--	--
Aβ+	-0.7 (1.2)	-0.3 (0.4)	-0.8 (0.4)	-0.9 (0.5)	-3.2 (1.4)*	0.001 (0.0)
Tau+	-1.3 (1.6)	-0.3 (0.5)	0.3 (0.6)	0.2 (0.6)	-2.5 (1.8)	-0.01 (0.0)
Aβ+/Tau+	-0.3 (1.6)	-0.02 (0.6)	-0.1 (0.7)	0.07 (0.7)	-1.3 (2.0)	0.02 (0.0)
Age each visit (center)	-0.3 (0.1)***	-0.1 (0.0)***	-0.004 (0.0)	-0.02 (0.0)	-0.7 (0.1)***	0.01 (0.0)***
Age each visit (center)^2	-0.002 (0.0)	-0.002 (0.0)	--	--	--	0.0001 (0.0)
Sex (male)	-6.8 (0.8)***	-1.8 (0.3)***	-2.1 (0.3)***	-2.1 (0.4)***	-3.4 (1.1)**	0.03 (0.0)*
Education (years)	0.4 (0.2)**	0.1 (0.1)*	0.4 (0.1)***	0.3 (0.1)***	0.5 (0.2)*	-0.004 (0.0)
Practice Effect	1.2 (0.1)***	0.2 (0.0)***	0.2 (0.1)**	0.3 (0.1)***	0.8 (0.2)***	-0.01 (0.0)***
Age each visit x Group						
Age x Aβ-/Tau-	--	--	--	--	--	--
Age x Aβ+	-0.2 (0.1)*	-0.1 (0.0)*	-0.1 (0.1)*	-0.2 (0.1)**	-0.4 (0.1)*	-.0003 (0.0)
Age x Tau+	0.1 (0.1)	-0.05 (0.0)	-0.1 (0.1)	0.02 (0.1)	0.1 (0.2)	.0004 (0.0)
Age x Aβ +/Tau+	-0.2 (0.2)	-0.05 (0.1)	-0.2 (0.1)**	-0.2 (0.1)**	-0.4 (0.2)*	-.0005 (0.0)
Age each visit^2 x Group						
Age^2 x Aβ-/Tau-	--	--	--	--	--	--
Age^2 x Aβ+	-0.01 (0.0)	-0.01 (0.0)*	--	--	--	.0004 (0.0)*
Age^2 x Tau+	0.01 (0.0)	-0.001 (0.0)	--	--	--	.0002 (0.0)
Age^2 x Aβ +/Tau+	-0.03 (0.0)*	-0.01 (0.0)*	--	--	--	.0002 (0.0)

RAVLT=Rey Auditory Verbal Learning Test; WMS-R LM=Wechsler Memory Scale-Revised Logical Memory Story A subtest; ***p≤.001; **p≤.01; *p<.05

Note: Quadratic terms were non-significant for LM and Digit Symbol measures. Final models with quadratic terms removed are reported here.

Figure Titles and Legends

Figure 1 Title. Biomarker groups and cognitive trajectories

Figure 1 Legend. Graphs depict neuropsychological performance on the y-axis for six cognitive measures and age at each visit (centered on mean age) on the x-axis. Each line depicts the estimated slope for the four biomarker groups, adjusting for covariates of sex, education, and practice effects. Higher scores equate better performance on all measures except TMT-B (higher scores = worse performance). Quadratic terms were retained for the RAVLT and TMT-B. Non-significant quadratic terms were removed for other outcomes and linear effects are depicted. Both the Aβ+ only group (orange) and the Aβ+/Tau+ group (green) exhibited significantly greater decline than the biomarker negative group (black). In contrast, the group with only tau+ (blue) did not differ from biomarker negative individuals.

Figure 2 Title. Relationships between Aβ42, tau, and longitudinal verbal memory performance.

Figure 2 Legend. Two-way interaction between age at each visit and Aβ42 (top) or ptau (bottom) on memory performance. Figures depict that although performance generally decreases with age, those with low Aβ42 (high brain amyloid) exhibit most rapid decline, whereas the association between age at each visit and memory performance does not vary by ptau. Facets depict biomarker level by quartile (1 = lowest quartile (0-25%), 2= 25-50%, 3 = 50-75%, 4 = highest quartile (75-100%)).

REFERENCES

1.   Soldan A, Pettigrew C, Cai Q, et al. Hypothetical Preclinical Alzheimer Disease Groups and Longitudinal Cognitive Change. JAMA Neurol 2016;73:698-705.
2.   Vos SJB, Gordon BA, Su Y, et al. NIA-AA staging of preclinical Alzheimer disease: discordance and concordance of CSF and imaging biomarkers. Neurobiol Aging 2016;44:1-8.
3.   Dubois B, Hampel H, Feldman HH, et al. Preclinical Alzheimer's disease: Definition, natural history, and diagnostic criteria. Alzheimers Dement 2016;12:292-323.
4.   Clark LR, Racine AM, Koscik RL, et al. Beta-amyloid and cognitive decline in late middle age: Findings from the Wisconsin Registry for Alzheimer's Prevention study. Alzheimer's & Dementia 2016;12:805-814.
5.   Lim YY, Maruff P, Pietrzak RH, et al. Effect of amyloid on memory and non-memory decline from preclinical to clinical Alzheimer's disease. Brain 2014;137:221-231.
6.   Sager MA, Hermann B, La Rue A. Middle-Aged Children of Persons With Alzheimer's Disease: APOE Genotypes and Cognitive Function in the Wisconsin Registry for Alzheimer's Prevention. Journal of Geriatric Psychiatry and Neurology 2005;18:245-249.
7.   Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's & Dementia 2011;7:270-279.
8.   McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's & Dementia 2011;7:263-269.
9.   Backman L, Jones S, Berger AK, Laukka EJ, Small BJ. Cognitive impairment in preclinical Alzheimer's disease: a meta-analysis. Neuropsychology 2005;19:520-531.
10.   Duke Han S, Nguyen CP, Stricker NH, Nation DA. Detectable Neuropsychological Differences in Early Preclinical Alzheimer's Disease: A Meta-Analysis. Neuropsychology review 2017.
11.   Schmidt M. Rey Auditory Verbal Learning Test: A handbook. Los Angeles, CA: Western Psychological Services, 1996.
12.   Wechsler. Wechsler Memory Scale-Revised. New York: The Psychological Corporation, Harcourt Brace Jovanovich, inc. for Psychological Corp., 1987.
13.   Reitan RM, & Wolfson, D. The Halstead–Reitan Neuropsychological Test Battery: Therapy and clinical interpretation. Tucson, AZ: Neuropsychological Press, 1985.
14.   Wechsler. Wechsler Adult Intelligence Scale - Revised. San Antonio: The Psychological Corporation, Harcourt brace & co for The Psychological Corporation, 1981.
15.   R: A language and environment for statistical computing [computer program]. Vienna, Austria: R Foundation for Statistical Computing, 2016.
16.   Robin X, Turck N, Hainard A, et al. pROC: an open-source package for R and S+ to analyze and compare ROC curves. BMC bioinformatics 2011;12:77.
17.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. 2015 2015;67:48.
18.   Jonaitis EM, Koscik RL, La Rue A, Johnson SC, Hermann B, Sager MA. Aging, practice effects, and genetic risk in the Wisconsin Registry for Alzheimer's Prevention. The Clinical neuropsychologist 2015;29:426-441.
19.   Bates DM, & Pinheiro, J.C. Mixed-Effects Models in S and S-PLUS. New York, NY: Springer, 2000: 81-82.

20.	Donohue MC, Sperling RA, Petersen R, et al. Association between elevated brain amyloid and subsequent cognitive decline among cognitively normal persons. Jama 2017;317:2305-2316.
21.	Vos SJ, Xiong C, Visser PJ, et al. Preclinical Alzheimer's disease and its outcome: a longitudinal cohort study. The Lancet Neurology 2013;12:957-965.
22.	Albert M, Soldan A, Gottesman R, et al. Cognitive Changes Preceding Clinical Symptom Onset of Mild Cognitive Impairment and Relationship to ApoE Genotype. Current Alzheimer Research 2014;11:773-784.
23.	Jack CR, Jr., Bennett DA, Blennow K, et al. A/T/N: An unbiased descriptive classification scheme for Alzheimer disease biomarkers. Neurology 2016;87:539-547.
24.	Gordon BA, Blazey T, Su Y, et al. Longitudinal beta-Amyloid Deposition and Hippocampal Volume in Preclinical Alzheimer Disease and Suspected Non-Alzheimer Disease Pathophysiology. JAMA Neurol 2016;73:1192-1200.