Genetic dissection of an ancient divergence in yeast thermotolerance

Carly V. Weiss[1], Jeremy I. Roop[1,2]*, Rylee K. Hackley[1,3,4], Julie Chuong[3],

Igor V. Grigoriev[1,5], Adam P. Arkin[6], Jeffrey M. Skerker[6], Rachel B. Brem[1,3]*

[1]Department of Plant and Microbial Biology, UC Berkeley, Berkeley, CA.

[2]Current address: Fred Hutchinson Cancer Research Center, Seattle, WA.

[3]Buck Institute for Research on Aging, Novato, CA.

[4]Current address: Duke University Program in Genetics & Genomics, Duke University, Durham,

NC.

[5]US Department of Energy Joint Genome Institute, Walnut Creek, CA.

[6]Department of Bioengineering, UC Berkeley, Berkeley, CA and Lawrence Berkeley National

Laboratory, Berkeley, CA.

*Correspondence to: jeremyianroop@gmail.com, rbrem@buckinstitute.org.

Abstract

Some of the most unique and compelling survival strategies in the natural world are fixed in

isolated species. To date, molecular insight into these ancient adaptations has been limited, as

classic experimental genetics has focused on interfertile individuals in populations. Here we use

a new mapping approach, which screens mutants in a sterile interspecific hybrid, to identify

eight housekeeping genes that underlie the growth advantage of *Saccharomyces cerevisiae*

over its distant relative *S. paradoxus* at high temperature. Pro-thermotolerance alleles at these

mapped loci were required for the adaptive trait in *S. cerevisiae* and sufficient for its partial

reconstruction in *S. paradoxus*. The emerging picture is one in which *S. cerevisiae* improved the

heat resistance of multiple components of the fundamental growth machinery in response to

selective pressure. This study lays the groundwork for the mapping of genotype to phenotype in

clades of sister species across Eukarya.

Geneticists since Mendel have sought to understand how and why wild individuals differ.

Studies toward this end routinely test for a relationship between genotype and phenotype via

linkage or association[1]. These familiar approaches, though powerful in many contexts, have an

important drawback—they can only be applied to interfertile members of the same species. This

rules out any case in which an innovation in form or function evolved long ago and is now fixed

in a reproductively isolated population.

As organisms undergo selection over long timescales, their traits may be refined by processes

quite different from those that happen early in adaptation[2,3]. We know little about these

mechanisms in the wild, expressly because when the resulting lineages become reproductively

incompatible, classic statistical-genetic methods cannot be used to analyze them[4]. To date, the

field has advanced largely on the strength of candidate-based studies that implicate a single

variant gene in an interspecific trait[5,6], with the complete genetic architecture often remaining

unknown. Against the backdrop of a few specialized introgression[7-10] and molecular-evolution[11]

techniques available in the field, dissection of complex trait differences between species has

remained a key challenge.

Here we develop a new genetic mapping strategy, based on the reciprocal hemizygosity

test[12,13], and use it to identify the determinants of a difference in high-temperature growth

between isolated *Saccharomyces* yeast species. We validate the contributions of the mapped

loci to the thermotolerance trait, and we investigate their evolutionary history.

Results

Species differences in thermotolerance

At high temperature, the yeast *Saccharomyces cerevisiae* grows qualitatively better than other members of its clade[14-16], including its closest relative, *S. paradoxus*, from which it diverged ~5 million years ago[17]. In culture at 39°C, *S. cerevisiae* doubled faster than *S. paradoxus* and accumulated more biomass over a timecourse, a compound trait that we call thermotolerance. The magnitude of differences in thermotolerance between species far exceeded that of strain variation within each species (Figure 1), whereas no such effect was detectable at 28°C (Figure S1). The failure by *S. paradoxus* to grow to high density at 39°C was, at least in part, a product of reduced survival relative to that of *S. cerevisiae,* as cells of the former were largely unable to form colonies after heat treatment (Figure S2). In microscopy experiments, *S. paradoxus* cells were almost uniformly visible as large-budded dyads after 24 hours at 39°C (Figure S3), suggestive of defects late in the cell cycle as a proximal cause of death[18]; no such tendency was apparent in *S. cerevisiae* at high temperature, or in either species at 28°C (Figure S3).

Massively parallel reciprocal hemizygosity testing by RH-seq

We set out to dissect the genetic basis of *S. cerevisiae* thermotolerance, using a genomic implementation of the reciprocal hemizygote test[12,13] (Figure 2A). For this purpose, we first mated *S. cerevisiae* strain DBVPG1373, a soil isolate from the Netherlands, with *S. paradoxus* strain Z1, an English oak tree isolate. The resulting sterile hybrid had a thermotolerance phenotype between those of its purebred parents (Figure S4). In this hybrid background we generated hemizygote mutants using a plasmid-borne, selectable PiggyBac transposon system[19]. We cultured the pool of mutants in bulk for ~7 generations at 39°C and, separately, at 28°C. From cells in each culture we sequenced transposon insertion locations[20] as a readout of the genotypes and abundance of mutant hemizygote clones present in the selected sample. In these sequencing data, at each of 4888 genes we detected transposon mutant clones in both

species' alleles in the hybrid (Figure S5), with transposon insertions distributed in a largely

unbiased manner across the genome (Figure S6). For a given gene, we tabulated the

abundances of mutants whose transposon insertion fell in the *S. cerevisiae* allele of the hybrid,

after high-temperature selection relative to the 28°C control, and we compared them to the

abundance distribution of mutants in the *S. paradoxus* allele (Figure 2A). Any difference in

abundance between these reciprocal hemizygote cohorts can be ascribed to variants between

the retained alleles at the respective locus; we refer to the comparison as reciprocal

hemizygosity analysis via sequencing (RH-seq). Integrating this approach with a quality-control

pipeline (Figure S5), in a survey of 3416 high-coverage genes we identified 8 top-scoring hits

(false discovery rate 0.01; Figure 2B). At each such locus, disruption of the *S. cerevisiae* allele

in the hybrid was associated with low clone abundance after selection at 39°C relative to 28°C

(Figure 2B), reflecting a requirement for the *S. cerevisiae* allele for thermotolerance. None of the

genes mapped by RH-seq had a known role in the yeast heat shock or stress responses. All

were annotated as housekeeping factors: *ESP1, DYN1, MYO1, CEP3, APC1,* and *SCC2*

function in chromosome segregation and cytokinesis, and *AFG2* and *TAF2* in

transcription/translation.

To evaluate the role in thermotolerance of genes that emerged from RH-seq, we first sought to

verify that growth differences between hemizygotes at a given locus were the consequence of

allelic variation and not an artifact of our genomic approach. As a complement to our analysis of

transposon mutants, for each RH-seq hit gene we engineered hemizygotes by targeted deletion

of each species' allele in turn in the hybrid. In growth assays, the strain lacking the *S. cerevisiae*

allele at each gene grew poorly at high temperature (Figure 2B), with little impact at 28°C

(Figure S7), consistent with fitness inferences from RH-seq. And in each case, the *S. paradoxus*

allele made no contribution to the phenotype of the hybrid, since deleting it had no effect

(Figures 2B and S7). We conclude that the loci emerging from RH-seq represent bona fide

determinants of thermotolerance in the hybrid.

Validation of RH-seq gene hits in purebreds

We expected that variation at our RH-seq hits, though mapped by virtue of their impact in the

hybrid, could also explain thermotolerance differences between purebred species. As a test of

this notion, for each mapped gene in turn, we replaced the two copies of the endogenous allele

in each purebred diploid with the allele from the other species. Growth assays of these

transgenics established the *S. cerevisiae* allele of each locus as necessary or sufficient for

biomass accumulation at 39°C, or both: thermotolerance in the *S. cerevisiae* background was

compromised by *S. paradoxus* alleles at 7 of the 8 genes and, in *S. paradoxus*, improved by *S.*

cerevisiae alleles at 6 of 8 loci (Figure 3). Allele replacements had little impact on growth at

28°C (Figure S8). These trends mirrored the direction of locus effects from hemizygotes in the

hybrid, though the magnitudes were often different. Most salient were the small effect sizes in *S.*

paradoxus relative to other backgrounds, indicative of strong epistasis in this poorly-performing

species (Figure S9). Thus, the loci mapped by RH-seq in an interspecies hybrid contribute

causally to thermotolerance in purebreds, with effect sizes that depend on the context in which

they are interrogated.

Avid growth at high temperature is a defining characteristic of *S. cerevisiae* as a species,

relative to other Saccharomycetes (refs. 14-16 and Figure 1). In principle, the loci mapped by

RH-seq could be unique to the genetic architecture of thermotolerance in our focal *S. cerevisiae*

strain, DBVPG1373, or be part of a mechanism common to many *S. cerevisiae* isolates. In

support of the latter model, transgenesis experiments showed that a diverse panel of *S.*

cerevisiae isolates all harbored alleles conferring modest but significant growth benefits at high

temperature, and alleles from multiple *S. paradoxus* isolates were deleterious (Figure S10A-B).

We detected no such impact at 28°C (Figure S10A-B). Similarly, we found elevated sequence

divergence from *S. paradoxus* to be a shared feature of *S. cerevisiae* strains at the loci mapped

by RH-seq (using the absolute divergence measure D_{xy}; Figure S10C). These findings indicate

that the *S. cerevisiae* population accumulated divergent, pro-thermotolerance alleles at the loci

mapped by RH-seq, consistent with a role in the trait for these genes across the species.

Likewise, in the yeast phylogeny, RH-seq hit genes were distinguished by accelerated evolution

along the branch leading to *S. cerevisiae,* as expected if the ancestral program has been

conserved among the other species in the clade (Figure S10C).

Discussion

In this work, we have developed the RH-seq method for genome-wide mapping of natural trait

variation, and we have used it to elucidate the genetics of thermotolerance in reproductively

isolated yeasts. Growth at high temperature is likely a derived character in *S. cerevisiae*[14-16],

and the mechanism by which evolution built the trait, after the split from *S. paradoxus*, has

remained unknown. In pursuing the genetics of this putative ancient adaptation, we complement

studies of younger, intra-specific variants that erode thermotolerance, in the few *S. cerevisiae*

strains that have lost the trait relatively recently[12,21]. We have sought to shed light on more

ancient evolutionary events by considering *S. paradoxus* as a representative of the ancestral

state, to which thermotolerant *S. cerevisiae* can be compared.

Using this approach, we have mapped eight loci at which *S. cerevisiae* alleles are necessary

and sufficient for thermotolerance. Six of these genes are essential for growth in standard

conditions[22], and all eight contribute to fundamental growth processes. *ESP1, DYN1, CEP3,*

APC1, MYO1, and *SCC2* mediate mitotic spindle assembly, chromatid cohesion and separation,

cytokinesis, and mitotic exit; *AFG2* regulates the release of maturation factors from the

ribosome; and *TAF2* encodes a TFIID subunit. In each case, our growth experiments in the

interspecific hybrid have shown that the *S. paradoxus* allele acts as a hypomorph at high

temperature. Our work leaves unanswered exactly how heat-treated *S. paradoxus* cells die in

the absence of these functions, though their large-budded morphology strongly suggests

regulated arrest or stochastic failure late in the cell cycle. Given that some but not all RH-seq hit

loci have roles in mitosis, it is likely only one of the choke points at which *S. paradoxus* alleles

are a liability at high temperature. Assuming that these heat-sensitive alleles also littered the

genome of the common ancestor with *S. cerevisiae*, thermotolerance would have evolved along

the *S. cerevisiae* lineage by resolving each of them, boosting the heat resistance of many

housekeeping processes. Such a mechanism would dovetail with the recent finding that, across

species, the limiting temperature for cell growth correlates with the melting temperatures of a

subset of essential proteins[23].

These insights into the evolution of a complex yeast trait serve as a proof of concept for RH-

seq. To date, the reciprocal hemizygosity test has led to landmark discoveries in a candidate-

gene framework, confirming the effects of variation at a given locus identified by other

means[12,13]. Schemes to scale up the test have generated a genome's worth of hemizygotes

from deletion-strain purebreds, which tend to harbor secondary mutations that come through

screens as false positives[24,25]. As such, a key advantage of RH-seq is that we carry out

mutagenesis in the hybrid, which ensures coverage of essential genes and obviates the use of

mutation-prone null genotypes. Furthermore, any secondary mutations that do arise in a given

hemizygote clone should have little influence in RH-seq mapping, because deep mutagenesis

generates many independent clones per gene that are analyzed together. One important caveat

of RH-seq, as in single-gene reciprocal hemizygote tests, is the assumption that no epistasis

unique to the hybrid will mask the effects of loci underlying a trait difference of interest between

the parents. In our case study, the genetic architecture of thermotolerance in the hybrid did bear

out as relevant for the purebreds, albeit with locus effect sizes that varied across the

backgrounds. More dramatic discrepancies may be particularly likely when the hybrid has a

heterotic (*i.e.* extreme) phenotype and is a poor model for the genetics of the parents[26]. The

choice of a non-heterotic hybrid in which to pursue RH-seq would be analogous to classical

linkage mapping in a cross whose progeny have, on average, phenotypes that are intermediate

between those of the parents.

In fact, although we have focused here on ancient divergence, the RH-seq method would be

just as applicable to individuals within a species, as a high-resolution alternative to linkage

analysis. We thus anticipate that RH-seq will accelerate the mapping of genotype to phenotype

in many systems, whether the parents of a cross are closely related or members of a species

complex that have been locally adapting for millions of years.

Figure 1. *S. cerevisiae* grows better at high temperature than *S. paradoxus.*

A, Each point reports cell density (OD_{600}) of the indicated wild isolate of *S. cerevisiae* (blue) or *S. paradoxus* (orange) over a timecourse of growth at 39°C. Each solid line shows a logistic population growth model fit to the respective cell density measurements. **B**, Each bar reports mean efficiency (*n* = 4) of the indicated strain at 39°C, defined as the difference between cell density at 24 hours of growth and that at the time of inoculation. *, $p = 3.78 \times 10^{-11}$; individual measurements are reported as open circles.

Figure 2. Mapping thermotolerance by RH-seq.

A, A transposon (rectangle) disrupts the allele from *S. cerevisiae* (blue) or *S. paradoxus*

(orange) of a gene (*YFG*) in an interspecific hybrid (green). Clones lacking the pro-

thermotolerance allele grow poorly at 39°C (dashed outlines), as measured by sequencing and

reported in smoothed histograms (traces, colored to indicate the species' allele that is not

disrupted). **B**, Each panel reports results from one RH-seq hit locus. In the main figure of a

given panel, the *x*-axis reports the $\log_2$ of abundance, measured by RH-seq after selection at

39°C, of a clone harboring a transposon insertion in the indicated species' allele, relative to the

analogous quantity for that clone from selection at 28°C. The *y*-axis reports the proportion of all

clones bearing insertions in the indicated allele that exhibited the abundance ratio on the *x*, as a

kernel density estimate. In insets, each bar reports the relative efficiency, calculated as the

mean growth efficiency at 39°C (n = 8-12) of the indicated targeted-deletion hemizygote

measured in liquid culture assays, normalized to the analogous quantity for the wild-type hybrid.

*, $p \leq 0.002$; individual measurements are reported as open circles.

Figure 3. *S. cerevisiae* thermotolerance alleles are necessary and sufficient for growth at

high temperature.

A, Each bar reports mean growth efficiency at 39°C, measured in liquid culture assays (*n* = 6-

12), of an *S. cerevisiae* strain harboring the *S. paradoxus* allele at the indicated RH-seq hit

locus, relative to the analogous quantity for wild-type *S. cerevisiae*. **B**, Data are as in **A**, except

that each bar reports results from a *S. paradoxus* strain harboring the *S. cerevisiae* allele at the

indicated locus, normalized to wild-type *S. paradoxus*. In a given panel, the top and bottom

dotted lines report the relative efficiency of wild-type *S. cerevisiae* and *S. paradoxus*,

respectively. *, $p \leq 0.036$; **, $p \leq 0.001$. Individual measurements are reported as open circles.

Materials and Methods

Strains

Strains used in this study are listed in Table S1. Homozygous diploid strains of *S. cerevisiae*

and *S. paradoxus* used as parents of the interspecific hybrid, and as the backgrounds for allele-

swap experiments, were homothallic DBVPG1373 and Z1, respectively. In the case of the

hybrid parents, each strain was rendered homozgyous null for *URA3* via homologous

recombination with a HYGMX cassette, then sporulated; a given mated spore from a dissected

tetrad was grown up into a diploid that was homozygous null at *URA3* and tested for the

presence of both genomes by PCR with species-specific primers.

piggyBac transposon machinery

For untargeted, genome-scale construction of reciprocal hemizygotes in the *S. cerevisiae* x *S.*

paradoxus hybrid, we adapted methods for piggyBac transposon mutagenesis[19] to develop a

system in which the transposon machinery was borne on a selectable and counter-selectable

plasmid lacking a centromere. We constructed this plasmid (final identifier pJR487) in three

steps. In step I we cloned the piggyBac transposase enzyme gene driven by the *S. cerevisiae*

TDH3 promoter (from plasmid p3E1.2, a gift from Malcolm Fraser, Notre Dame) into plasmid

pJED104 (which contains *URA3*, an ARS, and the *CEN6* locus, and was a gift from John

Dueber, UC Berkeley). For this cloning, the amplification used a forward and reverse primer

containing a BamHI and XhoI site, respectively, that upon restriction digest yielded sticky ends

for ligation to recipient BamHI and XhoI sites in digested pJED104. We used the resulting

plasmid as input into step II, removal of the *CEN6* sequence: we first amplified the entire

plasmid with primers that initiated outside of *CEN6* and were directed away from it, and

contained reciprocally complementary NheI sites; sticky ends of the linear PCR product were

then ligated together for re-circularization. We used the resulting plasmid as input into step III,

the cloning in of a construct comprised of the KANMX cassette flanked by long terminal arms

(328bp and 361bp) from the piggyBac transposon. We first amplified KANMX from pUG6[27] and

each transposon arm from p3E1.2, using primers that contained overlapping sequence on the

fragment ends that would ultimately be the interior of the construct, and XbaI sites on the

fragment ends that would ultimately be the 5' and 3'-most ends of the construct. We stitched the

three fragments together by overlap extension PCR, digested the resulting construct and the

plasmid from step II with XbaI, and annealed sticky ends of the two to yield the final pJR487

plasmid.

Untargeted hemizygote construction via transposon mutagenesis

For mutagenesis, pJR487 was gigaprepped using a column kit (Zymo Research) to generate

~11 mg plasmid. To prepare for transformation, JR507 (the *S. cerevisiae* DBVPG1373 x *S.*

paradoxus Z1 hybrid) was streaked from a -80°C freezer stock onto a yeast peptone dextrose

(YPD, 1% yeast extract [BD], 2% yeast peptone [BD], 2% D-glucose [Sigma]) agar plate and

incubated for 2 days at 26°C. A single colony was inoculated into 100 mL YPD and shaken at

28°C, 200rpm for ~24 hours. The next day, we transferred cells from this pre-culture, and YPD,

to each of four 1 L flasks at the volumes required to attain an optical density at 600 nm (OD_{600})

of 0.2 in 500 mL each. We cultured each for 6 hours at 28°C with shaking at 200rpm. Two of

these cultures were combined into 1 L of culture and two into a separate 1 L, and each such

culture was subjected to transformation (for a total of two transformations) as follows. The 1 L

was split into twenty 50-mL conical tubes. Each aliquot was centrifuged and washed with water

and then with 0.1 M lithium acetate (LiOAc, Sigma) mixed with 1X Tris-EDTA buffer (10 mM

Tris-HCl and 1.0 mM EDTA); after spin-down, to each tube was added a solution of 0.269 mg of

pJR487 mixed 5:1 by volume with salmon sperm DNA (Invitrogen), and then to each was added

3 mL of 39.52% polyethylene glycol, 0.12M LiOAc and 1.2X Tris-EDTA buffer (12 mM Tris-HCl

and 1.2 mM EDTA). Tubes were rested for 10 minutes at room temperature, then heat-shocked

in a water bath at 39°C for 26 minutes. Cells from all 20 tubes were then combined. We

transferred cells from this post-transformation culture, and YPD, to each of three 1 L flasks at

the volumes required to attain an OD_{600} of ~0.35-4 in 500 mL. Each such culture was recovered

by shaking at 28°C and 200 rpm for 2 hours. G418 (Geneticin, Gibco) was added to each at a

concentration of 300 µg/mL to select for those cells which had taken up the plasmid, and

cultures were incubated with 200 rpm shaking at 28°C for two days until each reached an OD_{600}

of ~2.3. All six such selected cultures across the two transformations were combined. We

transferred cells from this combined culture, and YPD + G418 (300 ug/mL), to each of two 1 L

flasks at the volumes required to attain an OD_{600} of 0.2 in 500 mL each. We cultured each flask

at 28°C and 200 rpm shaking overnight until reaching an OD_{600} of 2.18 and combined the two

cultures again to yield one culture. To cure transformants of the pJR487 URA+ plasmid, we

spun down a volume of this master culture and resuspended in water with the volume required

to attain a cell density of 1.85 OD_{600} units/mL. 12 mL of this resuspension were plated (1 mL per

24.1cm x 24.1cm plate) onto plates containing complete synthetic media with 5-fluoroorotic acid

(5-FOA) [0.2% drop-out amino acid mix without uracil or yeast nitrogen base (YNB) (US

Biological), 0.005% uracil (Sigma), 2% D-glucose (Sigma), 0.67% YNB without amino acids

(Difco), 0.075% 5-FOA (Zymo Research)]. After incubation at 28°C to enable colony growth,

colonies were scraped off all 12 plates and combined into water at the volume required to attain

40 OD_{600} units per 900 µL, yielding the final transposon mutant hemizygote pool. This was

aliquoted into 1 mL volumes with 10% DMSO and frozen at -80°C.

Thermotolerance phenotyping via selection of the hemizygote pool

One aliquot of the pool of transposon mutant hemizygotes in the JR507 *S. cerevisiae*

DBVPG1373 x *S. paradoxus* Z1 hybrid background was thawed and inoculated into 150 mL of

YPD in a 250 mL flask, and cultured for 7.25 hours at 28°C, with shaking at 200 rpm. We used

this timepoint as time zero of our thermotolerance experiment, and took four aliquots of 6.43 mL

(7 OD units) as technical replicates for sequencing of transposon insertion positions (see

below). 9.19 mL of the remaining culture was back-diluted to an OD_{600} of 0.02 in a total of 500

mL YPD in each of six 2L glass flasks for cultures that we call selections; three were grown at

28°C and three at 39°C (shaking at 200 rpm) until an OD_{600} of 1.9-2.12 was reached,

corresponding to about 6.5 doublings in each case. Four cell pellets of 7 OD_{600} units each were

harvested from each of these biological replicate flasks, for sequencing as technical replicates

(see below). In total, 28 pellets were subjected to sequencing: 4 technical replicates from the

time-zero culture; 3 biological replicates, 4 technical replicates each, from the 28°C selection;

and 3 biological replicates, 4 technical replicates each, from the 39°C selection.

Tn-seq library construction

To determine the abundance of transposon mutant hemizygote clones after selection, we first

sequenced transposon (Tn) insertions as follows. Each cell pellet from a time zero or selection

sample (see above) was thawed on ice, and its genomic DNA (gDNA) was harvested with the

ZR Fungal/Bacterial DNA MiniPrep kit (Zymo Research). gDNA was resuspended in DNA

elution buffer (Zymo) pre-warmed to 65°C and its concentration was quantified using a Qubit 3.0

fluorometer. Illumina Tn-seq library construction was as described[28]. Briefly, gDNA was

sonicated and ligated with common adapters, and for each fragment deriving from a Tn insertion

in the genome, a sequence containing a portion of the transposon and a portion of its genomic

context (the Tn-genome junction) was amplified using one primer homologous to a region in the

transposon, and another primer homologous to a region in the adapter. The indexed adapter-

specific primer was

CAAGCAGAAGACGGCATACGAGATNNNNNNGTGACTGGAGTTCAGACGTGTGCTCTTCCG

ATCT, where the six N's are a unique index used for multiplexing multiple libraries onto the

same Hiseq sequencing lane, and the transposon specific primer was

ATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT

NNNNNNAGCAATATTTCAAGAATGCATGCGTCAAT, where N's are random nucleotides.

Amplification used Jumpstart polymerase (Sigma) and the following cycling protocol: 94°C-2

min, [94°C-30 sec, 65°C-20 sec, 72°C-30 sec] X 25, 72°C-10 min. Sequencing of single-end

reads of 150 bp was done over eight lanes on a HiSeq 2500 at the Joint Genome Institute

(Walnut Creek, CA). Reads sequenced per library are reported in Table S2.

Tn-seq read-mapping and data analysis

For analysis of data from the sequencing of Tn insertion sites in pools of hemizygotes, we first

searched each read for a string corresponding to the last 20 base pairs of the left arm of the

piggyBac transposon sequence, allowing up to two mismatches. For each Tn-containing read,

we then identified the genomic location of the sequence immediately downstream of the Tn

insertion site, which we call the genomic context of the insertion, by mapping with BLAT

(minimum sequence identity = 95, tile size = 12) against a hybrid reference genome made by

concatenating amended *S. cerevisiae* DBVPG1373 and *S. paradoxus* Z1 genomes (see below).

These genomic-context sequence fragments were of variable length; any case in which the

sequence was shorter than 50 base pairs was eliminated from further analysis, as was any case

in which a genomic-context sequence mapped to more than one location in the hybrid

reference. The resulting data set thus comprised reads containing genomic-context sequences

specifically mapping to a single location in either *S. cerevisiae* DBVPG1373 or *S. paradoxus* Z1,

which we call usable reads. For a given library, given a cohort of usable reads whose genomic-

context sequence mapped to the same genomic location, we inferred that these reads

originated from clones of a single mutant with the Tn inserted at the respective site, which we

call an insertion. In cases where the genomic-context sequences from reads in a given library

mapped to positions within 3 bases of each other, we inferred that these all originated from the

same mutant genotype and combined them, assigning to them the position corresponding to the

single location to which the most reads mapped among those combined. For a given insertion

thus defined, we considered the number of associated reads n_{insert} as a measure proportional to

the abundance of the insertion clone in the cell pellet whose gDNA was sequenced. To enable

comparison of these abundances across samples, we tabulated the total number of usable

reads n_{pellet} from each cell pellet, took the average of this quantity across pellets, $<n_{pellet}>$, and

multiplied each n_{insert} by $<n_{pellet}>/ n_{pellet}$ to yield a_{insert}, the final normalized estimate of the

abundance of the insertion clone in the respective pellet. For any insertions that were not

detected in a given pellet's library ($n_{insert} = 0$) but detectable in another library of the data set, we

assigned $n_{insert} = 1$.

We evaluated, from the mapped genomic-context sequence of each insertion, whether it fell into

a gene according to the *S. cerevisiae* and *S. paradoxus* genome annotations[17,29], and we

retained for further analysis only those insertions that fell into genes that were syntenic in the

two species. For each such insertion, for each biological replicate corresponding to a selection

culture (at 28°C or 39°C), we averaged the normalized abundances a_{insert} across technical

replicates, yielding a single abundance estimate $<a_{insert}>_{technical}$ for the biological replicate. We

then calculated the mean of the latter quantities across all biological replicates of the selection,

to yield a final abundance estimate for the insertion in this selection, $<a_{insert}>_{total}$. Likewise, for

each insertion and selection experiment we calculated $CV_{insert,total}$, the coefficient of variation of

$<a_{insert}>_{technical}$ values across biological replicates.

To use Tn-seq data in reciprocal hemizygosity tests, we considered for analysis only genes

annotated with the same (orthologous) gene name in the *S. cerevisiae* and *S. paradoxus*

reference genomes. For each insertion, we divided the $<a_{insert}>_{total}$ value from the 39°C selection

by the analogous quantity from the 28°C selection and took the log_2 of this ratio, which we

consider to reflect thermotolerance as measured by RH-seq. For each gene in turn, we used a

two-tailed Mann-Whitney U test to compare thermotolerance measured by RH-seq between the

set of insertions falling into the *S. cerevisiae* alleles of the gene, against the analogous quantity

from the set of insertions falling into the *S. paradoxus* allele of the gene, and we corrected for

multiple testing using the Benjamini-Hochberg method.

We tabulated the number of inserts and genes used as input into the reciprocal hemizygote test,

and the number of top-scoring genes emerging from these tests, under each of a range of

possible thresholds for coverage and measurement noise parameter values (Figure S5). We

used in the final analysis the parameter-value set yielding the most extensive coverage and the

most high-significance hits: this corresponded to insertions whose abundances had, in the data

from at least one of the two selections (at 28°C or 39°C), $CV_{insert,total} \leq 1.5$ and $<a_{insert}>_{total} \geq 1.1$,

and genes for which this high-confidence insertion data set contained at least 5 insertions in

each species' allele. This final data set comprised 110,678 high-quality insertions (Table S3) in

3416 genes (Table S4).

Amended reference genome construction

We generated reference genomes for *S. cerevisiae* strain DBVPG1373 and *S. paradoxus* strain

Z1 as follows. Raw genome sequencing reads for each strain were downloaded from the

SGRP2 database (ftp://ftp.sanger.ac.uk/pub/users/dmc/yeast/SGRP2/input/strains). Reads were

aligned using bowtie2[30] with default options; DBVPG1373 reads were aligned to version

R64.2.1 of the reference sequence of the *S. cerevisiae* type strain S288C (Genbank Assembly

Accession GCA_000146045.2), and Z1 reads were aligned to the *S. paradoxus* strain CBS432

reference sequence[31]. Single nucleotide variants (SNPs) were called using a pipeline of

samtools[32], bcftools and bgzip, and were filtered for a quality score (QUAL) of >20 and a

combined depth (DP) of >5 and either <65 (*S. cerevisiae*) or <255 (*S. paradoxus*). We then

amended each reference genome with the respective filtered SNPs: we replaced the S288C

allele with that of DBVPG1373 at each filtered SNP using bcftools' consensus command with

default options (42,983 base pairs total), and amendment of the CBS432 sequence was carried

out analogously using Z1 alleles (15,126 base pairs total).

Targeted-deletion hemizygote construction by homologous recombination

A given targeted hemizygote for each RH-seq hit gene except *TAF2* was generated in the *S.*

cerevisiae DBVPG1373 x *S. paradoxus* Z1 hybrid (JR507) by knocking out the allele of the gene

from one species via homologous recombination with KANMX as described[33] with 70 base pairs

of homology on the 5' and 3' ends of the cassette; checking was via diagnostic PCR. Two or

more independent transformants were isolated and phenotyped for each hemizygote genotype

(Table S1).

Construction of allele replacement and targeted hemizgyote strains with Cas9

At each RH-seq hit gene, we constructed strains in wild-type homozygous diploid *S. cerevisiae*

DBVPG1373 in which both copies of the endogenous allele were replaced by the allele from an

S. paradoxus isolate (Z1 for Figures 3, S8, and S9, and other strains as indicated for Figure

S10), and likewise for replacement of alleles from *S. cerevisiae* (DBVPG1373 in Figures 3, S8,

and S9, and other strains as indicated for Figure S10) into *S. paradoxus* Z1. We call each such

strain an allele-replacement strain, and each was constructed using a dual-guide Cas9

transgenesis method[34] in which a linear PCR fragment from the donor species is incorporated

into the recipient genome by homology-directed repair of two chromosomal double-strand

breaks induced by Cas9. Briefly, for each allele of each gene, we designed two guide RNAs for

double-strand breaks by Cas9: one guide targeted a position ~1000 base pairs 5' to the coding

start or at the 3' end of the closest upstream gene, whichever was closer, and the other guide

targeted the region of the coding stop. The precise cut site of each was chosen to contain an

NGG immediately downstream of variants between the *S. cerevisiae* and *S. paradoxus* strains,

to avoid re-cutting of the donor allele by Cas9 after it had been introduced into the recipient

strain. We cloned the two guide RNAs, a KANMX cassette, and the gene encoding the *S.*

pyogenes Cas9 protein into a single plasmid as described[34]. The resulting plasmid was

propagated in DH5a *E. coli* and miniprepped with a column miniprep kit (Qiagen). Separately, to

generate the fragment to be used as the donor for DNA repair after Cas9 cutting, we PCR-

amplified the respective region from the donor strain, with primers whose 5'-most 70 base pairs

were homologous to the recipient and whose 3'-most 20 base pairs were homologous to the

donor, except in the case of transgenesis using *S. cerevisiae* donors other than DBVPG1373 or

S. paradoxus donors other than Z1 (for Figure S10), for which the 3'-most 20 base pairs were

homologous to DBVPG1373 or Z1, respectively. The homology region at the 5' end of the gene

ended at most 31 base pairs upstream of the 5' cut site, and the homology region at the 3' end

of the gene started at most 33 base pairs downstream of the 3' cut site. The donor fragment

product was purified with a column kit (Qiagen) and ethanol-precipitated. We then

simultaneously transformed, using the lithium acetate method, the donor fragment and dual-

guide Cas9 plasmid into the recipient strain, using donor:acceptor ratios of 0.38:10 to 1:5, with

0.5-26 ug of plasmid. In this transformation, heat shock was for 20-30 minutes at 39-42°C in

transformations of *S. cerevisiae* DBVPG1373, and 10-20 minutes at 37-39°C for transformations

of *S. paradoxus* Z1. Transformants were plated on YPD+G418 (300 µg/mL) to select for cells

that retained the plasmid. From this selection we patched single colonies onto YPD without

G418, under the expectation that by the time a lawn came up for each patch, its cells would

have lost the Cas9 plasmid. Each such strain was Sanger-sequenced at the junctions of the

recipient and donor sequence. Positive patches were streaked to single colonies on YPD plates,

and cells from each such colony were used to inoculate a patch on a YPD plate and, separately,

to inoculate a patch on a YPD+G418 plate. Those colonies whose patches grew on the former

but not on the latter were inferred to be cured of the plasmid and stored at -80°C. For all genes

except *DYN1*, 2-3 such strains from each transformation were retained for thermotolerance

assays and underwent Sanger sequencing of the entire locus to determine the exact swapped

region (Table S1). For *DYN1* allele-replacement in *S. paradoxus* Z1, the Cas9-based strategy

yielded a single verified clone in which the *S. paradoxus* Z1 allele of *DYN1* was replaced by that

of *S. cerevisiae* DBVPG1373, and likewise for *DYN1* allele-replacement in *S. cerevisiae*

DBVPG1373. In each case, we mated the single swap clone to a wild-type of the respective

species background, confirmed heterozygosity of the resulting diploid via allele-specific

diagnostic PCR at the *DYN1* locus, sporulated, and dissected tetrads, allowing each spore to

autodiploidize and grow up as a homozygote; we retained from one such tetrad the two spores

that were homozygous at the *DYN1* locus for the swapped allele, as confirmed by sequencing,

and stored these at -80°C.

Targeted-deletion hemizygote strains for *TAF2* were generated by knocking out the *S.*

cerevisiae or the *S. paradoxus* allele in the interspecific hybrid (JR507) using the above

methods for Cas9 cutting and repair, with the following differences. To generate the fragment to

be used as the donor for DNA repair after Cas9 cutting, we PCR-amplified the NATMX cassette

from pBC713 (a gift from John Dueber, constructed as in [35]) using primers whose 5'-most 70

base pairs were homologous to the recipient and whose 3'-most 20 base pairs were

homologous to the cassette. The homology region at the 5' end of the gene ended 22 base

pairs outside the 5' cut site (which was upstream of coding start), and the homology region at

the 3' end of the gene started 33 base pairs outside the 3' cut site. Positive strains were

confirmed by PCR. Two independent transformants were isolated and phenotyped for each

genotype (Table S1).

Growth assays

Growth measurements of wild-type SGRP strains.

For the growth timecourse of a given wild-type, purebred homozygote isolate from the SGRP

collection at 28°C, it was streaked from a -80°C freezer stock onto a YPD agar plate and

incubated at 26°C for 3 days. A single colony was inoculated into liquid YPD and grown for 24

hours at 28°C with shaking at 200 rpm. This culture was back-diluted into YPD at an OD_{600} of

~0.05 and grown for an additional 5.5 hours at 28°C, 200 rpm, until reaching logarithmic phase.

We transferred cells from each such pre-culture, and YPD, to each of 11 replicate wells of a 96-

well plate, with volumes sufficient to yield a total volume of 150 μL per well at an OD_{600} of 0.02.

The plate was covered with a gas-permeable membrane (Sigma) and incubated with orbital

shaking in an M200 plate reader (Tecan, Inc.) at 28°C for 24 hours. For curves in Figure S1A,

measurements for optical density at 595nm (OD_{595}) were taken every 30 minutes and for each

timepoint, the average was taken across replicate wells. To subtract background OD_{595} for the

resulting curve, we tabulated the mean of the five lowest values from all datapoints, excluding

the first two, and subtracted this value from that of each timepoint, setting any negative value to

0. To smooth the resulting curve, we first replaced each timepoint measurement by its average

with those of the timepoints immediately before and after it; then, for any timepoint whose

measurement was not greater than or equal to the previous one, we set it to be equal to that

previous data point. For Figure S1B, the efficiency for a given growth curve (from a single well)

was calculated as the difference between the OD_{595} measured at the last four smoothed and

averaged data points and that of the first four smoothed and averaged data points. Efficiencies

from all of the wells from every *S. cerevisia*e isolate were combined and compared to

efficiencies from all of the wells for every *S. paradoxus* isolate in a two-sample two-tailed *t*-test.

For the growth timecourse of a given SGRP strain at 39°C (Figure 1A), we used a large-volume

flask growth paradigm to avoid the influence of plate effects on growth measurements at high

temperature in the incubated microplate reader, as follows. Each strain was streaked from a -

80°C freezer stock onto a YPD agar plate and incubated at 26°C for 3 days. A single colony of a

given strain was inoculated into liquid YPD and grown for 24 hours at 28°C with shaking at 200

rpm. Each of these cultures was back-diluted into YPD at an OD_{600} of 0.05 and grown for an

additional 5.5-7.5 hours at 28°C, shaking at 200 rpm, until reaching logarithmic phase. We

transferred cells from each such pre-culture, and YPD, to a glass 250 mL flask at the volumes

required to attain an OD_{600} of 0.05 in 100 mL YPD, and incubated it at 39°C with shaking at 200

rpm. OD_{600} readings were taken every ~2 hours for ~18 hours. Figure 1A reports representative

data from one of three such independent timecourse experiments. For curve fits, we used the

getInitial and SSlogis functions in R to estimate starting values for the parameters of the logistic

equation, and the nls function to fit the final parameters.

For efficiency measurements of a given SGRP isolate at 39°C in the large-volume format

(Figure 1B), it was streaked from a -80°C freezer stock onto a YPD agar plate and incubated at

26°C for 3 days. Two single colonies of each isolate were each inoculated into liquid YPD and

grown for 24 hours at 28°C with shaking at 200 rpm to generate two replicate pre-cultures. Each

was back-diluted into YPD at an OD_{600} at 600 nm of 0.05 and grown for an additional 5.5 hours

at 28°C, shaking at 200 rpm, until reaching logarithmic phase. The two pre-cultures were each

again back-diluted into YPD in 1-inch diameter glass tubes with a target OD_{600} of 0.05; the

actual OD_{600} of each was measured, after which it was grown at 39°C with shaking at 200rpm

for 24 hours, and OD_{600} was measured again. The efficiency for each replicate was calculated

as the difference between these final and initial OD_{600} values. The pipeline from inoculation off

solid plates through pre-culture, two back-dilutions, and growth at 39°C we refer to as a day's

growth experiment. For each day's experiments, we calculated the average efficiency across

the replicates of each isolate $<e_{strain}>$. We carried out two days' worth of replicate growth

experiments for each isolate. For a given species, we used the complete cohort of

measurements of $<e_{strain}>$ from all isolates of each species, across all days, as input into a two-

sample, two-tailed *t*-test to evaluate whether the suite of e_{strain} values across isolates of *S.*

cerevisiae was significantly different from the analogous set of values from *S. paradoxus*.

Growth measurements of targeted-deletion hemizygotes and allele-replacement strains at 28°C.

For efficiency measurements of a given targeted-deletion hemizygote or allele-replacement

strain at 28°C (Figures S7, S8 and S10), pre-culture and plate reader assays were as for wild-

type SGRP strains, except that 6 or more replicate wells were cultured per strain. 2-3

independently isolated targeted-deletion hemizygotes or allele-replacement strains (Table S1)

were assayed for each genotype. Each timecourse of targeted-deletion hemizygote or allele-

replacement strains also included the wild-type hybrid (JR507) or parent (*S. cerevisiae*

DBVPG1373 or *S. paradoxus* Z1), respectively, with pre-culture and replication as above.

Efficiency for a given growth curve (from a single well) was calculated as the difference between

the OD_{600} measured at the last four smoothed and averaged datapoints and that of the first four

smoothed and averaged datapoints, with smoothing and averaging as detailed above. For

Figure S7, relative efficiency for a given well of a given targeted-deletion hemizygote strain at

28°C was tabulated as its efficiency divided by that of the average of all replicate wells of the

wild-type hybrid (JR507) grown in the same experiment. For a given gene, we used the

complete cohort of these measurements, from all isogenic hemizygotes, as input into a two-

sample, two-tailed *t*-test to evaluate whether the relative efficiency of the strain in which the *S.*

cerevisiae allele was knocked out was lower than the analogous quantity from the strain in

which the *S. paradoxus* allele was knocked out. In Figures S8 and S10, allele-replacement

strains for a given gene were analyzed analogously, with the relative efficiency calculated

against the respective wild-type parent, and with a one-sample, two-tailed *t*-test to evaluate

whether the relative efficiency was significantly different from 1.

Growth measurements of targeted-deletion hemizygotes and allele-replacement strains at 39°C.

For efficiency measurements of a given targeted-deletion hemizygote strain at 39°C in the large-

volume format (Figure 2B), each strain was streaked from a -80°C freezer stock onto a YPD

agar plate and incubated at 26°C for 3 days. Two single colonies of a given strain were each

inoculated into liquid YPD and grown for 24 hours at 28°C with shaking at 200 rpm. Each such

pre-culture at stationary phase, or a log-phase outgrowth of it (which we used in the case of

DYN1 and *TAF2*: the pre-culture and YPD were added at the volumes required to attain an

OD_{600} of 0.05 and grown for an additional 5.5 hours at 28°C, shaking at 200 rpm, until the

culture reached logarithmic phase) was used to inoculate YPD in 1-inch diameter glass culture

tubes with a target cell density corresponding to an OD_{600} of 0.05. The actual OD_{600} of each was

measured, after which it was grown at 39°C with shaking at 200 rpm for 24 hours, and OD_{600}

was measured again. The efficiency for each such replicate was then calculated as the

difference between the final and initial OD_{600} values. The pipeline from inoculation off solid

plates through pre-culture, back-dilution, and growth at 39°C we refer to as a day's growth

experiment for a targeted-deletion homozygote. In each such experiment, 2-3 independently

isolated targeted-deletion hemizygotes of a given gene in each direction were all assayed on

the same day, alongside the wild-type hybrid parent (JR507) with replicate structure and

methods as above. From each day's experiments, we calculated the average efficiency across

the replicates of the wild-type hybrid $<e_{hybrid}>$, and we used this quantity to normalize the

efficiency $e_{hemizyg}$ measured for each replicate of each hemizygote strain assayed on that day.

Thus, the final observable used for analysis for each replicate on a given day was

$e_{hemizyg}/<e_{hybrid}>$. We carried out 2-3 days' replicate growth experiments for each gene's

hemizygotes. For a given gene, we used the complete cohort of these measurements of

$e_{hemizyg}/<e_{hybrid}>$, from all days and all isogenic hemizygotes, as input into a two-sample, one-

tailed *t*-test to evaluate whether $e_{hemizyg}/<e_{hybrid}>$ of the strain in which the *S. cerevisiae* allele

was knocked out was lower than the analogous quantity from the strain in which the *S.*

paradoxus allele was knocked out.

For growth measurements of a given allele-replacement strain at 39°C in the large-volume

format (Figure 3, S7, S8 and S10), each strain was streaked from a -80°C freezer stock onto a

YPD agar plate and incubated at 26°C for 3 days. 1-2 single colonies of each strain were each

inoculated into liquid YPD and grown for 24 hours at 28°C with shaking at 200 rpm to generate

1-2 replicate pre-cultures. Each was back-diluted into YPD at an OD_{600} of 0.05 and grown for an

additional 5.5 hours at 28°C, shaking at 200 rpm, until reaching logarithmic phase. Each pre-

culture were each again back-diluted into YPD in 1-inch diameter glass tubes with a target

OD_{600} of 0.05 (for experiments using a single pre-culture, it was now split into two replicate pre-

cultures, each of the same OD_{600}); the actual OD_{600} of each was measured, after which it was

grown at 39°C with shaking at 200rpm for 24 hours, and OD_{600} was measured again. The

efficiency for each replicate was calculated as the difference between these final and initial

OD_{600} values. The pipeline from inoculation off solid plates through pre-culture, two back-

dilutions, and growth at 39°C we refer to as a day's growth experiment for an allele-swap strain.

In each such experiment, 2-3 independently isolated allele-swap strains targeting a given gene

in a given background were all assayed on the same day, alongside the respective wild-type

background strain (*S. cerevisiae* DBVPG1373 or *S. paradoxus* Z1) with replicate structure and

methods as above. For each day's experiments, we calculated the average efficiency across the

replicates of the wild-type parent $<e_{parent}>$, and we used this quantity to normalize the efficiency

e_{swap} measured for each replicate assayed on that day of each allele-swap strain in the

respective background. Thus, the final measurement used for analysis for each replicate on a

given day was $e_{swap}/<e_{parent}>$. We carried out 2-3 days' worth of replicate growth experiments for

each gene's allele-swap strains. For a given gene in a given background, we used the complete

cohort of measurements of $e_{swap}/<e_{parent}>$ from all days and all allele-swap strains as input into a

one-sample, one-tailed *t*-test to evaluate whether $e_{swap}/<e_{parent}>$ was significantly different from

1. For swaps of the *S. cerevisiae* allele of a given gene into *S. paradoxus*, we tested whether

$e_{swap}/<e_{parent}>$ was greater than 1 (*i.e.* that the swap strain grew better at 39°C than did its

parent), and for swaps of the *S. paradoxus* allele of a given gene into *S. cerevisiae*, we tested

whether $e_{swap}/<e_{parent}>$ was less than 1 (*i.e.* that the swap strain grew worse at 39°C than its

parent).

Testing survival of wild-type parent strains at 39°C.

To test the survival of heat-treated *S. paradoxus* Z1 and *S. cerevisiae* DBVPG1373 (Figure S2),

each strain was streaked from -80°C freezer stocks onto YPD agar plates and incubated at

26°C for 3 days. 1-2 single colonies of each parent were inoculated into liquid YPD and grown

for 24 hours at 28°C with shaking at 200 rpm to create 1-2 replicate pre-cultures. After 24 hours,

we transferred cells from each pre-culture, and YPD, to each of 2-4 tubes at the volumes

required to attain an OD_{600} of 0.05 in 11 mL YPD. Two tubes were incubated at 28°C and two at

39°C, all with shaking at 200 rpm for 24 hours. OD_{600} of each was measured; for each culture

grown at 28°C, 100 µL of a 1.0×10^{-5} serial dilution was plated to YPD, and for cultures grown at

39°C, *S. paradoxus* and *S. cerevisiae* were serially diluted to 10^{-1} and 5.0×10^{-5}, respectively.

Plates were incubated at 26°C for three days until single colonies appeared. Colonies on each

plate were counted, from which we tabulated the colony forming units per mL of culture plated,

and normalized by the optical density of the original plated culture. The pipeline from pre-culture

through treatment and colony counting we refer to as one day's worth of experiments. We used

the results of two days' worth of experiments (a total of four for each species and temperature)

as input into an ANOVA with species and temperature as factors, and took the *p*-value for the

interaction between the factors as the estimate of the significance of the difference between

species in the effect of temperature (of the 24-hour liquid culture) on cells' ability to grow into

colonies (at the permissive temperature).

Microscopy and quantification

For microscopy of *S. cerevisiae* DBVPG1373 and *S. paradoxus* Z1 (Figure S3), each species

was streaked from a -80°C freezer stock onto a YPD agar plate and incubated at 26°C for 3

days. Two single colonies of each strain were each inoculated into liquid YPD and grown for 24

hours at 28°C with shaking at 200 rpm to generate two replicate pre-cultures. Each was then

back-diluted into YPD at an OD_{600} of 0.05; one was grown at 39°C with shaking at 200rpm for

24 hours, and the other was grown at 28°C with shaking at 200rpm for 24 hours. After the 24

hour growth period, 0.5 OD units of each culture were harvested through centrifugation and

incubated in 66.5% ethanol for 1-4 hours at room temperature. Each sample was washed twice

with 1X Dulbecco's Phosphate Buffered Saline (DPBS, Gibco), resuspended in 0.5 mL of 1X

DPBS, and vortexed for 15 seconds on high. 5 µL of each sample was transferred to an

agarose pad made with 1% agarose and YPD. We observed samples on a Zeiss Axio Observer

inverted bright-field microscope at 100X magnification. Images were taken using a Hamamatsu

ORCA-Flash4.0 digital camera and visualized using ZEN software for image analysis. The

exposure of each image was set automatically through ZEN, and brightness was adjusted using

the "Min/Max" adjustment for black and white light. The pipeline from inoculation off solid plates

through pre-culture, growth at 39°C or 28°C, fixation, and imaging we refer to as a day's

experiment. We carried out 2 days' worth of experiments for each species, yielding a total of 17-

21 images. In each image, free-floating cells were scored manually as singlets or those with a

small, medium, or large bud (for a total of 31-151 scored cells per species and temperature).

The proportion of scored cells with large buds was tabulated for each day's experiment for each

species and temperature. We used these proportions (a total of two for each species and

temperature) as input into an ANOVA with species and temperature as factors, and took the *p*-

value for the interaction between the factors as the estimate of the significance of the difference

between species in the effect of temperature on the proportion of cells with large buds.

Sequence analysis

D_{xy} *analysis.*

To assess the divergence between *S. cerevisiae* and *S. paradoxus* at loci mapped by RH-seq,

we used the D_{xy} statistic[36], the average number of pairwise differences between all *S. cerevisiae*

strains and *S. paradoxus*, normalized for gene length, as follows. We downloaded *S. cerevisiae*

genomic sequences from the following sources: YJM978, UWOPS83-787, Y55, UWOPS05-

217.3, 273614N, YS9, BC187, YPS128, DBVPG6765, YJM975, L1374, DBVPG1106, K11,

SK1, 378604X, YJM981, UWOPS87-2421, DBVPG1373, NCYC3601, YPS606, Y12,

UWOPS05-227.2, and YS2 from http://www.yeastrc.org/g2p/home.do; Sigma1278b, ZTW1, T7,

and YJM789 from http://www.yeastgenome.org/; and RM11 from NCBI (accession

PRJNA13674). For each strain, we extracted the coding sequence of each gene in turn, and we

downloaded the *S. paradoxus* reference sequence for each orthologous coding region from [17].

Sequences were aligned using MAFFT[37] with default settings. Alignments that did not contain a

start and stop codon, or those that contained gaps at greater than 40% of sites were considered

poor quality and discarded. We tabulated D_{xy} for each gene. To evaluate whether the eight RH-

seq hit genes were enriched for elevated D_{xy}, we first tabulated $<D_{xy}>_{true}$, the mean value across

the eight RH-seq hit genes. We then sampled eight random genes from the set of 3416 genes

tested by RH-seq; to account for biases associated with lower rates of divergence among

essential genes, the resampled set contained six essential genes and two non-essential genes,

mirroring the breakdown of essentiality among the RH-seq hits. Across this random sample we

tabulated the mean D_{xy}, $<D_{xy}>_{resample}$. We repeated the resampling 5000 times and used as an

empirical *p*-value the proportion of resamples at which $<D_{xy}>_{resample} \leq <D_{xy}>_{true}$.

Phylogenetic analysis.

We downloaded orthologous protein coding regions for the type strains of *S. cerevisiae*, *S.*

paradoxus, and an outgroup, *S. mikatae*, from [17]. For each gene for which ortholog sequences

were available in all three species, we aligned the sequences with PRANK[38] utilizing the "-

codon" option for codon alignment. These alignments were used as input into the codeml

module of PAML[39], which was run assuming no molecular clock and allowing omega values to

vary for each branch in the phylogeny. From the resulting inferences, we tabulated the branch

length on the *S. cerevisiae* lineage for each gene. To evaluate whether sequence divergence of

the eight RH-seq hit genes showed signatures of rapid evolution along the *S. cerevisiae* lineage,

we used the resampling test detailed above.

Locus effect size

Locus effect sizes in Figure S9 were computed from the data in Figures 2B and 3 of the main

text as follows. For analyses in the hybrid background, for a given locus we calculated $m_{S.par}$, the

mean of all replicate measurements of $e_{hemizyg}/<e_{hybrid}>$ of hemizygote strains lacking the *S.*

paradoxus allele. We took, as independent estimates of the locus effect size (each

corresponding to an open circle on the respective grey bar in Figure S9), each measurement of

$e_{hemizyg}/<e_{hybrid}>$ of a hemizygote strain lacking the *S. cerevisiae* allele, as a ratio against $m_{S.par}$.

For analyses in the *S. cerevisiae* DBVPG1373 background, for a given locus we used each

measurement of $e_{swap}/<e_{parent}>$ as an independent estimate of the locus effect size (each

corresponding to an open circle on the respective orange bar in Figure S9). For analyses in the

S. paradoxus Z1 background, for a given locus we used each measurement of $<e_{parent}>/e_{swap}$ as

an independent estimate of the locus effect size (each corresponding to an open circle on the

respective blue bar in Figure S9).

Data availability

RH-seq data will be deposited in the Sequence Read Archive (SRA) upon publication.

Code availability

Custom Python and R scripts used for RH-seq data analysis are available upon request.

References

Flint, J. & Mott, R. Finding the molecular basis of quantitative traits: successes and pitfalls. *Nat Rev Genet* **2**, 437-445, doi:10.1038/35076585 (2001).

Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The dynamics of molecular evolution over 60,000 generations. *Nature*, doi:10.1038/nature24287 (2017).

Savolainen, O., Lascoux, M. & Merila, J. Ecological genomics of local adaptation. *Nat Rev Genet* **14**, 807-820, doi:10.1038/nrg3522 (2013).

Allen Orr, H. The genetics of species differences. *Trends Ecol Evol* **16**, 343-350 (2001).

Nadeau, N. J. & Jiggins, C. D. A golden age for evolutionary genetics? Genomic studies of adaptation in natural populations. *Trends Genet* **26**, 484-492, doi:10.1016/j.tig.2010.08.004 (2010).

Wray, G. A. Genomics and the Evolution of Phenotypic Traits. *Annual Review of Ecology, Evolution, and Systematics* **44**, 51-72 (2013).

Masly, J. P. & Presgraves, D. C. High-resolution genome-wide dissection of the two rules of speciation in Drosophila. *PLoS biology* **5**, e243, doi:10.1371/journal.pbio.0050243 (2007).

Greig, D. A screen for recessive speciation genes expressed in the gametes of F1 hybrid yeast. *PLoS genetics* **3**, e21, doi:10.1371/journal.pgen.0030021 (2007).

Eshed, Y. & Zamir, D. An introgression line population of Lycopersicon pennellii in the cultivated tomato enables the identification and fine mapping of yield-associated QTL. *Genetics* **141**, 1147-1162 (1995).

Lazzarano, S. *et al.* Genetic mapping of species differences via in vitro crosses in mouse embryonic stem cells. *Proc Natl Acad Sci U S A* **115**, 3680-3685, doi:10.1073/pnas.1717474115 (2018).

Roop, J. I., Chang, K. C. & Brem, R. B. Polygenic evolution of a sugar specialization trade-off in yeast. *Nature* **530**, 336-339, doi:10.1038/nature16938 (2016).

Steinmetz, L. M. *et al.* Dissecting the architecture of a quantitative trait locus in yeast. *Nature* **416**, 326-330, doi:10.1038/416326a (2002).

Stern, D. L. Identification of loci that cause phenotypic variation in diverse species with the reciprocal hemizygosity test. *Trends Genet* **30**, 547-554, doi:10.1016/j.tig.2014.09.006 (2014).

Goncalves, P., Valerio, E., Correia, C., de Almeida, J. M. & Sampaio, J. P. Evidence for divergent evolution of growth temperature preference in sympatric Saccharomyces species. *PLoS One* **6**, e20739, doi:10.1371/journal.pone.0020739 (2011).

Salvado, Z. *et al.* Temperature adaptation markedly determines evolution within the genus Saccharomyces. *Appl Environ Microbiol* **77**, 2292-2302, doi:10.1128/AEM.01861-10 (2011).

Sweeney, J. Y., Kuehne, H. A. & Sniegowski, P. D. Sympatric natural Saccharomyces cerevisiae and S. paradoxus populations have different thermal growth profiles. *FEMS Yeast Res* **4**, 521-525 (2004).

Scannell, D. R. *et al.* The Awesome Power of Yeast Evolutionary Genetics: New Genome Sequences and Strain Resources for the Saccharomyces sensu stricto Genus. *G3 (Bethesda)* **1**, 11-25, doi:10.1534/g3.111.000273 (2011).

Hartwell, L. H. Saccharomyces cerevisiae cell cycle. *Bacteriol Rev* **38**, 164-198 (1974).

Mitra, R., Fain-Thornton, J. & Craig, N. L. piggyBac can bypass DNA synthesis during cut and paste transposition. *EMBO J* **27**, 1097-1109, doi:10.1038/emboj.2008.41 (2008).

803   20   van Opijnen, T., Lazinski, D. W. & Camilli, A. Genome-Wide Fitness and Genetic
804   Interactions Determined by Tn-seq, a High-Throughput Massively Parallel Sequencing
805   Method for Microorganisms. *Curr Protoc Mol Biol* **106**, 7 16 11-24,
806   doi:10.1002/0471142727.mb0716s106 (2014).
807   21   Parts, L. *et al.* Revealing the genetic structure of a trait by sequencing a population
808   under selection. *Genome Res* **21**, 1131-1138, doi:10.1101/gr.116731.110 (2011).
809   22   Winzeler, E. A. *et al.* Functional characterization of the S. cerevisiae genome by gene
810   deletion and parallel analysis. *Science* **285**, 901-906 (1999).
811   23   Leuenberger, P. *et al.* Cell-wide analysis of protein thermal unfolding reveals
812   determinants of thermostability. *Science* **355**, doi:10.1126/science.aai7825 (2017).
813   24   Wilkening, S. *et al.* An evaluation of high-throughput approaches to QTL mapping in
814   Saccharomyces cerevisiae. *Genetics* **196**, 853-865, doi:10.1534/genetics.113.160291
815   (2014).
816   25   Neurological Disorders: Public Health Challenges. 232 (World Health Organization,
817   2006).
818   26   Sinha, H., Nicholson, B. P., Steinmetz, L. M. & McCusker, J. H. Complex genetic
819   interactions in a quantitative trait locus. *PLoS Genet* **2**, e13,
820   doi:10.1371/journal.pgen.0020013 (2006).
821   27   Guldener, U., Heck, S., Fielder, T., Beinhauer, J. & Hegemann, J. H. A new efficient
822   gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res* **24**, 2519-
823   2524 (1996).
824   28   Wetmore, K. M. *et al.* Rapid quantification of mutant fitness in diverse bacteria by
825   sequencing randomly bar-coded transposons. *MBio* **6**, e00306-00315,
826   doi:10.1128/mBio.00306-15 (2015).
827   29   Skelly, D. A. *et al.* Integrative phenomics reveals insight into the structure of phenotypic
828   diversity in budding yeast. *Genome Res* **23**, 1496-1504, doi:10.1101/gr.155762.113
829   (2013).
830   30   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
831   **9**, 357-359, doi:10.1038/nmeth.1923 (2012).
832   31   Liti, G. *et al.* Population genomics of domestic and wild yeasts. *Nature* **458**, 337-341,
833   doi:10.1038/nature07743 (2009).
834   32   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
835   2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
836   33   Roop, J. I. & Brem, R. B. Rare variants in hypermutable genes underlie common
837   morphology and growth traits in wild Saccharomyces paradoxus. *Genetics* **195**, 513-525,
838   doi:10.1534/genetics.113.155341 (2013).
839   34   Maurer, M. J. *et al.* Quantitative Trait Loci (QTL)-Guided Metabolic Engineering of a
840   Complex Trait. *ACS Synth Biol* **6**, 566-581, doi:10.1021/acssynbio.6b00264 (2017).
841   35   Lee, M. E., DeLoache, W. C., Cervantes, B. & Dueber, J. E. A Highly Characterized
842   Yeast Toolkit for Modular, Multipart Assembly. *ACS Synth Biol* **4**, 975-986,
843   doi:10.1021/sb500366v (2015).
844   36   Nei, M. *Molecular Evolutionary Genetics*. (Columbia University Press, 1987).
845   37   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
846   improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
847   doi:10.1093/molbev/mst010 (2013).
848   38   Loytynoja, A. & Goldman, N. webPRANK: a phylogeny-aware multiple sequence aligner
849   with interactive alignment browser. *BMC Bioinformatics* **11**, 579, doi:10.1186/1471-2105-
850   11-579 (2010).
851   39   Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-
852   1591, doi:10.1093/molbev/msm088 (2007).

End Notes

Supplementary Information is available in the online version of the paper.

Acknowledgements The authors thank Faisal AlZaben, Anna Flury, Gina Geiselman,

Justin Hong, Jake Kim, Matt Maurer, and Luke Oltrogge for technical assistance, Dave

Savage for his generosity with microscopy resources, and Ben Blackman, Sam

Coradetti, Avi Flamholz, Vinny Guacci, Doug Koshland, Chris Nelson, and Arjun

Sasikumar for discussions. This work was supported by R01 GM120430-A1 and by

Community Sequencing Project 1460 to RBB at the U.S. Department of Energy Joint

Genome Institute, a DOE Office of Science User Facility. The work conducted by the

latter was supported by the Office of Science of the U.S. Department of Energy

under Contract No. DE-AC02-05CH11231.

Author Contributions R.B.B. and J.I.R. developed the project design; C.V.W., J.I.R.,

R.H. and J.C. performed experiments; C.V.W. and J.I.R. and analyzed the data; J.M.S.

contributed to the development of mutagenesis and sequencing methods; A.P.A.

contributed mutagenesis and sequencing reagents; I.V.G. provided technical assistance

with sequencing; R.B.B. and C.V.W wrote the manuscript with input from all authors.

Author Information The authors do not declare any competing financial interests.

Correspondence and request for materials should be addressed to R.B.B.

(rbrem@buckinstitute.org) and J.I.R. (jeremyianroop@gmail.com).