Improved genome assembly and annotation for the rock pigeon (*Columba livia*)

Carson Holt[*,†], Michael Campbell[*,1], David A. Keays[‡], Nathaniel Edelman[‡], Aurélie

Kapusta[*,†], Emily Maclary[§], Eric Domyan[§,**], Alexander Suh[††], Wesley C. Warren[‡‡],

Mark Yandell[*,†], M. Thomas P. Gilbert[§§,***], Michael D. Shapiro[§]

* Department of Human Genetics, University of Utah, Salt Lake City, UT, USA

† USTAR Center for Genetic Discovery, University of Utah, Salt Lake City, UT, USA

‡ Research Institute of Molecular Pathology, Vienna, Austria

§ Department of Biology, University of Utah, Salt Lake City, UT, USA

** Department of Biology, Utah Valley University, Orem, UT, USA

†† Department of Evolutionary Biology (EBC), University of Uppsala, Uppsala, Sweden

‡‡ Genome Institute at Washington University, St. Louis, MO, USA

§§ Natural History Museum of Denmark, University of Copenhagen, Copenhagen,

Denmark

*** Norwegian University of Science and Technology, University Museum, 7491

Trondheim, Norway

1 Current address: Division of Plant Biology, Cold Spring Harbor Laboratory, Cold

Spring Harbor, NY, USA

Keywords:

Columba livia, rock pigeon, HiRise assembly, MAKER annotation

Author for Correspondence:

Michael D. Shapiro, Department of Biology, University of Utah, 257 S 1400 E, Salt Lake

City, UT 84108, shapiro@biology.utah.edu, +1 801 581 5690

Abstract

The domestic rock pigeon (*Columba livia*) is among the most widely distributed and

phenotypically diverse avian species. This species is broadly studied in ecology, genetics,

physiology, behavior, and evolutionary biology, and has recently emerged as a model for

understanding the molecular basis of anatomical diversity, the magnetic sense, and other

key aspects of avian biology. Here we report an update to the *C. livia* genome reference

assembly and gene annotation dataset (Cliv_1.0). Greatly increased scaffold lengths in

the updated reference assembly, along with an updated annotation set, provide improved

tools for evolutionary and functional genetic studies of the pigeon, and for comparative

avian genomics in general.

Introduction

Intensive selective breeding of the domestic rock pigeon (*Columba livia*) has resulted in

over 350 breeds with extreme differences in morphology and behavior (Levi 1986;

Domyan and Shapiro 2017). The large phenotypic differences among different breeds

make them a useful model for studying the genetic basis of radical phenotypic changes,

which are more typically found among different species rather than within a single

species.

In genetic and genomic studies of *C. livia*, linkage analysis is important for identifying

genotypes associated with specific phenotypic traits of interest (Domyan and Shapiro

2017); however, short scaffold sizes in the Cliv_1.0 draft reference assembly (Shapiro et

al. 2013) hinder computationally-based comparative analyses. Short scaffolds also make

it more difficult to identify structural changes, such as large insertions or deletions, that

are responsible for traits of interest (Domyan et al. 2014; Kronenberg et al. 2015).

Here we present the Cliv_2.0 reference assembly and an updated gene annotation set. The

new assembly greatly improves scaffold length over the previous draft reference

assembly, and updated gene annotations show improved concordance with both

transcriptome and protein homology evidence.

Methods & Materials

Genome sequencing and assembly

Genomic DNA from a female Danish tumbler pigeon (full sibling of the male bird used

for the original Cliv_1.0 assembly (Shapiro et al. 2013)) was used to produce long-range

sequencing libraries using the "Chicago" (Putnam et al. 2016) method by Dovetail

Genomics (Santa Cruz, CA). Two Chicago libraries were prepared and sequenced on the

Illumina HiSeq platform to a final physical coverage (1-50 kb pairs) of 390x (see Table

1).

Scaffolding was performed by Dovetail Genomics using HiRise assembly software and

the Cliv_1.0 assembly as input. Briefly, Chicago reads were aligned to the input assembly

to identify and mask repetitive regions, and then a likelihood model was applied to

identify mis-joins and score prospective joins for scaffolding. The final assembly was

then filtered for length and gaps according to NCBI submission specifications.

Genome annotation

The pre-existing reference Gnomon (Souvorov et al. 2010) derived gene models for the

Cliv_1.0 assembly (GCA_000337935.1) were mapped onto the updated Cliv_2.0

reference assembly using direct alignment of transcript FASTA entries. This was done

using the alignment workflow of the genome annotation pipeline MAKER (Cantarel et al.

2008; Holt and Yandell 2011), which first seeds alignments using BLASTN (Altschul et

al. 1990) and then polishes the alignments around splice sites using Exonerate (Slater and

Birney 2005). Results were then filtered to remove alignments that had an overall match

of less than 90% of the original model (match is calculated as percent identity multiplied

by percent end-to-end coverage).

For final annotation, MAKER was allowed to identify *de novo* gene models that did not

overlap the aligned Gnomon models. Protein evidence sets used by MAKER included

annotated proteins from *Pterocles gutturalis* (yellow-throated sandgrouse) (Zhang et al.

2014) and *Gallus gallus* (chicken) (International Chicken Genome Sequencing 2004)

together with all proteins from the UniProt/Swiss-Prot database (Bairoch and Apweiler

2000; UniProt 2007). The transcriptome evidence sets for MAKER included Trinity

(Grabherr et al. 2011) mRNA-seq assemblies from multiple *C. livia* breeds and tissues

(methods for transcriptome assembly are described below). Gene predictions were

produced within MAKER by Augustus (Stanke and Waack 2003; Stanke et al. 2008)

trained against the Cliv_1.0 Gnomon gene models. Repetitive elements in the genome

were identified using a custom repeat library.

Custom repeat library

A repeat library for *C. livia* was built by combining libraries from existing avian species

(Zhang et al. 2014) together with with repeats identified *de novo* for the Cliv_2.0

assembly. *De novo* repeat identification was performed using RepeatScout (Price et al.

2005) with default parameters (>3 copies) to generate consensus repeat sequences.

Identified repeats with greater than 90% sequence identity and a minimum overlap of 100

bp were assembled using Sequencher (Yokouchi et al. 1993). Repeats were classified into

transposable element (TE) families using multiple lines of evidence, including homology

to known elements, presence of terminal inverted repeats (TIRs), and detection of target

site duplications (TSDs). Homology-based evidence was obtained using RepeatMasker

(Smit et al. 1996), as well as the homology module of the TE classifying tool RepClass

(Feschotte et al. 2009). RepClass was also used to identify signatures of transposable

elements (TIRs, TSDs). We then eliminated non-TE repeats (simple repeats or gene

families), using custom Perl scripts (available at https://github.com/4ureliek/ReannTE).

In our custom repeat analysis, using the script ReannTE_FilterLow.pl, consensus

sequences were labeled as simple repeats or low complexity repeats if 80% of their length

could be annotated as such by RepeatMasker (the library was masked with the option -

noint). Next using the ReannTE_Filter-mRNA.pl script, consensus sequences were

interrogated against RefSeq (Pruitt et al. 2007) mRNAs (as of March 7th 2016) with

TBLASTX (Altschul et al. 1990). Sequences were eliminated from the library when: (i)

the e-value of the hit was lower than 1E-10; (ii) the consensus sequence was not

annotated as a TE; and (iii) the hit was not annotated as a transposase or an unclassified

protein. The script ReannTE_MergeFasta.pl was then used to merge our library with a

library combining RepeatModeler (Smit and Hubley 2008) outputs from 45 bird species

(Kapusta et al. 2017) and complemented with additional avian TE annotations

(International Chicken Genome Sequencing 2004; Warren et al. 2010; Bao et al. 2015).

Merged outputs were then manually inspected to remove redundancy, and all DNA and

RTE class transposable elements were removed and replaced with manually curated

consensus sequences.

Transcriptomics Methods

RNA was extracted from adult tissues (brain, retina, subepidermis, cochlear duct, spleen,

olfactory epithelium) of the racing homer breed, and one whole embryo each of a racing

homer and a parlor roller (approximately embryonic stage 25 (Hamburger and Hamilton

1951)). RNA-seq libararies were prepared and sequenced using 100-bp paired-end

sequencing on the Illumina HiSeq 2000 platform at the Research Institute of Molecular

Pathology, Vienna (adult tissues), and the Genome Institute at Washington University, St.

Louis (embryos). RNA-seq data generated for the Cliv_1.0 annotation were also

downloaded from the NCBI public repository for *de novo* re-assembly. Accession

numbers for the public data are SRR521357 (Danish tumbler heart), SRR521358 (Danish

tumbler liver), SRR521359 (Oriental frill heart), SRR521360 (Oriental frill liver),

SRR521361 (Racing homer heart), and SRR521362 (Racing homer liver).

Each FASTQ file was processed with FastQC (http://www.bioinformatics.babraham.ac.

uk/projects/fastqc/) to assess quality. When FastQC reported overrepresentation of

Illumina adapter sequences, we trimmed these sequences with fastx_clipper from the

FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). We used FASTX-Toolkit for

two additional functions: runs of low quality bases at the start of reads were trimmed with

fastx_trimmer when necessary (quality cutoff of -Q 33), and reads were then trimmed

with fastq_quality_trimmer (-Q 33). Finally, each pair of sequence files was assembled

with Trinity (Grabherr et al. 2011) version r20131110 using the --jaccard_clip option.

Linkage map construction and anchoring to current assembly

Genotyping by sequencing (GBS) data was generated, trimmed, and filtered as previously

described (Domyan et al. 2016). Reads were mapped to the Cliv_2.0 assembly using

Bowtie2 (Langmead and Salzberg 2012). Genotypes were called using Stacks (Catchen et

al. 2011), with a minimum read-depth cutoff of 10. Thresholds for automatic corrections

were set using the parameters –min_hom_seqs 10, --min_het_seqs 0.01, --max_het_seqs

0.15. Sequencing coverage and genotyping rate varied between individuals, and birds

with genotyping rates in the bottom 25% were excluded from map assembly.

Genetic map construction was performed using R/qtl (www.rqtl.org) (Broman et al.

2003). For autosomal markers, markers showing segregation distortion (Chi-square, p <

0.01) were eliminated. Sex-linked scaffolds were assembled and ordered separately, due

to differences in segregation pattern for the Z-chromosome. Z-linked scaffolds were

identified by assessing sequence similarity and gene content between pigeon scaffolds

and the Z-chromosome of the annotated chicken genome (Ensembl Gallus_gallus-5.0).

Pairwise recombination frequencies were calculated for all autosomal and Z-linked

markers. Missing data were imputed using "fill.geno" with the method "no_dbl_XO".

Duplicate markers were identified and removed. Within individual scaffolds, R/Qtl

functions "droponemarker" and "calc.errorlod" were used to assess genotyping error.

Markers were removed if dropping the marker led to an increased LOD score, or if

removing a non-terminal marker led to a decrease in length of >10 cM that was not

supported by physical distance. Individual genotypes were removed if they showed with

error LOD scores >5 (Lincoln and Lander 1992). Linkage groups were assembled from

2960 autosomal markers and 232 Z-linked markers using the parameters (max.rf 0.1,

min.lod 6). In the rare instance that single scaffolds were split into multiple linkage

groups, linkage groups were merged if supported by recombination frequency data; these

instances typically reflected large physical gaps between markers on a single scaffold.

Scaffolds in the same linkage group were manually ordered based on calculated

recombination fractions and LOD scores.

To compare the linkage map to the prior assembly (Cliv_1.0), each 90-bp locus

containing a genetic marker was parsed from the Stacks output file

"catalogXXX_tags.tsv" and queried to the Cliv_1.0 assembly using Nucleotide-

Nucleotide blast (v2.6.0+) with the parameters –max_target_seqs 1 – max hsps 1. 3175 of

the 3192 loci (99.47%) from the new assembly had a BLAST hit with an E-value < 4e-24

and were retained.

Data availability

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under

the accession AKCR00000000. The version described in this paper is version

AKCR02000000. RNA-seq data are deposited in the SRA database with the BioSample

accession numbers SAMN07417936-SAMN07417943. Assembly and RNA-seq data are

publicly available in NCBI databases under BioProject PRJNA167554.

Results and Discussion

The final reference assembly is 1,108,534,737 base pairs in length and consists of 15,057

scaffolds (Table 1). A total of 1,015 scaffolds contain a gene annotation. Completion

analysis of the assembly using BUSCO (Simao et al. 2015) suggests that Cliv_2.0 is

72.9-86.2% complete which is nearly identical to the Cliv_1.0 assembly estimate of 72.3-

86.4% (Table 2). Thus, we found no significant changes to assembly completeness

between the two assemblies. The major improvement to the Cliv_2.0 assembly is rather

an increase in scaffold length (Fig. 1a). Overall, the N50 scaffold length increased to 14.3

megabases compared to 3.15 megabases for the previous reference assembly, a greater

than 4-fold increase. Recently, Damas et al. (Damas et al. 2017) used computational

methods and universal avian bacterial artificial chromosome (BAC) probes to achieve

chromosome-level scaffolding using the Cliv_1.0 assembly as input material; however,

this assembly is currently unannotated.

The new assembly joins scaffolds that we knew were adjacent but were separated

previously (see Table S1 for full catalog of positions of the original assembly in the new

assembly, and Table S2 for full catalog of breaks in the original assembly to form the

new assembly). For example, we previously determined that Cliv_1.0 Scaffolds 70 and

95 were joined based on genetic linkage data from a laboratory cross (Domyan et al.

2016). These two sequences are now joined into a single scaffold in the Cliv_2.0

assembly (see Table S3 for positions of genetic markers in Cliv_1.0 and Cliv_2.0). At

least one gene model (RefSeq LOC102093126), which was previously split across two

contigs, has now been unified into a single model on a single scaffold.

The updated annotation set contains 15,392 gene models encoding 18,966 transcripts (see

Table 3). This represents only a minor update of the reference annotation set as 94.7% of

previous models were mapped forward nearly unmodified (90% exact match for 14,898

out of 15,724 previous gene models) and only 494 new gene models were added to the

Cliv_2.0 annotation set (see Table 4).

The updated annotation set shows a modest improvement in concordance with aligned

evidence datasets from mRNA-seq and cross species protein homology evidence relative

to the Cliv_1.0 set as measured by Annotation Edit Distance (AED) (Eilbeck et al. 2009;

Holt and Yandell 2011). As a result, transcript models in the Cliv_2.0 annotation tend to

have lower AED values than the Cliv_1.0 set (Figure 2; the CDF curve is shifted to the

left). Lower AED values indicate greater model concordance with aligned transcriptome

and protein homology data. Furthermore, the Cliv_2.0 dataset displays greater transcript

counts in every AED bin despite having slightly fewer transcripts overall compared to the

Cliv_1.0 dataset (Table S4). The higher bin counts indicate that lower AED values are

not solely a result of removing unsupported models from the annotation set, but rather

suggest that evidence concordance has improved overall.

The improved scaffold lengths as well as updated gene model annotations should further

empower ongoing studies to identify genes responsible for phenotypic traits of interest

and improve detection of regions under selection due to longer scaffolds. We also expect

to be able to better identify large deletions and other structural variants responsible for

specific phenotypes now that they can be more clearly mapped to longer scaffolds.

Finally, the new transcriptomic data provides tissue-specific expression profiles for

several adult tissue types and an important embryonic stage for the morphogenesis of

limbs, craniofacial structures, skin, and other tissues.

Acknowledgements

We thank Dovetail Genomics for their aid in scaffolding the assembly, and Julia

Carleton, and Anna Vickrey for technical support. This work was supported by National

Science Foundation grant DEB1149160 and National Institutes of Health (NIH) grant

R01GM115996 to MDS; NSF EAGER grant IOS 1561337 to MY; a European Research

Council starting grant 336724 and Austrian Science Fund (FWF) grant Y726 to DAK;

and European Research Council Consolidator grant 681396 to MTPG. We gratefully

acknowledge research support from Boehringer Ingelheim at the Research Institute of

Molecular Pathology, and support and resources from the Center for High Performance

Computing at the University of Utah.

References

Altschul, S. F., W. Gish, W. Miller, E. W. Meyers, and D. J. Lipman. 1990. Basic Local Alignment Search Tool. Journal of Molecular Biology 215:403-410.

Bairoch, A. and R. Apweiler. 2000. The SWISS-PROT protein sequence database and its supplement TrEMBL in 2000. Nucl. Acids Res. 28:45-48.

Bao, W., K. K. Kojima, and O. Kohany. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. Mob DNA 6:11.

Broman, K., H. Wu, S. Sen, and G. Churchill. 2003. R/qtl: QTL mapping in experimental crosses. Bioinformatics 19:889-890.

Cantarel, B. L., I. Korf, S. M. C. Robb, G. Parra, E. Ross, B. Moore, C. Holt, A. Sanchez Alvarado, and M. Yandell. 2008. MAKER: An easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res. 18:188-196.

Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait. 2011. Stacks: building and genotyping loci de novo from short-read sequences. G3 1:171-182.

Damas, J., R. O'Connor, M. Farre, V. P. E. Lenis, H. J. Martell, A. Mandawala, K. Fowler, S. Joseph, M. T. Swain, D. K. Griffin, and D. M. Larkin. 2017. Upgrading short-read animal genome assemblies to chromosome level using comparative genomics and a universal probe set. Genome Res 27:875-884.

Domyan, E. T., M. W. Guernsey, Z. Kronenberg, S. Krishnan, R. E. Boissy, A. I. Vickrey, C. Rodgers, P. Cassidy, S. A. Leachman, J. W. Fondon, 3rd, M. Yandell, and M. D. Shapiro. 2014. Epistatic and combinatorial effects of pigmentary gene mutations in the domestic pigeon. Curr Biol 24:459-464.

Domyan, E. T., Z. Kronenberg, C. R. Infante, A. I. Vickrey, S. A. Stringham, R. Bruders, M. W. Guernsey, S. Park, J. Payne, R. B. Beckstead, G. Kardon, D. B. Menke, M. Yandell, and M. D. Shapiro. 2016. Molecular shifts in limb identity underlie development of feathered feet in two domestic avian species. eLife 5:e12115.

Domyan, E. T. and M. D. Shapiro. 2017. Pigeonetics takes flight: Evolution, development, and genetics of intraspecific variation. Dev Biol 427:241-250.

Eilbeck, K., B. Moore, C. Holt, and M. Yandell. 2009. Quantitative measures for the management and comparison of annotated genomes. BMC Bioinformatics 10:67.

Feschotte, C., U. Keswani, N. Ranganathan, M. L. Guibotsy, and D. Levine. 2009. Exploring repetitive DNA landscapes using REPCLASS, a tool that automates the classification of transposable elements in eukaryotic genomes. Genome Biol Evol 1:205-220.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X. Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A. Gnirke, N. Rhind, F. di Palma, B. W. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman, and A. Regev. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nature biotechnology 29:644-652.

Hamburger, V. and H. L. Hamilton. 1951. A series of normal stages in the development of the chick embryo. Journal of Morphology 88:49-92.

Holt, C. and M. Yandell. 2011. MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. BMC Bioinformatics
12:491.
International Chicken Genome Sequencing, C. 2004. Sequence and comparative analysis
of the chicken genome provide unique perspectives on vertebrate evolution.
Nature 432:695-716.
Kapusta, A., A. Suh, and C. Feschotte. 2017. Dynamics of genome size evolution in birds
and mammals. Proceedings of the National Academy of Sciences 114:E1460-
E1469.
Kronenberg, Z. N., E. J. Osborne, K. R. Cone, B. J. Kennedy, E. T. Domyan, M. D.
Shapiro, N. C. Elde, and M. Yandell. 2015. Wham: Identifying Structural
Variants of Biological Consequence. PLoS Comput Biol 11:e1004572.
Langmead, B. and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9:357-359.
Levi, W. M. 1986. The Pigeon (Second Revised Edition). Levi Publishing Co., Inc.,
Sumter, S.C.
Lincoln, S. E. and E. S. Lander. 1992. Systematic detection of errors in genetic linkage
data. Genomics 14:604-610.
Price, A. L., N. C. Jones, and P. A. Pevzner. 2005. De novo identification of repeat
families in large genomes. Bioinformatics 21 Suppl 1:i351-358.
Pruitt, K. D., T. Tatusova, and D. R. Maglott. 2007. NCBI reference sequences (RefSeq):
a curated non-redundant sequence database of genomes, transcripts and proteins.
Nucleic Acids Res:D61 - 65.
Putnam, N. H., B. L. O'Connell, J. C. Stites, B. J. Rice, M. Blanchette, R. Calef, C. J.
Troll, A. Fields, P. D. Hartley, C. W. Sugnet, D. Haussler, D. S. Rokhsar, and R.
E. Green. 2016. Chromosome-scale shotgun assembly using an in vitro method
for long-range linkage. Genome Res 26:342-350.
Shapiro, M. D., Z. Kronenberg, C. Li, E. T. Domyan, H. Pan, M. Campbell, H. Tan, C. D.
Huff, H. Hu, A. I. Vickrey, S. C. Nielsen, S. A. Stringham, H. Hu, E. Willerslev,
M. T. Gilbert, M. Yandell, G. Zhang, and J. Wang. 2013. Genomic diversity and
evolution of the head crest in the rock pigeon. Science 339:1063-1067.
Simao, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov.
2015. BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. Bioinformatics 31:3210-3212.
Slater, G. and E. Birney. 2005. Automated generation of heuristics for biological
sequence comparison. BMC Bioinformatics 6:31.
Smit, A. F. and R. Hubley. 2008. RepeatModeler Open-1.0
http://www.repeatmasker.org/.
Smit, A. F., R. Hubley, and P. Green. 1996. RepeatMasker Open-3.0
http://www.repeatmasker.org/.
Souvorov, A., Y. Kapustin, B. Kiryutin, V. Chetvernin, T. Tatusova, and D. Lipman.
2010. Gnomon – NCBI eukaryotic gene prediction tool. NCBI.
Stanke, M., M. Diekhans, R. Baertsch, and D. Haussler. 2008. Using native and
syntenically mapped cDNA alignments to improve de novo gene finding.
Bioinformatics 24:637-644.

Stanke, M. and S. Waack. 2003. Gene prediction with a hidden Markov model and a new
356   intron submodel. Bioinformatics 19:ii215-225.
357   UniProt, C. 2007. The Universal Protein Resource (UniProt). Nucleic Acids Res:D193 -
358   197.
359   Warren, W. C., D. F. Clayton, H. Ellegren, A. P. Arnold, L. W. Hillier, A. Kunstner, S.
360   Searle, S. White, A. J. Vilella, S. Fairley, A. Heger, L. Kong, C. P. Ponting, E. D.
361   Jarvis, C. V. Mello, P. Minx, P. Lovell, T. A. Velho, M. Ferris, C. N.
362   Balakrishnan, S. Sinha, C. Blatti, S. E. London, Y. Li, Y. C. Lin, J. George, J.
363   Sweedler, B. Southey, P. Gunaratne, M. Watson, K. Nam, N. Backstrom, L.
364   Smeds, B. Nabholz, Y. Itoh, O. Whitney, A. R. Pfenning, J. Howard, M. Volker,
365   B. M. Skinner, D. K. Griffin, L. Ye, W. M. McLaren, P. Flicek, V. Quesada, G.
366   Velasco, C. Lopez-Otin, X. S. Puente, T. Olender, D. Lancet, A. F. Smit, R.
367   Hubley, M. K. Konkel, J. A. Walker, M. A. Batzer, W. Gu, D. D. Pollock, L.
368   Chen, Z. Cheng, E. E. Eichler, J. Stapley, J. Slate, R. Ekblom, T. Birkhead, T.
369   Burke, D. Burt, C. Scharff, I. Adam, H. Richard, M. Sultan, A. Soldatov, H.
370   Lehrach, S. V. Edwards, S. P. Yang, X. Li, T. Graves, L. Fulton, J. Nelson, A.
371   Chinwalla, S. Hou, E. R. Mardis, and R. K. Wilson. 2010. The genome of a
372   songbird. Nature 464:757-762.
373   Yokouchi, Y., M. Yamamoto, T. Toyota, H. Sasaki, and A. Kuroiwa. 1993. Regulatory
374   interaction of positional signalings on coordinate expression of homeobox genes
375   in developing limb buds. Limb Development and Regeneration. Wiley-Liss, Inc.
376   Zhang, G., B. Li, C. Li, M. T. Gilbert, E. D. Jarvis, J. Wang, and C. Avian Genome.
377   2014. Comparative genomic data of the Avian Phylogenomics Project.
378   Gigascience 3:26.
379

380 **FIGURES**

382 **Figure 1.** Assembly scaffolding contiguity and scaffolding library insert size

383 distributions. (a) Scaffolding comparison between Cliv_1.0 (input scaffolds) and Cliv_2.0

384 (final scaffolds) assemblies. (b) Distribution of Dovetail Genomics "Chicago" library

385 inserts.

390 **Figure 2.** Evidence support comparison of annotation sets. Annotation edit distance

391 (AED) support for gene models in Cliv_2.0 (red line) is improved over Cliv_1.0 (NCBI

392 Gnomon annotation, blue line).

     TABLES

Table 1. Assembly statistics for Cliv_2.0

Estimated Physical Coverage	389.7x
Total Length	1,108,534,737bp
Total scaffolds	15,057
Total scaffolds >1kb	4,062
Total scaffolds >10kb	848

Table 2. Assembly version comparison

	Cliv_1.0	Cliv_2.0
Total Length	1110.8Mb	1110.9Mb
N50 Length	3.15Mb and 82 scaffolds	14.3Mb and 17 scaffolds
N90 Length	0.618Mb and 394 scaffolds	1.56Mb and 113 scaffolds
Completeness Estimate	72.3-86.4%	72.9-86.2%

Table 3. Annotation statistics

	Genes	Transcripts
Total	15,392	18,966
match[a]	14,898	18,472
new	494	494

[a] Count that match Cliv_1.0 annotations with a value of at least 90% (match is calculated as % identity multiplied by % end-to-end coverage)

Table 4. Annotation version comparison

	Cliv_1.0	Cliv_2.0
Total Gene Models	15,724	15,392
coding	15,022	14,683
non-coding	702	709
Total Transcripts	19,585	18,966
coding	18,569	18,148
non-coding	1016	818

SUPPLEMENTAL TABLES

Table S1. Tab-delimited table describing positions of Cliv_1.0 scaffolds in the Cliv_2.0 scaffolds. The table has the following format: column 1, Cliv_2.0 scaffold name; column 2, Cliv_1.0 sequence name; column 3, starting base (zero-based) of the Cliv_1.0 sequence; column 4, ending base of the Cliv_1.0 sequence; column 5, orientation of the Cliv_1.0 sequence in the Cliv_2.0 scaffold, where (-) indicates that the Cliv_2.0 scaffold sequence is reverse complemented relative to the Cliv_1.0 assembly; column 6, starting base (zero-based) in the Cliv_2.0 scaffold; column 7, ending base in the Cliv_2.0 scaffold.

Table S2. Tab-delimited table describing positions of breaks made in the Cliv_1.0 assembly to create the Cliv_2.0 assembly. Data fields follow the same format that is used in Supplemental Table 1.

Table S3. Table describing the linkage map assembled from genotype-by-sequencing markers aligned to the Cliv_2.0 assembly, and relative positions of aligned markers within the Cliv_2.0 and Cliv_1.0 genomes. The table has the following format: column 1, Linkage map marker ID; column 2, Linkage group ID; column 3, Linkage map position; column 4, Cliv_2.0 scaffold name; column 5, starting base (zero-based) of the alignment in the Cliv_2.0 scaffold; column 6, alignment orientation in the Cliv_2.0 scaffold; column 7, Cliv_1.0 scaffold name; column 8, starting base (zero-based) of the alignment the Cliv_1.0 scaffold; column 9, alignment orientation in the Cliv_1.0 scaffold.

Table S4. Tab-delimited table describing transcript count and CDF binned by Annotation

Edit Distance (AED) values. AED is a modified sensitivity/specificity metric used to

compare annotation datasets to each other or to aligned transcriptome and protein

homology datasets. For calculating AED, sensitivity is defined as the fraction of a given

reference overlapping a prediction and measures false negative rates. For our purposes,

the prediction is a transcript model and the reference (or truth set) is a set of aligned

transcriptome and protein homology evidence. We calculate sensitivity using the formula

$SN = |p{\cap}r|/|r|$; where $|p{\cap}r|$ represents the number overlapping nucleotides between the

prediction and reference, and $|r|$ represents the total number of nucleotides in the

reference. Specificity is then defined as the fraction of a prediction overlapping a given

reference, and it measures false positive rates. We calculate specificity using the formula

$SP = |p{\cap}r|/|p|$. We then define concordance to be the average of sensitivity and specificity

$(C = (SN+SP)/2)$, and AED is 1 minus the concordance $(AED = 1- C)$. Transcript models

that have high AED values then show little concordance to aligned experimental

evidence, and models with low AED values show high concordance.