**Identification of USP7 as an essential component to maintain integrity and function of**

**non-canonical PRC1.1 in leukemia**

Henny Maat[1], Jennifer Jaques[1], Aida Rodríguez López[1], Marcel P. de Vries[2,3], Chantal Gravesteijn[1],

Annet Z. Brouwers-Vos[1], Gerwin Huls[1], Edo Vellenga[1], Vincent van den Boom[1] and Jan Jacob

Schuringa[1,*].

[1]Department of Experimental Hematology, Cancer Research Center Groningen, University Medical

Center Groningen, University of Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands;

[2]Department of Pharmacy, Interfaculty Mass Spectrometry Center, University of Groningen, A.

Deusinglaan 1, 9713 AV Groningen, The Netherlands;

[3]Department of Pediatrics, Center for Liver, Digestive, and Metabolic Diseases, University Medical

Center Groningen, University of Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

[*]Correspondence: j.j.schuringa@umcg.nl. Department of Experimental Hematology, Medical Center

Groningen, University of Groningen, Hanzeplein 1, DA13, 9713 GZ Groningen, The Netherlands.

**ABSTRACT**

Polycomb proteins are essential epigenetic regulators of gene transcription. KDM2B, the chromatin-binding moiety of non-canonical PRC1.1, is critically important for human leukemias. Here, we investigated the complete interactome of KDM2B in human leukemic cells and identified that the deubiquitinase USP7 is an essential component of PRC1.1 and required for its stability and function. USP7 inhibition results in disassembly of the PRC1.1 complex and consequently loss of binding to its target loci. PRC1.1 can be associated with active loci and loss of PRC1.1 binding coincided with loss of H2AK119ub, reduced H3K27ac levels and reduced gene transcription, whereas H3K4me3 levels remained unaffected. Survival was reduced in (primary) acute myeloid leukemia cells in both cycling as well as quiescent populations upon USP7 inhibition, also independent of the USP7-MDM2-p53 axis. Finally, we evaluated the efficacy of USP7 inhibition *in vivo* and find that progression of MLL-AF9-induced leukemia is delayed, although in a niche-dependent manner.

**INTRODUCTION**

Regulation of gene transcription is carefully coordinated to control stem cell self-renewal and differentiation. Polycomb group (PcG) proteins are important epigenetic regulators and critically involved in controlling gene transcription by mediating post-translational modifications of histone proteins and chromatin remodeling (Kouzarides, 2007; Mas & Di, 2016; Simon & Kingston, 2013). Genome-wide analyses of Polycomb target genes revealed the occupancy of PcG proteins at promoters of genes regulating cell fate, highlighting their importance for proper lineage specification (Boyer et al., 2006; Bracken, Dietrich, Pasini, Hansen, & Helin, 2006; T. I. Lee et al., 2006). Yet, how PcG proteins are recruited, recognize their target genes and regulate gene expression still remains poorly understood. Understanding these processes is important since deregulation of PcG proteins frequently contributes to cancer and hematopoietic malignancies, like leukemia (Bracken & Helin, 2009; D. Hu & Shilatifard, 2016; Piunti & Shilatifard, 2016).

PcG proteins form multi-protein chromatin modifying complexes of which Polycomb Repressive

Complex 1 (PRC1) and 2 (PRC2) are best characterized (Simon & Kingston, 2009). PRC2, by means of

EZH1/EZH2, catalyzes the trimethylation of histone H3 on lysine 27 (H3K27me3) which can be

recognized by the chromodomain protein (CBX) of the canonical PRC1 complex (Cao et al., 2002;

Kirmizis et al., 2004; Kuzmichev, Nishioka, Erdjument-Bromage, Tempst, & Reinberg, 2002). PRC1

monoubiquitinates histone H2A on lysine 119 (H2AK119ub) via either one of the two E3 ligases

RING1A/1B (de Napoles et al., 2004; H. Wang et al., 2004). Proteome analysis has identified the

composition of six major PRC1 complexes that all contain RING1A or RING1B and distinguished by the

presence of one of the six PCGF proteins (PCGF1-6) (Gao et al., 2012; van den Boom et al., 2016).

Canonical PRC1 complexes contain either PCGF2 or PCGF4 and one of the CBX paralogs

(CBX2/4/6/7/8), PHC (1/2/3) and SCML1/SCML2/SCMH1 subunits. Non-canonical PRC1 complexes

lack a CBX subunit and instead contain RYBP or YAF2 which all three compete for the same binding

site of RING1B (Gao et al., 2012; R. Wang et al., 2010). The exact function of individual subunits in the

PRC1 complex is not fully understood, though it is suggested that they are involved in maintaining

the integrity of the complex, in providing or controlling enzymatic activity or in targeting to

chromatin. For example, the H2AK119 E3 ligase activity is enhanced by the dimerization of RING1A/B

with either PCGF2 or PCGF4 (Buchwald et al., 2006; Cao, Tsukada, & Zhang, 2005; Elderkin et al.,

2007) or can be stimulated by the PCGF1-RYBP/YAF2 interaction in the case of non-canonical PRC1.1

(Gao et al., 2012; Rose et al., 2016). A shRNA approach for individual PRC1 subunits in hematopoietic

stem cells revealed a lack of functional redundancy, suggesting unique functions of distinct PRC1

complexes (van den Boom et al., 2013) and indeed PRC1 complex composition changes upon lineage

specification (Morey et al., 2015).

To better understand the function of PRC1 complexes in human leukemias, we recently identified an

essential role for non-canonical PRC1.1 proteins (van den Boom et al., 2016). PRC1.1 was first

identified by the purification of the BCOR protein, which was found to interact with RING1A/B, RYBP,

PCGF1, SKP1 and KDM2B (Gearhart, Corcoran, Wamstad, & Bardwell, 2006; Sanchez et al., 2007). A

potential oncogenic role of PRC1.1 is underlined by the fact that KDM2B is overexpressed in

leukemias, breast and pancreatic cancers where it functions as an oncogene, conversely knockdown

of KDM2B abrogated tumorigenicity (Andricovich, Kai, Peng, Foudi, & Tzatsos, 2016; He, Nguyen, &

Zhang, 2011; Kottakis et al., 2014; Ueda et al., 2015; van den Boom et al., 2016).

Via the ZF-CxxC domain of KDM2B, PRC1.1 is targeted to non-methylated CpG islands genome-wide

and contributes to PRC1.1 recruitment (Blackledge et al., 2014; Farcas et al., 2012; He et al., 2013;

van den Boom et al., 2016; Wu, Johansen, & Helin, 2013). The PRC1.1 complex assembles around a

core PCGF1-RING1B dimer (Junco et al., 2013; Rose et al., 2016; Wong et al., 2016). While Polycomb-

mediated silencing by canonical PRC1 and PRC2 complexes is thought to maintain compacted

chromatin, which is inhibitory to transcription (Francis, Kingston, & Woodcock, 2004; Muller &

Verrijzer, 2009), less is known about the function of PRC1.1 on gene transcription. In mouse

embryonic stem cells (mESCs) and mouse embryonic fibroblasts it is suggested that KDM2B functions

together with PRC2 to repress lineage-specific genes (Blackledge et al., 2014; He et al., 2013; Tzatsos,

Pfau, Kampranis, & Tsichlis, 2009). On the other hand, Boulard and colleagues find more DNA

methylation in KDM2B null mESC at CpG islands co-occupied by PRC2, suggesting that KDM2B keeps

promoters in an activated non-methylated state (Boulard, Edwards, & Bestor, 2015). We and others

have shown that PRC1.1/KDM2B can also be targeted to sites independent of PRC2/H3K27me3 and

notably is associated with 'permissive' or active chromatin suggested by the occupancy of H3K4me3,

H3K36me3, RNAPII (S5P) and H3K27ac (Engelen et al., 2015; Liang, He, & Zhang, 2012; Tavares et al.,

2012; van den Boom et al., 2016).

A better understanding of PRC1.1 function is required in order to understand its role in gene

regulation and to explore possibilities to target PRC1.1 in human diseases such as leukemia. Here, we

investigated KDM2B/PRC1.1 in more detail with the aim to identify alternative approaches to target

the non-canonical PRC1.1 complex in human leukemic cells. LC-MS/MS-based proteome studies were

used to investigate the complete interactome of KDM2B in human leukemic cells and we identified

that the deubiquitinase USP7 is an essential component of PRC1.1 and is required to maintain its

integrity and function. As a consequence of inhibiting USP7 deubiquitinase activity the PRC1.1

complex disassembles and loses binding to chromatin, with a concomitant reduction in gene

expression of target loci. Furthermore, our data show that USP7 is essential for leukemic cells and

suggests that targeting of USP7 might provide an alternative therapeutic approach for leukemia.

**RESULTS**

**Identification of the deubiquitinase USP7 as a subunit of PRC1.1**

We recently identified KDM2B as a critically important factor for the survival of human leukemic

stem cells (van den Boom et al., 2016). Therefore, we set out to study the KDM2B interactome in

detail in leukemic cells. Human K562 cells were transduced with KDM2B-EGFP followed by anti-EGFP

pull outs and LC-MS/MS analysis to identify interacting proteins. Thus, 406 KDM2B-interacting

proteins were identified involved in cellular processes like DNA/chromatin binding, protein binding,

RNA binding and RNA polymerase activity (Figure 1 – source data 1, Figure 1A). Gene Ontology

analyses revealed that this list contained proteins that associated with rRNA processing, mRNA

splicing, translation, mRNA processing, positive regulation of gene expression and DNA damage

response (Figure 1B). The most abundant KDM2B interaction partners were non-canonical PRC1.1

proteins (Figure 1C), including the ubiquitin-specific protease 7 (USP7). Since inhibitors are available

that inhibit its deubiquitinase activity, we questioned whether inhibition of USP7 could be an

alternative way to target leukemic stem cells, in part via inhibition of PRC1.1 and therefore we

continued to study the role of USP7 in further detail. To validate the interaction of USP7 with PRC1.1

proteins, besides KDM2B, we also performed EGFP pull outs on nuclear extracts of stable K562 cell

lines expressing PCGF1-EGFP and EGFP-RING1B followed by LC-MS/MS analysis (Figure 1 – source

data 2). Figure 1D shows the spectral counts (MS/MS counts) corrected for expected peptides based

on *in silico* protein digests. Clearly, USP7 was found to interact directly with KDM2B, PCGF1 and

RING1B and all other PRC1.1 proteins were co-purified. In addition, Western blots of independent

pull outs on canonical PRC1 (PCGF2, PCGF4, CBX2) and non-canonical PRC1.1 (PCGF1, RING1B) were

performed using antibodies against USP7 and RING1B (Figure 1E). As expected, RING1B precipitated

with both PRC1 and PRC1.1 proteins. The interaction of USP7 with RING1B and PCGF1 was confirmed

by Western blot while less interaction was observed with the canonical PRC1 proteins PCGF2, PCGF4

or CBX2 (Figure 1E).

**USP7 inhibition results in disassembly of the PRC1.1 complex**

Deubiquitinating enzymes (DUBs) exert a variety of important cellular functions, including the control

over protein stabilization or degradation, protein localization, protein activity or by modulating

protein-protein interactions (Leznicki & Kulathu, 2017). We therefore questioned whether inhibition

of the deubiquitinase USP7 might affect PRC1.1 stability and function. P22077 (1-(5-((2,4-

difluorophenyl)thio)-4-nitrothiophen-2-yl)ethanone) has been identified as a selective inhibitor of

USP7, as well as USP47, using activity-based chemical proteomics (Altun et al., 2011; Tian et al.,

2011). In general, DUB inhibitors increase overall protein polyubiquitination (Altun et al., 2011).

Similarly, we observed accumulation of polyubiquitinated proteins in our EGFP-RING1B and PCGF1-

EGFP K562 cells treated for 24h with P22077 (Figure 2A). Next, we investigated the effect of USP7

inhibition on the stability of PRC1.1. EGFP pull outs were performed on nuclear extracts from K562

PCGF1-EGFP and EGFP-RING1B cells treated with DMSO or P22077 followed by LC/MS-MS analysis

(Figure 2B, Figure 2 – source data 1). Volcano plots were generated using label-free quantification

(LFQ) intensities of potential interactors of EGFP-RING1B (left) and PCGF1-EGFP (right) plotted as fold

change difference (control/USP7i) against significance (t-test p-value). Interactions with several

PRC1.1 proteins, highlighted in orange, were significantly reduced in both EGFP-RING1B and PCGF1-

EGFP pull outs as a consequence of USP7 inhibition (Figure 2B). The ubiquitin protein, UBB, was

enriched in both PCGF1-EGFP and EGFP-RING1B pull outs upon P22077 treatment, suggesting that

these proteins or other Polycomb interaction proteins might be more ubiquitinated upon USP7

inhibition. In addition we analysed the intensity-based absolute quantification (iBAQ) values, as a

measure for protein abundance, relative to either RING1B or PCGF1 and normalized to control pull

outs (Figure 2C). Similarly, these data demonstrated a clear reduced interaction of PRC1.1 proteins

with RING1B and PCGF1 after P22077 treatment. Independent EGFP pull outs performed on PCGF1-

EGFP and EGFP-RING1B cell lines further confirmed that USP7 inhibition indeed resulted in reduced

interaction of PCGF1-EGFP with endogenous RING1B and EGFP-RING1B with endogenous PCGF1

(Figure 2D). Importantly, input samples did not reveal loss of protein stability and also the expression

of EGFP-RING1B and PCGF1-EGFP was not affected as quantified by FACS analysis (Figure 2E). Taken

together these data indicate that USP7 is essential for PRC1.1 complex integrity.

**PRC1.1 chromatin binding relies on functional USP7 deubiquitinase activity**

To determine the functional consequences of disassembly of the PRC1.1 complex upon USP7

inhibition, we performed ChIP to investigate whether PRC1.1 would still be recruited to target loci.

Previously, we identified PRC1 and PRC1.1 target loci by ChIP-seq for PCGF1/2/4, CBX2, RING1A/1B,

KDM2B, H2AK119ub and H3K27me3 in the leukemic cell line K562 (van den Boom et al., 2016). Here,

ChIP-qPCRs were performed for KDM2B, PCGF1 and RING1B, the core PRC1.1 subunits, on a selected

set of loci in the absence or presence of P22077 (30 μM). Inhibition of USP7 resulted in a complete

loss of KDM2B binding, with concomitant strong reductions in PCGF1 and RING1B chromatin binding

(Figure 3A). Importantly, USP7 inhibition did not result in reduced protein levels of KDM2B, PCGF1 or

RING1B (Figure 2D/2E and data not shown). Since RING1B mediates H2AK119 ubiquitination we then

analysed the levels of this histone modification upon USP7 inhibition. Similarly, H2AK119ub was lost

from PRC1.1 target loci. Since USP7 does not target H2AK119 directly, these data  suggest that USP7

inhibition severely impacts on PRC1.1 chromatin binding and as a consequence H2AK119ub levels.

Since non-canonical PRC1.1 can be recruited to active loci in leukemic cells, we also investigated the

levels of H3K4me3, but no changes were observed on this histone mark upon USP7 inhibition, further

highlighting that not all posttranslational histone modifications are affected as a consequence of

treatment (Figure 3B). To test whether the loss of H2AK119ub would indeed be the result of

impaired *de novo* ubiquitination by either one of the two E3 ligases RING1A/1B, cells were cross-

linked at 4h, 8h and 16h followed by a ChIP for H2AK119ub, KDM2B and RING1B (Figure 3C). The

time-dependent loss of H2AK119ub coincided with a loss of KDM2B and RING1B binding at the same

target loci, strongly suggesting that loss of *de novo* ubiquitination underlies these observations.

Again, H3K4me3 levels remained unaffected, while H3K27ac levels were reduced upon USP7

inhibition indicative for reduced transcriptional activity at these loci (Figure 3D).

**USP7 inhibition leads to downregulation of several PRC1.1 active target genes**

Given that USP7 inhibition severely impaired PRC1.1 occupancy to several target loci we investigated

whether this would also affect the expression of PRC1.1 target genes in leukemic cells. RNA-seq was

performed to compare gene expression in DMSO or USP7 inhibitor treated K562 cells (P22077, 30

µM) for 4h, 8h, 16h and 24h. Since PRC1.1 can be associated with active genes in leukemic cells we

focused on the genes that were downregulated upon USP7 inhibition. In total 297 genes were

downregulated more than 2 fold after 24h of USP7 inhibition (Figure 4 – source data 1). Since USP7

has also PRC1.1 independent functions, for instance by controlling MDM2/p53 (Li, Brooks, Kon, & Gu,

2004), we then analysed whether downregulated genes overlapped with previously identified PRC1.1

(Figure 4A, Figure 4 – source data 2). Twenty-four percent of these genes overlapped with PRC1.1

targets. Gene Ontology (GO) analysis revealed that this set was enriched for GO terms like

transcription, regulation of gene expression and chromatin modification, while PRC1.1 independent

downregulated genes were enriched for GO terms like mRNA splicing, protein folding and protein

polyubiquitination (Figure 4B). ChIP-seq profiles of TOP2B, SIN3A, CHD1 and MYC are shown in Figure

4C as representative examples for PRC1.1 target genes. The tracks show clear binding of PCGF1,

RING1A/1B and KDM2B and little occupancy of canonical PRC1 proteins (PCGF2, PCGF4, CBX2). These

loci were also enriched for H2AK119ub and active chromatin marks H3K4me3, H3K36me3, RNAPII

and H3K27ac but devoid of the repressive mark H3K27me3. Loss of PRC1.1 binding was confirmed by

ChIP-qPCRs for KDM2B, PCGF1 and RING1B on these 4 loci (Figure 4D). Again, a concomitant loss of

H2AK119ub and H3K27ac marks was seen, without any alterations in H3K4me3. Subsequently, the

observed downregulation based on RNA-seq data was validated by independent quantitative RT-

PCRs (Figure 4E). In the majority of cases we observed that loss of PRC1.1 binding strongly correlated

with loss of gene expression, with only a few exceptions. For instance PKM, KIF21B and SLC25A22

were not downregulated despite a clear loss of PRC1.1 binding at these loci, suggesting that

transcription is controlled by additional mechanisms as well. Taken together, these data suggest that

the presence of PRC1.1 is required to maintain the transcriptional activity of several target genes.

**Both quiescent and cycling primary AML  cell populations are sensitive towards USP7 inhibition**

To study the functional consequences of USP7 inhibition, we tested P22077 on the AML cell lines

MOLM-13, NB4, KG1a, K562 and HL60. USP7 inhibition severely impaired cell growth in all tested

AML cell lines (Figure 5A) associated with increased apoptosis. Also cell lines that do not express

functional p53 (HL60 and K562) showed sensitivity, indicating that at least in those cell lines the

effect of P22077 was independent of p53 (Figure 5 – source data 1). Next, we assessed whether USP7

inhibition also affected the survival of primary AML cells. The long-term proliferation of primary AML

CD34$^+$ cells (n=3) on MS5 co-culture was clearly reduced upon USP7 inhibition (Figure 5B). Patient

characteristics are shown in Figure 5 – source data 1. With current therapies for leukemia patients

the majority of the blast population is readily eradicated, while the rare quiescent population of

leukemic stem cells is more difficult to target. To examine the effects of USP7 inhibition on both

cycling as well as quiescent ($G_0$) cells, we performed Hoechst/PyroninY stainings. AML 2 was

characterized by a large CD34$^+$/CD38$^+$ population that is high in cycling activity, while the

CD34$^+$/CD38$^-$ population is predominantly in $G_0$ (Figure 5C). In both populations, the $G_0$ fraction was

equally or even more efficiently targeted by P22077 compared to the cycling fraction, indicating that

cycling cells are not necessarily more sensitive than quiescent cells. AML3 was low in CD34$^+$ and

therefore we gated on the whole blast population which also showed that both quiescent and cycling

cells were sensitive for P22077 (Figure 5D).

**Evaluation of USP7 inhibition in xenograft leukemia mouse model**

Next, we evaluated the effect of USP7 inhibition in our human CB MLL-AF9 xenograft mouse model (Figure 6A) (Horton et al., 2013; Sontakke et al., 2016). Importantly, these lymphoid MLL-AF9 cells grown on a co-culture of MS5 *in vitro* were sensitive towards USP7 inhibition (Figure 6 – figure supplement 1). Here 5 x 10$^4$ MLL-AF9 leukemic cells, from a primary leukemic mouse, were intravenously (IV) injected into secondary recipients (n=11). Three weeks after injection 10/11 mice showed engraftment of MLL-AF9 EGFP$^+$/CD45$^+$/CD19$^+$ cells in peripheral blood (Figure 6B). Subsequently, mice were divided into two groups that were treated either with DMSO as control (n=5) or 20 mg/kg P22077 (n=6) via intraperitoneal (IP) injections. Peripheral blood chimerism levels were monitored by regular blood sample analysis and mice were sacrificed when chimerism levels in the blood exceeded 40%. Mice were treated daily starting four weeks post-transplant (indicated with the arrow, Figure 6C). Two-and-a-half weeks after starting treatment, chimerism levels were significantly lower in P22077-treated mice compared to DMSO (Figure 6B and 6C). The chimerism levels for DMSO treated mice rapidly increased to 44% (average) within 6 weeks after injection, whereas the chimerism levels of P22077 treated mice were around 25% on average. Notably, 3/6 mice gave a better response to USP7 inhibition with lower chimerism levels of 18.3, 16.8 and 12% respectively at day 46 (Figure 6B). Upon treatment we followed the mice for some additional weeks and within 60 days all control mice exceeded 40% chimerism in the blood, indicative for a full blown leukemia and were sacrificed (Figure 6C and D). Bone marrow, spleen and liver analyses showed high levels (>90%) of chimerism (data not shown). USP7 inhibition delayed leukemia development and in particular in two mice a clear response to USP7 inhibition was observed and chimerism levels remained relatively stable between day 56-67 post-transplant, although ultimately mice did develop MLL-AF9 induced leukemia after day 77 (Figure 6D).

Since the efficacy of USP7 inhibition *in vitro* in AML cell lines and patient samples was rather high, in contrast to what we observed *in vivo* in our xenograft mouse model, we hypothesized that the bone marrow niche could potentially play a protective role in the survival of leukemic cells and dampen

the response to therapeutic agents (Greim et al., 2014; Karjalainen et al., 2017; Parmar et al., 2011).

Therefore we evaluated the efficacy of USP7 inhibition in a direct comparison of CB MLL-AF9 cells

grown in liquid culture or in a co-culture setting on murine MS5 bone marrow stromal cells. In both

settings P22077 impaired cell growth, but the efficacy was much higher in liquid culture conditions

(Figure 6E). Loss of KDM2B binding to PRC1.1 target loci was confirmed by ChIP-qPCRs, both under

liquid culture and MS5 co-culture conditions, although we did note that the efficacy was somewhat

higher under liquid culture conditions (Figure 6F).

**DISCUSSION**

In this study, we reveal that the deubiquitinase USP7 is an essential component of non-canonical

PRC1.1 in leukemia and is required for its stability and function. Inhibiting USP7 deubiquitinase

activity, by P22077 treatment, resulted in disassembly of the PRC1.1 complex. Consequently,

recruitment of the PRC1.1 complex to target loci was heavily lost coinciding with the loss of its

H2AK119ub catalytic activity. This causes repression of a subset of transcriptionally active PRC1.1

target genes and corresponds with a reduction in H3K27ac, highlighting an essential function for

PRC1.1 in maintaining gene transcription. Furthermore, USP7 inhibition strongly impaired leukemic

cell growth *in vitro* and delayed MLL-AF9 induced leukemia *in vivo*, although in a niche-dependent

manner.

The non-canonical PRC1.1 complex consists of several Polycomb proteins and although we begin to

learn more about the function of some of the individual proteins within the complex, the role that

others fulfill remains far from completely understood. PRC1.1 is recruited to non-methylated CpG

islands via KDM2B and a single point mutation in the ZF-CxxC domain (K643A) of KDM2B has been

shown to prevent this interaction (Farcas et al., 2012; He et al., 2013; Wu et al., 2013). The H2AK119

E3 ubiquitin ligase activity of both canonical PRC1.2/1.4 and non-canonical PRC1.1 is mediated by

RING1A/1B proteins and is supported by the dimerization with one of the PCGFs (PCGF1-6) (Dou,

Buetow, Sibbet, Cameron, & Huang, 2012; Endoh et al., 2008; Taherbhoy, Huang, & Cochran, 2015).

For example, non-canonical PCGF1/3/5/6-RING1 heterodimers show much higher E3 ligase activity

than canonical PCGF2/4-RING1 heterodimers (Taherbhoy et al., 2015). This difference possibly lies in

a structural difference in PCGF helix α3 residues between canonical and non-canonical PCGFs which

may limit ubiquitin transfer by the RING1 dimer. Furthermore, the interaction of RYBP/YAF2 with the

PCGF1-RING1B core dimer increased the relative H2AK119 E3 ligase activity of PRC1.1 in an *in vitro*

assay (Rose et al., 2016). KDM2B and BCOR contributed little or nothing to this activity.

A role for the deubiquitinase USP7 within PRC1.1 was suggested since USP7 stably interacts with

PRC1.1 proteins, KDM2B, PCGF1 and RING1B (Figure 1). Sánchez and colleagues first identified the

ubiquitin protease USP7 as an interactor of RING1B (Sanchez et al., 2007). Then, USP7 was also

identified in proteomics analysis of PCGF1, RYBP, YAF2 and RING1A/B pullouts (Gao et al., 2012; van

den Boom et al., 2016). Previously, USP7 has been shown to be associated with the canonical PRC1

protein BMI1 (PCGF4) and potentially also with MEL18 (PCGF2), although Lecona and colleagues

suggested that USP7 interacts directly with SCML2 and thereby bridges the interaction with PRC1.4

(Lecona, Narendra, & Reinberg, 2015; Maertens, El Messaoudi-Aubert, Elderkin, Hiom, & Peters,

2010). Similar to Gao and colleagues (Gao et al., 2012) however, in our canonical PCGF2, PCGF4 and

CBX2 proteome analyses we found little or no interaction with USP7 (van den Boom et al., 2016). We

did notice that several canonical RING1B-Polycomb interactions were also reduced upon USP7

inhibition, including CBX2/4/8, SCML2 and SCMH1, though the interaction with PCGF2 and PCGF4

was not affected. This might also suggest that USP7 impacts on PRC1 protein stability, although

further studies are required to clarify these issues. In line with that notion, a recent paper

demonstrated that another deubiquitinase, USP26, controls the stability of CBX4 and CBX6 and

thereby affects the complex composition of PRC1 during ESC reprogramming (Ning et al., 2017).

Maertens and colleagues demonstrated that also USP11 interacts with PRC1 proteins and affects the

stability of BMI1 (Maertens et al., 2010). Thus, DUBs reveal an interesting role on Polycomb stability

and complex composition.

Based on our observations, the complex composition of PRC1.1 is affected upon USP7 inhibition. This

suggest that controlling ubiquitin levels of Polycomb proteins is important for the integrity of the

protein-protein interactions within PRC1.1 and its chromatin binding. Upon USP7 inhibition an overall

increase in protein ubiquitination was observed (Figure 2). Importantly, USP7 inhibition did not lead

to degradation of PCGF1 or RING1B, allowing us to conclude that USP7 controls protein-protein

interactions rather than proteasomal degradation. In line with that notion, overexpression of USP7

has been shown to deubiquitinate RING1B thereby stabilizing RING1B and preventing it from

proteasomal degradation (de, Zaaroor-Regev, & Ciechanover, 2010). Furthermore, RING1B self-

ubiquitination is required for its ligase activity on histone H2A (Ben-Saadon, Zaaroor, Ziv, &

Ciechanover, 2006). This is supported by the observation that H2AK119ub is lost upon USP7

inhibition, most likely as a consequence of loss of *de novo* ubiquitination mediated via RING1B.

Similarly, knockout of USP7 reduced H2AK119ub levels coinciding with reduced levels of RING1B

(Lecona et al., 2015). Other studies also show that USP7 is not likely a DUB for H2AK119 (Maertens et

al., 2010; Sarkari et al., 2009; van der Knaap et al., 2005). Thus, regulating ubiquitin levels within the

PRC1.1 complex itself is critically important and mediated via USP7 deubiquitinase activity.

With the efficient loss of PRC1.1 binding and consequent loss of H2AK119ub from target loci upon

USP7 inhibition we were able to study PRC1.1 function in relation to gene regulation in more detail.

Previously we identified genes that are targeted by PRC1.1 independent of H3K27me3 and have a

transcriptionally active profile suggested by H3K4me3, RNAPII and H3K27ac occupancy close to the

transcription start site. Loss of PRC1.1 binding resulted in reduced gene expression coinciding with

reduced levels of H3K27ac in the majority of analysed cases. The mechanisms underlying differences

in gene expression when PRC1.1 binding is lost are currently unclear. We hypothesize that PRC1.1

creates a transcriptionally permissive and open chromatin state which enables transcription factors

to bind and initiate gene expression. Addressing a possible cross-talk between PRC1.1 proteins,

CBP/p300-linked H3K27ac, recruitment of transcription factors and accessibility of chromatin is

definitely a focus for future work. Since H3K4me3 levels remained unaffected following USP7

inhibition, this suggest that the H3K4 methyltransferase complex that is likely targeted to CpGs via

CFP1 protein is not targeted in a PRC1.1-dependent manner (J. H. Lee & Skalnik, 2005).

In ESCs PRC1.1 targets are usually described together with canonical PRC1/PRC2 signaling, known as

bivalent loci and shown to repress target genes (Blackledge et al., 2014; Farcas et al., 2012; He et al.,

2013). For example, knockdown of KDM2B leads to the up-regulation of a subset of poised target

genes, coinciding with some reduction in RING1B binding and H2AK119ub (Farcas et al., 2012). While

we find no clear interaction of USP7 with canonical PRC1 in leukemic cells, it was shown that USP7

inhibition does impact functionally on canonical PRC1 (Lecona et al., 2015). Inhibiting the catalytic

core by removal of RING1A/B in ESCs does not lead to loss of KDM2B binding (Farcas et al., 2012).

However, in leukemic cells we find that USP7 inhibition not only leads to reduced H2AK119ub but

also induces loss of PRC1.1/KDM2B binding as a result of complex disassembly, suggesting that

KDM2B-PCGF1-BCOR/L1/SKP1 interactions might be needed for targeting. This is in agreement with

data that suggest that dimerization of PCGF1 and BCOR(L1) is required for binding to KDM2B and

recruiting PRC1.1 to the chromatin (Wong et al., 2016).

PRC1.1 is essential for human leukemic cells (van den Boom et al., 2016). By inhibiting individual

Polycomb proteins using an shRNA approach we observed impaired cell growth of leukemic cells

upon knockdown of PCGF1, RING1B, KDM2B or BCOR. *In vivo* experiments in humanized niche

xenograft mice using primary AML patient samples (Antonelli et al., 2016) or lentiviral leukemia

models (Sontakke et al., 2016) demonstrated that leukemogenesis was delayed or even abrogated

upon knockdown of PRC1.1 proteins. While these genetic studies clearly highlight the importance of

PRC1.1 for the survival of leukemic cells, small molecule inhibitors would be easier to implement in a

therapeutic clinical setting. Various USP7 inhibitors have been developed, and most recently

selective USP7 inhibitors which destabilize USP7 substrates including MDM2 and thereby increase

p53-mediated apoptosis of cancer cells were generated (Kategaya et al., 2017; Turnbull et al., 2017).

We observed that leukemic cells were very sensitive to USP7 inhibition, both in model systems as

well as in primary patient samples. Importantly, also the quiescent leukemic cells, that are usually

more difficult to target, showed sensitivity to P22077. It is evident that USP7 can function in several

pathways, often through regulating protein stability of tumor suppressors or epigenetic regulators

(Carra et al., 2017; Felle et al., 2011; van der Horst et al., 2006) and it is particularly the p53 pathway

that is strongly controlled by USP7 (Colland et al., 2009; Fan et al., 2013; M. Hu et al., 2006; Ye et al.,

2015). While we cannot exclude the possibility that some of these pathways were also affected in

some of our studies, we have analysed the p53 status in our models and primary AML patient

samples and see sensitivity even in the absence of a normal p53 response. Furthermore, while USP7

knockout mice are embryonically lethal, deletion of p53 was not able to rescue this phenotype,

further highlighting that p53-independent pathways downstream of USP7 exist as well (Kon et al.,

2010). Although P22077 has shown antitumor effects *in vivo* in neuroblastoma models (Fan et al.,

2013; Tavana et al., 2016), the responses that we observed *in vivo* in our CB MLL-AF9 leukemia model

were more modest, despite very strong effects seen upon USP7 inhibition *in vitro*. We wondered

whether the presence of a protective bone marrow microenvironment might play a role in regulating

P22077 sensitivity and therefore performed experiments in the absence or presence of bone marrow

stroma. We indeed demonstrate that MLL-AF9 cells grown on MS5 stromal cells were less sensitive

towards USP7 inhibition compared to cells grown in liquid culture conditions. These observations are

in line with other data clearly supporting the notion that the niche plays a protective role and can

dampen the response to therapeutic agents (Greim et al., 2014; Karjalainen et al., 2017; Parmar et

al., 2011).

In conclusion, our data reveal an important role for USP7 deubiquitinase activity in the integrity of

the PRC1.1 complex. We provide insight into the recruitment of PRC1.1 to target loci and function in

gene regulation and show that PRC1.1 is a potential interesting therapeutic target in leukemia.

**MATERIALS AND METHODS**

**GFP-mediated pull outs**

Pull outs were performed on nuclear extracts from K562 cells stably expressing KDM2B-EGFP, PCGF1-

EGFP and EGFP-RING1B. At least $80x10^6$ cells were collected for each pull out, washed with PBS and

lysed in 3 ml buffer A including freshly added protease inhibitors (10 mM HEPES/KOH pH 7.9, 1.5 mM

MgCl2, 10 mM KCl, 1 mM DTT, 1x Chymostatin Leupeptin Antipain Pepstatin (CLAP) protease-

inhibitor cocktail (all 1 µg/ml) and 0.1 mM PMSF (all from Sigma-Aldrich). After 10 minutes

incubation on ice, 10% IGEPAL CA-630 (endconc. 0.5%, Sigma-Aldrich) was added, followed by 10

seconds vortexing and 1 minute 14000 rpm spin to recover the nuclei. Nuclei were resuspended in

0.7 ml high salt buffer (20 mM HEPES/KOH pH 7.9, 1.5 mM MgCl2, 420 mM NaCl, 0.2 mM EDTA, 25%

Glycerol, 1 mM DTT, 1x CLAP and 0.1 mM PMSF) and incubated on ice for 20 minutes with 10

seconds vortexing intervals every  5 minutes. Extracted nuclei were collected after 15 minutes at

14000 rpm at 4°C and 2 volumes of Dilution buffer (25 mM Tris/Cl pH 7.5, 0.45% IGEPAL CA-630, 1 x

CLAP, 0.1 mM PMSF) and 3 volumes Wash buffer (TBS, 0.3% IGEPAL CA-630, 1 x CLAP, 0.1 mM PMSF)

were added. Pre-clearing of cell lysate was done by adding 50 µl pre-equilibrated binding control

magnetic agarose beads (Chromotek) and incubated for 30 min at 4°C on a rotating platform. Then

pre-cleared lysate was incubated with 70 µl pre-equilibrated GFP-Trap magnetic agarose beads

(Chromotek) overnight at 4°C on a rotating platform. Beads were separated using a magnetic rack

and six times washed in Wash buffer. Bound (B) fractions were eluted from the beads by boiling for

10 min in 2x Leammli sample buffer.

**In gel trypsin digestion**

Bound fractions of KDM2B-EGFP, PCGF1-EGFP (control/USP7i) and GFP-RING1B (control/USP7i) were

loaded on a 4-12% pre-cast NuPAGE gel (Invitrogen) and were run briefly. Gels were stained with

Coomassie dye R-250 (Thermo Scientific) and subsequently destained with ultrapure water

overnight. Gel lanes were cut into one slice, further cut into small pieces and completely destained

using 70% 50 mM $NH_4HCO_3$ and 30% acetonitrile (ACN). Reduction and alkylation of cysteines was

performed by adding 10 mM DTT dissolved in 50 mM $NH_4HCO_3$ and incubated at 55°C for 30 min.

Next, 55 mM iodoacetamide in 50 mM $NH_4HCO_3$ was added and incubated for 30 min, in dark,  at

room temperature. Remaining fluid was removed and 50 mM $NH_4HCO_3$ was added with 10 min.

shaking. Then 100% ACN was added incubated for 30 min. while shaking. Fluid was removed and gel

pieces were dried for 15 min. at 55°C. Proteins were digested with adding 10 ng/µl sequencing-grade

modified trypsin (Promega) in 50 mM $NH_4HCO_3$ to the gel pieces and incubated overnight at 37°C.

Next day, peptides were extracted using 5% formic acid followed by second elution with 5% formic

acid in 75% acetonitrile. Samples were dried in a SpeedVac centrifuge and dissolved in 5% formic

acid.

**LC-MS/MS analysis**

Online chromatography of the extracted tryptic peptides was performed with the Ultimate 3000

nano-HPLC system (Thermo Fisher Scientific) coupled online to a Q-Exactive-Plus mass spectrometer

with a NanoFlex source (Thermo Fisher Scientific) equipped with a stainless steel emitter. Tryptic

digests were loaded onto a 5 mm × 300 µm i.d. trapping micro column packed with PepMAP100 5

µm particles (Dionex) in 0.1% FA at the flow rate of 20 µL/min. After loading and washing for 3

minutes, peptides were forward-flush eluted onto a 50 cm × 75 µm i.d. nanocolumn, packed with

Acclaim C18 PepMAP100 2 µm particles (Dionex). The following mobile phase gradient was delivered

at the flow rate of 300 nL/min: 2–50% of solvent B in 90 min; 50–80% B in 1 min; 80% B during 9 min,

and back to 2 % B in 1 min and held at 3% A for 19 minutes. Solvent A was 100:0 H2O/acetonitrile

(v/v) with 0.1% formic acid and solvent B was 0:100 H2O/acetonitrile (v/v) with 0.1% formic acid. MS

data were acquired using a data-dependent top-10 method dynamically choosing the most abundant

not-yet-sequenced precursor ions from the survey scans (300–1650 Th) with a dynamic exclusion of

20 seconds . Sequencing was performed via higher energy collisional dissociation fragmentation with

a target value of 2e5 ions determined with predictive automatic gain control. Isolation of precursors

was performed with a window of 1.6. Survey scans were acquired at a resolution of 70,000 at m/z

200. Resolution for HCD spectra was set to 17,500 at m/z 200 with a maximum ion injection time of

110 ms. The normalized collision energy was set at 28. Furthermore, the S-lens RF level was set at 60

and the capillary temperature was set at 250degr. C. Precursor ions with single, unassigned, or six

and higher charge states were excluded from fragmentation selection.

**Data analysis**

Raw mass spectrometry data were analysed using MaxQuant version 1.5.2.8 (Cox & Mann, 2008),

using default settings and LFQ/ iBAQ enabled, searched against the Human Uniprot/Swissprot

database (downloaded June 26, 2016, 20197 entries). Further data processing was performed using

Perseus software, version 1.5.6.0 (Tyanova et al., 2016). Network visualization of KDM2B interactome

was performed using Cytoscape software, version 3.5.1. For Gene Ontology (GO) analysis we used

DAVID Bioinformatics Resources (http://david.abcc.ncifcrf.gov/home.jsp). ChIP-seq tracks were

visualized and analysed using UCSC genome browser (http://genome.ucsc.edu). Accession number

for the Polycomb ChIP-seq data previously reported and used for analysis in this paper is GEO:

GSE54580 (van den Boom et al., 2016). Previously published ChIP-seq used for analysis include

H3K4me3, H3K36me3, H3K27ac and RNAPII/Pol2(b) from  ENCODE/Broad Institute (GSE29611).

**Patient samples**

AML blasts from peripheral blood or bone marrow from untreated patients were studied after

informed consent and the protocol was approved by the Medical Ethical Committee, in accordance

with the Declaration of Helsinki. Mononuclear cells were isolated by density gradient centrifugation

and CD34+ cells were selected automatically by using autoMACS (Miltenyi Biotec).

**Cell culture**

The AML cell lines K562, MOLM13, NB4 and HL-60 were cultured in RPMI 1640 (BioWhittaker, Lonza,

Verviers, Belgium) supplemented with 10% fetal bovine serum (FCS, HyClone Laboratories, Logan,

Utah, US) and 1% penicillin/streptomycin (p/s, PAA Laboratories).  KG1a cells were cultured in IMDM

(BioWhittaker), supplemented with 20% FCS and 1% p/s. MS5 murine stromal cells were cultured in

alpha-MEM with 200 mM glutamine (BioWhittaker) supplemented with 10% FCS and 1% p/s. Primary

AMLs were cultured in Gartner's medium consisting of alpha-MEM (BioWhittaker) supplemented

with 12.5% heat-inactivated FCS and 12.5% heat-inactivated Horse serum (Sigma-Aldrich, Saint Louis,

MO, USA), 1 µM hydrocortisone (Sigma-Aldrich), 57.2 µM β-mercaptoethanol (Sigma) and 1% p/s in

the presence of 20 ng/ml granulocyte colony-stimulating factor (G-CSF), thrombopoietin (TPO) and

IL-3. All AMLs were grown on a MS5 stromal layer. CB MLL-AF9 liquid and MS5 co-cultures under

myeloid permissive conditions were grown in Gartner's medium supplemented with 20 ng/ml Flt-3L,

stem cell factor (SCF) and IL-3 (Horton et al., 2013; Sontakke et al., 2016). For lymphoid CB-MLL-AF9

cells were grown on MS5 co-culture in Gartner's medium without Horse serum and hydrocortisone

and no IL-3 was added. Instead ascorbic acid (50 µg/ml, Sigma) and IL-7 (10 ng/ml) were added. All

cultures were kept at 37°C and 5% $CO_2$. For USP7 inhibition experiments, P22077 (1-(5-((2,4-

difluorophenyl)thio)-4-nitrothiophen-2-yl)ethanone) was purchased from Merck Millipore (662142)

(Billerica, MA, USA) and stock solution dissolved in DMSO then added 1:1000 to the culture.

**Generation of lentiviral vectors and transductions**

Lentiviral pRRL SFFV PCGF1-EGFP and EGFP-RING1B vectors were generated as described previously

(van den Boom, 2016). pRRL SFFV KDM2B EGFP was generated as follows. KDM2B was PCR amplified

from cDNA in two parts (from the ATG to the RsrII site [fragment 1] and from the RsrII site to end of

KDM2B [excluding the stop codon, fragment 2]). Both fragments were independently subcloned into

pJet1.2 resulting in the pJet1.2 KDM2B[1] and pJet1.2 KDM2B[2] plasmids and that were

subsequenty verified by sequencing. Next, KDM2B fragment 2 was isolated from pJet1.2 KDM2B[2]

using RsrII and XbaI digestion and ligated into pJet1.2 KDM2B[1] that was also digested with RsrII and

XbaI, resulting in a pJet1.2 plasmid with the full length KDM2B ORF but excluding the stop codon.

Finally, the KDM2B ORF was subcloned into the pRRL SFFV GFP vector using  AgeI digestion, resulting

in the pRRL SFFV KDM2B-GFP construct. A lentiviral pRRL SFFV-His-Ubiquitin-mBlueberry2 vector was

generated by PCR amplification of His-Ubiquitin from pCl His-Ubi (Addgene) plasmid using primers

including BamHI sites. The PCR product was first ligated into pJet1.2 using Blunt-End cloning

protocol. Subsequently His-Ubiquitin was isolated from pJet1.2 using BamHI digestion and ligated

into pRRL SFFV-IRES-mBlueberry2 vector also using BamHI digestion and verified by sequencing.

Generation of lentiviral viruses and transductions were performed as described previously (van

Gosliga et al., 2007).

**Flow cytometry analysis**

Flow cytometry analyses were performed on the BD LSR II (Becton Dickinson (BD) Biosciences) and

data were analysed using FlowJo (Tree Star Inc, Ashland, OR, USA). Cells were sorted on a MoFlo XDP

or MoFLo Astrios (Beckman Coulter). For Hoechst/PyroninY staining, cells were resuspended in

HPGM (Lonza, Leusden, The Netherlands) and stained with 5 ug/ml Hoechst 33342 (Invitrogen) at

37°C for 30-45 min. Then 1 ug/ml PyroninY (Sigma) was added and incubated for 30-45 min at 37C.

Upon FcR blocking (MACS miltenyi Biotec), cells were stained with CD34-APC (581, BD Biosciences)

and CD38-AlexaFluor 700 (HIT2, Biolegend) at 4°C for 30 min. Cells were washed in medium

containing Hoechst/PyroninY and analysed on the BD LSR II. *In vivo* engraftment levels were analysed

in peripheral blood (PB), bone marrow, liver and spleen. Prior to staining, cells were blocked with

anti-human FcR block (MACS miltenyi Biotec) and anti-mouse CD16/CD32 block (BD Biosciences) and

stained with CD45-BV421 (HI30), CD19-BV785 (HIB19) and CD33-APC (WM53) all from Biolegend at

4°C for 30 min.

**Western blotting**

For detecting His-tagged ubiquitinated proteins cells were lysed in 1 ml denaturing lysis buffer (6M

Guanidium-HCl, 100 mM $Na_2HPO_4$ x $2H_2O$, 10 mM Tris-HCl pH 8.0, 5 mM Imidazole, 10 mM β-

mercaptoethanol) and sonicated on ice. Lysates were pre-cleared by centrifugation at 14000 rpm at

4°C and five volumes denaturing lysis buffer was added. His-tagged ubiquitinated proteins were

purified by adding 75 μl pre-equilibrated Ni-NTA magnetic agarose beads (Jena Bioscience) and

incubated for 4h on a rotating wheel. Beads were separated using a magnetic rack and washed 1x in

denaturing lysis buffer without Imidazole, 1x in wash buffer pH 8.0 (8M Urea, 100 mM $Na_2HPO_4$ x

$2H_2O$, 10 mM Tris-HCl pH 8.0 and 10 mM β-mercaptoethanol), 1x in wash buffer pH 6.3 (8M Urea,

100 mM $Na_2HPO_4$ x $2H_2O$, 10 mM Tris-HCl pH 6.3, 10 mM β-mercaptoethanol) plus 0.2% Triton X-100,

and 1x in wash buffer pH 6.3 plus 0.1% Triton X-100. His-tagged ubiquitinated proteins were eluted

from the beads by adding 75 µl elution buffer (200 mM Imidazole, 150 mM Tris-HCl pH 6.7, 30%

glycerol, 5% SDS and 720 mM β-mercaptoethanol) and incubated for 20 min. on a rotating wheel.

Elution samples were diluted 2x in Laemmli sample buffer containing 10% β-mercaptoethanol and

immediately boiled for 5 min. prior to Western blot analysis.

GFP mediated pull out cell lysates, input, non-bound and bound fractions were boiled for 5-10 min. in

Laemmli sample buffer prior to SDS-polyacrylamide gel electrophoresis. Proteins were transferred to

fluorescence polyvinylidene difluoride (PVDF FL, Millipore) membrane by semidry blotting.

Membranes were blocked in Odyssey blocking buffer (Westburg). The following primary antibodies

were used: anti-USP7 (A300-033A, Bethyl Laboratories), anti-RING1B (ab 181140, Abcam), anti-GFP

(sc-9996, Santa Cruz), anti-Ubiquitin (FK2, Enzo Life Sciences), anti-PCGF1 (ab183499, Abcam).

Fluorescent secondary antibodies either goat anti-mouse IRDye 800 or goat anti-rabbit IgG (H+L)

Alexa Fluor 680 (Invitrogen) were used for detection. Membranes were scanned using the Odyssey

CLx Imaging System (Li-Cor Biosciences).

**Chromatin immunoprecipitation**

ChIP was essentially performed as described previously (Frank, 2001), with minor modifications.

Briefly, K562 cells stably expressing low levels of EGFP-fusion vectors encoding, PCGF1-EGFP, EGFP-

RING1B, KDM2B-EGFP or non-transduced K562 cells and CB-MLL-AF9 cells were treated with DMSO

or P22077 for indicated timepoints and subsequently cross-linked with 1% formaldehyde for 10 min.

Reactions were quenched with addition of Glycine at a final concentration of 125 mM. Cells were

lysed in SDS buffer including protease inhibitors (1x CLAP, 0.1 mM PMSF) and snap frozen in liquid

nitrogen and stored at -80°C. ChIP was continued by resuspending cells in cold IP buffer and

sonicated using a Bioruptor (Diagenode). Pre-clearing was performed by adding Protein G Dynabeads

(Invitrogen) and incubated for 15 min at 4°C on a rotating platform. Then ChIP reactions were

performed using the following antibodies: anti-GFP (ab290, Abcam), anti-KDM2B (ab137547, Abcam),

anti-H2AK119ub (D27C4, Cell Signaling Technology), anti-H3K4me3 (ab8580, Abcam), anti-H3K27ac

(C15410196, Diagenode) and IgG (I8141, Sigma) followed by overnight incubation at 4°C on a rotating

platform. Next day Protein G Dynabeads were added for 2-4h after beads were separated using a

magnetic rack and washed with 3x Mixed Micelle Buffer, 2x Buffer 500, 2x LiCl Detergent Wash

Buffer and 1x TE (pH 8.0). ChIP DNA was eluted and reverse cross-linked at 65°C in the presence of

1% SDS and 0.1M NaHCO$_3$. Samples were treated with RNase and Proteinase K and purified using

QIAquick PCR purification kit (Qiagen). ChIPs were analysed by qPCR as percentage of input, primer

sequences are available on request.

**RNA seq analysis and quantitative real-time PCR**

RNA samples for sequencing were prepared for DMSO and P22077 (30 µM) treated K562 cells at 4h,

8h, 16h and 24h. Total RNA was isolated using the RNeasy Mini Kit from Qiagen (Venlo, The

Netherlands) according to the manufacturer's recommendations. Initial quality check and RNA

quantification of the samples was performed by capillary electrophoresis using the LabChip GX

(Perkin Elmer). Sequence libraries were generated with 50 ng mRNA, using Lexogen Quantseq 3' prep

kit (Lexogen GmbH) according to the manufacturer's recommendations. The obtained cDNA

fragment libraries were sequenced on an Illumina NextSeq500 using default parameters (single

read). Bioinformatics were performed on the Strand Avadis NGS (v3.0) software (Strand Life Sciences

Pvt.Ltd). Sequence quality was checked for GC content, base quality and composition using FASTQC

and StrandNGS. Quality trimmed reads were aligned to build Human Hg19 transcriptome. Ensembl

Genes and transcripts (2014.01.02) was used as gene annotation database. Quantified reads were

normalized using the DESeq package. Reads with failed vendor QC, quality score less than 24

(average), mapping quality score below 50 and length less than 20 were all filtered out.

For quantitative RT-PCR, RNA was reverse transcribed using the iScript cDNA synthesis kit (Bio-Rad)

and amplified using SsoAdvanced SYBR Green Supermix (Bio-Rad) on a CFX384 Touch Real-Time PCR

Detection System (Bio-Rad). RPL27 was used as housekeeping gene. Primer sequences are available

on request.

**USP7 inhibition *in vivo***

Eight to ten week old female NSG (NOD.Cg-Prkdcscid ll2rgtm1Wjl/SzJ) mice were purchased from the

Centrale Dienst Proefdieren (CDP) breeding facility within the University Medical Center Groningen.

Mouse experiments were performed in accordance with national and institutional guidelines, and all

experiments were approved by the Institutional Animal Care and Use Committee of the University of

Groningen (IACUC-RuG). 24h prior to transplantations, mice were sub-lethally irradiated with a dose

of 1.0 Gy (X-RAD 320 Unit, PXINC 2010). After irradiation mice received Neomycin (3.5 g/l) in their

drinking water and soft food (RM Convalescence + BG SY (M); Special Diet Services; Witham,

England) for two weeks. For secondary transplantations, 5 x $10^4$ MLL-AF9 EGFP cells from primary

leukemic mice (CB MLL-AF9 xenograft mouse model, (Horton et al., 2013; Sontakke et al., 2016))

were injected IV (lateral tail vein). Peripheral blood chimerism levels were monitored by regular

blood sample analysis. Mice were randomly divided into two groups, weighted and treated with

DMSO as control (n=5) or 20 mg/kg P22077 (n=6) via intraperitoneal (IP) injections daily starting four

weeks post-transplant. Prior to injections, P22077 was dissolved in DMSO (or DMSO only as control)

and directly mixed with Cremophor EL (1:1). This solution was then diluted 1:4 in saline, to get an end

concentration of max. 10% DMSO. Mice were humanely terminated by cervical dislocation under

isoflurane anesthesia when chimerism levels in the blood exceeded 40%. Peripheral blood, bone

marrow, spleen and liver were analysed.

**Acknowledgments**

The authors thank Dr. S. Bergink and E. de Mattos for providing pCl His-Ubi plasmid and advice with ubiquitin experiments. We thank M. Elliot for generation of KDM2B-EGFP construct. We would like to thank Roelof-Jan van der Lei, Theo Bijma, Henk Moes en Geert Mesander for help with cell sorting. We acknowledge Prof. Robert E Campbell (Department of Chemistry, University of Alberta, Edmonton, Alberta, Canada) for providing the mBlueberry2 Fluorescent Protein. This work is supported by the European Research Council (ERC-2011-StG 281474-huLSCtargeting) and the Dutch Cancer Foundation (RUG 2014-6832).

**Competing interest**

The authors declare no competing interests.

Reference List

Altun, M., Kramer, H. B., Willems, L. I., McDermott, J. L., Leach, C. A., Goldenberg, S. J., . . . Kessler, B.
M. (2011). Activity-based chemical proteomics accelerates inhibitor development for
deubiquitylating enzymes. *Chem Biol, 18*(11), 1401-1412.
doi:10.1016/j.chembiol.2011.08.018
Andricovich, J., Kai, Y., Peng, W., Foudi, A., & Tzatsos, A. (2016). Histone demethylase KDM2B
regulates lineage commitment in normal and malignant hematopoiesis. *J. Clin. Invest*.
doi:84014 [pii];10.1172/JCI84014 [doi]
Antonelli, A., Noort, W. A., Jaques, J., de, B. B., de Jong-Korlaar, R., Brouwers-Vos, A. Z., . . . Schuringa,
J. J. (2016). Establishing human leukemia xenograft mouse models by implanting human
bone marrow-like scaffold-based niches. *Blood, 128*(25), 2949-2959. doi:blood-2016-05-
719021 [pii];10.1182/blood-2016-05-719021 [doi]
Ben-Saadon, R., Zaaroor, D., Ziv, T., & Ciechanover, A. (2006). The polycomb protein Ring1B
generates self atypical mixed ubiquitin chains required for its in vitro histone H2A ligase
activity. *Mol Cell, 24*(5), 701-711. doi:10.1016/j.molcel.2006.10.022
Blackledge, N. P., Farcas, A. M., Kondo, T., King, H. W., McGouran, J. F., Hanssen, L. L., . . . Klose, R. J.
(2014). Variant PRC1 complex-dependent H2A ubiquitylation drives PRC2 recruitment and
polycomb domain formation. *Cell, 157*(6), 1445-1459. doi:S0092-8674(14)00589-3
[pii];10.1016/j.cell.2014.05.004 [doi]
Boulard, M., Edwards, J. R., & Bestor, T. H. (2015). FBXL10 protects Polycomb-bound genes from
hypermethylation. *Nat. Genet, 47*(5), 479-485. doi:ng.3272 [pii];10.1038/ng.3272 [doi]
Boyer, L. A., Plath, K., Zeitlinger, J., Brambrink, T., Medeiros, L. A., Lee, T. I., . . . Jaenisch, R. (2006).
Polycomb complexes repress developmental regulators in murine embryonic stem cells.
*Nature, 441*(7091), 349-353.
Bracken, A. P., Dietrich, N., Pasini, D., Hansen, K. H., & Helin, K. (2006). Genome-wide mapping of
Polycomb target genes unravels their roles in cell fate transitions. *Genes Dev, 20*(9), 1123-
1136.
Bracken, A. P., & Helin, K. (2009). Polycomb group proteins: navigators of lineage pathways led astray
in cancer. *Nat. Rev. Cancer, 9*(11), 773-784. doi:nrc2736 [pii];10.1038/nrc2736 [doi]
Buchwald, G., van der Stoop, P., Weichenrieder, O., Perrakis, A., van, L. M., & Sixma, T. K. (2006).
Structure and E3-ligase activity of the Ring-Ring complex of polycomb proteins Bmi1 and
Ring1b. *EMBO J, 25*(11), 2465-2474. doi:7601144 [pii];10.1038/sj.emboj.7601144 [doi]
Cao, R., Tsukada, Y., & Zhang, Y. (2005). Role of Bmi-1 and Ring1A in H2A ubiquitylation and Hox gene
silencing. *Mol. Cell, 20*(6), 845-854. doi:S1097-2765(05)01817-4
[pii];10.1016/j.molcel.2005.12.002 [doi]
Cao, R., Wang, L., Wang, H., Xia, L., Erdjument-Bromage, H., Tempst, P., . . . Zhang, Y. (2002). Role of
histone H3 lysine 27 methylation in Polycomb-group silencing. *Science, 298*(5595), 1039-
1043.
Carra, G., Panuzzo, C., Torti, D., Parvis, G., Crivellaro, S., Familiari, U., . . . Morotti, A. (2017).
Therapeutic inhibition of USP7-PTEN network in chronic lymphocytic leukemia: a strategy to
overcome TP53 mutated/deleted clones. *Oncotarget, 8*(22), 35508-35522.
doi:10.18632/oncotarget.16348
Colland, F., Formstecher, E., Jacq, X., Reverdy, C., Planquette, C., Conrath, S., . . . Daviet, L. (2009).
Small-molecule inhibitor of USP7/HAUSP ubiquitin protease stabilizes and activates p53 in
cells. *Mol Cancer Ther, 8*(8), 2286-2295. doi:10.1158/1535-7163.MCT-09-0097
Cox, J., & Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-
range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol, 26*(12),
1367-1372. doi:nbt.1511 [pii];10.1038/nbt.1511 [doi]

de, B. P., Zaaroor-Regev, D., & Ciechanover, A. (2010). Regulation of the Polycomb protein RING1B
ubiquitination by USP7. *Biochem. Biophys. Res. Commun, 400*(3), 389-395. doi:S0006-
291X(10)01592-5 [pii];10.1016/j.bbrc.2010.08.082 [doi]
de Napoles, M., Mermoud, J. E., Wakao, R., Tang, Y. A., Endoh, M., Appanah, R., . . . Brockdorff, N.
(2004). Polycomb group proteins Ring1A/B link ubiquitylation of histone H2A to heritable
gene silencing and X inactivation. *Dev. Cell, 7*(5), 663-676.
Dou, H., Buetow, L., Sibbet, G. J., Cameron, K., & Huang, D. T. (2012). BIRC7-E2 ubiquitin conjugate
structure reveals the mechanism of ubiquitin transfer by a RING dimer. *Nat Struct Mol Biol,*
*19*(9), 876-883. doi:10.1038/nsmb.2379
Elderkin, S., Maertens, G. N., Endoh, M., Mallery, D. L., Morrice, N., Koseki, H., . . . Hiom, K. (2007). A
phosphorylated form of Mel-18 targets the Ring1B histone H2A ubiquitin ligase to chromatin.
*Mol. Cell, 28*(1), 107-120. doi:S1097-2765(07)00552-7 [pii];10.1016/j.molcel.2007.08.009
[doi]
Endoh, M., Endo, T. A., Endoh, T., Fujimura, Y., Ohara, O., Toyoda, T., . . . Koseki, H. (2008). Polycomb
group proteins Ring1A/B are functionally linked to the core transcriptional regulatory
circuitry to maintain ES cell identity. *Development, 135*(8), 1513-1524.
doi:10.1242/dev.014340
Engelen, E., Brandsma, J. H., Moen, M. J., Signorile, L., Dekkers, D. H., Demmers, J., . . . Poot, R. A.
(2015). Proteins that bind regulatory regions identified by histone modification chromatin
immunoprecipitations and mass spectrometry. *Nat. Commun, 6*, 7155. doi:ncomms8155
[pii];10.1038/ncomms8155 [doi]
Fan, Y. H., Cheng, J., Vasudevan, S. A., Dou, J., Zhang, H., Patel, R. H., . . . Yang, J. (2013). USP7
inhibitor P22077 inhibits neuroblastoma growth via inducing p53-mediated apoptosis. *Cell*
*Death Dis, 4*, e867. doi:10.1038/cddis.2013.400
Farcas, A. M., Blackledge, N. P., Sudbery, I., Long, H. K., McGouran, J. F., Rose, N. R., . . . Klose, R. J.
(2012). KDM2B links the Polycomb Repressive Complex 1 (PRC1) to recognition of CpG
islands. *Elife, 1*, e00205. doi:10.7554/eLife.00205 [doi];00205 [pii]
Felle, M., Joppien, S., Nemeth, A., Diermeier, S., Thalhammer, V., Dobner, T., . . . Langst, G. (2011).
The USP7/Dnmt1 complex stimulates the DNA methylation activity of Dnmt1 and regulates
the stability of UHRF1. *Nucleic Acids Res, 39*(19), 8355-8365. doi:10.1093/nar/gkr528
Francis, N. J., Kingston, R. E., & Woodcock, C. L. (2004). Chromatin compaction by a polycomb group
protein complex. *Science, 306*(5701), 1574-1577. doi:10.1126/science.1100576
Gao, Z., Zhang, J., Bonasio, R., Strino, F., Sawai, A., Parisi, F., . . . Reinberg, D. (2012). PCGF Homologs,
CBX Proteins, and RYBP Define Functionally Distinct PRC1 Family Complexes. *Mol. Cell, 45*(3),
344-356. doi:S1097-2765(12)00035-4 [pii];10.1016/j.molcel.2012.01.002 [doi]
Gearhart, M. D., Corcoran, C. M., Wamstad, J. A., & Bardwell, V. J. (2006). Polycomb group and SCF
ubiquitin ligases are found in a novel BCOR complex that is recruited to BCL6 targets. *Mol.*
*Cell Biol, 26*(18), 6880-6889. doi:26/18/6880 [pii];10.1128/MCB.00630-06 [doi]
Greim, H., Kaden, D. A., Larson, R. A., Palermo, C. M., Rice, J. M., Ross, D., & Snyder, R. (2014). The
bone marrow niche, stem cells, and leukemia: impact of drugs, chemicals, and the
environment. *Ann. N. Y. Acad. Sci, 1310*, 7-31. doi:10.1111/nyas.12362 [doi]
He, J., Nguyen, A. T., & Zhang, Y. (2011). KDM2b/JHDM1b, an H3K36me2-specific demethylase, is
required for initiation and maintenance of acute myeloid leukemia. *Blood, 117*(14), 3869-
3880. doi:blood-2010-10-312736 [pii];10.1182/blood-2010-10-312736 [doi]
He, J., Shen, L., Wan, M., Taranova, O., Wu, H., & Zhang, Y. (2013). Kdm2b maintains murine
embryonic stem cell status by recruiting PRC1 complex to CpG islands of developmental
genes. *Nat. Cell Biol, 15*(4), 373-384. doi:ncb2702 [pii];10.1038/ncb2702 [doi]
Horton, S. J., Jaques, J., Woolthuis, C., van, D. J., Mesuraca, M., Huls, G., . . . Schuringa, J. J. (2013).
MLL-AF9-mediated immortalization of human hematopoietic cells along different lineages
changes during ontogeny. *Leukemia, 27*(5), 1116-1126. doi:leu2012343
[pii];10.1038/leu.2012.343 [doi]

Hu, D., & Shilatifard, A. (2016). Epigenetics of hematopoiesis and hematological malignancies. *Genes*
*Dev, 30*(18), 2021-2041. doi:30/18/2021 [pii];10.1101/gad.284109.116 [doi]
Hu, M., Gu, L., Li, M., Jeffrey, P. D., Gu, W., & Shi, Y. (2006). Structural basis of competitive
recognition of p53 and MDM2 by HAUSP/USP7: implications for the regulation of the p53-
MDM2 pathway. *PLoS Biol, 4*(2), e27. doi:10.1371/journal.pbio.0040027
Junco, S. E., Wang, R., Gaipa, J. C., Taylor, A. B., Schirf, V., Gearhart, M. D., . . . Kim, C. A. (2013).
Structure of the Polycomb Group Protein PCGF1 in Complex with BCOR Reveals Basis for
Binding Selectivity of PCGF Homologs. *Structure*. doi:S0969-2126(13)00053-1
[pii];10.1016/j.str.2013.02.013 [doi]
Karjalainen, R., Pemovska, T., Popa, M., Liu, M., Javarappa, K. K., Majumder, M. M., . . . Heckman, C.
A. (2017). JAK1/2 and BCL2 inhibitors synergize to counteract bone marrow stromal cell-
induced protection of AML. *Blood, 130*(6), 789-802. doi:10.1182/blood-2016-02-699363
Kategaya, L., Di Lello, P., Rouge, L., Pastor, R., Clark, K. R., Drummond, J., . . . Wertz, I. E. (2017). USP7
small-molecule inhibitors interfere with ubiquitin binding. *Nature*. doi:10.1038/nature24006
Kirmizis, A., Bartley, S. M., Kuzmichev, A., Margueron, R., Reinberg, D., Green, R., & Farnham, P. J.
(2004). Silencing of human polycomb target genes is associated with methylation of histone
H3 Lys 27. *Genes Dev, 18*(13), 1592-1605. doi:10.1101/gad.1200204
Kon, N., Kobayashi, Y., Li, M., Brooks, C. L., Ludwig, T., & Gu, W. (2010). Inactivation of HAUSP in vivo
modulates p53 function. *Oncogene, 29*(9), 1270-1279. doi:10.1038/onc.2009.427
Kottakis, F., Foltopoulou, P., Sanidas, I., Keller, P., Wronski, A., Dake, B. T., . . . Tsichlis, P. N. (2014).
NDY1/KDM2B Functions as a Master Regulator of Polycomb Complexes and Controls Self-
Renewal of Breast Cancer Stem Cells. *Cancer Res, 74*(14), 3935-3946. doi:0008-5472.CAN-13-
2733 [pii];10.1158/0008-5472.CAN-13-2733 [doi]
Kouzarides, T. (2007). Chromatin modifications and their function. *Cell, 128*(4), 693-705. doi:S0092-
8674(07)00184-5 [pii];10.1016/j.cell.2007.02.005 [doi]
Kuzmichev, A., Nishioka, K., Erdjument-Bromage, H., Tempst, P., & Reinberg, D. (2002). Histone
methyltransferase activity associated with a human multiprotein complex containing the
Enhancer of Zeste protein. *Genes Dev, 16*(22), 2893-2905.
Lecona, E., Narendra, V., & Reinberg, D. (2015). USP7 COOPERATES WITH SCML2 TO REGULATE THE
ACTIVITY OF PRC1. *Mol. Cell Biol*. doi:MCB.01197-14 [pii];10.1128/MCB.01197-14 [doi]
Lee, J. H., & Skalnik, D. G. (2005). CpG-binding protein (CXXC finger protein 1) is a component of the
mammalian Set1 histone H3-Lys4 methyltransferase complex, the analogue of the yeast
Set1/COMPASS complex. *J Biol Chem, 280*(50), 41725-41731. doi:10.1074/jbc.M508312200
Lee, T. I., Jenner, R. G., Boyer, L. A., Guenther, M. G., Levine, S. S., Kumar, R. M., . . . Young, R. A.
(2006). Control of developmental regulators by Polycomb in human embryonic stem cells.
*Cell, 125*(2), 301-313. doi:10.1016/j.cell.2006.02.043
Leznicki, P., & Kulathu, Y. (2017). Mechanisms of regulation and diversification of deubiquitylating
enzyme function. *J Cell Sci, 130*(12), 1997-2006. doi:10.1242/jcs.201855
Li, M., Brooks, C. L., Kon, N., & Gu, W. (2004). A dynamic role of HAUSP in the p53-Mdm2 pathway.
*Mol Cell, 13*(6), 879-886.
Liang, G., He, J., & Zhang, Y. (2012). Kdm2b promotes induced pluripotent stem cell generation by
facilitating gene activation early in reprogramming. *Nat. Cell Biol, 14*(5), 457-466.
doi:ncb2483 [pii];10.1038/ncb2483 [doi]
Maertens, G. N., El Messaoudi-Aubert, S., Elderkin, S., Hiom, K., & Peters, G. (2010). Ubiquitin-specific
proteases 7 and 11 modulate Polycomb regulation of the INK4a tumour suppressor. *EMBO J,*
*29*(15), 2553-2565. doi:10.1038/emboj.2010.129
Mas, G., & Di, C. L. (2016). The role of Polycomb in stem cell genome architecture. *Curr. Opin. Cell*
*Biol, 43*, 87-95. doi:S0955-0674(16)30155-7 [pii];10.1016/j.ceb.2016.09.006 [doi]
Morey, L., Santanach, A., Blanco, E., Aloia, L., Nora, E. P., Bruneau, B. G., & Di, C. L. (2015). Polycomb
Regulates Mesoderm Cell Fate-Specification in Embryonic Stem Cells through Activation and
Repression Mechanisms. *Cell Stem Cell, 17*(3), 300-315. doi:S1934-5909(15)00364-1
[pii];10.1016/j.stem.2015.08.009 [doi]

Muller, J., & Verrijzer, P. (2009). Biochemical mechanisms of gene regulation by polycomb group
protein complexes. *Curr Opin Genet Dev, 19*(2), 150-158. doi:10.1016/j.gde.2009.03.001
Ning, B., Zhao, W., Qian, C., Liu, P., Li, Q., Li, W., & Wang, R. F. (2017). USP26 functions as a negative
regulator of cellular reprogramming by stabilising PRC1 complex components. *Nat. Commun,
8*(1), 349. doi:10.1038/s41467-017-00301-4 [doi];10.1038/s41467-017-00301-4 [pii]
Parmar, A., Marz, S., Rushton, S., Holzwarth, C., Lind, K., Kayser, S., . . . Gotze, K. S. (2011). Stromal
niche cells protect early leukemic FLT3-ITD+ progenitor cells against first-generation FLT3
tyrosine kinase inhibitors. *Cancer Res, 71*(13), 4696-4706. doi:10.1158/0008-5472.CAN-10-
4136
Piunti, A., & Shilatifard, A. (2016). Epigenetic balance of gene expression by Polycomb and COMPASS
families. *Science, 352*(6290), aad9780. doi:10.1126/science.aad9780
Rose, N. R., King, H. W., Blackledge, N. P., Fursova, N. A., Ember, K. J., Fischer, R., . . . Klose, R. J.
(2016). RYBP stimulates PRC1 to shape chromatin-based communication between Polycomb
repressive complexes. *Elife, 5*. doi:10.7554/eLife.18591 [doi]
Sanchez, C., Sanchez, I., Demmers, J. A., Rodriguez, P., Strouboulis, J., & Vidal, M. (2007). Proteomics
analysis of Ring1B/Rnf2 interactors identifies a novel complex with the Fbxl10/Jhdm1B
histone demethylase and the Bcl6 interacting corepressor. *Mol. Cell Proteomics, 6*(5), 820-
834. doi:M600275-MCP200 [pii];10.1074/mcp.M600275-MCP200 [doi]
Sarkari, F., Sanchez-Alcaraz, T., Wang, S., Holowaty, M. N., Sheng, Y., & Frappier, L. (2009). EBNA1-
mediated recruitment of a histone H2B deubiquitylating complex to the Epstein-Barr virus
latent origin of DNA replication. *PLoS Pathog, 5*(10), e1000624.
doi:10.1371/journal.ppat.1000624
Simon, J. A., & Kingston, R. E. (2009). Mechanisms of polycomb gene silencing: knowns and
unknowns. *Nat. Rev. Mol. Cell Biol, 10*(10), 697-708. doi:nrm2763 [pii];10.1038/nrm2763
[doi]
Simon, J. A., & Kingston, R. E. (2013). Occupying chromatin: Polycomb mechanisms for getting to
genomic targets, stopping transcriptional traffic, and staying put. *Mol. Cell, 49*(5), 808-824.
doi:S1097-2765(13)00141-X [pii];10.1016/j.molcel.2013.02.013 [doi]
Sontakke, P., Carretta, M., Jaques, J., Brouwers-Vos, A. Z., Lubbers-Aalders, L., Yuan, H., . . .
Schuringa, J. J. (2016). Modeling BCR-ABL and MLL-AF9 leukemia in a human bone marrow-
like scaffold-based xenograft model. *Leukemia, 30*(10), 2064-2073. doi:leu2016108
[pii];10.1038/leu.2016.108 [doi]
Taherbhoy, A. M., Huang, O. W., & Cochran, A. G. (2015). BMI1-RING1B is an autoinhibited RING E3
ubiquitin ligase. *Nat. Commun, 6*, 7621. doi:ncomms8621 [pii];10.1038/ncomms8621 [doi]
Tavana, O., Li, D., Dai, C., Lopez, G., Banerjee, D., Kon, N., . . . Gu, W. (2016). HAUSP deubiquitinates
and stabilizes N-Myc in neuroblastoma. *Nat Med, 22*(10), 1180-1186. doi:10.1038/nm.4180
Tavares, Λ., Dimitrova, E., Oxley, D., Webster, J., Poot, R., Demmers, J., . . . Brockdorff, N. (2012).
RYBP-PRC1 Complexes Mediate H2A Ubiquitylation at Polycomb Target Sites Independently
of PRC2 and H3K27me3. *Cell*.
Tian, X., Isamiddinova, N. S., Peroutka, R. J., Goldenberg, S. J., Mattern, M. R., Nicholson, B., & Leach,
C. (2011). Characterization of selective ubiquitin and ubiquitin-like protease inhibitors using a
fluorescence-based multiplex assay format. *Assay Drug Dev Technol, 9*(2), 165-173.
doi:10.1089/adt.2010.0317
Turnbull, A. P., Ioannidis, S., Krajewski, W. W., Pinto-Fernandez, A., Heride, C., Martin, A. C. L., . . .
Komander, D. (2017). Molecular basis of USP7 inhibition by selective small-molecule
inhibitors. *Nature*. doi:10.1038/nature24451
Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M. Y., Geiger, T., . . . Cox, J. (2016). The Perseus
computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods,
13*(9), 731-740. doi:nmeth.3901 [pii];10.1038/nmeth.3901 [doi]
Tzatsos, A., Pfau, R., Kampranis, S. C., & Tsichlis, P. N. (2009). Ndy1/KDM2B immortalizes mouse
embryonic fibroblasts by repressing the Ink4a/Arf locus. *Proc. Natl. Acad. Sci. U. S. A, 106*(8),
2641-2646. doi:0813139106 [pii];10.1073/pnas.0813139106 [doi]

Ueda, T., Nagamachi, A., Takubo, K., Yamasaki, N., Matsui, H., Kanai, A., . . . Honda, H. (2015). Fbxl10
overexpression in murine hematopoietic stem cells induces leukemia involving metabolic
activation and upregulation of Nsg2. *Blood*. doi:blood-2014-03-562694 [pii];10.1182/blood-
2014-03-562694 [doi]

van den Boom, V., Maat, H., Geugien, M., Rodriguez, L. A., Sotoca, A. M., Jaques, J., . . . Schuringa, J. J.
(2016). Non-canonical PRC1.1 Targets Active Genes Independent of H3K27me3 and Is
Essential for Leukemogenesis. *Cell Rep, 14*(2), 332-346. doi:S2211-1247(15)01458-8
[pii];10.1016/j.celrep.2015.12.034 [doi]

van den Boom, V., Rozenveld-Geugien, M., Bonardi, F., Malanga, D., van, G. D., Heijink, A. M., . . .
Schuringa, J. J. (2013). Non-redundant and locus-specific gene repression functions of PRC1
paralog family members in human hematopoietic stem/progenitor cells. *Blood, 121*(13),
2452-2461. doi:blood-2012-08-451666 [pii];10.1182/blood-2012-08-451666 [doi]

van der Horst, A., de Vries-Smits, A. M., Brenkman, A. B., van Triest, M. H., van den Broek, N.,
Colland, F., . . . Burgering, B. M. (2006). FOXO4 transcriptional activity is regulated by
monoubiquitination and USP7/HAUSP. *Nat Cell Biol, 8*(10), 1064-1073. doi:10.1038/ncb1469

van der Knaap, J. A., Kumar, B. R., Moshkin, Y. M., Langenberg, K., Krijgsveld, J., Heck, A. J., . . .
Verrijzer, C. P. (2005). GMP synthetase stimulates histone H2B deubiquitylation by the
epigenetic silencer USP7. *Mol Cell, 17*(5), 695-707. doi:10.1016/j.molcel.2005.02.013

van Gosliga, D., Schepers, H., Rizo, A., van der Kolk, D., Vellenga, E., & Schuringa, J. J. (2007).
Establishing long-term cultures with self-renewing acute myeloid leukemia stem/progenitor
cells. *Exp. Hematol, 35*(10), 1538-1549.

Wang, H., Wang, L., Erdjument-Bromage, H., Vidal, M., Tempst, P., Jones, R. S., & Zhang, Y. (2004).
Role of histone H2A ubiquitination in Polycomb silencing. *Nature, 431*(7010), 873-878.
doi:10.1038/nature02985 [doi];nature02985 [pii]

Wang, R., Taylor, A. B., Leal, B. Z., Chadwell, L. V., Ilangovan, U., Robinson, A. K., . . . Kim, C. A. (2010).
Polycomb group targeting through different binding partners of RING1B C-terminal domain.
*Structure, 18*(8), 966-975. doi:S0969-2126(10)00220-0 [pii];10.1016/j.str.2010.04.013 [doi]

Wong, S. J., Gearhart, M. D., Taylor, A. B., Nanyes, D. R., Ha, D. J., Robinson, A. K., . . . Kim, C. A.
(2016). KDM2B Recruitment of the Polycomb Group Complex, PRC1.1, Requires Cooperation
between PCGF1 and BCORL1. *Structure*. doi:S0969-2126(16)30222-2
[pii];10.1016/j.str.2016.07.011 [doi]

Wu, X., Johansen, J. V., & Helin, K. (2013). Fbxl10/Kdm2b Recruits Polycomb Repressive Complex 1 to
CpG Islands and Regulates H2A Ubiquitylation. *Mol. Cell, 49*(6), 1134-1146. doi:S1097-
2765(13)00049-X [pii];10.1016/j.molcel.2013.01.016 [doi]

Ye, M., Tang, Y., Tang, S., Liu, J., Wu, K., Yao, S., . . . Tan, W. (2015). STIP is a critical nuclear
scaffolding protein linking USP7 to p53-Mdm2 pathway regulation. *Oncotarget, 6*(33), 34718-
34731. doi:10.18632/oncotarget.5303

  **Figures and Figure legends**

**Figure 1. The deubiquitinase USP7 interacts with non-canonical PRC1.1 proteins**

(A) Visualization of KDM2B interactome using Cytoscape, categorized by Molecular Function as

annotated by DAVID.

(B) Gene Ontology analysis of KDM2B interactome shown as Biological Process.

(C) Schematic representation of the non-canonical PRC1.1 complex, including USP7. PRC1.1

preferentially binds to non-methylated CpG islands via the ZF-CxxC domain of KDM2B. The

ubiquitination of histone H2A on lysine 119 is mediated by RING1A/B. PRC1.1 can be recruited to

active genes in leukemic cells, indicated by H3K4me3.

(D) Identification of USP7 and other PRC1.1 proteins by LC-MS/MS in KDM2B-EGFP, PCGF1-EGFP and

EGFP-RING1B pull outs from nuclear extracts in K562 cells. The numbers indicate MS/MS counts for

each interacting protein corrected for expected peptides based on *in silico* digests (x10).

(E) Validation of USP7 and RING1B interactions as analysed by Western blot in canonical PRC1

(PCGF2/PCGF4/CBX2/RING1B) and non-canonical PRC1.1 (PCGF1/RING1B) pull outs. Input (I), non-

bound (NB) and bound (B) fractions are shown.

**Figure 2. USP7 is essential for PRC1.1 integrity**

(A) Purification of his-tagged ubiquitinated proteins under denaturing conditions in EGFP-RING1B and

PCGF1-EGFP cells treated with DMSO (-) or P22077 (+) for 24h followed by Western blot analysis.

(B) Label-free quantification (LFQ) intensities of EGFP pull outs on EGFP-RING1B (left) and PCGF1-

EGFP (right) interactions in control (DMSO, 0.1%) and USP7i treated cells are illustrated as Volcano

plot. The fold change difference (log2) of the EGFP pull out in control versus USP7i is plotted against

the –log10 t-test p-value (y-axis). The bait protein (RING1B/PCGF1) is highlighted in green, PRC1.1

proteins in orange and UBB protein in purple.

(C) Intensity based absolute quantification (iBAQ) values for several identified PRC1.1 proteins are

shown in USP7i/control EGFP pull outs relative to RING1B (left) and PCGF1 (right). Data are shown as

mean ± SD (n=2 or 3).

(D) Western blot of EGFP pull outs on PCGF1-EGFP and EGFP-RING1B in the absence (-) or presence

(+) of P22077 for 72h probed with antibodies for EGFP, RING1B and PCGF1. Input (I), non-bound (NB)

and bound (B) fractions are shown.

(E) Mean fluorescent intensity (MFI) analysis of EGFP-RING1B and PCGF1-EGFP K562 cells in control

or USP7i treated cells at 72h.

**Figure 3. Loss of PRC1.1 occupancy and H2AK119ub at target loci upon USP7 inhibition**

(A) ChIP-qPCRs on control or 72h P22077 (USP7i) treated K562 cells for endogenous KDM2B, PCGF1-

EGFP, EGFP-RING1B (B) H2AK119ub and H3K4me3 on several PRC1.1 loci previously identified by

ChIP-seq. Error bars represent SD of technical triplicates.

(C) ChIP-qPCRs on control or USP7i treated cells for 4h/8h/16h ( ➟ ) for H2AK119ub, KDM2B-

EGFP, EGFP-RING1B (D) H3K4me3 and H3K27ac on five PRC1.1 loci. Error bars represent SD of

technical triplicates.

**Figure 4. USP7 inhibition leads to downregulation of a subset of PRC1.1 active target genes**

(A) Venn diagram showing overlap of genes downregulated by more than two fold after 24h of USP7

inhibition with previously identified PRC1.1 target genes.

(B) Gene Ontology analysis of downregulated genes identified as PRC1.1 targets (73) or Polycomb

independent targets (224) shown as Biological Process.

(C) ChIP-seq profiles of TOP2B, SIN3A, CHD1 and MYC as representative examples for PRC1.1 target

genes (73). Our ChIP-seq tracks are shown for GFP (control), H2AK119ub, H3K27me3, PCGF1, PCGF2,

PCGF4, CBX2, RING1A, RING1B and KDM2B. ChIP-seq tracks for H3K4me3, H3K36me3, RNAPII and

H3K27ac were downloaded from ENCODE/Broad.

(D) ChIP-qPCRs in control or 72h USP7i for KDM2B, PCGF1-EGFP, EGFP-RING1B and H2AK119ub or

16h USP7i for H3K4me3 and H3K27ac.

(E) Validation of relative gene expression levels of PRC1.1 target genes by qRT-PCR in control and

P22077 treated cells (24h). Data are represented as mean ± SD.

**Figure 5: Sensitiviy of primary AMLs towards USP7 inhibition**

(A) Cumulative cell growth of different AML cell lines in control (DMSO) or P22077 treated (10 µM

and 20 µM) conditions.

(B) Cumulative cell growth of primary AML patient cells (n=3) grown on MS5 co-culture treated with

DMSO (control) or P22077 (20 µM) performed in duplicate.

(C) Analysis of both cycling and quiescent AML cells by Hoechst/PyroninY stainings within the

CD34[+]/CD38[+] and CD34[+]/CD38[-] populations (D) or whole population (AML3). Cells were treated with

DMSO (control) or P22077 (20 µM) for 3 days while AMLs were grown on MS5 co-culture. Absolute

cell numbers were calculated in $G_0$, $G_1$ and $S/G_2/M$ fractions.

**Figure 6: Evaluation of USP7 inhibition in xenograft leukemia mouse model**

(A) Experimental setup of our human CB MLL-AF9 xenograft mouse model. Here 5 x $10^4$ MLL-AF9

EGFP cells from a primary leukemic mouse were IV injected into secondary recipients (n=11).

(B) Peripheral blood analysis of MLL-AF9 EGFP/CD45$^+$/CD19$^+$ cells, three weeks after injection prior to

treatment (left) and 2.5 weeks following treatment (right). Mice were treated daily with either DMSO

as control (n=5) or 20 mg/kg P22077 (n=6).

(C) Peripheral blood chimerism levels of control and P22077 (20 mg/kg) treated mice over the course

of the experiment. Treatment was started at day 28 (4 weeks post-transplant) indicated with the

arrow.

(D) Kaplan-Meier curve showing the survival of DMSO (n=5) and P22077 (n=6) treated mice.

Statistical analysis was performed using a log-rank test.

(E) Cumulative cell growth of CB MLL-AF9 cells grown under myeloid permissive conditions in liquid

and co-culture on MS5, treated with DMSO (control) or P22077 (15 and 20 μM).

(F) ChIP-qPCR for KDM2B in CB MLL-AF9 cells treated with DMSO or P22077 (48h) under either liquid

culture or MS5 co-culture conditions.

A

936 **Figure supplement 1. Sensitivity of lymphoid MLL-AF9 cells grown on MS5 for P22077**

937 (A) Lymphoid MLL-AF9 (GFP) co-culture on murine stromal cells treated with DMSO or P22077 (20

938 µM), day 6. Lymphoid MLL-AF9 cells typically form cobblestone areas (left). Absolute cell numbers

939 corrected for human CD45[+] and GFP[+] are shown 10 days upon DMSO or P22077 (20 µm) treatment

940 (right).