Title: Emergence of binocular disparity selectivity through Hebbian learning

Author details:

Tushar Chauhan*[1,2] (Tushar.Chauhan@cnrs.fr)

Timothée Masquelier[1,2] (Timothee.Masquelier@cnrs.fr)

Alexandre Montlibert[1,2] (Alexandre.Montlibert@cnrs.fr)

Benoit R. Cottereau*[1,2] (Benoit.Cottereau@cnrs.fr)

1.   Université de Toulouse, Centre de Recherche Cerveau et Cognition, Toulouse, France

2.   Centre National de la Recherche Scientifique, Toulouse, France

*Corresponding authors

Acknowledgements: We thank Simon J. Thorpe and Lionel Nowak for their comments on the
study. This research was supported by the IDEX Emergence (ANR-11-IDEX-0002-02, BIS-
VISION) and ANR JCJC (ANR-16-CE37-0002-01, 3D3M) grants awarded to B.R.C., and the
FRM grant (FRM: SPF20170938752) awarded to T.C.

Abstract

Neural selectivity in the early visual cortex strongly reflects the statistics of our environment

(Barlow, 2001; Geisler, 2008). Although this has been described extensively in literature

through various encoding hypotheses (Barlow and Földiák, 1989; Atick and Redlich, 1992;

Olshausen and Field, 1996), an explanation as to how the cortex develops the structures to

support these encoding schemes remains elusive. Here, using the more realistic example of

binocular vision as opposed to monocular luminance-field images, we show how a simple

Hebbian coincidence-detector is capable of accounting for the emergence of binocular, disparity

selective, receptive fields. We propose a model based on spike-timing dependent plasticity

(STDP) which not only converges to realistic single-cell and population characteristics, but also

demonstrates how known biases in natural statistics may influence population encoding and

downstream correlates of behaviour. Furthermore, we show that the receptive fields we obtain

are closer in structure to electrophysiological data than those predicted by normative encoding

schemes (Ringach, 2002). We also demonstrate the robustness of our model to the input dataset,

noise at various processing stages, and internal parameter variation. Taken together, our

modelling results suggest that Hebbian coincidence-detection is an important computational

principle and could provide a biologically plausible mechanism for the emergence of selectivity

to natural statistics in the early sensory cortex.

1. Introduction

Sensory stimuli are not only the inputs to, but also shape the very process of, neural computation (Hebbs, 1949; Barlow, 1961). A modern, more rigorous extension of this idea is the efficient coding theory (Barlow and Földiák, 1989; Atick and Redlich, 1992), which postulates that the computational aim of early sensory processing is to use the least possible resources (neurones, energy) to code the most informative features of the stimulus (information efficiency). A direct corollary to the efficient coding hypothesis is that if the inputs signals are coded efficiently, the statistical consistencies in the stimuli should then be reflected in the organisation and structure of the early cortex (Geisler, 2008). The largest body of work on the efficient coding principle lies within the visual sensory modality. In the context of vision, a number of studies have shown that information-theoretic constraints do indeed predict localised, oriented and band-pass representations, akin to those reported in the early visual cortex (Olshausen and Field, 1996). Over the years, several studies have shown how properties of natural images can not only explain neural selectivity (Olshausen and Field, 1996, 1997; Furmanski and Engel, 2000; Karklin and Lewicki, 2009; Samonds et al., 2012; Okazawa et al., 2015), but also predict behaviour (Webster and Mollon, 1997; Howe and Purves, 2002; Geisler, 2008; Geisler and Perry, 2009; Girshick et al., 2011; Cooper and Norcia, 2014; Burge and Geisler, 2015; Sebastian et al., 2017).

Nearly all these studies, however, rely on images of natural scenes acquired using a single camera – effectively using a 2-D projection of a 3-D scene. Although this representation captures the luminance-field statistics of the scene; however, it does not fully reflect how visual data is acquired by the human brain. Humans are binocular, and use two simultaneous retinal projections which enable the sensing of disparity (differences in retinal projections in the two eyes). Disparity, in turn, can be used to make inferences about the 3-D structure of the scene such as calculations of distance, depth and surface-slant. Despite critical results in the analysis of luminance-field statistics, only a handful of studies (Hoyer and Hyvärinen, 2000; Burge and Geisler, 2014; Hunter and Hibbard, 2015; Goncalves and Welchman, 2017) have attempted to analyse the relationship between binocular projections of natural scenes and the properties of binocular neurones in the early visual system. Furthermore, these studies investigated the relationship between natural scenes and cortical selectivity in terms of a global optimisation problem which is *solved* in the adult brain, leading to cortical structures which encode relevant

statistics of the stimuli. However, the question as to how these encoding schemes might actually

emerge in the early sensory cortex remains, as yet, unanswered (Stanley, 2013). Here, we show

how simple coincidence detection based on spike-timing dependent plasticity (Bi and Poo,

1998; Caporale and Dan, 2008) (STDP) could offer a biologically plausible mechanism for

arriving at neural population responses close to those reported in the early visual system. We

endow a neural network with a Hebbian STDP rule, and find that an unsupervised exposure of

this network to natural stereoscopic stimuli leads to a converged population which shows

single-cell and population characteristics close to those reported in electrophysiological studies

(Anzai et al., 1999; Durand et al., 2002; Prince et al., 2002b; Sprague et al., 2015). Moreover,

the emergent receptive fields differ from those obtained by optimisation-based methods, and

are more representative of those reported in literature (Ringach, 2002). Thus, taken together,

our findings suggest that known rules of synaptic plasticity are sufficient to explain the

relationship between biases reported in the early visual system and the statistics of natural

stimuli. Furthermore, they also provide a compelling demonstration of how simple biological

coincidence-detection (König et al., 1996; Brette, 2012; Masquelier, 2017) could explain the

emergence of selectivity in early sensory and multi-modal neural populations, both during and

after development.

2. Materials and Methods

Datasets

To simulate binocular retinal input, we chose two datasets – the Hunter-Hibbard dataset and

KITTI. The main results are reported for the Hunter-Hibbard database of stereo-images (Hunter

and Hibbard, 2015), available at https://github.com/DavidWilliamHunter/Bivis under the MIT

license which guarantees free usage and distribution. This database consists of 139 stereo-

images of natural scenes relevant to everyday binocular experience, containing objects like

vegetables, fruit, stones, shells, flowers, and vegetation. The database was captured using a

realistic acquisition geometry close to the configuration of the human visual system while

fixating straight ahead with zero elevation (Figure 1A). The distance between the two cameras

was close to the human inter-ocular separation (65 mm), and the two cameras were focussed at

realistic fixation-points from 50 cm to several metres away from the cameras. The lenses had a

large field of view (20°), enabling them to capture binocular statistics across a sizeable part of

the visual field. Figure 1A also shows a red-cyan anaglyph of one representative scene from the

database, while Figure 1B shows the images captured by the left and right cameras. Due to the
geometry of the cameras, there are subtle differences in the two images (disparity) which
provide important information about the 3-D structure of the scene. The mimicking of realistic
acquisition geometry is crucial for capturing disparity statistics which resemble those
experienced by human observers. For comparison, we also report results using the KITTI
database (available at http://www.cvlibs.net/datasets/kitti/) which uses parallel cameras – a
geometry which does not correspond to the human visual system (see Figure 6C and related
discussion for more details).

Figure 1: Processing pipeline. **A.** Acquisition geometry for the Hunter-Hibbard database. The set-up mimicked the typical geometry of the human visual system (see more details in the text). This geometry is crucial for capturing realistic disparity statistics from natural scenes. A stereoscopic reconstruction (red-cyan anaglyph) of a sample from the dataset is also shown. **B.** Input images captured by the left and the right camera for the scene shown in **A**. The subtle differences in the images (disparity) provide important cues about the 3-D structure of the scene. Depending on the simulation, a region-of-interest (ROI) in the visual field was identified, and inputs were sampled only within this ROI (foveal ROI illustrated in purple). Each sample consisted of two patches with equal retinal coordinates in the left and the right eye (illustrated in red and green respectively). All simulations presented in this paper were run using $N = 100,000$ samples each. **C.** The processing pipeline. The samples were processed

sequentially, one per iteration. The processing of the i^{th} sample is shown in a dashed box. First, the ON/OFF

processing in the LGN was modelled using difference-of-Gaussian filters in the two eyes – leading to left-on, left-

off, right-on and right-off maps. The activations of the four maps were then converted to first-spike relative

latencies using a simple inverse operation. These first-spikes constituted the input to the V1 layer. The V1 layer

consisted of 300 neurones connected to the LGN maps through Hebbian synapses endowed with spike-timing-

dependent-plasticity (STDP). A lateral inhibition scheme was also implemented to ensure no more than one

neurone fired per iteration. The first-spikes from the i^{th} sample were propagated through the synapses, and this

was followed by a change in synaptic weights dictated by the STDP algorithm (see text for details about the

implemented STDP model). This concluded the processing of the i^{th} sample, and was followed by the $(i + 1)^{th}$

sample, which was processed in a similar fashion.

Input sampling

Random locations in the scenes were sampled to provide inputs to our model. In each

simulation, the sampling was restricted to a specific region of interest (ROI). In this paper, we

present results from four main ROIs – foveal (eccentricities less than 3°), peripheral

(eccentricities larger than 6°), upper hemi-field, and lower hemi-field. Figure 1B shows the

foveal ROI shaded in purple. Each sample consisted of two patches corresponding to the two

eyes, both centred at the same random retinal coordinates. Figure 1B shows a random sample,

with the left patch outlined in red and the right patch in green. The sizes of the patches varied

with the ROI and can be interpreted as the initial dendritic receptive field of the V1 neurones,

which was subsequently pruned by STDP. In the simulations presented in this paper, the foveal

patches were 3° × 3° while the peripheral patches were 6° × 6°. 100,000 input samples were

used in each simulation.

To simulate the fixation behaviour of typical strabismic amblyopes (Figure 3B), the left and

right patches of each sample were no longer constrained to be centred at the same retinal

coordinates, thereby introducing a misalignment between the two eyes (while still keeping

within the ROI in each eye). The retinal correspondence between the ON and OFF LGN layers

of the same eye was still maintained.

Modelling of the retinogeniculate pathway

The computational pipeline employed by the model is shown in Figure 1C. The 100,000

samples were presented sequentially. The first stage of the model implemented the processing

in the retinogeniculate pathway. The patches were convolved with ON and OFF centre-surround

kernels to simulate the computations performed by the retinal ganglion cells (RGCs) and the

lateral geniculate nucleus (LGN). The filters were implemented using Difference of Gaussian

(DoG) kernels, and their receptive field sizes were chosen to reflect the size of representative

LGN centre-surround magnocellular receptive fields (Solomon et al., 2002) – 0.3°/1°

(centre/surround) for foveal simulations and 1°/2° (centre/surround) for peripheral simulations.

The activity of each kernel can roughly be interpreted as a retinotopic contrast map of the

sample. Four maps were calculated, corresponding to ON- and OFF-centre populations in the

left and right monocular pathways. The activity of each sample was thresholded such that, on

average, only a small portion (10%) of the LGN units fired. This activity was converted to

relative first-spike latencies through a monotonically decreasing function (a token function $y =$

$1/x$ was chosen, but all monotonically decreasing functions are equivalent (Masquelier and

Thorpe, 2007)), thereby ensuring that the most active units fired first, while units with lower

activity fired later, or not at all. Latency-based encoding of stimulus properties has been

reported extensively in the early visual system (Celebrini et al., 1993; Gawne et al., 1996;

Albrecht et al., 2002; Gollisch and Meister, 2008; Shriki et al., 2012), and allows for fast and

efficient networks describing early visual selectivity (Thorpe et al., 2001; Masquelier and

Thorpe, 2010; Masquelier, 2012). These trains of first- spikes (represented by their latencies)

from the random samples constituted the input to the STDP network.

V1: STDP neural network

The samples were presented to the network sequentially (Figure 1C). For each sample, the first-

spikes from the LGN were propagated through plastic synapses to a V1 population of 300

integrate and fire neurones. The initial dendritic receptive fields of the neurones were three

times the size of the LGN filters (foveal: $3° \times 3°$; peripheral: $6° \times 6°$). At the start, each neurone

was fully connected to all LGN afferents within its receptive field through synapses with

randomly assigned weights between 0 and 1. The weights were restricted between 0 and 1

throughout the simulation. The non-negative values of the weights reflect the fact that thalamic

connections to V1 are predominantly excitatory in nature (Ferster and Lindström, 1983; Tanaka,

1985).

Figure 1C shows the processing for the i^{th} sample in a dashed box. After the LGN spike

propagation, the synaptic weights were updated using an unsupervised multiplicative STDP

rule(Gütig et al., 2003). For a synapse connecting a pair of pre- and post-synaptic neurones, this

rule is described by:

$$\Delta w = \begin{cases} -\alpha^- \cdot w^{\mu^-} \cdot K(\Delta t, \tau_-), & \Delta t \leq 0 \\ \alpha^+ \cdot (1-w)^{\mu^+} \cdot K(\Delta t, \tau_+), & \Delta t > 0 \end{cases}$$

Here, a pre- and post- synaptic pair of spikes with a time difference Δt introduce a change Δw

in the weight of the synapse which is given by the product of a temporal filter K and a power

function of its current weight w. When a post-synaptic spike occurs shortly after a pre-synaptic

spike ($\Delta t > 0$), there is a strengthening of the synapse – also called long-term potentiation

(LTP). Conversely, when the post-synaptic spike occurs before the pre-synaptic spike ($\Delta t \leq$

0), or in the absence of a pre-synaptic spike, there is a weakening of the synapse or a long-term

depression (LTD). The LTP and LTD are driven by their respective learning rates α^+ and α^-.

The learning rates are non-negative ($\alpha^\pm \geq 0$), and determine the maximum amount of change

in synaptic weights when $\Delta t \to \pm 0$. The parameters $\mu^\pm \in [0, 1]$ describe the degree of non-

linearity in the LTP and LTD update rules. In practice, a non-linearity ensures that the final

weights are graded and prevents convergence to bimodal distributions saturated at the upper

and lower limits (0 and 1 in our case).

For the network used in the present study, the learning rates were fixed with $\alpha^+ = 5 \times 10^{-3}$

and $\alpha^+/\alpha^- = (4/3)$. The rate ratio α^+/α^- is crucial to the stability of the network, and was

chosen based on previous work demonstrating STDP based visual feature learning (Masquelier

and Thorpe, 2007). Our simulations show that the results presented in this paper are robust for

a large range of this parameter (see Figure 6B for more details). Furthermore, we used a high

nonlinearity for the LTP process ($\mu^+ = 0.65$) to ensure that we were able to capture fading

receptive fields through continuous weights, and used an almost additive LTD rule ($\mu^- = 0.05$)

to ensure pruning of continuously depressed synapses. In both LTP and LTD updates the

weights were maintained in the range $w \in [0, 1]$. Since our network relies only on first-spike

latencies, the temporal filter K was approximated as a step-function ($\tau_\pm \to \infty$), making the

update rule very efficient and fast. Furthermore, lateral inhibition through a winner-take-all

scheme (Maass, 2000) was implemented such that, if any neurone fired during a certain

iteration, it simultaneously prevented other neurones in the population from firing until the next

sample was processed. This scheme leads to a sparse neural population where the probability

of any two neurones learning the same feature is greatly reduced.

Analysis of converged receptive fields

The post-convergence receptive fields of the STDP neurones were approximated by a linear

summation of the afferent LGN receptive fields weighted by the converged synaptic weights.

If a weight w_{ij} connected the i^{th} neurone to the j^{th} LGN afferent with receptive field $\boldsymbol{\psi}_j$, the

receptive field ξ_i of the neurone was estimated by:

220
$$\xi_i \approx \sum_{j \in LGN} w_{ij} \boldsymbol{\psi}_j$$

221 The receptive fields calculated using this method are, in principle, similar to point-wise

222 estimates of the receptive field calculated by electrophysiologists. To better characterise the

223 properties of these receptive fields, they were fitted by two dimensional Gabor functions. These

224 Gabor functions are given by:

225
$$G(k, f, \theta, \phi, \boldsymbol{x}, \boldsymbol{\sigma}) = k * \boldsymbol{S}(f, \theta, \phi) * \boldsymbol{G}(\boldsymbol{x}, \boldsymbol{\sigma}, \theta)$$

where, k is a measure of the sensitivity of the Gabor, $\boldsymbol{S}$ is a 2-D sinusoid propagating with

frequency f and phase ϕ in the direction θ, and $\boldsymbol{G}$ is a 2-D Gaussian envelope centred at $\boldsymbol{x}$, with

size parameter $\boldsymbol{\sigma}$, also oriented at an angle θ. A multi-start global search algorithm was used to

estimate optimal parameters in each eye for each neurone. The process was expedited by using

the frequency response of the cell (Fourier amplitude) to limit the search-space of the frequency

parameter. For each neurone, the goodness of the Gabor fit was characterised using the

maximum of the R^2 goodness-of-fit values in the two eyes:

233
$$R^2_{max} = \max \{R^2_{Left}, R^2_{Right}\}$$

This ensured that the goodness-of-fit calculations were independent of whether the unit was

monocular or binocular.

After the fitting procedure, the Gabor parameters were then used to characterise the structure

and symmetry of the receptive field in the dominant eye (defined as the eye with the higher

value of the Gabor sensitivity-parameter k). This was done by calculating the symmetry-

invariant coordinates proposed by Ringach (2002). These coordinates (referred to as Ringach
coordinates or RC from here on) are derived from the Gabor parameters by:

242
$$\begin{bmatrix} n_x \\ n_y \end{bmatrix} = \begin{bmatrix} \sigma_x \\ \sigma_y \end{bmatrix} f$$

where, σ_x is the spread of the Gaussian envelope along the direction of sinusoidal propagation,
σ_y is the Gaussian spread perpendicular to σ_x, and f is the frequency of the sinusoid. While n_x
is a measure of the number of lobes in the receptive field, n_y reflects the elongation of the
receptive field perpendicular to the sinusoidal carrier. Typical cells reported in cat and monkey
have both RC coordinates (n_x and n_y) less than 0.5 (Ringach, 2002).

The responses of the converged neurones to disparity (disparity tuning curves or DTCs) were
characterised using two methodologies: binocular correlation (BC) of the left and right
receptive fields, and estimation of responses to random dot stereograms (RDSs). In the BC
method, we estimated the DTCs by cross-correlating the receptive fields in the left and the right
eye along a given direction. In the RDS method, we estimated the DTCs by calculating
responses of the converged neurones to random dot stereograms (RDSs) at various disparities.
The RDS patterns used for testing were perfectly correlated, with a dot size of about $15'$ visual
angles, and a dot density of 0.24 (i.e., 24% of the stimulus was covered by dots, discounting
any overlap). The DTC was calculated by averaging the response over a large number of
stimulus presentations ($N = 10,000$ reported here).

The responses were further quantified by a Binocularity Interaction Index (Ohzawa and
Freeman, 1986; Prince et al., 2002b) (BII) given by:

262
$$BII = \frac{R_{max} - R_{min}}{R_{max} + R_{min}}$$

where R_{max} and R_{min} are maximum and minimum responses on the disparity tuning curve. BII
values close to zero indicate that the neural response is weakly modulated by disparity, while
higher values indicate a sharper tuning to specific disparity values. Furthermore, the phase and
position disparity were estimated by methods commonly applied in electrophysiological studies
– one-dimensional Gabor functions were fitted to the DTCs (for this estimation, we used the
BC estimates as they are less subject to noise), and the phase of the sinusoid and the offset of

the Gaussian envelope from zero-disparity were taken as estimates of the phase and position
disparities respectively.

Network convergence

The evolution of the synaptic weights with time was characterised by using a Convergence
Index (CI). If $\Delta w_{ij}(t)$ is the change in the weight of the synapse connecting the i^{th} neurone to
the j^{th} LGN afferent at time t, the convergence index was defined as:

$$CI(t) = \frac{1}{N_{LGN} N_{V1}} \sum_{i \in V1} \sum_{j \in LGN} \left| \Delta w_{ij}(t) \right|$$

where N_{LGN} and N_{V1} are the number of LGN units and the number of the V1 neurones
respectively. The CI can be interpreted as a measure of the change in the synaptic weight
distribution over time. Assuming a statistically stable input to a Hebbian network, the number
of synapses undergoing a change in strength decreases with time. Thus, if our network is driven
to convergence, CI should reduce asymptotically with time. To report the convergence for a
single unit (say the i^{th} V1 neurone), a modified form of the above index (CI_i) was defined as:

$$CI_i(t) = \frac{1}{N_{LGN}} \sum_{j \in LGN} \left| \Delta w_{ij}(t) \right|$$

Robustness analysis

Biological systems show a considerable resilience to factors such as noise and internal
parameter variations (Burge and Jaini, 2017). We tested the robustness of our model using three
approaches – 1) by introducing timing jitter at both the input and the LGN level, 2) by varying
a key internal parameter of our network (the ratio of the LTP rate to the LTD rate), and 3) by
comparing our results obtained using the Hunter-Hibbard dataset (realistic acquisition
geometry) to results obtained using a dataset with non-realistic acquisition geometry (parallel
cameras).

The robustness of the model to noise was tested by adding external (image) and internal (LGN)

noise to the system. This simulated timing-jitter in the firing latencies from both the retina and

the LGN. Gaussian noise with an SNR varying between -3dB ($\sigma_{signal} = 0.5 \times \sigma_{noise}$) to 10dB

($\sigma_{signal} = 10 \times \sigma_{noise}$) was used, and the performance of the network at various noise levels

was characterised using two metrics – the convergence index (CI), and the R^2_{max} of Gabor

fitting. While the CI can be used to evaluate the stability of the converged synaptic connectivity,

the R^2_{max} shows how well the receptive fields can be characterised by Gabor functions. Taken

together, the CI and R^2_{max} characterise the robustness of the network in terms of both the

synaptic and the functional convergence.

The ratio of LTD to LTP rates ($\lambda = \alpha^-/\alpha^+$) is a critical parameter for the stability of the model

as it determines the number of pruned synapses after convergence. If the probabilities of LTD

and LTP events are p^- and p^+ respectively, the learning rate γ can be approximated as

308
$$\gamma \propto [\, p^+\alpha^+ + p^-\alpha^- \,]$$

Given our initial simulation values of the LTD and LTP rates α_0^- and α_0^+ respectively, we

defined $\alpha^- = (\alpha_0^-/\beta)$ and $\alpha^+ = \beta\alpha_0^+$, where the factor β was introduced to ensure that the

two rates changed such that the overall learning rate γ remained stable. The effect of the LTP

to LTD ratio on the convergence and functional stability of the network was then tested by

running simulations for λ in logarithmic steps from 0.01 to 100 and estimating both the CI and

R^2_{max} parameters. This simulated models where the LTD rates were from 1/100 to 100 times

the LTP rates.

After testing the robustness of our model to noise and internal parameter variations, the aim

was to test the model on a second dataset. As mentioned earlier, the Hunter-Hibbard dataset

used in all our simulations was chosen because it has an acquisition geometry close to the

human visual system. Thus, for this analysis we chose the KITTI dataset which was collected

using a markedly different acquisition geometry. This dataset uses parallel cameras mounted 54

cm. apart, leading to a highly non-ecological, zero-vergence geometry. The aim of this analysis

was to verify that the main results presented in this paper are not specific to the dataset, and that

the model is capable of capturing natural disparity statistics whist being robust to acquisition

geometry. Of course, in this case we did not expect the tunings of the converged neurones to

match those reported in human electrophysiology (see Figure 6C and related discussion for

more details).

Decoding of disparity

The ability of the converged network to encode disparity was tested through a decoding
paradigm. Random dot stereogram (RDS) stimuli at 11 equally spaced disparities between
$-1.5°$ and $1.5°$ were presented to the converged model, and the activity of the V1 layer was
used to train, and subsequently test, linear and quadratic discriminant classifiers. We chose RDS
stimuli because they do not contain any other information except disparity. A total of 10,000
stimuli per disparity were used, with a 70/30 training/testing split. 25-fold cross-validation
testing was performed to ensure robust results. In this article, we report the detection probability
(probability of correct identification) at each disparity. It must be noted that here, decoding
through classifiers is only an illustrative representation of perceptual responses as it is based on
inputs from very early visual processes and completely ignores important interactions such as
cortico-cortical connections and feedback.

Code accessibility

343   On acceptance of the manuscript for publication, the code will be freely made available on a
344   public server (ModelDB and gitlab). During the review process, the editor and the reviewers
345   will be given access to the code upon request.

3.   Results

In this section, we present the results from simulations (10^5 iterations per simulation) where
binocular stimuli from specific ROIs were presented to our model. In the model, retinal inputs
were filtered through an LGN layer, which was in turn connected to a V1 layer consisting of
300 integrate-and-fire neurones through STDP-driven synapses (see *Materials and Methods* for
details). The main results are shown for the foveal ROI, while other ROIs (peripheral, upper-
hemifield and lower-hemifield) are used to illustrate retinotopic biases.

Foveal population

356   In Figure 2A, we present a representative unit from a foveally trained population of 300 V1

neurones. The first row of Figure 2A shows the initial receptive field of the neurone. Since the weights were set to random values, the resulting receptive field has no specificity in terms of structure or excitatory/inhibitory sub-fields. The second row of the figure shows the receptive field of the same neurone after STDP learning from 10^5 samples (see also, Movie 1). The converged receptive field is local, bandpass, oriented. Clear on and off regions can be observed, and the receptive field closely resembles classical Gabor-like receptive fields reported for simple cells (Hubel and Wiesel, 1962; Jones and Palmer, 1987). Figure 2A also shows the Convergence Index (CI) of the neurone (bottom-left panel) which characterises the stability of its synaptic weights over the learning period. We find that the fluctuations in the average synaptic strength decrease over time to a very small value − indicating that the weight distribution has converged.

Figure 2: **A.** A representative neurone from the simulated population. Receptive fields in the left and the right eye before (first row), and after (second row) presentation of 10^5 samples. The responses are scaled such that the maximum response is 1. The initial receptive fields are random, while the converged receptive fields show Gabor-like structure. The bottom-left panel shows the Convergence Index (CI), which measures the mean change in synaptic strength over time. Estimates of the Disparity tuning curve (DTC) for horizontal disparity are shown in the bottom-right panel. The neurone shown here is tuned to uncrossed (or positive) disparities. The DTC estimates using two techniques are presented − binocular correlation of the left and right receptive fields (red solid line), and averaging responses to multiple ($N = 10^4$) presentations of random dot stereogram (RDS) stimuli (grey dotted line). The standard error for the RDS DTCs is shown as a grey envelope around the average responses. The Binocular Integration Index (BII) is also indicated within the graph. BII is a measure of the sensitivity of the

neuronal response to disparity variations, with values close to 0 indicating low sensitivity, and values close to 1

indicating higher disparity sensitivity. See *Materials and Methods* for more details on the calculation of the

receptive fields, DTCs and BII. **B.** Converged receptive fields and DTCs for neurones showing zero-tuned (first

row), tuned near (second row) and phase-modulated (third row) responses to horizontal disparity.

In electrophysiology, a neurone with a binocular receptive field is often characterised in terms

of its response to binocular disparity (Freeman and Ohzawa, 1992; Prince et al., 2002b, 2002a),

i.e. its disparity tuning curve (DTC). The bottom-right panel of Figure 2A shows the DTC

estimates for the selected neurone, estimated using both the BC (solid red line) and the RDS

(dotted grey line, with standard error marked as a grey envelope) methods. The DTCs obtained

from BC were very similar to those obtained through RDS stimuli. As the BC DTCs are, in

general, less noisy, they were used to calculate all the disparity results presented in the article.

This neurone shows a selectivity for positive or uncrossed disparities, indicating that it is more

likely to respond to objects further away from the observer than fixation. It also has a high

Binocularity Interaction Index (Ohzawa and Freeman, 1986; Prince et al., 2002b) (BII) of 0.83,

indicating that there is a substantial modulation of its neuronal activity by binocular disparity

(BII values close to 0 indicate no modulation of neuronal response by disparity, while values

close to 1 indicate high sensitivity to disparity variations). Furthermore, Figure 2B shows three

other examples of converged receptive fields and the corresponding DTCs (one neurone per

row). The first neurone shows a tuning to zero-disparity (fixation), the second neurone to

crossed or negative disparity (objects closer than the fixation), and the third neurone shows a

tuning curve which is typically attributed to phase-tuned receptive fields. The receptive fields

have a variety of orientations and sizes, and closely resemble two-dimensional Gabor functions.

Receptive field properties

Since most converged receptive fields in the foveal simulation show a Gabor-like spatial

structure, we investigated how well an ideal Gabor function would fit these receptive fields.

Figure 3A shows the goodness-of-fit R^2 parameter for the left and the right eye for all 300

neurones in the foveal population, with the colour code corresponding to the value $R^2_{max} =$

$\max\{R^2_{left}, R^2_{right}\}$. This value is maximal of the R^2 parameter across the two eyes, and

indicates how well the Gabor model fits the receptive field irrespective of whether it is

monocular or binocular. A majority of the neurones lie along the diagonal in the upper right

quadrant, showing an excellent fit in both eyes; with only few monocular neurones lying in the

upper-left and bottom-right quadrants. It is also interesting to note that irrespective of the degree

of fit, the R^2 values for the left and the right eye are tightly correlated. This binocular correlation

is non-trivial and cannot simply be attributed to learning from a stereoscopic dataset, as the

inter-ocular correlations in the stimuli compete with intra-ocular correlations. To demonstrate

this, in Figure 3B we show neurones from two simulated populations – the normal foveal

population from Figure 3A (red dots), and a simulated amblyopic population (blue dots). Only

the neurones with a reasonably good Gabor-fit ($R^2_{max} \geq 0.5$) in each population are shown. The

neurones in the amblyopic population occupy the top-left and bottom-right quadrants

(indicating the neurones are monocular), as opposed to the binocular neurones from the normal

population in the top-right quadrant. In the amblyopic population, the projections in the two

eyes were sampled from different foveal locations within the same image (see *Materials and*

Methods), thereby mimicking the oculomotor behaviour exhibited by strabismic amblyopes.

Here, although neurones learn from left and right image-sets with the same widefield statistics,

due to the lack of local inter-eye correlations they primarily develop monocular receptive fields

– choosing features from either the left or the right eye.

Figure 3: Characterising a simulated population of 300 foveal V1 neurones. **A.** The goodness-of-fit R^2 criteria for Gabor-fitting in the left and the right eye. Each dot represents a neurone, while the filled colour indicates the quantity $R^2_{max} = \max\{R^2_{left}, R^2_{right}\}$, a measure of how well a Gabor function describes the receptive field irrespective of whether the neurone is monocular or binocular. **B.** Neurones ($R^2_{max} \geq 0.5$) from normal (red dots) and amblyopic (blue dots) populations trained on foveal stimuli (see *Materials and Methods* for more details). **C.** The orientation parameter for the Gabors fitted to the left and the right eye receptive fields. Only neurones with binocular receptive fields defined by the criterion $\min\{R^2_{left}, R^2_{right}\} \geq 0.5$ are shown. Note that the range of the angles is $[0°, 180°]$, with $0°$ being the same as $180°$. **D.** DTCs for the binocular neurones from **C.** were estimated by the binocular cross-correlation of the left and right-eye receptive fields under horizontal displacement. Gabors were fitted to the resulting curves, thereby giving a measure of the position and phase disparity in the receptive fields(Prince et al., 2002b, 2002a). This figure shows the position disparity plotted against the relative phase disparity. It also shows the histogrammes for the two types of disparities.

Figure 3C shows the receptive field orientations in the left and the right eye of binocular

neurones in the foveal population. Here, we define a binocular unit by using the criterion:

$\min\{R^2_{left}, R^2_{right}\} \geq 0.5$, i.e., both eyes fit the Gabor-model with an R^2 of at least 0.5. We see

that there is a strong correspondence between the orientation in the two eyes (0° and 180° are

equivalent orientations), with a large number of neurones being oriented along the cardinal

directions, i.e., either horizontally or vertically. Though not shown here, a similar

correspondence is found between the Gabor frequencies in the two eyes (the frequency range

is limited between 0.75 and 1.25 cpd, due to the use of fixed LGN receptive field sizes), with

an octave bandwidth of approximately 1.5 – a value close to those measured for simple-cells

with similar frequency peaks (De Valois et al., 1982). Furthermore, Figure 3D shows the

position and phase encoding of the horizontal disparities as derived by fitting Gabor functions

to DTCs estimated by the BC method (see legend and *Materials and Methods* for details). Both

position and phase are found to code for disparities between $-0.5°$ and $0.5°$ visual angle (phase

disparities were converted to absolute disparities using the frequency of the fitted Gabor), with

most units coding for near-zero disparity.

In addition to edge-detecting Gabor-like receptive fields, more compact, blob-like neurones

have also been reported in cat and monkey recordings (Jones and Palmer, 1987; Ringach, 2002).

These receptive fields are not satisfactorily predicted by efficient approaches such as ICA and

sparse coding (Ringach, 2002). In Figure 4, we present the distribution of Ringach-coordinates

(RC, a symmetry-invariant descriptor for simple-cell receptive-fields, see *Materials and*

Methods for details) for the dominant eye of our converged foveal population. The n_x and n_y

axes approximately code for the number of lobes and the elongation of the receptive-field

perpendicular to the periodic function (see *Materials and Methods* for more details). Most of

the receptive fields in our converged population lie within 0.5 units on the two axes (blue shaded

region) – two typical receptive fields in this region are shown (cyan markers). These receptive

fields are not very elongated and typically contain 1-1.5 cycles of the periodic function. We also

observe blob-like receptive fields (magenta markers) which display a larger invariance to

orientation. In addition, we show three receptive fields which lie further away from the main

cluster – along the n_x axis with multiple lobes (brown markers), and along the n_y axis with an

elongated receptive field (purple marker). This RC distribution is close to those reported in cat

and monkey recordings (Jones and Palmer, 1987; Ringach, 2002); while typically, RC

distributions found by optimisation-based coding schemes such as ICA and sparse coding tend

to lie to the right of this distribution (Ringach, 2002).

Figure 4: Ringach-coordinate (RC) distribution for the simulated foveal population (dominant eye). RC is a symmetry-invariant size descriptor for simple-cell receptive-fields (see *Materials and Methods* for a detailed description). High values along the n_x axes convey large number of lobes in the receptive field, while high values along the n_y axes describe elongated receptive fields. Receptive-fields for some representative neurones are also shown. The pink neurones are examples of blob-like receptive fields, while cyan markers illustrate 2-3 lobed neurones which constitute the majority of the converged population. Brown markers show neurones with a higher number of lobes, while the purple marker shows a neurone with an elongated receptive field. Most of the converged neurones lie within the square bounded by 0 and 0.5 along both axes (shaded in blue), something that is also observed in electrophysiological recordings in monkey and cat (Ringach, 2002).

Biases in natural statistics and neural populations

The disparity statistics of any natural scene show certain characteristic biases which are often reflected in the neural encoding in the early visual pathway. In this section, we investigate whether such biases are also found in our neural populations. Figure 5A shows a histogram of the horizontal disparity in two simulated populations of neurones, one trained on inputs from

the central field of vision (eccentricity ≤ 3°, shown in purple), and the other on inputs from the

periphery (6° ≤ eccentricity ≤ 10°, shown in green). The peripheral population shows a clear

broadening of its horizontal disparity tuning with respect to the central population (Wilcoxon

rank-pair test: $p = 0.046$). As one traverses from the fovea to the periphery, the range of

encoded horizontal disparities has indeed been shown to increase, both in V1 (Durand et al.,

2007), and in natural scenes (Sprague et al., 2015).

Figure 5: Retinotopic simulations and population biases in horizontal. **A.** Histograms and estimated probability

density functions for horizontal disparity in two simulated populations of 300 V1 neurones each. The first

population was trained on inputs sampled from the fovea (purple) while the second population was trained on

inputs from the periphery (green). **B.** Histogram and probability density estimates of two populations of 300

neurones each, trained on inputs from the upper (blue) and lower (red) hemifield. **D.** A meta-analysis(Sprague et

al., 2015) of results from electrophysiological recordings in monkey V1 neurones (820 neurones in total) by five

separate groups. The neurones were split in two groups – neurones with upper-hemifield receptive fields are in

blue, while those with receptive fields in the lower hemifield are in red.

Another statistical bias reported in retinal projections of natural scenes (Sprague et al., 2015) is

that horizontal disparities in the lower hemifield are more likely to be crossed (negative) while

the upper hemifield is more likely to contain uncrossed (positive) disparities. This bias is not

altogether surprising considering that objects in the lower field of vision are likely to be closer

to the observer than objects in the upper field (Hibbard and Bouzit, 2005; Sprague et al., 2015).

A meta-analysis (Sprague et al., 2015) of 820 V1 neurones from five separate

electrophysiological studies indicates that this bias could also be reflected in the binocular

neurones in the early visual system. Figure 5C shows the results of this meta-analysis with the

probability density for the neurones in the lower hemifield (in red) showing a bias towards

crossed disparities while the upper hemifield (in blue) shows a bias towards uncrossed

disparities. In comparison, Figure 5B presents histograms of the horizontal disparity in

binocular neurones from two simulated populations – one trained on samples from the lower

hemifield (red), and the other on the upper hemifield (blue). A comparison of Figure 5B and

Figure 5C shows that both the range and the peak-tuning of the two simulated populations

closely resemble those reported in electrophysiology. The peaks in Fig5B are at 0.17° for the

upper hemifield and −0.02° for the lower hemifield (Wilcoxon rank-pair test: $p =$

2.15×10^{-14}), while the meta-analysis by Sprague et al. (2015) shown in Fig5D reports the

values as 0.14° and −0.09° respectively.

Robustness to internal and external noise

In the results reported so far, the only source of noise are the camera sensors used to capture the

stereoscopic dataset. To further test the robustness of our model we introduced additive white

Gaussian noise $\sim N(0, \sigma_{noise})$ at two stages of the visual pipeline – external noise in the image,

and internal noise in the LGN layer. The noise was manipulated such that the SNR varied in

steps of approximately 3 dB from -3 dB ($\sigma_{noise} = 2 \times \sigma_{signal}$) to 10 dB ($\sigma_{noise} =$

$0.1 \times \sigma_{signal}$). Since the noise in these simulations induces timing jitter at the retinal and LGN

stages, they are especially relevant for our model which relies on first-spike latencies. In Figure

6A we show the behaviour of the system under internal (in green colours) and external noise

(red colours). The left and middle panels show the convergence index (CI) of the network,

plotted against the iteration number (training duration) for the internal and external noise

conditions respectively. The lightness of the colour indicates the level of noise (the darker the

colour, the higher the noise). The right panel shows the mean R_{max}^2, plotted against the SNR.

We see that the CI approaches zero asymptotically in all conditions, indicating that the weights

converge to a stable distribution for all levels of noise. The mean R_{max}^2, as expected, increases

with the SNR (the higher the noise, the less Gabor-like the receptive fields). We see that the

receptive fields converge to Gabor-like functions for a large range of internal and external noise,

with the R_{max}^2 dropping below 0.5 only at SNR levels below 3 dB ($\sigma_{noise} \geq 0.5 \times \sigma_{signal}$).

Figure 6: Robustness analysis. **A.** Internal and external noise. Additive white Gaussian noise (SNR levels between -3 and 10 dB) was introduced in the LGN activity (internal, in green colours), and the retinal input (external, in red colours). The left and middle panels show the Convergence Index (CI, see *Materials and Methods*) for the internal and external noise conditions respectively, plotted against the number of iterations (training period). In both graphs, the lightness of the curve codes for the SNR (the darker the curve, the higher the noise). For a stable synaptic convergence, the CI should approach zero asymptotically. The right panel shows the mean R^2_{max} for the simulations (internal noise in green and external noise in red), plotted against the SNR. The higher the value of the mean R^2_{max}, the closer the converged receptive fields are to Gabor-like receptive fields. **B.** Parameter variation. The ratio of the LTD rate (α^-) to the LTP rate (α^+), say λ, is a crucial parameter which determines the relative contributions of synaptic potentiation and depression to the overall learning rate. This parameter was varied logarithmically from 10^{-2} to 10^2 (i.e., $\alpha^- = \alpha^+/100$ to $\alpha^- = \alpha^+ \times 100$) to determine the robustness of the model to internal parameter variation. The left panel presents the CI for each level of λ, with purple colours coding for $\alpha^- \leq \alpha^+$ and orange/brown colours coding for $\alpha^- \geq \alpha^+$. The darker the colour, the closer the α^- to the α^+, with black being $\lambda = 1$. The right panel shows the mean R^2_{max} for the simulations plotted against λ, with the symbols

following the same colour coding as the left panel. An additional grey dot is added to indicate the value of λ (=

3/4) used for all the simulations presented in the paper (except this figure). The higher the mean R^2_{max}, the more

Gabor-like the converged receptive fields. **C.** A second dataset. The model was tested using a second dataset

(KITTI, see *Materials and Methods* for details). The left panel shows the acquisition geometry of the KITTI

dataset, and a cyan-red anaglyph reconstruction of a representative scene. The cameras were parallel and separated

by 54 cm. – a geometry that does not resemble the human visual system. The middle panel shows the R^2 for the

Gabor fit procedure on the left and right receptive fields of the converged neurones, with the colour of the symbol

coding for the R^2_{max} (compare to Figure 3A for the Hunter-Hibbard dataset). The right panel shows horizontal

disparity tunings for two populations of neurones trained on the upper (blue) and lower (red) hemi-fields of the

KITTI database. For comparison, the curves obtained using the Hunter-Hibbard database (Figure 5B) are plotted

in dashed lines using the same colour scheme.

Overall, these analyses demonstrate that our approach is robust to a large range of noise induced

timing-jitter at the input and LGN stages. While low noise conditions produce a synaptically

converged network with Gabor-like receptive fields, high noise conditions produce a network

which still converges, albeit to receptive fields which are no longer well-modelled by Gabor

functions.

Robustness to parameter variation

One of the key parameters of a Hebbian STDP model is the ratio (say, λ) of the LTD and LTP

rates (α^- and α^+ respectively). This parameter determines the balance between the weight

changes induced by potentiation and depression, and thus directly influences the number of

pruned synapses after convergence. The simulations presented in this article use $\lambda = 3/4$, a

value based on STDP models previously developed in the team (Masquelier and Thorpe, 2007).

In this section, we test the robustness of the model to variations in this important parameter.

The value of λ was varied between 0.01 and 100 in log-steps (see *Materials and Methods* for

the details). Figure 6B shows the results of these simulations. The left panel shows the CI,

plotted against the iteration number (training period). The purple colours code for $\alpha^- \leq \alpha^+$ and

the brown/orange colours code for $\alpha^- \geq \alpha^+$, with the intensity of the colour coding for the

deviation from $\lambda = 1$. Once again, the synaptic weights converge for all values of the parameter,

with convergence being quicker for higher values of λ. The right panel shows the mean R^2_{max}

plotted against λ. The same colour code as the left panel is used, and an additional grey symbol

indicates the performance of the model at $\lambda = 3/4$. The model produces Gabor-like receptive

fields for a very broad range of λ values (for $R^2_{max} \geq 0.5$ this range is roughly between 0.05

and 2), with the performance deteriorating for small values of λ (α^- is very low compared to

α^+) and saturating for $\alpha^- \geq 6\alpha^+$. This analysis shows that our model shows stable synaptic

convergence for all tested values of λ (Figure 6B, left panel), and a functional convergence to

Gabor-like receptive fields for a large range of the parameter (Figure 6B, right panel).

Robustness to dataset acquisition geometry

All our simulations (except this section) use the Hunter-Hibbard database which was collected

using an acquisition geometry close to the human visual system. The emergent receptive fields

are disparity-sensitive, and closely resemble those reported in the early visual system. In this

analysis, we prove that this emergence is not only a property of the dataset, but also of the

human-like acquisition geometry. To do this, the model was tested on another dataset (KITTI,

available at http://www.cvlibs.net/datasets/kitti/) which was collected using two widely-spaced

(54 cm apart), parallel cameras – an acquisition-geometry which does not resemble the human

visual system at all (left panel, Figure 6C). In the middle and right panel of Figure 6C, we

present two results from this series of simulations. The middle panel shows the R^2 of Gabor-

fitting in the two eyes (symbols colour coded by R^2_{max}) for a population trained on the foveal

region of the KITTI dataset. We observe both monocular (high R^2 in one eye, top-left or bottom-

right quadrants) and binocular (high R^2 in both eyes, top-right quadrant) neurones. In

comparison, the converged receptive fields for the Hunter-Hibbard dataset are mostly binocular

(Figure 3A). This is not surprising as the large inter-camera distance (54 cm as compared to 6.5

cm for the Hunter-Hibbard dataset) combined with the parallel geometry results in less overlap

in the left and right images of the KITTI dataset, which further leads to weaker binocular

correlations.

The right panel shows population disparity tunings for two separate populations trained on the

upper (blue) and lower (red) hemi-fields of the KITTI dataset. As expected, the lower-hemifield

population is tuned to disparities which are more crossed (negative) than the upper-hemifield

population (see Figure 3C and the section *Biases in natural statistics and neural populations*).

For comparison, the curves for the Hunter-Hibbard dataset (from Figure 3C) are plotted as

dotted lines. The parallel geometry of the cameras in the KITTI dataset results in predominantly

crossed disparities (the fixation point is at infinity, and hence all disparities are, by definition,

crossed). Thus, while the upper vs. lower hemifield bias (see section *Biases in natural statistics*

and neural populations) still emerges from our model, the population tunings no longer
correspond to those reported in electrophysiological recordings(Sprague et al., 2015) (see
Figure 3D).

Decoding disparity through V1 responses

Recently, the effect of natural 3D statistics on behaviour has been modelled using labelled
datasets in conjunction with task-optimised filters (Burge and Geisler, 2014) and feedforward
convolutional networks (Goncalves and Welchman, 2017). Typically, such models are
evaluated by how well they perform at classifying/detecting previously unseen data. In Figure
7A, we show the result (detection probability) of training two very simple classifiers on the
activity of the converged V1 layer from our model, using RDS stimuli at 11 equally spaced
disparities between −1.5° and 1.5° (see *Materials and Methods* for details). RDS stimuli were
used because they contain no other meaningful information except disparity. The first, a decoder
based on linear discriminant analysis (green), performs well above chance (dashed red),
especially for realistic values of disparity between −0.5° and 0.5°. Interestingly, when the
complexity of the decoder is increased slightly by including quadratic terms (blue), one
observes a substantial increase in discrimination performance. This is not surprising as our
neurones are linear units similar to simple cells, and a non-linear processing of their activity
makes the decoding units conceptually closer to sharply tuned complex cells (Ohzawa et al.,
1997; Cumming and DeAngelis, 2001; Henriksen et al., 2016).

Figure 7: Decoding disparity through random-dot stereograms (RDS). **A.** Output spikes of the V1 layer (foveal
simulation) to RDS stimuli at various disparities were used to train linear (green) and quadratic (blue) discriminant
classifiers. The RDS stimuli were chosen as they do not contain any other information except disparity. The
performance of these classifiers was then tested using different sets of labelled RDS stimuli (10000 total RDS

stimuli; 70% used for training, 30% used for testing; cross-validation using 25 folds). This plot shows the detection

probability (probability of correct identification), plotted against the disparity. The quadratic classifier performs

better than the linear classifier, while both classifiers perform above chance-level (in dashed red). **B.** Two

populations of V1 neurones trained on the upper (blue) and lower (red) hemi-fields were used to test a linear

classifier using the same stimuli as **A**. The detection probability of the two classifiers reflects the population bias

induced by the disparity statistics of the upper and lower hemifields (Figure 5B).

We have previously seen that V1 populations trained on the upper and lower hemi-fields show

systematic biases for uncrossed and crossed disparities respectively (see Figure 5B). In Figure

7B, we show the performance of two linear classifiers, trained on V1 populations from the upper

(blue) and lower (red) hemi-fields using the same method as Figure 7A. The crossed/uncrossed

disparity biases in the V1 layer are also transferred to the decoding performance of the

classifiers. Thus, our model predicts that biases in early binocular neural populations can drive

downstream disparity discrimination biases too – something which has indeed been reported in

human subjects (Sprague et al., 2015).

4. Discussion

We have demonstrated that unsupervised Hebbian learning from natural stereoscopic scenes

leads to a population of neurones which are capable of encoding binocular disparity statistics

in natural scenes. Our findings provide evidence that known mechanisms of neuronal Hebbian

plasticity (such as STDP) could play a key role in the encoding of the statistics of natural scenes

in the early visual pathway.

Starting from random receptive-fields, our neural population converges to Gabor-like receptive

fields found in V1 simple cells (see Movie 1 and Figure 2). This convergence is achieved by a

Hebbian pruning of the initial densely connected dendritic receptive field of size $3° \times 3°$. The

mean size of the converged binocular receptive fields when described using a Gabor fit is about

$1°$ (average 1σ of the Gaussian envelope). Although in the reported foveal population we started

with a $3° \times 3°$ receptive field, simulations with an initial receptive field size of $6° \times 6°$ yielded

very similar results. Thus, the pruning process is independent of the size of the initial dendritic

tree, and yields converged receptive fields which correspond to about 3 times the size of the

central ON/OFF region of the retinal ganglion receptive fields.

We find that the converged receptive fields in the two eyes show strong correlations for properties such as the orientation, size and frequency (Figure 3C shows the orientation, but similar results were found for size and frequency). This suggests that the selectivity of the neurones is driven by the inter-eye correlations, despite competition from within-eye correlations. Although correlation-based refinement of binocular receptive-fields has previously been proposed at a theoretical level (Berns et al., 1993), our results prove that natural scenes and the geometry of the human visual system are indeed capable of providing the strength of inter-ocular correlation required for binocularity to occur. This point is further illustrated by our simulation of the oculomotor behaviour in strabismic amblyopes where the inputs are binocular, but the inter-ocular correlations are weakened due to uncoordinated eye fixations. Whilst this population of neurones also converges to Gabor-like receptive fields, a majority of the neurones are found to be monocular (Figure 3B). This is not altogether surprising as STDP learning from monocular images is known to produce oriented Gabor-like receptive fields (Delorme et al., 2001; Masquelier, 2012). These results also echo findings in monkeys reared with amblyopic deficits (Movshon et al., 1987; Kiorpes et al., 1998) where the binocularity of V1 neurones is found to be severely restricted.

The converged neurones show selectivity to zero, near and far disparities between −0.5° and 0.5° (Figure 3D shows this for position disparity, and a similar range is observed when relative phase values are converted to visual angles), which is in line with electrophysiological findings (Poggio et al., 1985; Prince et al., 2002a), and gives the population the theoretical capacity to encode a large range of realistic disparity values (Sprague et al., 2015). Through retinotopic simulations, we give two further examples which show that when allowed to learn from specific regions of the visual field, our model produces population biases which correspond closely to retinotopic biases reported in electrophysiology. In the first example (Figure 5A), we demonstrate that neurones trained on stimuli from peripheral eccentricities are selective to a larger range of horizontal disparities (Durand et al., 2002) when compared with neurones trained on foveal stimuli. In the second example (Figure 5B) we use two neural populations (upper and lower hemi-field) to show how Hebbian learning can lead to neural populations which reflect the intuitive bias that objects in the lower hemi-field are more likely to be closer than fixation and objects in the upper hemifield are likely to be further away. This bias has been

demonstrated through free-viewing tasks, and is also reflected in neural responses at both the

single-unit (Sprague et al., 2015) and macroscopic (Nasr and Tootell, 2016) levels. Through

these simulations, we show that retinotopic learning based on a Hebbian approach could explain

how biases in natural statistics can transfer to neural populations in the early visual system.

Few studies have previously characterised the relationship between natural statistics and

binocular disparity responses in the visual cortex. Based on their approach to learning, they can

roughly be classified under two main headings – unsupervised strategies (Hoyer and Hyvärinen,

2000; Hunter and Hibbard, 2015) which learn from unlabelled stimuli, and supervised

approaches (Burge and Geisler, 2014; Goncalves and Welchman, 2017) which use labelled data

to learn specific properties of the scene. Whilst results from unsupervised learning can be

interpreted as possible encoding schemes for the stimuli, supervised approaches optimise model

performance on specific tasks such as disparity or contrast discrimination. Both these

approaches solve an optimisation problem which may involve global properties like entropy

and mutual information, or task-specific metrics such as classification accuracy. Although our

model does not require supervision, it differs critically from traditional, objective-function

based unsupervised approaches as it does not necessarily converge towards a static global

optimum - instead, it can be interpreted as a self-regulating coincidence detector based on causal

Hebbian updates. In our case, the convergence towards an optimum is a result of the stability

of natural statistics across samples.

A commonly employed unsupervised approach to studying the encoding of 3D natural statistics

is the Independent Component Analysis (ICA) (Hoyer and Hyvärinen, 2000; Hunter and

Hibbard, 2015), a technique which maximises the mutual information between the stimuli and

the activity of the encoding units. Here, we mention three critical differences which separate

our results from those obtained using ICA. First, Hebbian learning could potentially be sub-

optimal when viewed in terms of the mutual-information based objective functions used for

ICA. For example, Hebbian learning leads to blobby receptive fields which are highly feature

invariant, and a very inefficient choice for encoding natural images. Second, Hebbian learning

is usually unable to converge to features which do not recur in the input, whereas efficient

coding approaches learn the optimal basis representations of the data which could be very

different from any individual stimulus. As an example, the converged population in our model

contains very few neurones with antiphase receptive fields in the two eyes (Figure 3D). This is

in line with electrophysiological data collected in monkeys, cats and the barn owl (Prince et al.,

2002a). This result is also very intuitive because in everyday experience, the probability of a

given object projecting antiphase features in the two eyes is very low. In contrast, ICA models

usually predict a higher number of antiphase receptive fields in the neural population (Hunter

and Hibbard, 2015). Third, the receptive fields obtained through the Hebbian model are closer

in structure to electrophysiological data than those obtained by ICA (Figure 4) – for example,

they have fewer lobes and show broader orientation tuning. Thus, although ICA representations

offer a highly optimal encoding of the data in terms of information transfer, they are inconsistent

with electrophysiological findings which often include neurones which are sub-optimal in terms

of any particular coding strategy (Ringach, 2002).

As initially pointed out by Barlow (1961), although encoding schemes offer explanations as to

how scene statistics may be represented in early processing, the precise nature of the encoding

must be capable of driving upstream processes which ultimately determine behaviour. We

demonstrate (Figure 7) that the activity of the converged V1 population in our model is capable

of driving simple classifiers to an above chance performance. This suggests that disparity

encoding in the early visual system does not necessarily need supervised training, and an

unsupervised feedforward Hebbian process can lead to neural units whose responses can be

interpreted in terms of 3D percept through downstream processing. It must be noted that here,

decoding through a classifier is only presented as a simplified illustration of perceptual

responses, which may involve other processes such as cortico-cortical interactions and

feedback.

We have shown that a simple Hebbian learning rule expressed through STDP could provide a

plausible mechanism for the encoding of natural statistics in the early visual system. This form

of learning through coincidence detection could present a more intuitive explanation for the

emergence of representations found in the early visual system and address crucial discrepancies

between efficiency based approaches and electrophysiological data (Ringach, 2002). Although

this study demonstrates the relevance of Hebbian plasticity in binocular vision, this approach

could easily be extended to other sensory modalities, and even multi-modal representations in

the early sensory cortex. Finally, we would like to point out that although the present work

focuses on the encoding properties of model, this simple plasticity rule also makes it possible to study the time-course of changes in the connectivity of individual units/neurones. For computational neuroscientists, this offers a critical advantage over conventional optimisation-based approaches as it allows for models which track cortical connectivity over time – such as studies on development, critical-period plasticity, and perceptual learning.

References

Albrecht D, Geisler W, Frazor R, Crane A (2002) Visual Cortex Neurons of Monkeys and Cats: Temporal Dynamics of the Contrast Response Function. J Neurophysiol 88 Available at: http://jn.physiology.org/content/88/2/888.short [Accessed September 11, 2017].

Anzai A, Ohzawa I, Freeman RD (1999) Neural Mechanisms for Processing Binocular Information I. Simple Cells. J Neurophysiol 82:891–908 Available at: http://jn.physiology.org/content/82/2/891.full-text.pdf+html [Accessed May 3, 2016].

Atick JJ, Redlich AN (1992) What Does the Retina Know about Natural Scenes? Neural Comput 4:196–210 Available at: http://www.mitpressjournals.org/doi/10.1162/neco.1992.4.2.196 [Accessed February 21, 2017].

Barlow H (1961) Possible Principles Underlying the Transformations of Sensory Messages. In: Sensory Communication, pp 217–234. The MIT Press. Available at: http://mitpress.universitypressscholarship.com/view/10.7551/mitpress/9780262518420.001.0001/upso-9780262518420-chapter-13.

Barlow H (2001) The exploitation of regularities in the environment by the brain. Behav Brain Sci 24:602–607 Available at: http://www.journals.cambridge.org/abstract_S0140525X01000024 [Accessed October 21, 2016].

Barlow H, Földiák P (1989) Adaptation and decorrelation in the cortex. In: The computing neuron, pp 54–72. Addison-Wesley Longman Publishing Co., Inc.

Berns G, Dayan P, Sejnowski T (1993) A correlational model for the development of disparity selectivity in visual cortex that depends on prenatal and postnatal phases. Proc Natl Acad Sci 90.

Bi GQ, Poo MM (1998) Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. J Neurosci 18:10464–10472 Available at: http://www.ncbi.nlm.nih.gov/pubmed/9852584 [Accessed June 27, 2016].

Brette R (2012) Computing with Neural Synchrony Sporns O, ed. PLoS Comput Biol 8:e1002561 Available at: http://dx.plos.org/10.1371/journal.pcbi.1002561 [Accessed November 9, 2017].

Burge J, Geisler W (2014) Optimal disparity estimation in natural stereo images. J Vis 14:1 Available at: http://jov.arvojournals.org/article.aspx?articleid=2121400 [Accessed January 15, 2016].

Burge J, Geisler W (2015) Optimal speed estimation in natural image movies predicts human performance. Nat Commun 6:7900 Available at: http://www.nature.com/articles/ncomms8900 [Accessed March 26, 2018].

Burge J, Jaini P (2017) Accuracy Maximization Analysis for Sensory-Perceptual Tasks: Computational Improvements, Filter Robustness, and Coding Advantages for Scaled Additive Noise Kording KP, ed. PLOS Comput Biol 13:e1005281 Available at: http://dx.plos.org/10.1371/journal.pcbi.1005281 [Accessed April 19, 2018].

Caporale N, Dan Y (2008) Spike Timing–Dependent Plasticity: A Hebbian Learning Rule. Annu Rev Neurosci 31:25–46 Available at: http://www.annualreviews.org/doi/10.1146/annurev.neuro.31.060407.125639 [Accessed March 23, 2017].

Celebrini S, Thorpe S, Trotter Y, Imbert M (1993) Dynamics of orientation coding in area V1 of the awake primate. Vis Neurosci 10:811–825 Available at: http://www.journals.cambridge.org/abstract_S0952523800006052 [Accessed September 11, 2017].

Cooper EA, Norcia AM (2014) Perceived depth in natural images reflects encoding of low-level luminance statistics. J Neurosci 34:11761–11768 Available at: http://www.jneurosci.org/content/34/35/11761.short [Accessed February 23, 2016].

Cumming BG, DeAngelis GC (2001) The physiology of stereopsis. Annu Rev Neurosci 24:203–238 Available at: http://www.annualreviews.org/doi/abs/10.1146/annurev.neuro.24.1.203 [Accessed April 29, 2016].

De Valois R, Albrecht D, Thorell L (1982) Spatial frequency selectivity of cells in macaque visual cortex. Vision Res 22:545–559 Available at: https://www.sciencedirect.com/science/article/pii/0042698982901134 [Accessed May 11, 2018].

Delorme A, Perrinet L, Thorpe SJ (2001) Networks of integrate-and-fire neurons using Rank Order Coding B: Spike timing dependent plasticity and emergence of orientation selectivity. Neurocomputing 38–40:539–545 Available at: http://www.sciencedirect.com/science/article/pii/S0925231201004039 [Accessed January 8, 2016].

Durand J-B, Celebrini S, Trotter Y (2007) Neural Bases of Stereopsis across Visual Field of the Alert Macaque Monkey. Cereb Cortex 17:1260–1273 Available at: https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bhl050 [Accessed April 3, 2017].

Durand J-B, Zhu S, Celebrini S, Trotter Y (2002) Neurons in Parafoveal Areas V1 and V2 Encode Vertical and Horizontal Disparities. J Neurophysiol 88:2874 LP-2879 Available at: http://jn.physiology.org/content/88/5/2874.abstract.

Ferster D, Lindström S (1983) An intracellular analysis of geniculo-cortical connectivity in area 17 of the cat. J Physiol 342:181–215 Available at: http://doi.wiley.com/10.1113/jphysiol.1983.sp014846 [Accessed September 11, 2017].

Freeman R, Ohzawa I (1992) Development of binocular vision in the kitten's striate cortex. J Neurosci 12.

Furmanski C, Engel S (2000) An oblique effect in human primary visual cortex. Nat Neurosci 3:535–536 Available at: http://www.nature.com/articles/nn0600_535 [Accessed March 26, 2018].

Gawne T, Kjaer T, Richmond B (1996) Latency: another potential code for feature binding in striate cortex. J Neurophysiol 76 Available at: http://jn.physiology.org/content/76/2/1356 [Accessed September 11, 2017].

Geisler W (2008) Visual perception and the statistical properties of natural scenes. Annu Rev Psychol 59:167–192 Available at: http://www.annualreviews.org/doi/abs/10.1146/annurev.psych.58.110405.085632 [Accessed October 14, 2015].

Geisler W, Perry J (2009) Contour statistics in natural images: Grouping across occlusions. Vis Neurosci 26:109 Available at: http://www.journals.cambridge.org/abstract_S0952523808080875 [Accessed March 26, 2018].

Girshick A, Landy M, Simoncelli E (2011) Cardinal rules: visual orientation perception reflects knowledge of environmental statistics. Nat Neurosci 14:926–932 Available at: http://www.nature.com/articles/nn.2831 [Accessed March 26, 2018].

Gollisch T, Meister M (2008) Rapid Neural Coding in the Retina with Relative Spike Latencies. Science (80-) 319 Available at: http://science.sciencemag.org/content/319/5866/1108 [Accessed September 11, 2017].

Goncalves N, Welchman A (2017) "What Not" Detectors Help the Brain See in Depth. Curr Biol Available at: http://www.sciencedirect.com.insb.bib.cnrs.fr/science/article/pii/S0960982217304049 [Accessed May 16, 2017].

Gütig R, Aharonov R, Rotter S, Sompolinsky H (2003) Learning Input Correlations through Nonlinear Temporally Asymmetric Hebbian Plasticity. J Neurosci 23:3697–3714.

Hebbs D (1949) The organization of behavior. New York, New York, USA: Wiley and Sons.

Henriksen S, Cumming BG, Read JCA (2016) A Single Mechanism Can Account for Human Perception of Depth in Mixed Correlation Random Dot Stereograms. PLoS Comput Biol 12:e1004906 Available at: http://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1004906#sec011 [Accessed June 1, 2016].

Hibbard P, Bouzit S (2005) Stereoscopic correspondence for ambiguous targets is affected by elevation and fixation distance. Spat Vis 18:399–411 Available at: http://booksandjournals.brillonline.com/content/journals/10.1163/1568568054389589 [Accessed October 16, 2017].

Howe C, Purves D (2002) Range image statistics can explain the anomalous perception of length. Proc Natl Acad Sci U S A 99:13184–13188 Available at: http://www.ncbi.nlm.nih.gov/pubmed/12237401 [Accessed March 26, 2018].

Hoyer P, Hyvärinen A (2000) Independent component analysis applied to feature extraction from colour and stereo images. Netw Comput Neural Syst 11:191–210 Available at: http://www.tandfonline.com/doi/full/10.1088/0954-898X_11_3_302 [Accessed July 2, 2017].

Hubel D, Wiesel T (1962) Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. J Physiol 160:106–154 Available at: http://doi.wiley.com/10.1113/jphysiol.1962.sp006837 [Accessed May 15, 2018].

Hunter D, Hibbard P (2015) Distribution of independent components of binocular natural images. J Vis 15:6 Available at: http://jov.arvojournals.org/article.aspx?doi=10.1167/15.13.6 [Accessed January 9, 2017].

Jones J, Palmer L (1987) An evaluation of the two-dimensional Gabor filter model of simple receptive fields in cat striate cortex. J Neurophysiol 58:1233–1258.

Karklin Y, Lewicki MS (2009) Emergence of complex cell properties by learning to generalize in natural scenes. Nature 457:83–86 Available at: http://dx.doi.org/10.1038/nature07481 [Accessed May 31, 2016].

Kiorpes L, Kiper DC, O'Keefe LP, Cavanaugh JR, Movshon JA (1998) Neuronal Correlates of Amblyopia in the Visual Cortex of Macaque Monkeys with Experimental Strabismus and Anisometropia. J Neurosci 18:6411 LP-6424 Available at: http://www.jneurosci.org/content/18/16/6411.abstract.

König P, Engel A, Singer W (1996) Integrator or coincidence detector? The role of the cortical neuron revisited. Trends Neurosci 19:130–137 Available at: https://www.sciencedirect.com/science/article/pii/S0166223696800191 [Accessed November 9, 2017].

Maass W (2000) On the Computational Power of Winner-Take-All. Neural Comput 12:2519–2535 Available at: http://www.mitpressjournals.org/doi/10.1162/089976600300014827 [Accessed May 16, 2018].

Masquelier T (2012) Relative spike time coding and STDP-based orientation selectivity in the early visual system in natural continuous and saccadic vision: a computational model. J Comput Neurosci 32:425–441 Available at: http://www.ncbi.nlm.nih.gov/pubmed/21938439 [Accessed January 8, 2016].

Masquelier T (2017) STDP allows close-to-optimal spatiotemporal spike pattern detection by single coincidence detector neurons. Neuroscience Available at: http://linkinghub.elsevier.com/retrieve/pii/S0306452217304372 [Accessed November 9, 2017].

Masquelier T, Thorpe SJ (2007) Unsupervised learning of visual features through spike timing dependent plasticity. PLoS Comput Biol 3:e31 Available at: http://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.0030031 [Accessed April 11, 2016].

Masquelier T, Thorpe SJ (2010) Learning to recognize objects using waves of spikes and Spike Timing-Dependent Plasticity. In: The 2010 International Joint Conference on Neural Networks (IJCNN), pp 1–8. IEEE. Available at: http://ieeexplore.ieee.org/lpdocs/epic03/wrapper.htm?arnumber=5596934 [Accessed January 8, 2016].

Movshon J, Eggers H, Gizzi M, Hendrickson A, Kiorpes L, Boothe R (1987) Effects of early unilateral blur on the macaque's visual system. III. Physiological observations. J Neurosci 7:1340–1351 Available at: http://www.ncbi.nlm.nih.gov/pubmed/3572484 [Accessed October 20, 2017].

Nasr S, Tootell R (2016) Visual field biases for near and far stimuli in disparity selective
columns in human visual cortex. Neuroimage Available at:
http://www.sciencedirect.com/science/article/pii/S1053811916304736.

Ohzawa I, Deangelis G, Freeman R (1997) Encoding of Binocular Disparity by Complex Cells
in the Cat's Visual Cortex. J Neurophysiol 77:2879–2909.

Ohzawa I, Freeman R (1986) The binocular organization of simple cells in the cat's visual
cortex. J Neurophysiol 56:221–242 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/3746398 [Accessed May 3, 2018].

Okazawa G, Tajima S, Komatsu H (2015) Image statistics underlying natural texture selectivity
of neurons in macaque V4. Proc Natl Acad Sci U S A 112:E351-60 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25535362 [Accessed October 20, 2016].

Olshausen BA, Field DJ (1996) Emergence of simple-cell receptive field properties by learning
a sparse code for natural images. Nature 381:607–609 Available at:
http://dx.doi.org/10.1038/381607a0 [Accessed October 1, 2015].

Olshausen BA, Field DJ (1997) Sparse coding with an overcomplete basis set: A strategy
employed by V1? Vision Res 37:3311–3325.

Poggio GF, Motter BC, Squatrito S, Trotter Y (1985) Responses of neurons in visual cortex (V1
and V2) of the alert macaque to dynamic random-dot stereograms. Vision Res 25:397–406
Available at: http://www.sciencedirect.com/science/article/pii/0042698985900653.

Prince SJD, Cumming BG, Parker AJ (2002a) Range and Mechanism of Encoding of Horizontal
Disparity in Macaque V1. J Neurophysiol 87:209 LP-221 Available at:
http://jn.physiology.org/content/87/1/209.abstract.

Prince SJD, Pointon AD, Cumming BG, Parker AJ (2002b) Quantitative Analysis of the
Responses of V1 Neurons to Horizontal Disparity in Dynamic Random-Dot Stereograms.
J Neurophysiol 87:191–208 Available at: http://jn.physiology.org/content/87/1/191.short
[Accessed January 20, 2016].

Ringach D (2002) Spatial Structure and Symmetry of Simple-Cell Receptive Fields in Macaque
Primary Visual Cortex. J Neurophysiol 88 Available at:
http://jn.physiology.org/content/88/1/455.short [Accessed June 22, 2017].

Samonds JM, Potetz BR, Lee TS (2012) Relative luminance and binocular disparity preferences
are correlated in macaque primary visual cortex, matching natural scene statistics. Proc
Natl Acad Sci 109:6313 LP-6318 Available at:
http://www.pnas.org/content/109/16/6313.abstract.

Sebastian S, Abrams J, Geisler W (2017) Constrained sampling experiments reveal principles
of detection in natural scenes. Proc Natl Acad Sci U S A 114:E5731–E5740 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/28652323 [Accessed March 26, 2018].

Shriki O, Kohn A, Shamir M (2012) Fast Coding of Orientation in Primary Visual Cortex
Graham LJ, ed. PLoS Comput Biol 8:e1002536 Available at:
http://dx.plos.org/10.1371/journal.pcbi.1002536 [Accessed October 10, 2017].

Solomon SG, White AJR, Martin PR (2002) Extraclassical Receptive Field Properties of
Parvocellular, Magnocellular, and Koniocellular Cells in the Primate Lateral Geniculate
Nucleus. J Neurosci 22:338 LP-349 Available at:
http://www.jneurosci.org/content/22/1/338.abstract.

Sprague W, Cooper E, Tosic I, Banks M (2015) Stereopsis is adaptive for the natural
environment. Sci Adv 1:e1400254–e1400254 Available at:
http://advances.sciencemag.org/content/1/4/e1400254.abstract [Accessed October 13,
2015].

Stanley G (2013) Reading and writing the neural code. Nat Neurosci 16:259–263 Available at:
http://www.nature.com/doifinder/10.1038/nn.3330 [Accessed November 9, 2017].

Tanaka K (1985) Organization of geniculate inputs to visual cortical cells in the cat. Vision Res
25:357–364 Available at: http://linkinghub.elsevier.com/retrieve/pii/0042698985900604
[Accessed September 11, 2017].

Thorpe S, Delorme A, Van Rullen R (2001) Spike-based strategies for rapid processing. Neural
Networks 14:715–725.

Webster M, Mollon J (1997) Adaptation and the color statistics of natural images. Vision Res
37:3283–3298   Available   at:
http://www.sciencedirect.com/science/article/pii/S0042698997001259   [Accessed
December 13, 2015].

1003