1 **Title:** Tn-Core: context-specific reconstruction of core metabolic models using Tn-seq data

2 **Running head:** Tn-Core: Tn-seq and metabolic reconstruction

3 **Authors:** George C diCenzo[*], Alessio Mengoni, Marco Fondi[1*]

4 **Affiliation:** Department of Biology, University of Florence, 50019, Sesto Fiorentino, FI, Italy

6 *Correspondence: georgecolin.dicenzo@unifi.it, marco.fondi@unifi.it

8   **ABSTRACT**

Motivation: Tn-seq (transposon mutagenesis and sequencing) and constraint-based metabolic

modelling represent highly complementary approaches. They can be used to probe the core

genetic and metabolic networks underlying a biological process, revealing invaluable

information for synthetic biology engineering of microbial cell factories. However, while

algorithms exist for integration of –omics data sets with metabolic models, no method has been

explicitly developed for integration of Tn-seq data with metabolic reconstructions.

Results: We report the development of Tn-Core, a Matlab toolbox designed to generate gene-

centric, context-specific core reconstructions consistent with experimental Tn-seq data.

Extensions of this algorithm allow: i) the generation of context-specific functional models

through integration of both Tn-seq and RNA-seq data; ii) to visualize redundancy in core

metabolic processes; and iii) to assist in curation of *de novo* draft metabolic models. The utility

of Tn-Core is demonstrated primarily using a *Sinorhizobium meliloti* model as a case study.

Availability   and   implementation: The   software   can   be   downloaded   from

https://github.com/diCenzo-GC/Tn-Core. All results presented in this work have been obtained

with Tn-Core v. 1.0.

Contact: georgecolin.dicenzo@unifi.it, marco.fondi@unifi.it

Supplementary information: Supplementary data are available at Bioinformatics online.

INTRODUCTION

The chemical complexity of biological entities hampers a full understanding of life and,

consequently, its characterization is one of the strongest motivations in systems biology.

Constraint-based metabolic modelling (CBMM) [1] is a well-established tool to formally

represent cellular metabolism at the genome-scale level (by means of Genome Scale Metabolic

Reconstructions, GSMRs) and to derive reliable predictions [2]. Despite this approach having

shown remarkable predictive capabilities over the years [3], there are constant efforts aimed at

improving and customizing the procedures of CBMM analyses.

It is increasingly recognized that the complexity of modern GSMRs often masks their

utility in various applications [4], and that most studies to date only focus on the core metabolic

pathways of the organism [5, 6]. Furthermore, due to the scaling of computational complexity,

many stoichiometric (e.g. elementary flux modes enumeration [7]) and/or dynamic approaches

(e.g. kinetic modelling [8]) cannot be applied to GSMRs embedding thousands of reactions. As a

result, algorithms have been implemented to reduce a GSMR to a core set of reactions necessary

to produce a pre-defined phenotype(s) [4, 9-11]. These algorithms share a similar overall

approach: they are reaction-centric and require a user-defined list of reactions, metabolites,

and/or phenotypes that must remain in the core model. However, by not directly incorporating

experimental data, the biological accuracy of these core models cannot be guaranteed.

As GSMRs generally incorporate as much of the cell's metabolism as possible, regardless

to the activity of the reaction in a given environment, additional constraints are required to

accurately represent environment-specific metabolism. This can be accomplished by constraining

GSMRs with –omics data sets. This most commonly involves integrating gene expression data,

constraining the allowable flux across each reaction based on the expression level(s) of the

corresponding gene(s) [12, 13]. Similarly, tools exist for combining GSMRs with proteomics

[14], fluxomics [15], and metabolomics data [16]. Ultimately, these applications have a common

goal: reducing a GSMR to a smaller model with only the reactions active in the specific

condition.

High-throughput transposon mutagenesis and sequencing (Tn-seq) generates a genome-

wide list of genes essential in a given environment [17]. Arguably, these data sets are the best

experimental representation of which reactions are active in a given environmental condition.

Combining core metabolic networks and Tn-seq can allow deep functional refinement of GSMRs

to account for only those (core) reactions and genes active under the tested conditions.. From a

synthetic biology viewpoint, the central metabolism of an organism is of paramount importance

as it i) produces the precursors for all natural chemicals and ii) has a high capacity of pathway

fluxes; as such, central metabolism can be exploited as a chassis for production of industrially

important molecules [18, 19]. Consequently, a Tn-seq curated core metabolic model is of high

value for synthetic biology attempts at engineering designing cell factories. Indeed, genome

streamlining, i.e., the construction of cells with minimal genomes, is known to generate cells

with improved biotechnological properties, including increased protein or metabolite production

[20-24]. However, despite the highly complementary nature of Tn-seq and CBMM, we are

unaware of a tool for generating context-specific models through the automated incorporation of

Tn-seq data with GSMRs.

Here, we report the development of Tn-Core, a MATLAB toolbox for use with COBRA

formatted metabolic models. Tn-Core is designed for the generation of gene-centric, context-

specific core metabolic models consistent with experimental gene fitness data produced through

Tn-seq experiments, or through both Tn-seq and RNA-seq data. Tn-Core can further be used to:

i) evaluate potential redundancy in core metabolism (does not require Tn-seq data); ii) identify

which of the alternate pathway(s) contributes to higher flux through the objective function; and

iii) perform Tn-seq-guided refinement of the Gene-Protein-Reaction rules (GPRs) in a GSMR.

77   **IMPLEMENTATION**

Tn-Core was developed to facilitate the generation of context-specific core metabolic

models through the integration Tn-seq data, then expanded to further allow the integration of

RNA-seq data and to examine core metabolic redundancy in the presence or absence of these

data. The toolbox is written in Matlab and uses COBRA formatted models and the COBRA

Toolbox [25]. Tn-Core is available as Supplementary Materials S1, and the current and future

versions will be available through GitHub (https://github.com/diCenzo-GC/Tn-Core).The

functionality of the entire toolbox has been validated on four machines, running three versions of

Matlab (R2015b, R2016b, R2017a) and three distinct COBRA toolbox setups (openCOBRA

downloaded between 12/2016 and 08/2017), suggesting that Tn-Core should work in a broad

range of computing environments.

Generation of core metabolic models.

The pseudocode for Tn-Core is given in Algorithm 1, the main workflow is depicted in the

flowchart of Figure 1, and a detailed manual describing its usage is provided in Supplementary

Materials S1. The minimum input is a COBRA-formatted metabolic model. Optionally, the user

may provide: (i) Tn-seq data for all genes in the genome; (ii) RNA-seq data for all genes in the

genome, and/or (iii) a list of pre-determined core/essential genes. Tn-Core begins with the

optional step (Figure 1a) of producing a list of model genes to be protected during the generation

of

Algorithm 1. The Tn-Core algorithm

Input: n is the number of iterations; *model* is the initial GSMR. **Other variables and lists:** G_m is the list of genes in *model*, T is the Tn-seq data, L is the RNA-seq data, M_i is the final array of core metabolic reconstructions, t is the threshold for objective function. **Functions**: *detectDeadEnds, deleteModelGene, findRxnsFromMets, singleGeneDeletion*, and *optimizeCbModel* are part of the COBRA Toolbox. All the other functions are implemented as Matlab code (see Supplementary Material S1).

1: D = *detectDeadEnds*(*model*)

2: R_D = *findRxnsFromMets(D)*

3: m_{red} = **remove reactions and unused genes** (*model*, R_D)

4: E_{model} = *singleGeneDeletion*(*model*)

5: (E, S, W) = **get essential, strong, and weak growth promoting Genes** (*T*)

6: L_H = **get highly expressed genes** (*L*)

7: $U_m = (G_m \sim ((E \cap G_m) \cup E_{model} \cup (L_H \cap G_m)))$

8: for i = 1 **to** n

9: $m = m_{red}$

10: U_m^* = shuffle(U_m)

11:   **for** j = 1 **to** length(U_m^*)

12: m' = *deleteModelGene*(*m*, $U_m^*(j)$)

13: φ = *optimizeCbModel*(*m'*)

14:   **if** $\varphi > t$

15: $m = m'$

16:   **end if**

17:   **end for**

18: $M(i) = m$

19: $G_{M(i)}$ = **get the genes in** $M(i)$

20: $O(i)$ = *optimizeCbModel*($M(i)$)

21: $\{N_E(i); N_S(i); N_W(i)\}$ = {**length**($G_{M(i)} \cap E$); **length**($G_{M(i)} \cap S$); **length**($G_{M(i)} \cap W$)}

22: end for

23: M_{core} = $M($**max**(N_E)$)$

24: if length(M_{core}) > 1

25: M_{core} = $M_{core}($**max**(N_S)$)$

26: M_{core} = $M_{core}($**max**(N_W)$)$

27: M_{core} = $M_{core}($**max**(O)$)$

28: end if

29: return M_{core}

random core models. This list is based on: (i) all user-defined core genes, (ii) highly expressed

genes if RNA-seq are provided, and (iii) essential genes based on Tn-seq data (optional even if

Tn-seq data are provided). Next, Tn-Core produces a reduced GSMR by iteratively removing all

reactions that produce dead-end metabolites (and associated genes, if they are not in the GPR of

another reaction). Additionally, all GPRs not assigned to a coding sequence (e.g. gap-filling

reactions) are removed. As the order in which reactions are added/removed from a model might

alter the predictive capability of the reconstruction, randomized core models (M, Algorithm 1)

are then generated from the reduced model (Figure 1b). Importantly, this step can be parallelized,

reducing the running time. This involves first preparing a list of all non-protected model genes

(U_m), and randomly shuffling their order at each iteration (U_m^*). All genes (and corresponding

reactions) from each shuffled set are individually deleted from the model and growth is tested. If

the objective function flux (φ) stays above the threshold (t), the gene is excluded from the model;

otherwise, the gene is put back to the model. The result is a population of models (M) each

containing the initially protected genes (optional), and a minimal amount of additional genes

required to maintain objective function flux φ above the threshold t. If Tn-seq data is provided,

the objective function flux of each core model is recorded, genes are classified into four

categories from 'essential' to 'non-essential' based on the Tn-seq data (Figure S1), and the

number of core model genes in each category is recorded (Figure 1c).

Finally, the core reconstruction that maximizes the number of essential Tn-seq genes is

chosen as the reconstruction most consistent with the Tn-seq data (M_{core}). If two or more models

embed the same number of essential genes, the reconstruction maximizing the number of 'strong

growth promoting' and then 'weak growth promoting' genes is selected as the output. If multiple

models still remain, the model with the highest objective reaction flux is returned as the core

metabolic model most consistent with the gene essentiality data (Figure 1d). Independently, the

core model with the highest objective function flux is returned as the fastest growing core model

(Figure 1d); if multiple models have the same maximal objective function flux, the model most

consistent with the gene essentiality data is chosen. In some cases, it may be desirable to obtain

other core models produced during the running of Tn-Core, such as the slowest growing core

model. The output of Tn-Core additionally includes a cell array of the objective function flux for

all produced core models, as well as a binary presence/absence cell array indicating which genes

are included in each of the core models. By using the latter cell array with the *tncore_reconstruct*

function, it is possible to rebuild any of the core models produced during the running of Tn-Core.

Analysis of variation across the core metabolic models.

164   The redundancy embedded within GSMRs means that each of the models in the core

model population may contain a different set of genes and/or reactions. Tn-Core includes

functions to explore this redundancy, whether Tn-seq data is provided or not (Figure 1e). Two or

three primary matrixes are returned, and can display either gene or reaction information. A

binary presence/absence matrix is given, which indicates, for each model, whether each feature is

present or absent; only features embedded in at least one core model are included (Figure 2a, 2b).

A co-occurrence matrix is also provided; for each feature variably present in the core model

population, a Chi-squared statistics is reported to indicate which feature pairs are more likely

than chance to appear, or not appear, in the same core models (Figure 2c-2e). If the core models

are generated multiple times, for example, using different objective flux thresholds, a matrix can

be produced that indicates, for each population of core models, what percentage of models

contains each of the features (Figure 2f, 2g).

Refinement of genome-scale metabolic network reconstructions.

Finally, an extension is provided to use Tn-seq data to assist in the automated curation of

GSMRs (Figure 1f). First, Tn-seq essential genes are determined, and these genes are protected

during core model generation. The core model most consistent with the Tn-seq data is collected,

and where appropriate, 'or' statements in the GPRs are replaced with 'and' statements; if any Tn-

seq essential genes in the model have no effect when deleted, and if any occur in the same

reaction(s) and only the same reaction(s), and the GPR currently lacks an 'and' statement, the

'or' statements of the GPR are replaced with 'and' statements. The implementation of this

section of the code is rather strict in order to avoid artificially converting non-essential genes to

essential genes. Finally, for any core model reaction with a Tn-seq essential gene, the

corresponding GPRs of the original GSMR are replaced with those of the core reconstruction.

189   **RESULTS AND DISCUSSION**

Validation of Tn-Core.

Tn-Core was validated by extracting context-specific core models from the

Sinorhizobium meliloti iGD1575 GSMR [26]. Two core models were produced, each using a

growth threshold of 50% the full model, with 50,000 iterations, and with Tn-seq essential genes

pre-identified. In one Tn-Core run, only Tn-seq data [27] was used; in the second run, the same

Tn-seq data plus RNA-seq data [28] was included. The sizes of both models are summarized in

Table 1, and the inclusion of RNA-seq data resulted in a somewhat larger core model.

The ability of the core models to capture context-specific core metabolism was examined

by predicting the essentiality of central carbon metabolic genes (Figure 3). Results were

compared to both the full iGD1575 model and to the manually constructed *S. meliloti* iGD726

core metabolic reconstruction [27]. The entire set of central carbon metabolic pathways was

predicted to be non-essential in iGD1575 presumably due to network redundancy. In contrast,

most of central carbon metabolism was essential in the manually prepared iGD726 core model

(*gnd* and *tal* are correctly predicted as non-essential). Using only Tn-seq data, Tn-Core extracted

a core model largely consistent with iGD726, although the ATP synthase pump was missing.

However, by also including RNA-seq data in the pipeline, the extracted core model even better

reflected context-specific metabolism. This is highlighted by the lower half of the Embden-

Meyerhof-Parnas pathway. In particularly, mutation of *pgk* was experimentally shown to result

in a 40% growth rate decrease when grown with glucose [29]. Whereas *pgk* was essential in the

first core model, deletion of *pgk* in the core model extracted using Tn-seq and RNA-seq data

resulted in a growth rate decrease of 30%. Taken together, these results demonstrate the ability

of Tn-Core to produce highly accurate context-specific core metabolic models, and illustrates

how integrating both Tn-seq and RNA-seq data sets can lead to high precision fitness

predictions.

We subsequently implemented in Tn-Core the option to employ the Minimization of

Metabolic Optimization (MOMA) algorithm during core model generation instead of FBA.

Using MOMA instead of FBA is significantly slower, had little effect on the size of the core

models (Table 1), and, at least in central carbon metabolism (Figure 3), did not produce more

accurate core reconstructions. We have also found that the core models returned when using the

MOMA implementation are not guaranteed to grow. This appears to be due to certain core

models growing when using the *MOMA* function of the COBRA toolbox, but not growing when

using the *optimizeCbModel* function of the COBRA toolbox. We therefore suggest that the FBA

implementation should be used for most purposes.

The functionality of Tn-Core was further confirmed using the *Pseudomonas aeruginosa*

iPae1146 GSMR [30] and published Tn-seq data [31]. These results are reported in

Supplementary Material S2.

Benchmarking of Tn-Core.

There is currently no tool explicitly comparable to Tn-Core as none consider

experimental Tn-seq data during core model identification. Nevertheless, we compared Tn-Core

to two algorithms design for the extraction of core reconstructions: FASTCORE [10] and

minNW [11]. Both algorithms are reaction-centric, and require as input a set of reactions, not

genes, to be protected in the output model. To adapt these algorithms for use with Tn-seq data,

we set the protected reactions as those reactions that are constrained upon deletion of the Tn-seq

essential genes. Additionally, in both cases, a consistent model derived from iGD1575, generated

with FASTCC [10], was used as the starting model. For both FASTCORE and minNW, the

output models had similar or fewer reactions and metabolites, but a larger complement of genes,

than the models produced with Tn-Core (Table 1), which is related to its reaction-centric nature.

More importantly, although faster than Tn-Core, the accuracy of FASTCORE and minNW was

far exceeded by Tn-Core using central carbon metabolism as a proxy (Figure 3). This result

validates that Tn-Core fulfills a function that is currently lacking among the available algorithms.

The output of Tn-Core was also compared to the gene-centric TIGER implementation of

the GIMME algorithm [32, 33]. GIMME generates context-specific models based on expression

data, and is therefore not directly comparable to Tn-Core that primarily uses essentiality data.

GIMME initially failed to return a functional model using iGD1575 and the provided RNA-seq

data, but a working model could be recovered using a custom extension (see Supplementary File

S2). Overall, the models returned by GIMME and Tn-Core displayed high consistency, with the

central carbon metabolism extracted by GIMME of similar accuracy to those extracted by Tn-

Core (Figure 3). Additionally, the GIMME model and Tn-Core model produced with Tn-seq and

RNA-seq data (FBA implementation) share > 87% of their genes. Thus, at least in *S. meliloti*

where essential genes tend to be highly expressed [27], both Tn-Core and GIMME perform

similarly and the choice of algorithm would be driven primarily by the type of data being

incorporated with the GSMR.

Tn-Core performance.

In order for Tn-Core to be accurate, a sufficiently large population of core models must

be generated to ensure the optimal core model is represented. There are therefore two primary

factors contributing to the speed of Tn-Core: (i) running time per iteration (i.e., per core model

produced), and (ii) the number of iterations. To test the effect of starting model and parameter

settings on the performance of Tn-Core, we generated 25,000 core models for five different

GSMRs with varying parameter settings. A summary of these runs are provided in Table 2, and a

detailed description of is reported in Supplementary File S2. 25,000 iterations did not guarantee

the presence of all possible core models in any of the runs. However, the number of variably

present genes gives an indication of the number of iterations required to cover all possibilities;

the square of the variably present genes represents the theoretical maximum number of

genetically unique core models. Considering that the variability among core models is highly

dependent on the starting GSMR and the parameter settings, we recommend users first perform a

test run of 10,000 iterations, and use the gene variability to approximate how many iterations

must be performed. Additionally, if Tn-Core is being used to produce a core model and not only

to explore redundancies in the core network, we recommend setting Tn-Core to pre-determine

the essential genes prior to core model generation and to use a growth threshold of at least 50%.

Characterization of redundancy and growth promoting pathways with Tn-Core.

As is evident from Table 2, significant redundancy can exist in core metabolic pathways.

Tn-Core produces a series of matrixes to summarize this variability (Figure 2), which can be

easily imported into graphing tools to visualize the data (e.g. [34]). Here, we briefly illustrate the

usefulness of these matrixes in uncovering biologically interesting data. We note that the same

trends were observed for *S. meliloti* using the FBA (Figure 2) or MOMA (Figure S2)

implementation, and also when using GSMRs for *Eschericha coli*, *P. aeruginosa*, *Pseudomonas*

haloplanktis, and *Acinitobacter baumannii* (Figures S3-S6), demonstrating that these results are

not specific to a single model (Figure S6).

Gene/reaction presence matrixes (Figures 2a, 2b) provide an overview of the variability

of the models. In the case of *S. meliloti*, the core models contain an average of 434 genes, of

which 286 genes (~ 66%) are invariably present and the rest are from a set of 777 variably

present genes. In other words, a third of core *S. meliloti* metabolic genes can be functionally

replaced by alternative genes or pathways, consistent with recent experimental work [27]. The

variable and invariable core genes were mapped to KEGG pathways [35] using eggNOG-mapper

[36] to identify functional biases. Significant redundancy was observed in a diversity of

pathways, including carbon, amino acid and nucleotide metabolism. In contrast, the most

fundamental cellular processes appeared to lack redundancy, such as transcription, translation,

and aminoacyl-tRNA biosynthesis.

Gene/reaction co-occurrence matrixes summarize the frequency that two genes or

reactions occur in the same model relative to chance (Figures 2c-2d). This can identify modules

that work together (likely to co-occur), and genes or biochemical pathways that are functionally

redundant (unlikely to co-occur). For all GSMRs used in this work, clear modules and redundant

genes/pathways could be observed in the matrixes (Figure 2, Figures S2-S6). Known

redundancies could be detected in the *S. meliloti* iGD1575 reaction co-occurrence matrix. For

example, the two pathways for L-proline biosynthesis [37] were unlikely to occur in the same

model, as were thiamine transport and thiamine biosynthesis. These observations confirm that

these matrixes could be useful in detecting metabolic redundancy in core bacterial metabolism.

Finally, core models were generated using growth thresholds of 10% and 99% (of the

original objective function flux), and a scatterplot was used to compare the frequency of each

gene/reaction in the resulting core model populations (Figures 2f, 2g). In all cases, some

genes/reactions were found to be enriched in one of the two core model populations, and the use

of the MOMA algorithm increased the incidence of such genes/reactions (Figures 2 and S2).

When using the FBA algorithm, biases in the occurrence of genes in the two core model

populations were particularly prevalent in the *E. coli* iJO1366 model (Figure S5). Intriguingly,

some genes, such as *b2417* (glucose-specific enzyme IIA component of PTS, glycolysis), *b2342*

and *b3845* (both acetyl-CoA acyltransferase, fatty acid degradation), were ~ 5-fold more

prevalent in the core models generated with a 99% growth threshold compared to a 10% growth

threshold (differences statistically significant based on Fisher exact tests, p-value < 2.2e-16).

Yet, despite the importance of the pathways these genes are involved in, none of them had a

predicted effect on growth rate when deleted in the full iJO1366 model (using either FBA or

MOMA), likely due to the redundancy in the complete GSMR. Hence, Tn-Core may facilitate

the identification of genes contributing to optimal growth in core metabolic networks, including

genes not readily detected as important in the full GSMR.

Refinement of GSMRs using Tn-Core.

Automated metabolic network reconstruction methods are expected to incorrectly assign

multiple genes to the same core metabolic reaction. In the absence of experimental data, it can be

difficult to correct such errors. We therefore implemented a function for using Tn-Core to assist

in model refinement using Tn-seq data. We tested this pipeline using the *S. meliloti* iGD1575

model, as well as with a draft *S. meliloti* model prepared using the Kbase automated

reconstruction pipeline. This process resulted in the modification of the GPRs of 60 reactions in

iGD1575, with 69 genes removed from the model. Similarly, 107 GPRs (over 6% of reactions)

were modified in the draft model following this process, with 57 genes deleted from the model.

These results demonstrate that Tn-seq data and Tn-Core can play a valuable role in curation of

metabolic models, although it certainly does not replace the need of an accurate manual curation.

327   **CONCLUSIONS**

Here, we presented Tn-Core, a new tool for the generation of core metabolic network

reconstructions. The unique feature of Tn-Core is the ability to consider experimental Tn-seq

data, as well as both Tn-seq and RNA-seq data, for producing a core model that best represents

the true metabolism of the cell in a given physiological condition. Despite that this pipeline may

run slower than existing algorithms for the generation of core or context-specific models, Tn-

Core remains advantageous due to: i) its high accuracy; ii) its ability to consider both functional

genomics (Tn-seq) and transcriptomics data (RNA-seq); iii) its ease of use with little pre-

processing of the data required; and iv) its gene-centric approach.

METHODS

All data generated with Tn-Core (except for the timing of Table 2) was done using

Matlab 2016a (Mathworks), the COBRA Toolbox (downloaded December 9, 2016 from the

openCOBRA repository) [25], and using the Gurobi 6 solver (gurobi.com), SBMLToolbox 4.1.0

[38], and libSBML 5.11.8 [39]. All other computations were performed in Matlab 2017a using

the Gurobi 7.0.2 solver, SBMLToolbox 4.1.0, libSBML 5.15.0, scripts from the COBRA

Toolbox (downloaded May 12, 2017 from the openCOBRA repository), and the TIGER Toolbox

v.1.2.0-beta [33]. For running minNW, the iLOG CPLEX Studio 12.7.1 solver (ibm.com) was

used. Gene essentiality was determined using the *singleGeneDeletion* function and the MOMA

algorithm. In order to ensure that core model generation with Tn-Core did not occasionally fail

when using the MOMA algorithm, the MOMA.m script of the COBRA Toolbox was modified at

line 216 to to treat unbounded solutions the same as infeasible solutions. Additionally, the

solveCobraQP.m script of the COBRA Toolbox was modified to work with the Gurobi 6 solver.

Detailed usage, and modifications, of FASTCORE [10], minNW [11], and GIMME [32, 33] are

provided in Supplementary Materials S2.

The *S. meliloti* iGD1575 [26], *P. haloplanktis* iMF721 [40], *A. baumannii* iLP844 [41],

E. coli iJO1366 [42], and *P. aeruginosa* iPae1146 [30] models were previously published. Prior

to using iLP844, the genes 'Unknown1' through 'Unknown160' were replaced with a single

gene called 'Unknown'. The draft *S. meliloti* GSMR was generated using Kbase (kbase.us) as

described in Supplementary Materials S1.

Scripts to repeat all benchmarking, as well as all output data generated in this work, are

available at https://github.com/diCenzo-GC/Tn-Core. The complete Tn-Core toolbox, together

with a reference manual, are provided as Supplementary Materials S1. Tn-Core is also freely

available at https://github.com/diCenzo-GC/Tn-Core, and future releases of the toolbox will be

available through the same link.

364   **ACKNOWLEDGEMENTS**

365   GCD was funded by a Natural Sciences and Engineering Research Council (NSERC) of

366   Canada Postdoctoral Fellowship.

368   **REFERENCES**

[1] A. Varma, B.O. Palsson, Stoichiometric flux balance models quantitatively predict growth

and metabolic by-product secretion in wild-type Escherichia coli W3110, Appl Environ

Microbiol, 60 (1994) 3724-3731.

[2] A. Bordbar, J.M. Monk, Z.A. King, B.O. Palsson, Constraint-based models predict metabolic

and associated cellular functions, Nature reviews. Genetics, 15 (2014) 107-120.

[3] C.B. Milne, P.J. Kim, J.A. Eddy, N.D. Price, Accomplishments in genome-scale in silico

modeling for industrial and medical biotechnology, Biotechnology journal, 4 (2009) 1653-1670.

[4] M. Ataman, D.F. Hernandez Gardiol, G. Fengos, V. Hatzimanikatis, redGEM: Systematic

reduction and analysis of genome-scale metabolic reconstructions for development of consistent

core metabolic models, PLoS computational biology, 13 (2017) e1005444.

[5] K.C. Soh, L. Miskovic, V. Hatzimanikatis, From network models to network responses:

integration of thermodynamic and kinetic properties of yeast genome-scale metabolic networks,

FEMS yeast research, 12 (2012) 129-143.

[6] M.T. Alam, M.H. Medema, E. Takano, R. Breitling, Comparative genome-scale metabolic

modeling of actinomycetes: the topology of essential core metabolism, FEBS letters, 585 (2011)

2389-2394.

[7] C.T. Trinh, A. Wlaschin, F. Srienc, Elementary mode analysis: a useful metabolic pathway

analysis tool for characterizing cellular metabolism, Applied microbiology and biotechnology,

81 (2009) 813-826.

[8] N. Jahan, K. Maeda, Y. Matsuoka, Y. Sugimoto, H. Kurata, Development of an accurate

kinetic model for the central carbon metabolism of Escherichia coli, Microbial cell factories, 15

(2016) 112.

[9] P. Erdrich, R. Steuer, S. Klamt, An algorithm for the reduction of genome-scale metabolic

network models to meaningful core models, BMC systems biology, 9 (2015) 48.

[10] N. Vlassis, M.P. Pacheco, T. Sauter, Fast reconstruction of compact context-specific

metabolic network models, PLoS computational biology, 10 (2014) e1003424.

[11] A. Rohl, A. Bockmayr, A mixed-integer linear programming approach to the reduction of

genome-scale metabolic networks, BMC bioinformatics, 18 (2017) 2.

[12] A.S. Blazier, J.A. Papin, Integration of expression data in genome-scale metabolic network

reconstructions, Frontiers in physiology, 3 (2012) 299.

[13] D. Machado, M. Herrgard, Systematic evaluation of methods for integration of

transcriptomic data into constraint-based models of metabolism, PLoS computational biology, 10

(2014) e1003580.

[14] R. Grosseholz, C.C. Koh, N. Veith, T. Fiedler, M. Strauss, B. Olivier, B.C. Collins, O.T.

Schubert, F. Bergmann, B. Kreikemeyer, R. Aebersold, U. Kummer, Integrating highly

quantitative proteomics and genome-scale metabolic modeling to study pH adaptation in the

human pathogen Enterococcus faecalis, NPJ systems biology and applications, 2 (2016) 16017.

[15] N. Zamboni, E. Fischer, U. Sauer, FiatFlux--a software for metabolic flux analysis from

13C-glucose experiments, BMC bioinformatics, 6 (2005) 209.

[16] K. Yizhak, T. Benyamini, W. Liebermeister, E. Ruppin, T. Shlomi, Integrating quantitative

proteomics and metabolomics with a genome-scale metabolic network model, Bioinformatics, 26

(2010) i255-260.

[17] M.C. Chao, S. Abel, B.M. Davis, M.K. Waldor, The design and analysis of transposon

insertion sequencing experiments, Nature reviews. Microbiology, 14 (2016) 119-128.

[18] P. Jouhten, Metabolic modelling in the development of cell factories by synthetic biology,

Computational and structural biotechnology journal, 3 (2012) e201210009.

[19] J. Nielsen, It is all about metabolic fluxes, Journal of bacteriology, 185 (2003) 7031-7035.

[20] M. Juhas, D.R. Reuss, B. Zhu, F.M. Commichau, Bacillus subtilis and Escherichia coli

essential genes and minimal cell factories after one decade of genome engineering,

Microbiology, 160 (2014) 2341-2351.

[21] S. Lieder, P.I. Nikel, V. de Lorenzo, R. Takors, Genome reduction boosts heterologous gene

expression in Pseudomonas putida, Microbial cell factories, 14 (2015) 23.

[22] Y. Li, X. Zhu, X. Zhang, J. Fu, Z. Wang, T. Chen, X. Zhao, Characterization of genome-

reduced Bacillus subtilis strains and their application for the production of guanosine and

thymidine, Microbial cell factories, 15 (2016) 94.

[23] D. Zhu, Y. Fu, F. Liu, H. Xu, P.E. Saris, M. Qiao, Enhanced heterologous protein

productivity by genome reduction in Lactococcus lactis NZ9000, Microbial cell factories, 16

(2017) 1.

[24] J.H. Lee, B.H. Sung, M.S. Kim, F.R. Blattner, B.H. Yoon, J.H. Kim, S.C. Kim, Metabolic

engineering of a reduced-genome strain of Escherichia coli for L-threonine production,

Microbial cell factories, 8 (2009) 2.

[25] J. Schellenberger, R. Que, R.M. Fleming, I. Thiele, J.D. Orth, A.M. Feist, D.C. Zielinski, A.

Bordbar, N.E. Lewis, S. Rahmanian, J. Kang, D.R. Hyduke, B.O. Palsson, Quantitative

prediction of cellular metabolism with constraint-based models: the COBRA Toolbox v2.0,

Nature protocols, 6 (2011) 1290-1307.

[26] G.C. diCenzo, A. Checcucci, M. Bazzicalupo, A. Mengoni, C. Viti, L. Dziewit, T.M. Finan,

M. Galardini, M. Fondi, Metabolic modelling reveals the specialization of secondary replicons

for niche adaptation in Sinorhizobium meliloti, Nature communications, 7 (2016) 12219.

[27] G.C. diCenzo, A.B. Benedict, M. Fondi, G.C. Walker, T.M. Finan, A. Mengoni, J.S.

Griffitts, Robustness encoded across essential and accessory replicons in an ecologically

versatile bacterium, bioRxiv, (2017).

[28] G.C. diCenzo, Z. Muhammed, M. Osteras, S.A.P. O'Brien, T.M. Finan, A Key Regulator of

the Glycolytic and Gluconeogenic Central Metabolic Pathways in Sinorhizobium meliloti,

Genetics, (2017).

[29] G.C. diCenzo, T.M. Finan, Genetic redundancy is prevalent within the 6.7 Mb

Sinorhizobium meliloti genome, Molecular genetics and genomics : MGG, 290 (2015) 1345-

1356.

[30] J.A. Bartell, A.S. Blazier, P. Yen, J.C. Thogersen, L. Jelsbak, J.B. Goldberg, J.A. Papin,

Reconstruction of the metabolic network of Pseudomonas aeruginosa to interrogate virulence

factor synthesis, Nature communications, 8 (2017) 14631.

[31] K.H. Turner, A.K. Wessel, G.C. Palmer, J.L. Murray, M. Whiteley, Essential genome of

Pseudomonas aeruginosa in cystic fibrosis sputum, Proceedings of the National Academy of

Sciences of the United States of America, 112 (2015) 4110-4115.

[32] T. Shlomi, M.N. Cabili, M.J. Herrgard, B.O. Palsson, E. Ruppin, Network-based prediction

of human tissue-specific metabolism, Nature biotechnology, 26 (2008) 1003-1010.

[33] P.A. Jensen, K.A. Lutz, J.A. Papin, TIGER: Toolbox for integrating genome-scale

metabolic models, expression data, and transcriptional regulatory networks, BMC systems

biology, 5 (2011) 147.

[34] T. Galili, A. O'Callaghan, J. Sidi, C. Sievert, heatmaply: an R package for creating

interactive cluster heatmaps for online publishing, Bioinformatics, (2017).

[35] M. Kanehisa, Y. Sato, M. Kawashima, M. Furumichi, M. Tanabe, KEGG as a reference

resource for gene and protein annotation, Nucleic acids research, 44 (2016) D457-462.

[36] J. Huerta-Cepas, K. Forslund, L.P. Coelho, D. Szklarczyk, L.J. Jensen, C. von Mering, P.

Bork, Fast Genome-Wide Functional Annotation through Orthology Assignment by eggNOG-

Mapper, Molecular biology and evolution, 34 (2017) 2115-2122.

[37] G.C. diCenzo, M. Zamani, A. Cowie, T.M. Finan, Proline auxotrophy in Sinorhizobium

meliloti results in a plant-specific symbiotic phenotype, Microbiology, 161 (2015) 2341-2351.

[38] S.M. Keating, B.J. Bornstein, A. Finney, M. Hucka, SBMLToolbox: an SBML toolbox for

MATLAB users, Bioinformatics, 22 (2006) 1275-1277.

[39] B.J. Bornstein, S.M. Keating, A. Jouraku, M. Hucka, LibSBML: an API library for SBML,

Bioinformatics, 24 (2008) 880-881.

[40] M. Fondi, I. Maida, E. Perrin, A. Mellera, S. Mocali, E. Parrilli, M.L. Tutino, P. Lio, R.

Fani, Genome-scale metabolic reconstruction and constraint-based modelling of the Antarctic

bacterium Pseudoalteromonas haloplanktis TAC125, Environmental microbiology, 17 (2015)

751-766.

[41] L. Presta, E. Bosi, L. Mansouri, L. Dijkshoorn, R. Fani, M. Fondi, Constraint-based

modeling identifies new putative targets to fight colistin-resistant A. baumannii infections,

Scientific reports, 7 (2017) 3706.

[42] J.D. Orth, T.M. Conrad, J. Na, J.A. Lerman, H. Nam, A.M. Feist, B.O. Palsson, A

comprehensive genome-scale reconstruction of Escherichia coli metabolism--2011, Molecular

systems biology, 7 (2011) 535.

Table 1. Summary of the sizes of the produced core models relative to the parent model

(iGD1575) and the manually prepared core model (iGD726).

Model	Genes	Reactions	Metabolites
iGD1575	1577	1828	1579
iGD726	728	681	703 *
Core model A (without RNA-seq, FBA)	488	574	578
Core model B (with RNA-seq, FBA)	532	614	601
Core mdoel C (without RNA-seq, MOMA)	490	581	584
Core model D (with RNA-seq, MOMA)	532	602	590
FASTCORE	732	555	544
minNW	650	487	509
GIMME	546	1211 †	1165 †

483 * As iGD726 contains an updated biomass with a more complex membrane lipid composition,

484 this model is expected to have more metabolites than core models produced from iGD1575.

485 † The high number of reactions/metabolites is at least partially due to the presence of the

486 complete complement of exchange reactions.

Table 2. Parameters and summary statistics for Tn-Core runs.

Model	Gene Count *	Reaction Count *	Metabolite Count *	Growth Thresh	Pre-set EGs	RNA-seq	Method	Unique Gene Sets †	Unique Reaction Sets †	Variable Genes ¥	Variable Reactions ¥	Iteration run time (s) ø
iGD1575	1577 (1130)	1828 (920)	1579 (710)	10	No	No	FBA	25,000	25,000	777	416	26.7
				25	No	No	FBA	25,000	25,000	776	417	24.0
				50	No	No	FBA	25,000	25,000	773	413	23.8
				75	No	No	FBA	25,000	25,000	773	415	23.7
				90	No	No	FBA	25,000	25,000	771	412	23.9
				99	No	No	FBA	25,000	25,000	763	389	24.1
				10	Yes	No	FBA	25,000	25,000	471	296	20.6
				10	No	Yes	FBA	25,000	24,999	399	262	18.5
				10	Yes	Yes	FBA	25,000	24,995	265	163	17.1
				50	Yes	No	FBA	25,000	25,000	472	295	19.6
				50	No	Yes	FBA	25,000	25,000	402	265	18.4
				50	Yes	Yes	FBA	25,000	24,995	292	192	17.3
				10	No	No	MOMA	25,000	25,000	837	456	79.7
				50	No	No	MOMA	25,000	25,000	837	455	79.9
				99	No	No	MOMA	25,000	25,000	773	384	76.6
				50	Yes	No	MOMA	25,000	25,000	531	328	64.6
				50	Yes	Yes	MOMA	25,000	24,999	389	280	58.2
iPAE1160	1148 (808)	1496 (888)	1284 (643)	10	No	No	FBA	25,000	25,000	470	364	16.8
				50	No	No	FBA	25,000	25,000	476	362	17.1
				99	No	No	FBA	25,000	25,000	415	310	16.9
				10	Yes	No	FBA	25,000	24,997	319	258	14.8
				50	Yes	No	FBA	25,000	24,999	321	259	14.6
iJO1366	1367 (1255)	2583 (2333)	1805 (1578)	10	No	No	FBA	25,000	25,000	607	814	60.7
				50	No	No	FBA	25,000	25,000	510	719	59.4
				99	No	No	FBA	25,000	25,000	363	381	57.6
iLP844	887 (618)	1628 (816)	1518 (589)	10	No	No	FBA	25,000	25,000	340	303	11.3
				50	No	No	FBA	25,000	25,000	337	300	11.4
				99	No	No	FBA	25,000	25,000	304	263	11.1
iMF721	723 (611)	1324 (921)	1134 (688)	10	No	No	FBA	25,000	25,000	329	397	11.0
				50	No	No	FBA	25,000	25,000	338	399	10.7
				99	No	No	FBA	25,000	25,000	300	361	10.2

* The first set of numbers are based on the full starting model, while those in parentheses are based on the reduced model (following dead-end removal) that is used in the core model generation.

† Following 25,000 iterations, how many unique sets of genes or reactions were present in the core models.

¥ Following 25,000 iterations, how many genes or reactions were found to be variably present or absent in the core models.

ø Length of time in seconds required to produce a single core model. Total running time is approximately equal to the run time per iteration multiplied by the number of iterations and divided by the number of parallel pools used.

500 **Figure 1. Schematic representation of the Tn-Core pipeline.** Dashed lines represent optional

501 steps.

Figure 2. Evaluation of core metabolic redundancy with Tn-Core. The six primary matrixes generated by Tn-Core are shown. Tn-Core was run using the *S. meliloti* iGD1575 genome-scale metabolic reconstruction, with 25,000 iterations, a growth threshold of 10%, without essential gene pre-identified, and without RNA-seq data. Gene (**a**) and reaction (**b**) presence matrixes are shown for 1,000 of the randomly produced core models. Blue indicates the gene/reaction is present, white indicates the gene/reaction is absent. Gene (**c**) and reaction (**d**) co-occurrence matrixes are shown for the genes/reactions variably present in the 25,000 core models. (**e**) The legend for the co-occurrence matrixes is shown. The scale represent a Chi-squared statistic that summarizes if the gene or reaction pair is more (yellow) or less (blue) likely to occur in the same core model than by chance. Gene (**f**) and reaction (**g**) scatter plots displaying the correlation between the percentage of core models containing the gene/reaction when made using a growth threshold of 10% or 99%. Genes/reactions either present in all models or in no models are not included.

Figure 3. Comparison of central carbon metabolism of full and core metabolic models. This figure represents the full *S. meliloti* genome-scale metabolic reconstruction (iGD1575), the manually produced core metabolic reconstruction (iGD726), four core models produced from iGD1575 using Tn-Core (Core Model A [with Tn-seq, without RNA-seq, FBA algorithm], Core Model A [with Tn-seq, with RNA-seq, FBA algorithm], Core Model A [with Tn-seq, without RNA-seq, MOMA algorithm], Core Model A [with Tn-seq, with RNA-seq, MOMA algorithm]), and core models derived from iGD1575 using the FASTCORE, minNW, or GIMME algorithms. Representative genes from central carbon metabolism and the ATP synthase are shown. For each gene, a circle is shown if the gene is present in the model, and the circle is coloured according to the effect of deleting the gene on the growth rate of the model (determined using the MOMA algorithm); a value of 100 means no growth impact, a value of 0 means the gene deletion is lethal.