Title:

Naked mole-rat transcriptome signatures of socially-suppressed sexual maturation and links

of reproduction to aging

Authors:

Martin Bens[a,1], Karol Szafranski[a,1], Susanne Holtze[b], Arne Sahm[a], Marco Groth[a], Hans A.

Kestler[a,c], Thomas B. Hildebrandt[b,2], Matthias Platzer[a,2]

Affiliations:

[a] Leibniz Institute on Aging - Fritz Lipmann Institute, Beutenberg Str. 11, 07743 Jena, Germany

[b] Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Straße 17, 10315 Berlin, Germany

[c] Institute of Medical Systems Biology, Ulm University, James-Franck-Ring, 89069 Ulm, Germany

1 authors contributed equally to this work

2 shared senior authors

Corresponding Author:

Martin Bens (martin.bens@leibniz-fli.de)

Keywords:

reproduction, sexual maturation, aging, RNA-seq, naked-mole rat, eusociality

Abstract

Naked mole-rats (NMRs) are eusocially organized in colonies. Although breeders carry the additional metabolic load of reproduction, they are extremely long-lived and remain fertile throughout their lifespan. Comparative transcriptome analysis of ten organs from breeders and non-breeders of the eusocial long-lived NMR and the polygynous shorter-lived guinea pig provide comprehensive and unbiased molecular evidence that sexual maturation in NMR is socially suppressed. After transition into breeders, transcriptomes are markedly sex-specific, show pronounced feedback signaling via gonadal steroids and have similarities to reproductive phenotypes in African cichlid fish. Further, NMRs show functional enrichment of status-related expression differences associated with aging. Lipid metabolism and oxidative phosphorylation – molecular networks known to be linked to aging – were identified among most affected gene sets. Further, a transcriptome pattern associated with longevity is reinforced in NMR breeders contradicting the disposable soma theory of aging and potentially contributing to their exceptional long life- and healthspan.

Introduction

The naked mole-rat (NMR, *Heterocephalus glaber*) has become increasingly popular as an animal model in a variety of research fields due to its unique biology. This includes an exceptionally long lifespan and resistance to cancer [1, 2]. According to The Animal Ageing and Longevity Database (AnAge) [3] the maximum recorded lifespan is 31 years, i.e. 368% of the prediction based on body mass. NMRs stay fertile throughout their long and healthy life, i.e. show an extraordinary long life- and healthspan [4]. This lifelong fertility becomes even more astonishing, considering the extreme reproductive skew in NMR colonies. Like eusocial insects, NMRs are socially organized in colonies consisting of a pair of reproducing animals (breeders, queen and pasha) and up to 300 subordinates (non-breeders, female and male workers) [5]. However, although workers are in principle capable of reproduction [6, 7], sexual maturation is suppressed through the behavior of the dominating queen [8]. Non-breeding animals of both sexes are the backbone of the social organization of the colony and take care of foraging, brood care, colony defense and digging [9].

Naturally, new NMR colonies originate from fissioning of existing colonies or formation of new ones by dispersers that leave their natal colony [10, 11]. When under laboratory conditions non-breeders are removed from the colony and paired with the opposite sex, they have the capability to ascend into breeders. This process is accompanied with physiological and behavioral changes, and results in the formation of a new colony [5, 6]. Remarkably, despite the queen's enormous metabolic load of producing a large litter every three months and being exclusively in charge of lactation [12], data from wild and laboratory NMR colonies indicate that breeders live longer than their non-breeding counterparts [13, 14]. This contrasts the disposable soma theory of aging, which hypothesizes that energy is either allocated to body maintenance or reproduction [15].

In NMR, the reproductive suppression in female non-breeders is mediated through inhibition

of gonadotropin-releasing hormone (GnRH) secretion from the hypothalamus [6]. This in

turn leads to an inhibition of follicle stimulating hormone (FSH) and luteinizing hormone

(LH) released by the pituitary gland and causes a block of ovulation. Also for male non-

breeders, reproductive suppression is caused by inhibition of GnRH secretion, which results

in lower levels of urinary testosterone and plasma LH [7]. The impact, however, is less

profound compared to females as spermatogenesis is attenuated, but not entirely suppressed

[16]. Nevertheless, weight of testis and number of active spermatozoa is higher in breeders [7,

17]. The role of GnRH in mediating environmental cues to allow or block reproduction is well

described in a variety of species [18, 19].

The NMR can be regarded as a neotenic species and the prolonged retention of juvenile

features has been linked to its longevity [20]. In comparison to mice, e.g. postnatal NMR brain

maturation occurs at slower rate [12] and puberty is delayed. Female and male NMRs may

reach sexual maturity at 7.5 to 12 months of age [21]. In the colony, however, the queen

suppresses sexual maturation in both non-breeding males and females by aggressive social

behavior [8] and can delay – independently of neoteny - the puberty of female workers for life

[22]. Thus, sexual dimorphism is almost absent among non-breeding NMRs [23, 24]. Both

sexes show almost no difference in morphology, including body mass, body size and even

external genitalia, as well as no behavioral differences, in the sense that non-breeders

participate and behave equally in all colony labors [25]. Nevertheless, these features are

correlated with colony rank. Most profound differences can be observed comparing NMR

queens vs. non-breeders, reflected in morphological differences, such as an elongated spine

and higher body mass of queens, as well as behavioral differences, such as increased

aggressiveness, copulation and genital nuzzling [25].

In this work, we characterized the transcript signature of reproductive status (breeder vs. non-breeder) in tissue samples of ten organs or their substructures (hereinafter called for simplicity "tissues") from both sexes using RNA-seq. We contrast the NMR results with the transcript profiles of corresponding samples of guinea pig (GP, *Cavia porcellus*), a closely related, polygynous, not long-lived rodent species (AnAge: 12 years maximal longevity, 89% of the prediction based on body mass). We specifically focused our analyses on transcriptome signatures of the socially-suppressed sexual maturation in NMR as well as on differentially expressed genes (DEGs) that may contribute to the exceptional long life- and healthspan of NMR breeders.

Results

To gain molecular insights into the fascinating combination of NMR phenotypes, in particular their eusocial reproduction, lifelong fertility, extraordinary healthspan and longevity, we aimed to collect a comprehensive set of tissue samples for male and female breeders and non-breeders of NMR and GP - six biological replicates each. Towards this, NMR non-breeders were removed from their natal colony, paired with an unrelated animal of the opposite sex from a second colony and thereby turned into breeders. Respective male and female litter siblings remained in the two colonies as non-breeder controls. Time to first litters averaged in 6.5±4.9 (±SD) months and duration of pregnancies was approximately 70 days. GP breeders and non-breeders were housed as pairs of opposite or same sex, respectively. For this species, the time to first litters was 4.1±0.8 month and the pregnancies lasted about 68 days.

Female breeders gave birth to two litters each, with two exceptions. One NMR female was pregnant at least twice (ultrasonographically verified), but never gave birth to live offspring, and another gave birth to three litters, due to a pregnancy fathered by one of her sons. At

time of sampling, NMRs and GPs reached an age of 3.4±0.5 and 0.9±0.1 years, respectively

(Table S1).

Tissue and species are the major determinants of transcriptomes

To compare gene expression between reproductive statuses (breeder vs. non-breeder) in

NMR and GP, we performed RNA-seq of ten different tissue samples (heart - Hrt, skin - Skn,

liver - Lvr, kidney - Kid, cerebellum - Cer, hypothalamus - Hyp, pituitary - Pit, thyroid - Thy,

adrenal - Adr, and gonads – Gon, represented by either ovary – Ova or testis – Tes) from

24 animals for each species (six males, six females per status; Fig. S1). Seven of the 480

samples (1.5%) had to be excluded for different reasons (Tables S2, S3). On average (±SD), we

obtained per sample 27.6±3.6 million high-quality reads with 84.1±16.1% unique mapping rate

(Table S4). The grand mean of pairwise Pearson correlation within the 40 replicate groups (2

statuses x 2 sexes x 10 tissues per sex) was 0.981±0.013 and 0.984±0.01 for NMR and GP,

respectively, indicating high consistency between replicate samples (Table S5).

Based on these data, unsupervised hierarchical clustering gave a similar cluster hierarchy of

tissues for both species (Fig. S2). Brain tissues are grouped (Pit as a sister group to Cer and

Hyp); Kid and Thy are sister groups to the cluster of Adr and Ova. The results are confirmed

by principle component analysis, separating tissues by the first and species by the second

component (Figure 1). At this level of analysis, ovary was the only tissue, which showed a

separation of samples with respect to breeding status. Together, this indicates that: (i) tissue

source is dominant over other biological variables such as species, sex and status and (ii) the

impact of sex and status on transcriptome profiles is subtle.

Figure 1: Principle component (PC) analysis of groups based on mean expression levels (four groups per tissue and species: 2 sexes x 2 statuses, except gonads). Tissues are separated by PC1 and PC3, species by PC2.

Cross-species DEGs are enriched in aging-related genes

To further characterize the species differences between the long-lived NMR and the shorter-lived GP, we determined gene expression differences based on orthologous transcribed regions that show high sequence similarity. This filtering method avoided potentially misleading signals that may arise from assembly artifacts or the comparison of different transcript isoforms and identified 10,127 genes suitable for further analyses.

Figure 2: (A) Ras Related GTP Binding B (RRAGB) is consistently differentially expressed between species across all tissues. RRAGB is known to interact with mTORC1 complex [26, 27]. **(B)** Fibroblast Growth Factor Receptor 2 (FGFR2) shows opposing direction of expression between NMR and GP in breeders vs. non-breeders.

Across all tissues, we identified 18,000 significant expression differences (EDs; 9,651/8,349 higher/lower expressed in NMR) in 5,951 out of 10,127 genes (FDR<0.01, |log$_2$FC|>2; Table S6, Supplemental Data S1-S11). Among genes that are consistently differentially expressed across all tissues, we identified aging-related candidates. E.g. *RRAGB* (Ras related GTP binding B, Figure 2A) and *TMEM8C* (transmembrane protein 8C) are higher expressed in NMRs. *RRAGB* interacts with mTORC1 complex [26, 27]. *TMEM8C* is essential for muscle regeneration [28] and might be linked to the resistance to muscle loss in aged NMRs [29, 30].

To further assess the association of cross-species DEGs with aging, we examined their overlap with aging-related genes of human and mouse obtained from the Digital Ageing Atlas (DAA) [31]. This test revealed a significant overlap with DAA containing 1,056 genes (17.74% of DEGs; p=0.006, Fisher's exact test (FET); Table S7). The enrichment analysis of shared aging-related genes (Table S8) reveals that the top ranked GO term set is associated with lipid biosynthetic process (GO:0008610) (Fig. S3), a finding referring to existing links between lipid metabolism and lifespan [32, 33].

Figure 3: Euler diagrams showing overlaps of DEGs. **(A)** Female vs. male. **(B)** Breeder vs. non-breeder. **(C)** Gonads of NMR breeder vs. non-breeder and aging-related genes from the Digital Aging Atlas (DAA).

Sexual differentiation and maturation in NMR are delayed until transition from worker to breeder

The DEGs between sexes were determined within the groups of non-breeders and breeders for each tissue and species (Table 1, Table S9, Supplemental Data S12-S47). GP non-breeder females vs. males (GP-N-FvM) show over all tissues except gonads 1,713 significant expression differences (EDs) in 1,634 genes (FDR<0.01, Table 1), primarily in Adr (858 DEGs), Lvr (383), Thy (347) and Kid (109). Between female and male GP breeders (GP-B-FvM) 3,654/3,398 EDs/DEGs were observed. These transcriptome data confirm a clear sexual differentiation among sexually mature GPs that further increases after onset of breeding. Breeders have 790 DEGs in common with non-breeders (p<2.2x10^{-16}, FET; Figure 3A). Functional enrichment analysis of these shared genes reveals among the highest ranked GO sets immune system related terms (Fig. S5, Table S10).

192 **Table 1:** Numbers of DEGS identified in the different comparisons (FDR<0.01).

| | female vs. male | | | | breeder vs. non-breeder | | | |
| | non-breeder | | breeder | | females | | males | |
Tissue	GP	NMR	GP	NMR	GP	NMR	GP	NMR
Hrt	4	10	89	13	2	3	0	0
Skn	6	6	3	5	5	1	0	223
Lvr	383	4	235	10	71	0	9	1
Kid	109	6	106	21	1	4	0	0
Cer	1	5	2	25	0	15	0	0
Hyp	2	8	1	11	0	5	2	0
Pit	347	9	307	47	114	114	2	1
Thy	3	0	2,087	1,791	675	285	1	0
Adr	858	4	824	533	0	201	1	4
Gon	-	-	-	-	18	502	3	381
ED*	1,713	52	3,654	2,456	886	1,130	18	610
NR†	1,634	22	3,398	2,360	883	1,078	14	598

* significant expression differences across tissues

† non-redundant set of significant expression differences across tissues

Similarly to GP-B-FvM, NMR-B-FvM showed 2,456/2,360 sex-related EDs/DEGs (FDR<0.01),

mostly in Thy (1,791) and Adr (533). The overlap with GP-B-FvM is with 514 DEGs

considerable but does not reach significance (p=0.062, FET; Figure 3A). Nevertheless, these

data indicate basic similarities in sexual differentiation among breeders of both species.

Surprisingly, only 22 NMR-N-FvM EDs/DEGs were detected across all tissues (Table S11),

indicating an only minor sex differentiation between NMR non-breeding females and males

on the transcriptional level, consistent with the almost absent sexual dimorphism among non-

breeding NMRs [23, 24].

Status change of NMRs is accompanied by major changes in the endocrine system

The DEGs between breeders and non-breeders were determined within the same sex for each

species (Table 1; Table S12, Supplemental Data S48-S87). Females showed a similar amount of

EDs/DEGs in both species (GP-F-BvN: 886/883, NMR-F-BvN: 1,130/1,078) but have only 46

DEGs in common (Figure 3B). This is less than expected by chance, although not reaching

significance (p=0.075, FET for depletion) and indicates that the molecular signature of the

transition from female non-breeder to breeder is different in both species. E.g. in GP-F-BvN,

only 18 DEGs are observed in Ova and none in Adr while in NMR-F-BvN, these tissues show

most of the differences with 502 and 201 DEGs, respectively. Functional enrichment analysis

of DEGs in NMR Ova identifies reproductive structure development (GO:0048608) as the

highest ranked category (Fig. S6, Table S13). The same analysis in Adr revealed an obvious

directionality in expression changes. DEGs are preferentially upregulated among highest

ranked GO term sets (Fig. S7, Table S14), e.g. in reproduction (GO:0000003, 24 of 28) and

endocrine system development (GO:0035270, 17 of 18). Tallying with this, Cer DEGs are

enriched and upregulated in steroid metabolic process (GO:0008202, 6 of 6 upregulated) and

response to hormones (GO:0009725, 7 of 7) (Fig. S8, Table S15).

In male GPs, status-related differences (GP-M-BvN) were almost absent across all tissues

(only 18 EDs/14 DEGs). In contrast, NMR-M-BvN showed 610/598 EDs/DEGs, predominantly

in Tes (381) and Skn (223). NMRs share 55 status-related DEGs in both sexes (p=0.008, FET;

Table S16), while the few status-related changes in male GPs showed no overlap with those in

females (Figure 3B). Among shared DEGs in NMRs, 10 genes involved in endocrine signaling

were identified, including *SSTR3* (somatostatin receptor), *TAC4* (tachykinin), *PRDX1*

(peroxiredoxin 1) and ACPP (acid phosphatase, prostate), as well as in general signaling via

cAMP signaling (three genes) and through G-protein coupled receptors (four genes) further

underlining that social status transition in NMR is associated with changes in the endocrine

system.

Mitochondrial genes show opposed expression changes in Tes and Skn after status change

of NMR males

In NMR-M-BvN Tes, functional enrichment analysis revealed as highest ranked metabolism-

and energy-related GO sets. DEGs enriched therein are mostly upregulated, e.g. lipid

biosynthetic process (GO:0008610, 75 of 82 genes) and oxidation-reduction process

(GO:0055114, 64 of 64) (Fig. S9, Table S17). Furthermore, we observed an upregulation of

response to stimulus (GO:0050896, 79 of 106), in line with an upregulation of steroid

metabolic process (GO:0008202, 28 of 28) included in lipid biosynthetic process set. In

accordance with the dominance of energy-related processes, DEGs are enriched and

preferentially upregulated in the top GO cellular component terms mitochondria

(GO:0005739, 61 of 62) and peroxisomes (GO:0005777, 13 of 13) (Table S18). Together, this

indicates increased demands of energy, e.g. to produce steroid hormones in Tes of NMR

breeders.

Similar to Tes, Skn showed enrichment of energy-related processes (Table S19, Fig. S10).

However, GO term sets in Skn are mostly downregulated, including energy derivation by

organic compounds (GO:0015980, 36 of 38 genes) and oxidation-reduction process

(GO:0055114, 42 of 46). Consistently, this includes genes associated with mitochondria

(GO:0044429, 56 of 57) and respiratory chain (GO:0070469, 11 of 11).

Figure 4: Expression changes of nuclear genes encoding for mitochondrial respiratory chain complexes in status change of male NMRs (top) and GPs (bottom). DEGs are indicated by asterisk (*: FDR<0.05, **: FDR<0.01). Only NMRs show significant expression differences (skn: 46 genes, mean fold-change 0.76; tes 46 genes, mean fold-change 1.89).

The overlap between mitochondrial DEGs in Tes and Skn comprises 6 genes (p=7.27x10^{-8}, FET; Figure 4). Among common genes *PINK1* (PTEN induced putative kinase 1) is 1.5-fold upregulated in Tes and 2.5-fold downregulated in Skn, indicating a role in regulation of mitophagy [34] in both tissues.

To follow up the mitochondria-related findings, the 'mitonuclear transcript ratio' was determined as the read count ratio of mitochondrial-encoded genes versus nuclear-encoded genes. It differs largely between tissues and species (Table S20). Hrt showed the highest mitonuclear ratio, with a minor difference between species (NMR 30.1%, GP 30.4%). Tes showed the lowest ratio, particularly in GPs (NMR 5.5%, GP 0.9%), and a 43.1% increase in NMR-M-BvN (Fig. S11). This increase is accompanied with an upregulation of nuclear genes encoding for mitochondrial respiratory chain complexes (Figure 4; Table S21). The expected increase in ROS is compensated by an on average 1.59-fold upregulation of eight antioxidant

DEGs (Table S22). Consistently with functional enrichment analysis mentioned above, an opposing effect in Skn of NMR male breeders was observed, which showed a decline in mitonuclear ratio together with a downregulation of nuclear genes of the respiratory complexes (Figure 4). In line with the downregulation of the oxidative phosphorylation pathway (OXPHOS), a downregulation of antioxidant enzymes SOD2 (superoxide dismutase 2, 2.64-fold) and PRDX3 (peroxiredoxin 3, 2.14-fold) was observed. In general, downregulation of OXPHOS is linked to an extended lifespan [32, 35–37].

NMR status-related DEGs are enriched in aging-related genes

Guided by the exceptional long life- and healthspan of NMR breeders, we searched our transcriptome data for molecular signatures relating to this phenomenon. First, we found that only NMRs show status-related DEGs that are significantly enriched for aging-related genes from DAA (Table S23): males in Skn (55 genes; q=0.0012, FET) and Tes (80 genes, q=0.01), and females in Ova (127 genes; q=1.2×10^{-7}), Thy (59 genes, q=0.033) and Adr (43 genes, q=0.038). The significant overlap of 22 DEGs between NMR-F-BvN and NMR-M-BvN in Gon (p=0.0035, FET) contains nine aging-related genes (p=0.004, Figure 3C). In GP, only the non-redundant set of DEGs in GP-F-BvN showed a tendency of enrichment (160 genes, q=0.051), in contrast to NMRs, which showed enrichment in males (134 genes, q=4.5×10^{-5}) and females (245 genes, q=7×10^{-9}).

Second, we hypothesized that reproduction impacts the life expectancy of NMR and GP differently. Therefore, we searched for status-related DEGs that are shared in both species, but show opposing direction of expression. These genes might mark different coping mechanisms with the metabolic load of reproduction. As described above, the overlap of DEGs between species is very low (Table S24). Nevertheless, opposing direction of expression

change can be observed in Ova (1 of 2 shared DEGs), and female thyroid (8/8) and testis (1/1).

E.g. the fibroblast growth factor receptor 2 gene (*FGFR2*), linked to aging (AgeFactDB) [38], is

in Ova downregulated in NMRs, but upregulated in GPs (Figure 2B).

Third, and based on the assumption that status-related EDs with the greatest interspecies

difference (regardless of directionality) have an impact also on species-specific aging

trajectories of breeders, we determined enrichment of aging-related genes in the upper

quintile of those genes (Table S25). In males, we found significant enrichments of DAA genes

in Skn (q=7.04x10^{-7}) and Tes (q=0.0025); in females in Skn (q=0.0064), Hrt (q=0.0024), Pit

(q=0.0053) and Ova (q=0.0013). Further functional enrichment analysis of these aging-related

gene sets reveals differences between sexes. In males, the non-redundant set of genes shows

enrichment for lipid metabolism (GO:0006629), energy metabolism (energy derivation by

oxidation of organic compounds, GO:0015980; mitochondrial ATP synthesis coupled proton

transport, GO:0042776), glutathione metabolic process (GO:0006749) and immune system

(innate immune response-activating signal transduction, GO:0002758) (Table S26, Fig. S12).

Females showed enrichment in positive regulation of tumor necrosis factor production

(GO:0032760) and negative regulation of programmed cell death (GO:0043069) (Fig. S13,

Table S27).

Status related changes in NMR contradict the disposable some theory

Moreover, we assessed the connection of cross-species DEGs with expression changes that are

associated with status change in each species. Based on DEGs shared in two comparisons

(NMR vs. GP and breeder vs. non-breeder, FDR<0.05), we performed a correlation analyses of

fold changes in each species. We hypothesized – in agreement with the disposable soma

theory of aging – a negative impact of reproduction on lifespan for GP and – in contradiction

to this theory – an inverse effect for NMR. This was confirmed by opposing correlations

(combined p=4.2x10^{-9} (Lancaster procedure [39]), negative correlation for GP and positive

correlation for NMR (Figure 5). This means that DEGs with higher expression in NMR than

GP are preferentially upregulated in NMR breeders compared to non-breeders, and vice

versa; in other words, a transcriptome pattern associated with longevity (NMR vs. GP) is

reinforced by the status transition (breeder vs. non-breeder) and thus may contribute to the

exceptional long life- and healthspan of NMR breeders.

Figure 5: DEGs (FDR<0.05) occurring in both comparisons: cross-species (NMR vs. GP, y-

axes) and status change GPs (breeder vs. non-breeder, x-axes) separately for (A) NMRs and

(B). Correlation analysis between species shows opposing correlation (Lancaster procedure

[39], p=4.2x10^{-9}), while status-related DEGs in NMR are positively correlated with cross-

species DEGs (DEGs=3,820; spearman correlation=0.17, p=3.2×10^{-27}), status-related DEGs in

GP show a negative correlation (1,434; -0.1; 1.8×10^{-4}).

Discussion

In our comparative study of breeders vs. non-breeders of the eusocial long-lived NMR and

the polygynous and shorter-lived GP, we accumulated a comprehensive set of transcriptome

data to gain insights into molecular networks underlying naturally evolved interspecies differences in sexual maturation and links between reproduction and aging. Both species are able to breed year-round and produce four to five litters per year [40, 41]. Both species have a similar average gestation period of 70 days, which is long compared to similarly sized species. Notably, NMRs produce on average 10.5 offspring per litter, which is twice the number of offspring produced by similarly sized rodents and more than three times higher compared to GPs (average 3.2 offspring) [40, 41]. This underscores the apparent contradiction of the NMR queen's enormous metabolic load and extraordinary long life- and healthspan [4] to the disposable soma theory of aging [15], indicating that natural ways to extended healthspan remains to be uncovered in NMRs.

A first study to identify adaptations to unique NMR traits at the transcriptome level compared liver gene expression of young adult non-breeding male NMRs and mice [42]. Higher NMR transcript levels were observed for genes associated with oxidoreduction and mitochondria. This present study, more comprehensive in several aspects (sex, breeding status, numbers of animals and tissue samples), is based on a comparison of NMR vs. GP, which are phylogenetically closer than NMR and mouse. It revealed that between NMR and GP 58.8% of the analyzed genes are differentially expressed and that these DEGs are significantly enriched for aging-related genes. Among the latter, the main functional commonality is their association with lipid metabolism. Links of this molecular network to longevity of NMR were also obtained by a parallel proteome comparison of NMR and GP livers [43]. This study also shows that NMR liver mitochondria exhibit an increased capacity to utilize fatty acids.

In respect to sex-specific molecular signatures among either breeders or non-breeders in NMR and GP, the main finding is a nearly complete absence of significant transcriptional

differences between sexes in non-breeding NMRs. This fits their grossly identical morphology

and identical behavior in stable colonies [25]. This is in stark contrast to non-breeding GPs of

an even younger age where we observed more than a thousand DEGs. GP non-breeders and

breeders share a large and highly significant number of sex-related DEGs. Among others,

these DEGs are enriched in GO terms related to steroid metabolism and immune system. The

effect of gonadal steroids on the immune system is well described in GPs and other mammals

[44]. After separation of NMR non-breeders from their colony, sexes became not only

distinguishable by morphology and behaviour [22, 25], but also by gene expression. This

differentiation on transcriptional level provides further molecular support for the previously

described suppression of sexual maturation in non-breeding adult NMRs by social stress [8,

22] and identified major changes in the endocrine system after status change in NMRs but not

in GPs (Supplement Text S1). Notably, we found no significant differences in gonadotropin-

related genes (Supplement Text S2), indicating similar transcript turnovers in both non-

breeders and breeders, and in line with previous results indicating that LH is stored in non-

breeders, ready to be released upon GnRH signalling [45].

Glucocorticoids have been linked to stress, reproduction and social behavior in a variety of

species, including members of muroidae, primates and cichlids [46–48]. In NMRs, however,

correlation between social status and urinary cortisol is not clear and seems to depend on

colony stability [49, 50]. Here, we observed a significant upregulation of *NR3C1*

(glucocorticoid receptor) in Tes of NMR male and Thy of female breeders (Supplemental

Data S55/67). Interestingly, this is in line with elevated expression of glucocorticoid receptor

in Tes in African cichlid breeders. Males can reversibly change between dominant and

subordinate phenotypes [51]. Similar to NMRs, only dominant phenotypes are reproductively

active. In male African cichlids, moreover, aggression is negatively correlated with expression

of *SSTR3* (somatostatin receptor 3) in Tes. Similarly, we observed in NMR breeder Tes

significant downregulation of *SSTR3* (Supplemental Data S67). This indicates that *SSTR3* may be associated with social dominance in NMRs as well.

As our study was primarily motivated by the exceptional long life- and healthspan of NMR breeders, we searched for evidence indicating that NMR status change has an impact on genes involved in aging. We found enrichment of aging-related genes in the non-redundant DEG sets of males and females, as well as enrichment in most tissues with at least 50 DEGs (male Skn and Tes; female Ova, Thy and Adr). This contrasts with our observations in GPs, which showed only a tendency of aging-relation for the non-redundant set of status-related DEGs in females.

Further, we observed significant tissue-specific changes in OXPHOS of male NMR breeders. While Tes showed an upregulation of nuclear-encoded mitochondrial genes and a respective increase in mitonuclear transcript ratio, Skn showed the opposite. Moreover, we observed significant enrichments of genes involved in fatty acid metabolism among status-related DEGs in both NMR tissues. Consistent with the role of mitochondria in lipid homeostasis and the observed directionality of changes in OXPHOS, fatty acid metabolism DEGs in Tes were preferentially up- and in Skn downregulated. While the increased mitochondrial activity in Tes probably complies with demands of energy for the production of sex steroids and their anabolic effect on physiology, such as growth of testis [17], the observed changes in Skn may indicate a link to the extraordinary healthspan of NMR breeders. Previously, it was observed that inhibition of complex I activity during adult life prolongs lifespan and rejuvenates the tailfin transcriptome in short-lived fish [36]. Enhanced lipid metabolism and reduced mitochondrial respiration were also linked to NMR longevity in a parallel liver proteome study comparing NMR vs. GP and old vs. young NMR [43].

Finally, we performed a correlation analysis between species (NMR vs. GP) and status (breeder vs. non-breeder) EDs confirming the basic hypothesis of the present work: in contrast to GP and in line with recent demographic studies performed in NMRs [14], the transition into breeders results in molecular signatures linked to extended life- and healthspan only in NMRs. Genes which are higher or lower expressed in NMR compared to GP are also preferentially up- or downregulated in NMR breeders (positive correlation) opposite to GPs (negative correlation). In other words, the positive correlation in NMR contradicts the disposable soma theory of aging, as EDs contributing to a long lifespan (higher/lower expression in NMR than GP) are preferentially increased in NMR breeders compared to non-breeders, while diminished in GPs as expected by this theory.

Taken together, our comparative transcriptome analysis of breeders versus non-breeders of the eusocial, long-lived NMR versus the polygynous and shorter-lived GP identifies molecular networks underlying socially regulated sexual maturation and naturally evolved extended life- and healthspan that encourage further functional and mechanistic investigations of these extraordinary NMR phenotypes.

Materials & Methods

Animals

Naked mole-rats. NMR colonies were kept inside a climatized box (2x1x1 m) in artificial burrow systems, consisting of eight cylindrical acrylic glass containers (diameter: 240 mm height: 285 or 205 mm). The latter functioned as variable nest boxes, food chambers or toilet areas, and were interconnected with acrylic tubes having an inner diameter of 60 mm. Husbandry conditions were stabile during the entire experimental period of 22 months. Temperature and humidity were adjusted to 27.0±2.0°C and 85.0±5.0%, respectively. In

general, the NMR colonies were kept in darkness except for 2 to 4 hours of daily husbandry

activities. Fresh vegetable food was provided daily and *ad libitum*. In addition, commercial rat

pellets (Vita special, Vitakraft GmbH, Bremen, Germany) were fed as an additional source of

protein and trace elements.

To turn them into breeders, randomly selected non-breeding animals derived from two long-

term (>4 years) established colonies of more than 50 individuals were separated and paired

with the opposite sex. As non-breeder controls, litter siblings of paired animals remained in

their colonies as workers. After the lactation period of the second set of live offspring the

tissue sampling was scheduled. To avoid further pregnancies in the females, male partners

were removed and euthanized 8-10 days postpartum. The tissue collection in the females took

place 40-50 days after the end of last pregnancy.

Guinea pigs. GPs (breed: Dunkin Hartley HsdDhl:DH, Harlan Laboratories, AN Venray,

Netherlands) were housed in standardized GP cages (length: 850 mm, width: 470 mm, height:

450 mm) in breeding pairs plus offspring or in same-gender pairs of two. Commercial guinea

pig pellets and commercial pet food hay (Hellweg Zooland GmbH, Berlin, Germany) were

provided together with vitamin C enriched water *ad libitum*. Housing temperature and

humidity were 18.0 ± 2.0°C and 45.0 ± 5.0%, respectively. A 12h light/dark regime was provided.

After an initial adaption period of 6 to 8 weeks the GPs were randomly divided in breeding

pairs or in same-gender pairs of two. The offspring were separated from their parents after

weaning (~3 weeks postpartum). Tissue collection was scheduled after the lactation period of

the second set of live offspring. To avoid further pregnancies in the females, male partners

were removed between eleven days before and seven days after birth of the second litter. The

tissue collection in the females took place 42-83 days after the end of last pregnancy.

For tissue collection, all animals were anaesthetized by 3% isoflurane inhalation anaesthesia

(Isofluran CP, CP-Pharma, Burgdorf, Germany) and euthanized by surgical decapitation.

Animal housing and tissue collection at the Leibniz Institute for Zoo and Wildlife Research

was compliant with national and state legislation (breeding allowance #ZH 156; ethics

approval G 0221/12 "Exploring long health span", Landesamt für Gesundheit und Soziales,

Berlin).

Sample collection, RNA Isolation and Sequencing

For *de novo* transcriptome assembly, animals were euthanized and ten tissue samples (heart –

Hrt (NMR only), skin - Skn, liver - Lvr, kidney - Kid, cerebellum - Cer, hypothalamus - Hyp,

pituitary - Pit, thyroid - Thy, adrenal - Adr, and gonads - Gon (testis – Tes /ovaries - Ova))

were collected from NMR and GP individuals, as described previously [52]. Strand-specific

RNA-seq were prepared using the TruSeq Stranded RNA LT Kit (Illumina), and 200-nt reads

were obtained using a HiSeq2500 (Illumina), as described previously [52].

For expression analysis, the same ten tissues were collected from NMR and GP breeders and

non-breeders. RNA was purified as described above. Library preparation was done using

Illumina's TruSeq RNA Library Prep Kit v2 kit following the manufacturer's description.

Quantification and quality check of the libraries was done using Agilent's Bioanalyzer 2100 in

combination with a DNA 7500 Kit (both Agilent Technologies). Sequencing was done on a

HiSeq 2500 running the machine in 51 cycle, single-end, high-output mode by multiplexing

seven samples per lane. Demultiplexing and extraction of read information in FastQ format

was done using the tool bcl2astq v1.8.4 (provided by Illumina).

Data Analysis

De novo transcriptome assembly and annotation for GP was performed as described in [52].

Briefly, overlapping paired-end reads were joined into single fragments and then assembled

by Trinity [53]. Gene symbols were assigned to the assembled transcripts by similarity to

human transcripts using FRAMA [52].

As a reference for RNA-seq data mapping the public NMR (Bioproject PRJNA72441) [54] and

GP genomes (UCSC, cavpor3) were used. Reference transcript sets of NMR and GP were

mapped to the corresponding genome in two steps: BLAT (v36) [55] was used to identify the

locus and then SPLIGN (v1.39.8) [56] was applied to splice align the transcript sequence

within BLAT locus. RNA-seq data were aligned to the corresponding reference genome

utilizing STAR (v2.4.1d) [57] with a maximum mismatch of 6% and a minimum aligned

length of 90%. Reads mapped to multiple loci were discarded. Gene expression was

quantified using HTSEQ (v0.6.1p1) [58] based on the aligned reference transcripts. The

pairwise Pearson correlation between biological replicates was calculated based on 16,339 and

16,009 genes in NMR and GP, respectively (Table S5).

The 'mitonuclear transcript ratio' was calculated as the read count ratio of 13 mitochondrial-

encoded genes versus all nuclear-encoded genes.

PosiGene was applied to the transcriptome of human, NMR and GP with the parameter

'-prank=0 -max_anchor_gaps_hard=100 -rs=NMR' to determine orthologous transcribed

regions in NMR and GP having a protein identity >70%. RNA-seq data were aligned to the

corresponding transcriptomes utilizing bowtie2 (2.2.9) [59] with the parameter

'-very-sensitive-local'.

DESeq2 (v1.6.3) [60] was used to identify DEGs. A false discovery rate <0.01 (FDR; Benjamini

Hochberg corrected p values, [61]) was used for the identification of significant DEGs.

Gene Ontology analysis was performed using the web interface of GoMiner (Database build

2011-01) based on the functional annotation of human genes (UniProt) [62]. A FDR<0.05 was

used for the identification of significant GO terms that were summarized by REVIGO

(parameter SimRel=0.5) into non-redundant GO term sets [63]. GO term sets were then

ranked by number of summarized GO terms and number of changed genes. KEGG analysis

was performed using Fisher's exact test and significant pathways were identified using

FDR<0.05. KEGG results were redundant to Gene Ontology analysis and therefore not shown.

Overlap between gene sets was determined with Fisher's exact test (FET) using the one-sided

option. Generally, we tested for enrichment if not otherwise stated.

We obtained 3,009 aging-related genes in human and mouse from the Digital Ageing Atlas

(DAA) [31]. The corresponding counterparts in the NMR (2,588) and GP (2,539) were used for

enrichment analysis and results were corrected for multiple testing (FDR). P-values corrected

for multiple testing are indicated by q and nominal p-values by p.

To examine the connection between reproduction and aging in both species, we determined

the difference in log2-fold-change (breeders vs. non-breeders) of NMR and GP. For fold-

changes moving in opposite directions between species, we calculated the absolute difference

($| \log_2 NMR_{BvsN} - \log_2 GP_{BvsN} |$), and for fold-changes moving in the same direction, higher

fold-changes in NMR-BvN were rewarded ($|\log_2 NMR_{BvsN}| - |\log_2 GP_{BvsN}|$). The 20%-

quantile of genes having the greatest difference was determined separately for (i) the

complete gene set and for genes showing (ii) opposing and (iii) unidirectional fold-changes.

All sets were tested for enrichment of aging-related genes.

Statistical analyses were performed in R (version v3.1.2).

Data Availability

RNA-seq data for gene expression profiling were deposited at Gene Expression Omnibus

(GSE98719). RNA-seq data for *de novo* assembly were deposited at Sequence Read Archive

(SRP104222, SRP061363) and the corresponding gene collection is available as a gff3-file

(ftp://genome.leibniz-fli.de/pub/nmr2017/).

Acknowledgements

We like to acknowledge Angelika Kissmann and Jette Ziep for their professional help with

the animal husbandry. We also thank Michaela Wetzel for her logistic help during tissue

sampling and storage as well as for her support with the animal husbandry. We are grateful

to Ivonne Görlich for excellent technical assistance as well as Debra Weih and Steve

Hoffmann for critical reading of the manuscript.

Funding

This research was supported by the Leibniz-Gemeinschaft through the

Senatsausschusswettbewerb (SAW), grant SAW-2012-FLI-2, and the Deutsche

Forschungsgemeinschaft (DFG), grant PL 173/8-1.

Competing interests

None declared.

Author Contributions

TBH, MP and KS conceived the project. SH, TBH, MB, MG and KS performed the animal study, sampling and sequencing experiments. Data analysis and interpretation were performed by MB, KS, SH, AS, HK, MP and TBH. The manuscript was written by MB and KS, and revised by all authors. The manuscript submitted by MB; all authors reviewed and approved the submitted manuscript. TBH and MP are joint senior authors.

Abbreviations

DEG	differentially expressed gene
ED	expression differences
FDR	false discovery rate
FET	Fisher's exact test
GO	Gene Ontology
GP	guinea pig
GP-B-FvM	comparison of GP Breeder Females vs. Males
GP-F-BvN	comparison of GP Female Breeders vs. Non-breeders
GP-M-BvN	comparison of GP Male Breeders vs. Non-breeders
GP-N-FvM	comparison of GP Non-breeder Females vs. Males
NMR	naked mole-rat
NMR-BvN	comparison of NMR Breeders vs. Non-breeders (both sexes)
NMR-B-FvM	comparison of NMR Breeder Females vs. Males
NMR-F-BvN	comparison of NMR Female Breeders vs. Non-breeders
NMR-M-BvN	comparison of NMR Male Breeders vs. Non-breeders
NMR-N-FvM	comparison of NMR Non-breeder Females vs. Males

References

1. Kim EB, Fang X, Fushan A a., Huang Z, Lobanov A V, Han L, et al. Genome sequencing reveals insights into physiology and longevity of the naked mole rat. Nature. 2011;479:223–7.

2. Tian X, Azpurua J, Hine C, Vaidya A, Myakishev-Rempel M, Ablaeva J, et al. High-

molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. Nature.

2013;499:346–9. doi:10.1038/nature12234.

3. Tacutu R, Craig T, Budovsky A, Wuttke D, Lehmann G, Taranukha D, et al. Human Ageing

Genomic Resources: Integrated databases and tools for the biology and genetics of ageing.

Nucleic Acids Res. 2013;41:D1027–33. doi:10.1093/nar/gks1155.

4. Buffenstein R, Jarvis JUM. The naked mole rat--a new record for the oldest living rodent.

Sci aging Knowl Environ. 2002;2002:pe7. doi:10.1126/sageke.2002.21.pe7.

5. Jarvis JU. Eusociality in a mammal: cooperative breeding in naked mole-rat colonies.

Science. 1981;212:571–3. doi:10.1126/science.7209555.

6. Faulkes CG, Abbott DH, Jarvis JU. Social suppression of ovarian cyclicity in captive and

wild colonies of naked mole-rats, Heterocephalus glaber. Reproduction. 1990;88:559–68.

doi:10.1530/jrf.0.0880559.

7. Faulkes CG, Abbott DH, Jarvis JUM. Social suppression of reproduction in male naked

mole-rats, Heterocephalus glaber. J Reprod Fertil. 1991;91:593–604. doi:10.1530/jrf.0.0910593.

8. Smith TE, Faulkes CG, Abbott DH. Combined olfactory contact with the parent colony and

direct contact with nonbreeding animals does not maintain suppression of ovulation in

female naked mole-rats (Heterocephalus glaber). Horm Behav. 1997;31:277–88.

doi:10.1006/hbeh.1997.1384.

9. Burda H, Honeycutt RL, Begall S, Locker-Grutjen O, Scharff A. Are naked mole-rats

eusocial and if so, why? Behav Ecol Sociobiol. 2000;47:293–303.

10. Braude S. Dispersal and new colony formation in wild naked mole-rats: evidence against

inbreeding as the system of mating. Behav Ecol. 2000;11:7–12.

11. O'Riain MJ, Jarvis JU, Faulkes CG. A dispersive morph in the naked mole-rat. Nature.

1996;380:619–21.

12. Orr ME, Garbarino VR, Salinas A, Buffenstein R. Extended Postnatal Brain Development

in the Longest-Lived Rodent: Prolonged Maintenance of Neotenous Traits in the Naked Mole-

Rat Brain. Front Neurosci. 2016;10:504. doi:10.3389/fnins.2016.00504.

13. Hochberg ME, Noble RJ, Braude S. A Hypothesis to Explain Cancers in Confined Colonies

of Naked Mole Rats. bioRxiv. 2016.

14. Ruby JG, Smith M, Buffenstein R. Naked mole-rat mortality rates defy Gompertzian laws

by not increasing with age. Elife. 2018;7:e31157. doi:10.7554/eLife.31157.

15. Kirkwood TB. Evolution of ageing. Nature. 1977;270:301–4.

16. van der Horst G, Maree L, Kotzé SH, O'Riain MJ. Sperm structure and motility in the

eusocial naked mole-rat, Heterocephalus glaber: a case of degenerative orthogenesis in the

absence of sperm competition? BMC Evol Biol. 2011;11:351.

17. Faulkes CG, Trowell SN, Jarvis JU, Bennett NC. Investigation of numbers and motility of

spermatozoa in reproductively active and socially suppressed males of two eusocial African

mole-rats, the naked mole-rat (Heterocephalus glaber) and the Damaraland mole-rat

(Cryptomys damarensis). J Reprod Fertil. 1994;100:411–6. doi:10.1530/jrf.0.1000411.

18. Abbott DH, Hodges JK, George LM. Social status controls LH secretion and ovulation in

female marmoset monkeys (Callithrix jacchus). J Endocrinol. 1988;117:329–39.

doi:10.1677/joe.0.1170329.

19. White S, Nguyen T, Fernald RD. Social regulation of gonadotropin-releasing hormone. J

Exp Biol. 2002;205:2567–81.

20. Skulachev VP, Holtze S, Vyssokikh MY, Bakeeva LE, Skulachev M V., Markov A V., et al.

Neoteny, Prolongation of Youth: From Naked Mole Rats to "Naked Apes" (Humans). Physiol

Rev. 2017;97:699–720. doi:10.1152/physrev.00040.2015.

21. Sherman PW, Jarvis JUM, Alexander RD. Reproduction in Naked Mole-Rats. In: The

Biology of the Naked Mole-rat. Princeton University Press; 1991. p. 384–425.

22. Dengler-Crish CM, Catania KC. Phenotypic plasticity in female naked mole-rats after

removal from reproductive suppression. J Exp Biol. 2007;210 Pt 24:4351–8.

23. Pinto M, Jepsen KJ, Terranova CJ, Buffenstein R. Lack of sexual dimorphism in femora of

the eusocial and hypogonadic naked mole-rat: a novel animal model for the study of delayed

puberty on the skeletal system. Bone. 2010;46:112–20. doi:10.1016/j.bone.2009.08.060.

24. Peroulakis ME, Goldman B, Forger NG. Perineal muscles and motoneurons are sexually

monomorphic in the naked mole-rat (Heterocephalus glaber). J Neurobiol. 2002;51:33–42.

25. Sherman PW, Jarvis JUM, Alexander RD. An Ethogram for the Naked Mole-Rat: Nonvocal

Behaviors. In: The Biology of the Naked Mole-rat. Princeton University Press; 1991. p. 209–42.

26. Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L, et al. The Rag

GTPases Bind Raptor and Mediate Amino Acid Signaling to mTORC1. Science (80-).

2008;320:1496–501. http://science.sciencemag.org/content/320/5882/1496.abstract.

27. Zoncu R, Efeyan A, Sabatini DM. mTOR: from growth signal integration to cancer,

diabetes and ageing. Nat Rev Mol Cell Biol. 2011;12:21–35. doi:10.1038/nrm3025.

28. Millay DP, O/'Rourke JR, Sutherland LB, Bezprozvannaya S, Shelton JM, Bassel-Duby R,

et al. Myomaker is a membrane activator of myoblast fusion and muscle formation. Nature.

2013;499:301–5. http://dx.doi.org/10.1038/nature12343.

29. Stoll EA, Karapavlovic N, Rosa H, Woodmass M, Rygiel K, White K, et al. Naked mole-

rats maintain healthy skeletal muscle and Complex IV mitochondrial enzyme function into

old age. Aging (Albany NY). 2016. doi:10.18632/aging.101140.

30. Holtze S, Eldarov CM, Vays VB, Vangeli IM, Vysokikh MY, Bakeeva LE, et al. Study of

age-dependent structural and functional changes of mitochondria in skeletal muscles and

heart of naked mole rats (Heterocephalus glaber). Biochem. 2016;81:1429–37.

doi:10.1134/S000629791612004X.

31. Craig T, Smelick C, Tacutu R, Wuttke D, Wood SH, Stanley H, et al. The Digital Ageing

Atlas: integrating the diversity of age-related changes into a unified resource. Nucleic Acids

Res. 2014;43:D873--D878. doi:10.1093/nar/gku843.

32. Cellerino A, Ori A. What have we learned on aging from omics studies? Seminars in Cell

and Developmental Biology. 2017.

33. Bustos V, Partridge L. Good Ol' Fat: Links between Lipid Signaling and Longevity. Trends

in Biochemical Sciences. 2017.

34. Narendra DP, Jin SM, Tanaka A, Suen D-F, Gautier CA, Shen J, et al. PINK1 Is Selectively

Stabilized on Impaired Mitochondria to Activate Parkin. PLOS Biol. 2010;8:e1000298.

http://dx.doi.org/10.1371%2Fjournal.pbio.1000298.

35. Safdar A, deBeer J, Tarnopolsky MA. Dysfunctional Nrf2-Keap1 redox signaling in

skeletal muscle of the sedentary old. Free Radic Biol Med. 2010;49:1487–93.

36. Baumgart M, Priebe S, Groth M, Hartmann N, Menzel U, Pandolfini L, et al. Longitudinal

RNA-Seq Analysis of Vertebrate Aging Identifies Mitochondrial Complex I as a Small-

Molecule-Sensitive Modifier of Lifespan. Cell Syst. 2016;2:122–32.

doi:10.1016/j.cels.2016.01.014.

37. Miwa S, Jow H, Baty K, Johnson A, Czapiewski R, Saretzki G, et al. Low abundance of the

matrix arm of complex I in mitochondria predicts longevity in mice. Nat Commun. 2014;5.

38. Hühne R, Thalheim T, Sühnel J. AgeFactDB - The JenAge Ageing Factor Database -

Towards data integration in ageing research. Nucleic Acids Res. 2014;42:892–6.

39. Dai H, Leeder JS, Cui Y. A modified generalized Fisher method for combining

probabilities from dependent tests. Front Genet. 2014;5:32. doi:10.3389/fgene.2014.00032.

40. Roellig K, Drews B, Goeritz F, Hildebrandt TB. The long gestation of the small naked

mole-rat (Heterocephalus glaber Rüppell, 1842) studied with ultrasound biomicroscopy and

3D-ultrasonography. PLoS One. 2011;6:e17744.

41. Peaker M, Taylor E. Sex ratio and litter size in the guinea-pig. Reproduction. 1996;108:63–

7. doi:10.1530/jrf.0.1080063.

42. Yu C, Li Y, Holmes A, Szafranski K, Faulkes CG, Coen CW, et al. RNA sequencing reveals

differential expression of mitochondrial and oxidation reduction genes in the long-lived

naked mole-rat when compared to mice. PLoS One. 2011;6:e26729.

43. Heinze I, Bens M, Calzia E, Holtze S, Dakhovnik O, Sahm A, et al. Species comparison of

liver proteomes reveals enhanced lipid metabolism, reduced mitochondrial respiration and

enhanced expression of detoxifying enzymes in the long-lived naked mole-rat. bioRxiv. 2017.

doi:https://doi.org/10.1101/220343.

44. Grossman CJ. Interactions between the gonadal steroids and the immune system. Science.

1985.

45. Faulkes CG, Abbott DH, Jarvis JU, Sherriff FE. LH responses of female naked mole-rats,

Heterocephalus glaber, to single and multiple doses of exogenous GnRH. Reproduction.

1990;89:317–23. doi:10.1530/jrf.0.0890317.

46. Gesquiere LR, Learn NH, Simao MCM, Onyango PO, Alberts SC, Altmann J. Life at the

Top: Rank and Stress in Wild Male Baboons. Science (80-). 2011;333:357–60.

doi:10.1126/science.1207120.

47. O'Connor CM, Rodela TM, Mileva VR, Balshine S, Gilmour KM. Corticosteroid receptor

gene expression is related to sex and social behaviour in a social fish. Comp Biochem Physiol

A Mol Integr Physiol. 2013;164:438–46. doi:10.1016/j.cbpa.2012.12.003.

48. Crump CJ, Chevins PF. Prenatal stress reduces fertility of male offspring in mice, without

affecting   their   adult   testosterone   levels.   Horm   Behav.   1989;23:333–43.

http://www.ncbi.nlm.nih.gov/pubmed/2793076. Accessed 21 Mar 2017.

49. Clarke FM, Faulkes CG. Dominance and queen succession in captive colonies of the

eusocial naked mole-rat, Heterocephalus glaber. Proc R Soc London B Biol Sci. 1997;264:993–

1000. doi:10.1098/rspb.1997.0137.

50. Clarke FM, Faulkes CG. Hormonal and behavioural correlates of male dominance and

reproductive status in captive colonies of the naked mole-rat, Heterocephalus glaber. Proc

Biol Sci. 1998;265:1391–9. doi:10.1098/rspb.1998.0447.

51. Maruska KP, Fernald RD. Plasticity of the reproductive axis caused by social status change

in an african cichlid fish: II. Testicular gene expression and spermatogenesis. Endocrinology.

2011;152:291–302.

52. Bens M, Sahm A, Groth M, Jahn N, Morhart M, Holtze S, et al. FRAMA: from RNA-seq

data to annotated mRNA assemblies. BMC Genomics. 2016;17:54. doi:10.1186/s12864-015-

2349-8.

53. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.

2011;29:644–52.

54. Keane M, Craig T, Alfoldi J, Berlin a. M, Johnson J, Seluanov A, et al. The Naked Mole Rat

Genome Resource: facilitating analyses of cancer and longevity-related adaptations.

Bioinformatics. 2014;30:3558–60. doi:10.1093/bioinformatics/btu579.

55. Kent WJ. BLAT - The BLAST-like alignment tool. Genome Res. 2002;12:656–64.

56. Kapustin Y, Souvorov A, Tatusova T, Lipman D. Splign: algorithms for computing spliced

alignments with identification of paralogs. Biol Direct. 2008;3:20.

57. Dobin A, Davis C a., Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: Ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

58. Anders S, Pyl PT, Huber W. HTSeq-A Python framework to work with high-throughput

sequencing data. Bioinformatics. 2015;31:166–9.

59. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9:357–9. doi:10.1038/nmeth.1923.

60. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15:550. doi:10.1186/s13059-014-0550-8.

61. Kasen S, Ouellette R, Cohen P. Mainstreaming and postsecondary educational and

employment status of a rubella cohort. Am Ann Deaf. 1990;135:22–6. doi:10.2307/2346101.

62. Zeeberg BR, Feng W, Wang G, Wang MD, Fojo AT, Sunshine M, et al. GoMiner: a resource

for biological interpretation of genomic and proteomic data. Genome Biol. 2003;4:R28.

doi:10.1186/gb-2003-4-4-r28.

63. Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO summarizes and visualizes long lists of

gene ontology terms. PLoS One. 2011;6:e21800. doi:10.1371/journal.pone.0021800.

Supplementary Table Legends

Table S1. Age at death for NMRs and GPs.

Table S2. Number of analyzed biological replicates per group.

Table S3. Reason for exclusion of samples from further analysis.

Table S4. Number of uniquely aligned RNA-seq reads.

Table S5. Mean pairwise Pearson correlation coefficients between biological replicates in (A) naked mole-rat and (B) guinea pig.

Table S6. Number of differentially expressed genes (FDR < 0.01) between NMR and GP (A) without logFC threshold and (B) with logFC threshold ($|logFC| > 2$).

Table S7. DEGs in cross-species comparison that show an overlap with DAA.

Table S8. Gene set enrichment analysis for cross-species DEGs (FDR < 0.01, $|logFC| > 2$) that are age-related (DAA).

Table S9. Number of sex-related (female vs. male) differentially expressed genes in (A) non-breeder and (B) breeder (FDR < 0.01).

 Table S10. Functional enrichment analysis of 790 DEGs shared between GP-B-FvM and GP-N-FvM.

Table S11. Gene description of sex-related DEGs in NMR-N-FvM.

 Table S12. Number of status-related (breeder vs. non-breeder) differentially expressed genes in (A) females and (B) males (FDR < 0.01).

Table S13. Functional gene set enrichment analysis of DEGs in Ovr of NMR-F-BvsN.

Table S14. Functional gene set enrichment analysis of DEGs in Adr of NMR-F-BvsN.

Table S15. Functional gene set enrichment analysis of DEGs in Cer of NMR-F-BvsN.

Table S16. DEGs in intersection between NMR-F-BvsN and NMR-M-BvsN.

Table S17. Functional gene set enrichment analysis of DEGs in Tes of NMR-M-BvsN.

Table S18. Functional enrichment analysis in cellular components of DEGs in Tes

NMR-M-BvsN.

Table S19. Functional gene set enrichment analysis of DEGs in Skn of NMR-M-BvsN.

Table S20. Proportion of mitochondrial transcriptonal output.

Table S21. Differentially expressed nuclear genes (FDR<0.05) associated with mitochondrial

respiratory chain complexes in Tes and Skn of NMR-M-BvsN.

Table S22. Antioxidant enzymes differentially expressed in Tes and Skn of

NMR-M-BvsN (FDR<0.05).

Table S23. Enrichment of age-related genes (Digital Ageing Atlas) in status-related DEGs.

Only tissues having at least 50 DEGs were tested for enrichment.

Table S24. Shared status-related genes between NMR and GP.

Table S25. FDR values (Fisher's exact test) for overlap between Digital Ageing Atlas and

upper quintile of status-related expression differences having the greatest interspecies

difference. Three groups were investigated: (i) opposing and (ii) same direction of expression

in both species as well as (iii) independent of direction. Further, in each group direction in

each species was tested separately.

Table S26. Functional enrichment analysis of the non-redundant set of aging-related

20%-quantiles that show the greatest interspecies difference in males (Skn, Tes).

Table S27. Functional enrichment analysis of the non-redundant set of aging-related

20%-quantiles that show the greatest interspecies difference in females (Hrt, Pit, Ovr).

Supplementary Figure Legends

Figure S1: Collected tissues exemplified in schematic figure of NMR.

Figure S2: Hierarchical clustering of gene expression profiles. The clustering tree shows a

clear separation of tissues in both species, but less pronounced differences between sex and

breeding status.

Figure S3: Top 15 highest ranked GO sets based on enrichment analysis of cross-species

DEGs between NMR and GP. GO sets are ranked by number of summarized GO terms (x-

axes) and number of DEGs (alongside bar).

Figure S4: Top 15 highest ranked GO sets based on enrichment analysis of sex-related DEGs

that are shared between GP breeder and non-breeder. GO sets are ranked by number of

summarized GO terms (x-axes) and number of DEGs (alongside bar).

Figure S5: Top 15 highest ranked GO sets based on enrichment analysis of sex-related DEGs

in NMR breeder. GO sets are ranked by number of summarized GO terms (x-axes) and

number of DEGs (alongside bar, together with number of up- and downregulated genes).

Figure S6: Top 15 highest ranked GO sets based on enrichment analysis of status-related

DEGs in ovary of NMR females. GO sets are ranked by number of summarized GO terms

(x-axes) and number of DEGs (alongside bar, together with number of up- and

downregulated genes).

Figure S7: Top 15 highest ranked GO sets based on enrichment analysis of status-related

DEGs in adrenal gland of NMR females. GO sets are ranked by number of summarized GO

terms (x-axes) and number of DEGs (alongside bar, together with number of up- and

downregulated genes).

Figure S8: Top 15 highest ranked GO sets based on enrichment analysis of status-related

DEGs in cerebellum of NMR females. GO sets are ranked by number of summarized GO

terms (x-axes) and number of DEGs (alongside bar, together with number of up- and

downregulated genes).

Figure S9: Top 15 highest ranked GO sets based on enrichment analysis of status-related

DEGs in testis of NMR males. GO sets are ranked by number of summarized GO terms (x-

axes) and number of DEGs (alongside bar, together with number of up- and downregulated

genes).

Figure S10: Top 15 highest ranked GO sets based on enrichment analysis of status-related

DEGs in skin of NMR males. GO sets are ranked by number of summarized GO terms (x-

axes) and number of DEGs (alongside bar, together with number of up- and downregulated

genes).

Figure S11: Mitonuclear ratios in non-breeders and breeders per sex, tissue and species.

Boxplots shows median, 2nd/3rd quartiles, whiskers extend to 1.5 the interquartile range and

dots values outside this range. P-values were calculated using a two-tailed t-test; *: $p<0.05$,

**: $p<0.01$.

Figure S12: Top 15 highest ranked GO sets based on enrichment analysis of the non-

redundant set of aging-related 20%-quantiles that show the greatest interspecies difference in

males (Tes, Skn). GO sets are ranked by number of summarized GO terms (x-axes) and

number of DEGs (alongside bar).

Figure S13: Top 15 highest ranked GO sets based on enrichment analysis of the non-

redundant set of aging-related 20%-quantiles that show the greatest interspecies difference in

females (Hrt, Pit, Ovr). GO sets are ranked by number of summarized GO terms (x-axes) and

number of DEGs (alongside bar).

Supplemental Text S1

In NMR-M-BvsN Tes, this is indicated by a significant upregulation of receptors for

androgens (AR, 1.3-fold) and gonadotrophic hormones (LHCGR, 1.8-fold) as well as an

increase in steroid production. Specifically, we observe an upregulation of genes involved in

the transport of cholesterol into mitochondria (STAR, steroidogenic acute regulatory protein,

2.8-fold; TSPO, translocator protein, 1.5-fold; CAV1, caveolin 1, 1.4-fold), the rate limiting step

of steroidogenesis (1), and conversion of cholesterol into pregnenolone (CYP11A1,

cytochrome P450 family 11 subfamily A member 1, 1.8-fold), which is essential for all

steroids. Genes encoding for enzymes that produce testosterone (HSD17B1, hydroxysteroid

17-beta dehydrogenase 1, 1.4-fold), the more potent metabolite dihydrotestosterone (SRD5A1,

steroid 5 alpha-reductase 1, 1.3-fold) and cortisol (CYP51A1, cytochrome P450 family 51

subfamily A member 1, 2-fold) are upregulated as well (Supplement Figure 10). The

upregulation of steroidogenesis is accompanied by a significant increase in the mitonuclear

ratio and enrichment of nuclear DEGs associated with mitochondria, including OXPHOS.

Consequently, the load of reactive oxygen species (ROS) is presumably higher in NMR Tes,

but is compensated by increase in antioxidant enzymes. NMR-F-BvsN Ova shows more

subtle changes in steroidogenesis by increased estradiol production as indicated by

significant upregulation of CYP19A1 (aromatase, 2.5-fold) and increased sensitivity to

androgens and progesterogens as indicated by upregulation of corresponding receptors (Ova,

AR, 1,4-fold; Pit, PGR, 1.6-fold). We find, however, substantial expression changes in two

genes involved in the regulation of follicle-stimulating hormone secretion (2), INHA (inhibin

alpha subunit, 19.2-fold) and INHBA (inhibin beta A subunit, 4.4-fold), indicating that ovaries

of non-breeding NMRs lack preovulatory follicles. Further, enrichment of steroid metabolic

process, response to hormones and prostate gland growth indicates active feedback signalling

also in females and an impact on developmental processes. Increase in signalling is further

supported by significant enrichment of gene products located in secretory vesicle (secretory

granule lumen, GO:0034774) and on cell surface (GO:0009986).

References

1.   Stocco DM, Clark BJ (1996) Regulation of the acute production of steroids in

steroidogenic cells. Endocr Rev 17(3):221–244.

2.   Wijayarathna R, de Kretser DM (2016) Activins in reproductive biology and beyond.

Hum Reprod Update 22(3):dmv058.

Supplemental Text S2

Although well expressed, we found no significant differences for gonadotropin-related genes

(GNRH1, FSHB, CGA, LHB). This indicates a similar intracellular transcript turnover in non-

breeders and breeders, and is consistent with previous results indicating that LH is stored in

non-breeders, ready to be released upon GnRH signalling (1). Notably, GnIH (NPVF), a

negative regulator of reproductive function suggested to have a role in reproductive

immaturity in NMRs due to its decreased protein expression in brain of NMR breeder (2),

shows tendentially decreased mRNA expression in NMR breeders (males: 0.37, females

0.13). Also, previously elevated diencephalon mRNA levels in NMR were reported for AR

(androgen receptor) in male breeders and for CYP19A1 (aromatase), ESR1 (estrogen receptor

1) and PGR (progesterone receptor) in female breeders (3). Our corresponding Hyp data do

not support those increases (log2FC<0.1). Nevertheless, we identified significantly elevated

transcript levels of these genes both in male (Tes: AR; Supplemental Data S67) and female

breeders (Pit: PGR, Supplemental Data S54; Ova: AR, CYP19A1, ESR1, PGR, Supplemental

Data S57) supporting a complex status-related function of these genes. Furthermore, steroid

feedback and GnRH secretion are integrated by brain GABA and glutamate signalling in

mammals and cichlids (4). Although, we do not observe equivalent significant differences in

the NMR brain, the data show an upregulation of genes coding for three GABA receptor

subunits (GABRB3, GABRG1, GABRP) and one glutamate receptor subunit (GRIK2) in Tes

(Supplemental Data S67), as well as two of three differentially expressed GABA (GABRB2,

GABRG3) and three glutamate receptor subunits (GRIK1, GRIK2, GRIK4) in Ova and female

Pit of NMR breeders (Supplemental Data S54, S57). Together, this suggests increased

neuronal plasticity and/or activity predominantly in gonads of NMRs after becoming

breeders.

In the testis of immature chicken, expression of ADIPOR1/2 (adiponectin receptor 1/2) is

significantly lower than in mature animals (5). It has been hypothesised that these genes are

involved in supporting the higher metabolic activity related to spermatogenesis, testicular

steroid hormone production, and transport of spermatozoa and testicular fluid. In line with

these observations, our results show a significant upregulation of ADIPOR2 in Tes of NMR

breeders (1.7-fold; Supplemental Data S67).

The renin-angiotensin system predominantly involved in cardiovascular control has also been

associated with reproduction in mice and human (6). Signaling through receptors coded by

AGTR1 (angiotensin II receptor type 1) in males, and AGTR1/2 in females has been associated

with fertility and stimulation of reproduction. In accordance, we observed a significant

upregulation of AGTR1 and AGTR2 in gonads of NMR male and female breeders,

respectively (Supplemental Data S67,S57).

References

1. Faulkes CG, Abbott DH, Jarvis JU, Sherriff FE (1990) LH responses of female naked

mole-rats, Heterocephalus glaber, to single and multiple doses of exogenous GnRH.

Reproduction 89(1):317–323.

2. Peragine DE, et al. (2017) RFamide-related peptide-3 (RFRP-3) suppresses sexual

maturation in a eusocial mammal. Proc Natl Acad Sci 3:201616913.

3. Swift-Gallant A, Mo K, Peragine DE, Monks DA, Holmes MM (2015) Removal of

reproductive suppression reveals latent sex differences in brain steroid hormone receptors in

naked mole-rats, Heterocephalus glaber. Biol Sex Differ 6(1):31.

4. Renn SCP, Aubin-Horth N, Hofmann H a (2008) Fish and chips: functional genomics

of social plasticity in an African cichlid fish. J Exp Biol 211(Pt 18):3041–3056.

5. Ocón-Grove OM, Krzysik-Walker SM, Maddineni SR, Hendricks GL, Ramachandran

R (2008) Adiponectin and its receptors are expressed in the chicken testis: influence of sexual

maturation on testicular ADIPOR1 and ADIPOR2 mRNA abundance. Reproduction

136(5):627–38.

6. Pan PP, Zhan QT, Le F, Zheng YM, Jin F (2013) Angiotensin-converting enzymes play

a dominant role in fertility. Int J Mol Sci 14(10):21071–21086.