# A simple rule for the evolution of fast dispersal at the edge of expanding populations

Maxime Deforet[1], Carlos Carmona-Fontaine[2], Kirill S. Korolev[3], Joao B. Xavier[1]

[1]Program in Computational and Systems Biology, Memorial Sloan-Kettering Cancer Center, New York, New York

[2]Center for Genomics and Systems Biology, New York University, New York

[3]Department of Physics and Graduate Program in Bioinformatics, Boston University, Boston

Corresponding: deforetm@mskcc.org, xavierj@mskcc.org

**Abstract**

Evolution by natural selection is commonly perceived as a process that favors those that replicate faster to leave more offspring; nature, however, seem to abound with examples where organisms forgo some replicative potential to disperse faster. When does selection favor invasion of the fastest? Motivated by evolution experiments with swarming bacteria we searched for a simple rule. In experiments, a fast hyperswarmer mutant that pays a reproductive cost to make many copies of its flagellum invades a population of mono-flagellated bacteria by reaching the expanding population edge; a two-species mathematical model explains that invasion of the edge occurs only if the invasive species' expansion rate, $v_2$, which results from the combination of the species growth rate and its dispersal speed (but not its carrying capacity), exceeds the established species', $v_1$. The simple rule that we derive, $v_2 > v_1$, appears to be general: less favorable initial conditions, such as smaller initial sizes and longer distances to the population edge, delay but do not entirely prevent invasion. Despite intricacies of the swarming system, experimental tests agree well with model predictions suggesting that the general theory should apply to other expanding populations with trade-offs between growth and dispersal, including non-native invasive species and cancer metastases.

## Introduction

Natural selection is commonly perceived as a process that favors species that grow in numbers by reproducing faster and surviving better. In expanding populations, however, species face a choice between allocating resources to growth or to disperse faster [1]. Expanding populations consume local resources; the resource availability is highest outside the population range, which creates an advantage to being at the population margin [2]. Fast-dispersing species take advantage of this spatial heterogeneity: they take over the edge, cutting-off competitors' access to growth-limiting resources [1, 3]. The role of resource limitation in evolutionary processes, such as gene surfing [4] and the founder effect [5], has been analyzed before; however, crucial questions remain: When does natural selection favor dispersal over growth? How much growth can a fast-dispersing species forgo and still be favored by natural selection?

Invasiveness in expanding populations has global implications at many levels [6]: non-native species invade due to human introduction and climate change [7], expanding tumors become invasive cancers [8] and antibiotic-resistant bacteria spread due to global travel [9]. Predicting the evolution of expanding populations, however, is a non-trivial problem. Dispersal and growth are sometimes under an allocation trade-off where investing in one trait leaves fewer resources available for the other [10-13]. Mathematical modeling could help address these complex questions by defining the conditions favoring fast dispersal.

Testing model predictions with field studies in traditional ecology is challenging and experimental manipulation of natural ecosystems is often impractical. The shortage of robust empirical systems to test mathematical models stalls the development of predictive theory. Laboratory experiments with microbes can provide rigorous tests of mathematical models [5, 14-17]; microbes have large populations sizes, short generation times, affordable DNA sequencing and—in many cases—tools for genetic engineering. We recently discovered that experimental evolution in swarming colonies of the bacterium *Pseudomonas aeruginosa* leads to the spontaneous evolution of hyperswarmers [18]. Hyperswarmer mutants have a single point mutation in a gene called *fleN;* their cells are multiflagellated, and therefore more dispersive. Importantly, the many flagella come at the cost of a slower growth (table S1). *P. aeruginosa* wild-type outcompete hyperswarmers in well-mixed liquid media; hyperswarmers, on the other hand, swarm faster on semi-solid surfaces [19] and outcompete the wild-type in swarming competitions [18].

Here we exploited the differences in growth rate and dispersal between the wild-type *P. aeruginosa* and its hyperswarmer mutant to motivate, and then experimentally test, a mathematical model of invasion in expanding populations. We based our analysis on a well-established principle of spatial expansion in growing populations: the traveling wave derived from the Fisher-Kolmogorov-Petrovsky-Piscunov (F-KPP) equation. The F-KPP equation, in its original form, describes a 1-D monospecies population [20-22]. We expanded the F-KPP equation to investigate the conditions favoring invasion of a species with different dispersal and growth rates and solving that two-species system produced a simple rule to determine the outcome of invasion. Somewhat surprisingly, this simple rule had not been proposed before to

68 the best of our knowledge, despite much theoretical and experimental work in this field.
69 Simulations reveal the conditions at which the rule is applicable, and the time-scales necessary
70 for invasion in biologically relevant situations. Quantitative experiments with *P. aeruginosa*
71 hyperswarmers indicated that our model may predict the evolution of expanding populations in a
72 wide range of biological scenarios.
73
**Results**
*Modeling swarming in* P. aeruginosa *with the F-KPP equation*
*P. aeruginosa* populations swarm across agar gels containing nutrients and form
branched colonies. Bacteria at the branch tips travel at a nearly constant rate by dividing and
dispersing (19). Each growing tip consumes resources in its vicinity and thus forms a nutrient
gradient (16) that drives a resource-limited growth similar to the F-KPP model. Knowing that
cell sizes have a positive correlation with growth (23) we compared the size of cells collected
from the tip of a branch with the size of cells collected behind the tip; cells at the tip were longer
indicating faster growth at the edge of the population (Fig. S1). We then modeled *P. aeruginosa*
swarming as a one-dimensional expanding population according to the F-KPP equation:

84
$$\frac{\partial u}{\partial t} = ru\left(1 - \frac{u}{K}\right) + D\frac{\partial^2 u}{\partial x^2} \tag{1}$$

where $x$ is space, $t$ is time, $u$ is the local population density, $K$ is the carrying capacity, $r$ is the
maximum per-capita growth rate and $D$ quantifies dispersal. Growth and dispersal can obey
different laws in nature; for generality the F-KPP equation assumes logistic growth, where the
per-capita growth rate decreases linearly as the population density increases, leading to a
decrease in resource availability, and assumes Fickian diffusion for dispersal. The F-KPP
equation suits the common scenario where regions with excess of nutrients lie outside the
population and determine the direction of expansion. The resource availability, represented by
$1 - \frac{u}{K}$, is highest outside the population range and growth, represented by $ru(1 - \frac{u}{K})$, occurs
mostly at the edge of the population. Eq. (1) has a traveling wave solution, $u(x, t) = u_0(x - vt)$,
where the population front travels at a constant expansion rate $v = 2\sqrt{rD}$ and its density
increases from the edge with a length-scale $\lambda = \sqrt{D/r}$ (Movie S1 and Fig. S1) (4, 24). In
summary, the F-KPP equation—where low-density at the leading edge drives population
expansion—has a traveling wave solution where the expansion rate depends on both growth and
dispersal, but is independent of the carrying capacity.
99
*A simple rule for the evolution of faster dispersal*
Hyperswarmers grow slower in well-mixed liquid media due the cost of synthesizing
multiple flagella, but—thanks to their faster dispersal on semi-solid surfaces—outcompete the
wild-type in spatially structured environments (18, 19). In semi-solid surfaces lacking spatial
structure hyperswarmers are outcompeted, as expected (Fig. S2). At the micrometer scale, an
expanding population of hyperswarmers displays patterns of active turbulence typical of dense
bacterial suspensions, which is different from the wild-type where cells remain nearly static even

at the tips of swarming tendrils (Movie S2). To gain a better understanding of the competition
dynamics in expanding swarming colonies we mixed wild-type bacteria (labeled with a red
fluorescent protein) with hyperswarmers (labeled with a green fluorescent protein) at 10:1 ratio;
we then used time-lapsed florescence imaging to film the swarming competition (Fig. 1A). The
time-lapse showed that hyperswarmers quickly reached the population edge, increasing their
dominance as the colony expanded to win the competition (Movie S3).
113

**Figure 1: Invasion dynamics in an expanding population of swarming *P. aeruginosa* bacteria.** A: Fluorescence
time-lapse imaging shows a swarming competition in a mixed population of wild-type and hyperswarmers at 10:1
ratio and on an soft-agar gel. Leftmost panels: Monoclonal swarming colonies of wild-type (top) and hyperswarmer
(bottom) imaged at $T = 12$h (scale bars 1 cm). B: Snapshots of numerical simulations of Eq. 2 modeling
competition of *P. aeruginosa* (red lines) and the hyperswarmer mutant (green lines) with initial ratio 10:1, using
parameters extracted from experiments (Table S1). Left panel shows time points represented in panel A. Right panel
shows later time points, where the wild-type population stalls while the hyperswarmer population keeps expanding.
C: Ratio of hyperswarmer biomass over wild-type biomass. The experimental data is extracted from fluorescence
signals in panel A (black squares represent time points shown in A).

To determine the conditions favoring evolution of faster dispersal we used an extension
of the F-KPP equation for a two-species system. Species 1—representing the wild-type—has
density function $u_1(x,t)$, disperses with coefficient $D_1$ and grows with a rate $r_1$; species 2—
representing the hyperswarmer—has $u_2(x,t)$, $D_2$, and $r_2$. For simplicity, and according to data
for the hyperswarmer system (18) (Fig. S3), we assumed that both species have the same
carrying capacity, which we normalized to 1. We assumed that their dispersal rates, determined
by $D_1$ and $D_2$, are independent.
In this model, the only traits that may be improved in species 2 are its dispersal ($D_2$) and
growth ($r_2$). The two-species system has two coupled F-KPP equations (25-28):

$$\frac{\partial u_1}{\partial t} = r_1 u_1 (1 - u_1 - u_2) + D_1 \frac{\partial^2 u_1}{\partial x^2}$$

$$\text{(2)}$$

$$\frac{\partial u_2}{\partial t} = r_2 u_2 (1 - u_1 - u_2) + D_2 \frac{\partial^2 u_2}{\partial x^2}$$

Species 1 and 2 interact only by competing for the same resources, a feature implemented by the factor $1 - u_1 - u_2$. This framework is well established and it was used previously to investigate competition in various contexts of range expansion (17, 25-27, 29-36), including a linear trade-off between dispersal and growth (28). However, previous studies did not continue to derive a general rule to determine the invasion outcome for all possible values of dispersal and growth.

To derive that rule, we first investigated the conditions that allow an introduced population to invade and replace the resident population at the expansion front. We could determine analytically that invasion at the edge occurs only if the expansion rate of species 2 exceeds that of species 1 (see mathematical demonstration in *SI Appendix 2: Analytical solution for the condition of invasion*):

$$v_2 > v_1 \tag{3}$$

where $v_1 = 2\sqrt{r_1 D_1}$ and $v_2 = 2\sqrt{r_2 D_2}$ are the expansion rates of each species when grown alone.

Eq. 3 is a simple rule for invasion of the expansion edge. The intuition behind invasion dynamics is well illustrated in a simulation of the competition between an established species (species 1) and an invader with faster dispersal but slower growth (species 2), which we simulated (Fig. 1B) by numerically solving the system in Eq. 2 with parameters corresponding to the hyperswarmer system ($r_2/r_1 = 0.9$ and $D_2/D_1 = 2$, Table S1). Species 2, initially homogeneously mixed with species 1, outcompetes species 1 once it reaches the leading edge. The invasion succeeds despite the growth disadvantage of species 2 because its faster dispersal enables it to reach the low-density edge where it can take advantage of the resources available. Once species 2 dominates the edge, species 1 is left behind in the high-density region where growth has stopped. Over time, the global frequency of species 1—blocked by species 2 from reaching the edge and incapable of growing further—decreases in frequency whereas species 2 keeps increasing thanks to its edge domination (Fig. 1C, left). These simulation results are consistent with experimental tests conducted here (Fig. 1C, right) and also with the original experiment that led to evolution of hyperswarmers (18), which clearly showed that *fleN* mutants would outcompete the wild-type to extinction given sufficient competition time on swarming plates.

**Figure 2. Faster-dispersing species invade expanding populations despite growing slower as long as $v_2 > v_1$.**
A: Phase diagram of invasion in $r_2/r_1$, $D_2/D_1$ space, with 2 subdomains of interest in the invasive domain: (1) is the domain of invasion with lower growth rate and faster dispersal. (2) is the domain of invasion with faster dispersal but lower growth rate. The (+) symbol represents the hyperswarmer phenotype with respect to wild-type phenotype. The (*) symbol represents the wild-type phenotype with respect to the hyperswarmer phenotype. B: Fixation probability obtained from the stochastic model with death rate (stochastic simulations performed with $S = 1$, $K = 100$, $L = 2\lambda_1$). C: Invasion time obtained from the deterministic model (simulations performed with $L = 2\lambda_1$ and $S = 0.2$). The empty orange circles represent absence of invasion within the duration of the simulation ($r_1t = 180$). In A-C, the red dot depicts the reference point ($D_2 = D_1$ and $r_2 = r_1$).

The invasion rule obtained from the two species system (Eq. 3) states that the evolution of an expanding population is entirely determined from the growth and dispersal rates. Importantly, and similar to the expansion rate obtained for a monospecies traveling wave, the invasion rule is independent of the carrying capacity of each species (See Fig. S4 for confirmation with numerical simulations).

The invasion rule leads to a diagram that delineates the invasion outcome in a growth-dispersal space (Fig. 2A). This diagram shows two trivial domains: when both growth and dispersal of species 2 are lower ($D_2 < D_1$ and $r_2 < r_1$), invasion does not occur because $v_2$ is always lower than $v_1$. Numerical simulations explain that for very low values of $D_2$ and $r_2$ species 1 continues to expand and travel at constant expansion rate whereas species 2 spreads out and stalls (Movie S4, bottom left panel, and Fig. S5A). When growth and dispersal of species 2 are greater ($D_2 > D_1$ and $r_2 > r_1$), species 2 invades because $v_2$ is always higher than $v_1$: species 2 grows rapidly, moves to the front where it reaches the active layer and outcompetes species 1 (Movie S4, top right panel, and Fig. S5B).

The two domains where one trait is higher and the other is lower are less trivial, but arguably more relevant because of the trade-off between dispersal and growth commonly found in nature (6). Domain 1 of Fig. 2A shows that invasion occurs if species 2 disperses slower than species 1 as long as its growth rate is sufficiently higher ($r_2 > r_1 D_1/D_2$, Movie S4, top left panel, and Fig. S5C). Domain 2 shows that species 2 invades even if—like the hyperswarmer—it grows slower than species 1 as long as it disperses fast enough such that $D_2 > D_1 r_1/r_2$. A fast-

dispersing invader reaches the active layer where it can gain better access to growth-limiting
resources and outcompete the established species (Movie S4, bottom left panel, and Fig. S5D).
The simple invasion rule, therefore, makes quantitative predictions of invasion outcome
that hold true in our experimental system despite intricacies such as the large-scale branching
and small-scale turbulence. Hyperswarmers have a ~100% increase in dispersal ($D_2/D_1 \sim 2$) that
comes at a ~10% growth rate cost ($r_2/r_1 \sim 0.9$) (Movie S2 and Table S1). Therefore the
experimental system falls into domain 2 of the invasion diagram (cross symbol in Fig. 2A).
We measured the frequency of hyperswarmers within the first millimeter of the colony
from video frames and saw that it increased exponentially with a rate $\tilde{r} = 0.39 \pm 0.08\,\text{h}^{-1}$,
which is in quantitative agreement with the mathematical model (Fig. S6, see Eq. S6 of *SI*
*Appendix 2: Analytical solution for the condition of invasion*). Hyperswarmers introduced into an
expanding wild-type colony spread within a wild-type branch (Fig. S7), reach the tip of the
branch, and take over the population (Movie S5A) resembling our simulations (Fig. S5D).
Hyperswarmers evolved from a wild-type swarming colony (18) and could take over the
ancestral population thanks to a greater dispersal. To test the invasion rule, we asked whether
this process was reversible: Could wild-type cells invade the edge of an expanding a
hyperswarmer colony thanks to their greater growth rate? Our model predicted that wild-type
cells would be unable to invade the hyperswarmer population edge since in this case $v_2 < v_1$
($D_2/D_1 = 0.5$ and $r_2/r_1 = 1.1$, see star symbol in Fig. 2A). This was confirmed experimentally:
wild-type cells introduced in a hyperswarmer colony simply spread out and were rapidly
outpaced at the edge by the hyperswarmers (Movie S5B).

*Invasion rule valid despite phenotypic variability*
Even in mono-species systems, individuals with identical and defined genotypes can still
display phenotypic variation (19, 37). To study whether such variation had an effect on invasion
outcome, we introduced non-heritable fluctuations in birth and death events as well as in the
dispersal processes. These phenotypic variations were modeled as stochastic distributions around
the mean population value, which is determined by the strain's genotype (see *SI Appendix 4:*
*Stochastic Modeling*).
Our simulation results show that the invasion rule, $v_2 > v_1$, despite having been derived
from deterministic assumptions, holds valid even in stochastic situations. The transition at
$v_2 = v_1$ was, however, more gradual (Fig. 2B and S8): the zone of transition broadened as
stochasticity increased because, as expected from other stochastic studies (38, 39), stochasticity
allowed for a non-zero probability of deleterious mutants ($v_2 < v_1$) to invade and beneficial
mutants ($v_2 > v_1$) had a non-zero probability of failing to invade. Larger carrying capacities
lessened the stochastic effects and sharpened the transition zone, again as expected from
previous stochastic analyses (38, 39); importantly, however, the rule $v_2 > v_1$ could still predict
invasion of the population edge even with different carrying capacities (Fig. S9).
We confirmed the generality of the invasion rule further by carrying out evolutionary
simulations where mutations randomly arise at division. We considered two schemes: i)

mutations that change growth and/or dispersal relative to the ancestor phenotype but do so in an
uncorrelated way; ii) mutations that change growth and dispersal considering that the two traits
are linearly correlated (linear trade-off) but independent of the ancestor phenotype. In the case of
uncorrelated mutations, populations evolved—on average—towards a greater expansion rate
$v = 2\sqrt{rD}$ (Fig. S10C). In contrast, when the two phenotypes were constrained by a trade-off,
evolution converged to the value along the trade-off line which maximized the expansion rate
$v = 2\sqrt{rD}$ (Fig. S10D-G). In summary, we conducted several types of stochastic simulations
that all showed that invasion of the population edge obeys the simple rule $v_2 > v_1$.
241

242     *The role of spatial structure and founder effect*
243           We then investigated whether our model would account for other factors that can affect
invasion and competition in biologically relevant scenarios. For example, in most evolutionary
scenarios where competition starts with a mutation the size of the invader population is initially
very low (1 individual) whereas a competing species introduced by external processes (e.g.
human intervention) can start at higher densities. Also, the initial location of the invader species
matters because resources are not evenly distributed in nature, and an invading species may
invade faster if it is introduced in the resource-rich leading edge than if it is introduced in
deprived regions where it will take longer to grow to domination. In summary, species 2 should
take longer to invade (i) when it is introduced further from the edge where resources are already
limited or (ii) when its initial size is small. The invasion rule determines whether species 2 can
invade (Eq. 3, Fig. 2A) but does not give us the time necessary for invasion.
254           To investigate how the time to invasion depends on the location and initial size of the
invading population we modeled the introduction of species 2 into a traveling wave formed by
species 1. We assumed an initial density $S$ across a small interval at a distance $L$ from the edge
for species 2 *(SI Appendix 3: Time of invasion)*, and we determined the time needed to
outnumber species 1 at the front. Numerical simulations revealed that the general rule, $v_2 > v_1$
holds for all initial conditions given sufficient time (Fig. 2C). The time required, however,
depends on the initial conditions, increasing approximately linearly with the distance $L$ from the
front and decreasing sub-linearly with the initial density $S$ (Fig. S11 and S12).
262           To better distinguish the factors that influence the time-scale of invasion we considered
two steps: first, we considered that species 2 disperses until it reaches the active layer. The time
for species 2 to reach the edge depends on the distance from the introduction point to the front—
a distance that increases constantly because species 1 is itself advancing—and also on the initial
width of species 2. Second, once species 2 reaches the active layer it must grow to outnumber
species 1. When the introduction is sufficiently far from the edge the time of invasion, $t_i$, is:
$$t_i \sim \alpha L - \beta \log(S) \qquad\qquad (4)$$
where $\alpha$ and $\beta$ depend on the parameters $D_1, D_2, r_1$ and $r_2$ (see *SI Appendix 3: Time of invasion*).
This analysis confirms simulation results that the time to invasion depends linearly on $L$ but only
sub-linearly on $S$ (Fig. S11 and S12), highlighting that the distance to the edge is key to invasion.
272           We then tested these findings in our experimental system. We manipulated the distance to

the edge ($L$) and the density ($S$) of a small population of hyperswarmers introduced into an expanding wild-type population, and we compared the experimental results to the corresponding simulations. In simulations, the invasion outcome was calculated as the frequency $f$ of species 2 at the edge of the population 6 hours after implantation (Fig. 3A). In the experiments, we ranked the invasion outcome after 6 h of expansion as no invasion, scarce invasion, abundant invasion and complete invasion according to the amount of hyperswarmers visible at the edge (Fig. 3B, see details in *SI Appendix 1: Materials and Methods*). The experiments confirmed the dominant role of $L$ compared to $S$ in determining the time of invasion (Eq. 4), which is evident from the concave shape of the invasion scores (Fig. 3C, see *SI Appendix 5: Statistical analysis*).

The intuition behind the concave shape is that when the initial distance from the edge is too long then the invasive species may not be able to invade within biologically relevant time, even if its initial size is large. The shape of the iso-frequency contour lines can be calculated from the simplified two-step model of invasion described above and is given by

$$S \sim e^{L/L_0} \tag{5}$$

where $L_0 = 4D_2 v_1 / (3v_1^2 + v_2^2)$ is the characteristic length of theses lines. In the case of *P. aeruginosa* and its hyperswarmers $L_0 = 1.1 \pm 0.07$ mm (SD). The results from our hyperswarmer experiments agree with the theoretical model (Fig. 3A and C; compare to lines of constant mutant frequency), indicating that the two-species F-KPP model, in spite of its simplifying assumptions and despite any intricacies of the experimental system (e.g. the swarming population is tri-dimensional; bacterial cells tend to lose motility as they lose access to resources inside the population, which freezes the spatial organization), is sufficiently general to describe invasion in swarming colonies.

**Discussion**

313
Since the modern synthesis unified principles of natural selection and genetics in the
same mathematical framework, simple rules have provided important solutions to evolutionary
puzzles (40). Many cornerstone concepts of evolutionary theory can be expressed as such rules,
and their logic can be followed from simplifying assumptions. Real systems are often more
complicated, but simple rules still provide useful "rules of thumb" to guide intuition. For
instance, Hamilton's rule of kin selection (41), which among other feats determines conditions
for the evolution of altruism, drives despite its simplifying assumptions an important body of
work both theoretical and experimental in social evolution (42). The F-KPP is one such simple
equation to describe the contributions of growth and dispersal to the spatial expansion of a
mono-species population (20, 21). The F-KPP also defines conditions for a traveling wave which
resembles expanding populations observed in nature, and allows calculating the traveling wave
speed from the growth and dispersal rates.

Here we showed that extending the F-KPP equation to a multi-species situations produces
a simple mathematical rule to predict the outcome of invasion in an expanding edge depending
on the growth and dispersal rates of the competing species. The problem of invasion in an
expanding population had been investigated, both theoretically (e.g. 43, 44) and empirically (e.g.
7). But its simple solution, the inequality $v_2 > v_1$, had not been—to the best of our knowledge—
presented this way before, a fact that somewhat surprises us.

This simple rule derived analytically from the deterministic two-species F-KPP equation
establishes the mutual contribution of growth and dispersal for the invasion outcome.
Importantly, the outcome can be independent of the system's carrying capacity, particularly
because the competition dynamics relies especially on the low-density of the population at the
expanding edge. Further away from the edge, where population is denser, spatial structure and
dispersal are less relevant; the evolutionary fate of an invader might then be determined by high-
density dynamics where additional factors such as kin competition become important (36, 45).
The combined effect of growth and dispersal was also tested experimentally using a laboratory
system of swarming bacteria. The rule determined that a slower growing species could invade an
established species as long as its dispersal rate is sufficiently faster and that was indeed the case
in our experimental system. Hyperswarmer mutants paid a growth cost for synthesizing their
multiple flagella but—without affecting their competitive ability—dispersed faster than wild-
type bacteria (domain 2 of Fig. 2A).

Our experimental tests with hyperswarmers showed, then, that the invasion rule $v_2 > v_1$
applies to an empirical system despite intricacies that could violate the simplifying assumptions
of the F-KPP. This is important, because nature abounds with examples like our experiments
where invasive species speed up dispersal but at a cost to their growth rate: the invasion of the
cane toads in Australia, a human-introduced species, is led by faster long-legged individuals with
lower birth rates (46); the South African mountain fynbos is threatened by invasive pine trees
with lighter pine seeds that disperse better (47) but produce weaker seedlings (48); metastatic
cancer cells are more invasive due to a loss of contact inhibition of locomotion (49) that also

lowers their cellular proliferation rates (50-53). Additional examples include other plants (54-
56), freshwater ciliates (57), bugula (58), fish (59), crickets (60), beetles (61, 62) and butterflies
(63). Our invasive rule may potentially be applied to these systems as well, despite their specific
intricacies, and used to make quantitative predictions of invasion outcomes.

We should note two exceptions to our findings. First, while our model predicts that a
species with faster growth but slower dispersal should be able to invade (domain 1 of Fig. 2A),
we did not observe examples of this in our experimental system. And, besides observations by
another group in laboratory experiments with *Escherichia coli* (64), we could not find examples
in nature either. The reason for not finding invasion of rapid growers that disperse slower may be
population history: Empirical and theoretical studies of range expansions suggest that only
dispersal can be improved in expanding populations (65-68); invasion of new niches is possibly a
rare event, whereas competition within a confined, but relatively homogenous environment is
more common. In such situations, the selection is not on dispersal but on growth, which means
that—in most species—growth rates may already be close to their physiological limit (Fig. 4A).
Individuals challenged to overcome spatial structure may only have dispersal-related traits left to
improve. Second, while the trade-off between growth and dispersal may be found and seem
logical, a comparative analysis of dispersal in terrestrial and semi-terrestrial animals suggested
that dispersal and fecundity may be positively correlated (69).

The trade-off between growth and dispersal likely results from molecular, cellular and
physiological constraints, whereas the invasion rule $v_2 > v_1$ specifies the outcome of invasion.
The rule determines the maximal cost in growth $|\Delta r|_{\max}$ that the invader can afford to pay for
faster dispersal and still be able to dominate the edge of the expanding population (Fig. 4B):

375
$$|\Delta r|_{\max} = r_1 \left(1 - \frac{D_1}{D_1 + \Delta D}\right) \tag{7}$$

where $\Delta D$ is the increase of dispersal in species 2 relatively to species 1. The evolutionary
experiment that originally created the hyperswarmers always produced mutations in *fleN*, a gene
involved in the regulation of flagellar synthesis (18). It is possible that other mutations could
increase dispersal even more, but were not favored if they carried costs higher than $|\Delta r|_{\max}$.

The invasion rule $v_2 > v_1$ could be combined with quantitative knowledge on the
physiological mechanisms of a trade-off between growth and dispersal and used to better predict
the evolution of expanding populations. For example, if the trade-off can be represented by a line
in $(r, D)$ space the slope of that line, $dr/dD$, represents the cost of dispersal. When the trade-off
is subtle, the slope is shallow and we predict that the population will evolve to disperse faster
with a lower growth rate (Fig. 4C). Conversely, when the trade-off is strong, the slope is steep
and we predict that the population will evolve a higher growth rate and slower dispersal (Fig.
4D). According to this model, the *P. aeruginosa* system has a subtle trade-off: the improved
dispersal advantage of hyperswarmers is ~100% but costs only ~10 % of their growth rate
relatively to wild-type (18).

Using the simple invasion rule to make predictions about real world scenarios must take
into account the appropriate time scales. Our analysis showed that in some conditions where the
model predicts invasion, the invasion might take so long that it becomes irrelevant; other

processes—working on shorter time scales—would likely alter the system in the meantime. For
example, the hyperswarmers may never invade a real world population of the wild-type if
conditions change such that the bacteria need to suddenly form a biofilm. This is particularly
important because the same *fleN* mutations that give multiple flagella to hyperswarmers also
decreased their ability to form biofilm communities (18). Like Hamilton's rule and other
cornerstone concepts of evolutionary theory, if the conditions change dramatically before a fast
invader has the chance to dominate the edge of the expanding population then obviously the
model predictions will not hold.

401
402 **Figure 4. Evolutionary predictions from the invasion diagram in four scenarios**. A: the growth rate is bounded
403 by physiology. B: illustration of the maximal cost of an increased dispersal. C and D: linear trade-off between
404 growth and dispersal that corresponds to a low cost of dispersal (C) and high cost of dispersal (D).
405

In summary, we found a simple rule for invasion that, expanded with trade-off constraints
observed for each particular system, could be used to predict the fate of expanding populations.
Every model requires simplifying assumptions, and ours is certainly not an exception. In those
systems where the rule does hold, the invasion rule could be used to help control expanding
populations. Systems in this category may include the growth of cancer tumors and invasion of
non-native species in ecosystems.
412
413
414
415

**References**

1.    Phillips BL, Brown GP, Shine R. Life-history evolution in range-shifting populations. Ecology. 2010;91(6):1617-27.

2.    Murray JD. Mathematical biology. 3rd ed. New York: Springer; 2002.

3.    Nadell CD, Foster KR, Xavier JB. Emergence of spatial structure in cell groups and the evolution of cooperation. PLoS computational biology. 2010;6(3):e1000716.

4.    Hallatschek O, Nelson DR. Gene surfing in expanding populations. Theoretical population biology. 2008;73(1):158-70.

5.    Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding frontiers promotes gene segregation. Proceedings of the National Academy of Sciences of the United States of America. 2007;104(50):19926-30.

6.    Chuang A, Peterson CR. Expanding population edges: theories, traits, and trade-offs. Global change biology. 2016;22(2):494-512.

7.    Phillips BL, Brown GP, Webb JK, Shine R. Invasion and the evolution of speed in toads. Nature. 2006;439(7078):803.

8.    Waclaw B, Bozic I, Pittman ME, Hruban RH, Vogelstein B, Nowak MA. A spatial model predicts that dispersal and cell turnover limit intratumour heterogeneity. Nature. 2015;525(7568):261-4.

9.    Davies J, Davies D. Origins and evolution of antibiotic resistance. Microbiology and molecular biology reviews : MMBR. 2010;74(3):417-33.

10.   Bonte D, Van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, et al. Costs of dispersal. Biological reviews of the Cambridge Philosophical Society. 2012;87(2):290-312.

11.   Kneitel JM, Chase JM. Trade-offs in community ecology: linking spatial scales and species coexistence. Ecol Lett. 2004;7(1):69-80.

12.   Livingston G, Matias M, Calcagno V, Barbera C, Combe M, Leibold MA, et al. Competition-colonization dynamics in experimental bacterial metacommunities. Nature communications. 2012;3:1234.

13.   Tilman D. Competition and Biodiversity in Spatially Structured Habitats. Ecology. 1994;75(1):2-16.

14.   Dai L, Korolev KS, Gore J. Slower recovery in space before collapse of connected populations. Nature. 2013;496(7445):355-8.

15.   Jessup CM, Kassen R, Forde SE, Kerr B, Buckling A, Rainey PB, et al. Big questions, small worlds: microbial model systems in ecology. Trends in ecology & evolution. 2004;19(4):189-97.

16.   Mitri S, Clarke E, Foster KR. Resource limitation drives spatial organization in microbial groups. The ISME journal. 2016;10(6):1471-82.

17.   Gandhi SR, Yurtsev EA, Korolev KS, Gore J. Range expansions transition from pulled to pushed waves as growth becomes more cooperative in an experimental microbial population. Proceedings of the National Academy of Sciences. 2016;113(25):6922-7.

18.   van Ditmarsch D, Boyle KE, Sakhtah H, Oyler JE, Nadell CD, Deziel E, et al. Convergent evolution of hyperswarming leads to impaired biofilm formation in pathogenic bacteria. Cell reports. 2013;4(4):697-708.

19.   Deforet M, van Ditmarsch D, Carmona-Fontaine C, Xavier JB. Hyperswarming adaptations in a bacterium improve collective motility without enhancing single cell motility. Soft matter. 2014;10(14):2405-13.

20.   Fisher RA. The wave of advance of advantageous genes. Annals of Eugenics. 1937;7(4):355-69.

21.   Kolmogorov A, Petrovskii I, Piscunov N. A study of the equation of diffusion with increase in the quantity of matter, and its application to a biological problem. Byul Moskovskogo Gos Univ. 1937;1(6):1--25.

22.   Giometto A, Rinaldo A, Carrara F, Altermatt F. Emerging predictable features of replicated biological invasion fronts. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(1):297-301.

23.    Deforet M, van Ditmarsch D, Xavier JB. Cell-Size Homeostasis and the Incremental Rule in a Bacterial Pathogen. Biophysical journal. 2015;109(3):521-8.

24.    Murray J. Mathematical Biology: I. An Introduction (Interdisciplinary Applied Mathematics) (Pt. 1): Springer; 2007.

25.    Bénichou O, Calvez V, Meunier N, Voituriez R. Front acceleration by dynamic selection in Fisher population waves. Physical Review E. 2012;86(4):041908.

26.    Korolev KS. Evolution Arrests Invasions of Cooperative Populations. Physical review letters. 2015;115(20):208104.

27.    Pigolotti S, Benzi R, Perlekar P, Jensen MH, Toschi F, Nelson DR. Growth, competition and cooperation in spatial population genetics. Theoretical population biology. 2013;84:72-86.

28.    Reiter M, Rulands S, Frey E. Range Expansion of Heterogeneous Populations. Physical review letters. 2014;112(14):148103.

29.    Ramanantoanina A, Ouhinou A, Hui C. Spatial assortment of mixed propagules explains the acceleration of range expansion. PloS one. 2014;9(8):e103409.

30.    Bouin E, Calvez V, Meunier N, Mirrahimi S, Perthame B, Raoul G, et al. Invasion fronts with variable motility: Phenotype selection, spatial sorting and wave acceleration. Cr Math. 2012;350(15-16):761-6.

31.    Lehe R, Hallatschek O, Peliti L. The Rate of Beneficial Mutations Surfing on the Wave of a Range Expansion. PLoS computational biology. 2012;8(3):e1002447.

32.    King JR, McCabe PM. On the Fisher-KPP equation with fast nonlinear diffusion. P Roy Soc a-Math Phy. 2003;459(2038):2529-46.

33.    Holzer M, Scheel A. Accelerated Fronts in a Two-Stage Invasion Process. Siam J Math Anal. 2014;46(1):397-427.

34.    Berestycki H, Coulon AC, Roquejoffre JM, Rossi L. The effect of a line with nonlocal diffusion on Fisher-KPP propagation. Math Mod Meth Appl S. 2015;25(13):2519-62.

35.    Lewis MA, Li B, Weinberger HF. Spreading speed and linear determinacy for two-species competition models. Journal of Mathematical Biology. 2002;45(3):219-33.

36.    Jong-Shenq G, Chang-Hong W. Dynamics for a two-species competition–diffusion model with two free boundaries. Nonlinearity. 2015;28(1):1.

37.    Waite AJ, Frankel NW, Dufour YS, Johnston JF, Long J, Emonet T. Non‐genetic diversity modulates population performance. Molecular Systems Biology. 2016;12(12).

38.    Otto SP, Whitlock MC. The probability of fixation in populations of changing size. Genetics. 1997;146(2):723-33.

39.    Gillespie JH. Population genetics: a concise guide: JHU Press; 2010.

40.    Servedio MR, Brandvain Y, Dhole S, Fitzpatrick CL, Goldberg EE, Stern CA, et al. Not Just a Theory—The Utility of Mathematical Models in Evolutionary Biology. PLOS Biology. 2014;12(12):e1002017.

41.    Hamilton WD. The genetical evolution of social behaviour. I. Journal of Theoretical Biology. 1964;7(1):1-16.

42.    Foster K, Wenseleers T, Ratnieks F. Kin selection is the key to altruism. Trends in ecology & evolution. 2006;21(2):57-60.

43.    Burton OJ, Phillips BL, Travis JM. Trade-offs and the evolution of life-histories during range expansion. Ecol Lett. 2010;13(10):1210-20.

44.    Phillips BL. Evolutionary processes make invasion speed difficult to predict. Biol Invasions. 2015;17(7):1949-60.

45.    Kubisch A, Fronhofer EA, Poethke HJ, Hovestadt T, Associate Editor: Sean HR, Editor: Troy D. Kin Competition as a Major Driving Force for Invasions. The American Naturalist. 2013;181(5):700-6.

46.    Hudson CM, Phillips BL, Brown GP, Shine R. Virgins in the vanguard: low reproductive frequency in invasion-front cane toads. Biol J Linn Soc. 2015;116(4):743-7.

47.    Richardson DM, Cowling RM, Le Maitre DC. Assessing the risk of invasive success in Pinus and
Banksia in South African mountain fynbos. Journal of Vegetation Science. 1990;1(5):629-42.
48.    Reich PB, Oleksyn J, Tjoelker MG. Seed Mass Effects on Germination and Growth of Diverse
European Scots Pine Populations. Can J Forest Res. 1994;24(2):306-20.
49.    Carmona-Fontaine C, Matthews HK, Kuriyama S, Moreno M, Dunn GA, Parsons M, et al.
Contact inhibition of locomotion in vivo controls neural crest directional migration. Nature.
2008;456(7224):957-61.
50.    Gerlee P, Nelander S. The impact of phenotypic switching on glioblastoma growth and invasion.
PLoS computational biology. 2012;8(6):e1002556.
51.    Biddle A, Liang X, Gammon L, Fazil B, Harper LJ, Emich H, et al. Cancer stem cells in
squamous cell carcinoma switch between two distinct phenotypes that are preferentially migratory or
proliferative. Cancer research. 2011;71(15):5317-26.
52.    Widmer DS, Cheng PF, Eichhoff OM, Belloni BC, Zipser MC, Schlegel NC, et al. Systematic
classification of melanoma cells by phenotype-specific gene expression mapping. Pigment cell &
melanoma research. 2012;25(3):343-53.
53.    Kim IS, Heilmann S, Kansler ER, Zhang Y, Zimmer M, Ratnakumar K, et al. Microenvironment-
derived factors driving metastatic plasticity in melanoma. Nature communications. 2017;8:14343.
54.    Ganeshaiah KN, Shaanker RU. Seed Size Optimization in a Wind Dispersed Tree Butea-
Monosperma - a Trade-Off between Seedling Establishment and Pod Dispersal Efficiency. Oikos.
1991;60(1):3-6.
55.    Huang FF, Peng SL, Chen BM, Liao HX, Huang QQ, Lin ZG, et al. Rapid evolution of dispersal-
related traits during range expansion of an invasive vine Mikania micrantha. Oikos. 2015;124(8):1023-30.
56.    Williams JL, Kendall BE, Levine JM. Rapid evolution accelerates plant population spread in
fragmented experimental landscapes. Science. 2016;353(6298):482-5.
57.    Fronhofer EA, Altermatt F. Eco-evolutionary feedbacks during experimental range expansions.
Nature communications. 2015;6:6844.
58.    Lange R, Marshall DJ. Propagule size and dispersal costs mediate establishment success of an
invasive species. Ecology. 2016;97(3):569-75.
59.    Agostinho AA, Suzuki HI, Fugi R, Alves DC, Tonella LH, Espindola LA. Ecological and life
history traits of Hemiodus orthonops in the invasion process: looking for clues at home. Hydrobiologia.
2015;746(1):415-30.
60.    Simmons AD, Thomas CD. Changes in dispersal during species' range expansions. Am Nat.
2004;164(3):378-95.
61.    Ochocki BM, Miller TE. Rapid evolution of dispersal ability makes biological invasions faster
and more variable. Nature communications. 2017;8:14315.
62.    Weiss-Lehman C, Hufbauer RA, Melbourne BA. Rapid trait evolution drives increased speed and
variance in experimental range expansions. Nature communications. 2017;8:14303.
63.    Hughes CL, Hill JK, Dytham C. Evolutionary trade-offs between reproduction and dispersal in
populations at expanding range boundaries. Proceedings Biological sciences / The Royal Society.
2003;270 Suppl 2:S147-50.
64.    Fraebel DT, Mickalide H, Schnitkey D, Merritt J, Kuhlman TE, Kuehn S. Environment
determines evolutionary trajectory in a constrained phenotypic space. eLife. 2017;6.
65.    Phillips BL, Brown GP, Shine R. Evolutionarily accelerated invasions: the rate of dispersal
evolves upwards during the range advance of cane toads. Journal of evolutionary biology.
2010;23(12):2595-601.
66.    Perkins TA, Phillips BL, Baskett ML, Hastings A. Evolution of dispersal and life history interact
to drive accelerating spread of an invasive species. Ecol Lett. 2013;16(8):1079-87.
67.    Hallatschek O, Fisher DS. Acceleration of evolutionary spread by long-range dispersal.
Proceedings of the National Academy of Sciences of the United States of America. 2014;111(46):E4911-
9.
68.    Travis JMJ, Dytham C. Dispersal evolution during invasions. Evol Ecol Res. 2002;4(8):1119-29.

69.      Stevens VM, Whitmee S, Le Galliard JF, Clobert J, Bohning-Gaese K, Bonte D, et al. A
comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial animals. Ecol Lett.
2014;17(8):1039-52.
568
569