Title: Stoichiometric homeostasis of wetland vegetation along a nutrient gradient in a

subtropical wetland. Understanding stoichiometric mechanisms of nutrient retention in wetland

macrophytes.

Running Header: Wetland homeostasis

Paul Julian II[1], Stefan Gerber[2], Rupesh K. Bhomia[2], Jill King[3], Todd Z. Osborne[4,2], Alan L.
Wright[1],

[1] University of Florida, Soil and Water Sciences Department, Ft. Pierce, FL 34945
*Corresponding Author: <u>pjulian@ufl.edu;</u> ORCID: 0000-0002-7617-1354
[2] University of Florida, Soil and Water Sciences Department, Gainesville, FL, 32611

[3] South Florida Water Management District, Water Quality Treatment Technologies, West Palm
Beach, FL, 33406

[4] University of Florida, Whitney Laboratory for Marine Bioscience, St Augustine, FL 32080

Abstract (limit 250 words)

Nutrient homeostasis relates ambient stoichiometric conditions in an environment to the stoichiometry of

living entities of the ecosystem. In wetland ecosystems, vegetation can be a large, highly variable and

dynamic sink of nutrients. This study investigated stoichiometric homeostasis of dominant emergent and

submerged aquatic vegetation (EAV and SAV, respectively) within two treatment flow-ways (FW) of

Everglades Stormwater Treatment Area 2 (STA-2). These FW encompass a large gradient in plant

nutrient availability. The hypotheses of this study is that wetland vegetation is non-homeostatic relative to

ambient nutrients and consequently nutrient resorption will not vary along the nutrient gradient. We

developed a framework to investigate how vegetation uptake and resorption of nutrients contribute

separately to homeostasis. Overall, the wetland vegetation in this study was non-homeostatic with respect

to differential uptake of nitrogen (N) vs. phosphorus (P). Resorption evaluated for EAV was high for P

and moderate for N, resorption efficiency did not significantly vary along the gradient and therefore did

not affect overall homeostatic status. Nutrient addition experiments may help to compensate for some of

the limitation of our study, especially with respect to resolving the primary nutrient source (organic vs.

inorganic sources, water vs. soil compartment) and nutrient utilization rates.

Keywords: Phosphorus, Everglades, resorption, sink strength, Stormwater Treatment Area

"The environment not only determines the conditions under
which life exists, but the organisms influence the
conditions prevailing in their environment."
- (Redfield 1958)

A tenet of ecological stoichiometry is that abundance of carbon (C), nitrogen (N), phosphorus (P) or other elements is regulated by reciprocal interaction between organisms and their environment (Redfield 1958), and this relationship forms the underlying framework of stoichiometric homeostasis. In homeostatic organisms, nutrient stoichiometry is relatively fixed while non-homeostatic organisms display flexibility with their respective nutrient stoichiometry. Homeostatic organisms regulate their elemental composition by adjusting bulk material uptake (i.e. ingestion rate), degree of food selectivity for growth and post-assimilation processes to regulate the acquisition of elements. Elemental uptake by non-homeostatic organisms, is mostly controlled by supply (i.e. nutrient availability) rather than demand (i.e. relative growth rate) with limits to nutrient loss processes such as excretion in animals and senescence in plants (Killingbeck 1996; McGroddy et al. 2004).

The concept of stoichiometric homeostasis was formulated as a means to understand patterns of C, N and P in zooplankton to evaluate food quality and nutrient recycling (Jotaro Urabe et al. 1995; Elser and Urabe 1999; J. Urabe and Sterner 2001; Sterner and Elser 2002; Malzahn et al. 2007; Mulder 2007; Mulder and Bowden 2007). Later, the concept has been broadened to understanding food web stoichiometric dynamic (Persson et al. 2010), specific stream food web stoichiometric interactions (Feijoó et al. 2014), resource dependence of bacteria (Makino et al. 2003), and grassland community level responses to climate change (Gu et al. 2017). The stoichiometric homeostasis conceptual model is thus of particular interest with respect to nutrient

enrichment and biotic uptake processes due to selective uptake of nutrients during nutrient

enriched conditions (Sterner and Elser 2002; Small et al. 2009).

Stoichiometric flexibility or homeostatic plasticity (i.e. non-homeostasis) can manifest across

different spatial scales from organisms to ecosystems. At the organismal level, stoichiometric

flexibility can occur through changes in nutrient allocation to tissue types or biochemical

changes in cellular composition (Sistla and Schimel 2012; Rivas-Ubach et al. 2012). At the

ecosystem level, changes in stoichiometry can occur through changes in ecosystem composition

including changes in consumer and producer species composition and encroachment of invasive

species (Sistla et al. 2015). Therefore, stoichiometric flexibility is driven by species-specific

nutrient biomass storage, physiological plasticity, interspecific competition and the variability in

nutrient supply versus demand (Elser et al. 2010; Sardans et al. 2012; Sistla et al. 2015).

Heterotrophs (i.e. herbivores) are constantly faced with stoichiometric mismatch but can combat

it to maintain constant elemental composition through strategies of selective consumption or

elimination of particular elements (Filipiak and Weiner 2014). Meanwhile, autotrophs such as

plants have internal mechanisms to continue building up biomass and stoichiometric composition

through selective uptake and internal nutrient retention processes such as nutrient resorption.

Nutrient resorption, is a process in which nutrients are withdrawn from senescing leaves before

abscission to be used for new growth or stored for later use (Van Heerwaarden et al. 2003). It is

assumed to be one of the most important strategy used by plants to conserve nutrients

(Rejmánková 2005).

Most nutrient resorption studies focus on terrestrial (upland) evergreen, deciduous, forb and

graminoid species (Killingbeck 1996; Aerts and Chapin 1999; Van Heerwaarden et al. 2003;

Peng et al. 2016) with only few studies investigating wetland plant species (Miao 2004;

Rejmánková 2005; Rejmánková and Snyder 2008). Rejmánková (2005) observed relatively high

nutrient resorption efficiencies for wetland plant species including cattail (*Typha domingensis*)

for both N and P across a variety of nutrient limiting conditions. In a nutrient addition study,

Rejmánková and Snyder (2008) observed high N and P resorption across P-limiting conditions in

cattail and spike rush (*Eleocharis* spp.). Meanwhile, Miao (2004) observed high P and low N

resorption in two wetland macrophytes, sawgrass (*Cladium jamaicense*) and cattail in P-limiting

conditions. These studies suggest that P resorption is typically high regardless of P limitation

status while N resorption can vary depending on N availability and can contribute significantly to

stoichiometric homeostatic.

While limited to a few studies, prior work suggest that plants range in their degree of

homeostasis from very weak to non-homeostatic (Yu et al. 2011; Gu et al. 2017). However, these

studies focused on terrestrial grasses and shrub species and not wetland plant species distributed

along a nutrient gradient. In contrast to upland ecosystems biogeochemical processes in wetlands

are influenced by specific non-biotic (inundation, geochemical precipitation of nutrients) and

biotic (i.e. plant, periphyton, algae, etc.) interactions that can alter the oxidation-reduction and

pH environment (Reddy and DeLaune 2008; Kadlec and Wallace 2009). These interactions pose

additional challenges to producers with respect to nutrient supply and demand. Earlier work in

wetlands shows a decrease of plant P concentration with decreasing ambient P concentration

(Richardson et al.,1990, Julian et al., 2019). This suggests non-homeostatic behavior although

the author did not test for homeostasis. Therefore, the first objective of this study was to evaluate

the strength of stoichiometric homeostasis in dominant wetland species along a nutrient gradient

in two flow-ways (FW) within the Everglades Stormwater Treatment Areas (STAs; Fig. 1).

These treatment wetlands use emergent and submerged aquatic vegetation (EAV and SAV,

respectively) to remove nutrients via biotic uptake. Prior studies of nutrient resorption with

respect to emergent aquatic vegetation (EAV) suggest that EAV species use both "conservation

of use" and "enhanced acquisition" of P to fulfill nutrient demand in P-limited systems (Miao

2004; Rejmánková 2005; Rejmánková and Snyder 2008). Moreover these studies indicate that

EAV utilize the "conservation of use" strategy making resorption dependent upon nutrient

availability. Sterner and Elser (2002) suggest that autotrophic organism's stoichiometry is not

solely dictated by nutrient availability but also biophysical, physiological adaptions and optimal

nutrient utilization. The first objective of this study was to determine the degree of stoichiometric

homeostasis of wetlands plants along a strong nutrient gradient with the hypothesis that EAV and

SAV are generally homeostatic relative to the ambient environment due to biological adaptions

in nutrient uptake, utilization and storage. The second objective of this study was to evaluate

nutrient resorption dynamics of EAV species within the treatment wetland and with the

hypotheses that resorption efficiency (RE) significantly differs as a function of nutrient

limitation status (i.e. N or P limitation) and the degree of stoichiometric homeostasis observed

among wetland plants is caused by differences in nutrient resorption.

Methods

Study Area

A total of six Everglades STAs with an approximate area of 231 km^2 are located south of Lake

Okeechobee in the southern portion of the Everglades Agricultural Area (Fig. 1). The Everglades

STAs are established on land which earlier had natural wetlands and/or agricultural farmlands

under sugarcane production. The primary source of inflow water to the STAs is agricultural

runoff originating from approximately 284 km^2 of farmland upstream. Everglades STA treatment

cells or FWs are comprised of a mixture of EAV and SAV communities in several configurations

including EAV and SAV treatment cells arranged in parallel or in series (Chen et al. 2015).

Stormwater Treatment Area-2 (STA-2) has been in operation since June 1999 with an effective

treatment area of approximately 63 km^2 divided into eight treatment cells. This study was

conducted in two cells, FWs 1 and 3. The vegetative community of FW 1 is comprised

predominately of EAV vegetation including *Typha domingensis* Pers. (cattail) and *Cladium*

jamaicense Crantz (sawgrass) while FW 3 is dominantly SAV including *Chara* spp. (muskgrass),

Potamogeton spp. (pondweed) and *Najas guadalupensis* Spreng (southern naiad) with

approximately a quarter of the FW occupied by EAV species. Furthermore, prior to STA-2

construction FW 1 was a historic natural wetland while approximately three-quarters of FW 3

was previously farmed and is now managed as an SAV system (Juston and DeBusk 2006).

Data Source

Data used in this study were collected by South Florida Water Management District (SFWMD)

and University of Florida as a part of a larger project within the overall SFWMD's Restoration

Strategies Science Plan to evaluate performance and optimization of the Everglades STAs (South

Florida Water Management District 2013). Data from the Science Plan study: Evaluation of P

Sources, Forms, Flux and Transformation Processes in the STAs was used for this investigation

(UF-WBL 2017; Villapando and King 2018). We used weekly surface water grab samples

collected at monitoring locations along a nutrient gradient from inflow to outflow within FWs 1

and 3 in STA-2 to characterize changes in nutrient concentrations during prescribed/semi-

managed flow events (Fig. 1). Samples were collected just under the water surface at 10-15 cm

depth during prescribed flow events that cycled through various flow/no-flow sequences for FWs

1 and 3. For this study we used the water column parameters total P (TP), total N (TN), and

dissolved organic C (DOC) (Villapando and King 2018). In the Everglades system, the surface

water organic C pool is predominately composed of the DOC fraction with low particulate OC

concentrations (Julian et al. 2017), therefore total OC is assumed to be equivalent to DOC

concentrations. We also made use of soil samples collected along the flow transects twice during

the dry and wet seasons between 2015 and 2016 and analyzed for TP, TN, and TC. Vegetation

tissue were collected from live and senescent above-ground biomass (AGB) vegetation in FW 1

while only live-AGB was sampled from FW 3 at the end of the 2015 (November 2015) and 2016

(September 2016) wet seasons. The ability to collect senescent SAV is extremely difficult as it

decomposes rapidly and becomes part of the "litter" immediately, therefore no samples were

available for analysis. Vegetation samples were collected from inflow, mid and outflow regions

of the FWs within close proximity to the surface water and soil monitoring locations (Fig. 1).

Only co-located surface water, soil and vegetation sampling locations were considered for this

study (Fig. 1). Nutrient content of vegetation was determined from dried ground and

homogenized samples. For conducting analyses and summary statistics, data reported as less than

method detection limit (MDL; Table S1) were assigned a value of one-half the MDL, unless

otherwise noted.

Data Analysis

We categorized foliar TN:TP molar ratios for EAV along the FW 1 transect into potential N-

limitation (molar TN:TP<14), potential N and P co-limitation (14 < TN:TP < 16) and P

limitation (TN:TP > 16). For SAV along the FW 3 transect we categorized foliar TN:TP molar

ratios into potential N-limitation (TN:TP < 27) and potential P-limitation (TN:TP > 27). These

potential nutrient limiting categories were consistent with other EAV studies (Koerselman and

Meuleman 1996; Rejmánková 2005) and SAV studies (Duarte 1992; Fernández-Aláez et al.

1999). We compared nutrient limitation categories between FW regions (i.e. inflow, mid and

outflow) using chi-squared (RxC) analysis.

We evaluated P and N resorption efficiency (RE_{TP} and RE_{TN}, respectively) along the FW

transect by comparing living (X_{GR}) to senescent (X_{SEN}) EAV AGB nutrient tissue concentrations.

Resorption efficiency was estimated using Eq. 1, consistent with prior studies (Killingbeck 1996;

Reed et al. 2012).

$$RE_x = \frac{X_{GR} - X_{SEN}}{X_{GR}} \times 100 \qquad \text{Eq. 1}$$

We compared RE_{TP} and RE_{TN} values between P-limiting categories identified above for aquatic

vegetation type specific foliar TN:TP molar ratios using the Kruskal-Wallis rank sum test.

Resorption efficiency values for each site was compared with fractional distance from inflow

using Spearman's rank correlation. We also compared RE_{TP} and RE_{TN} between stations using

Dunn's multiple comparison rank sum test ('dunn.test' R package; Dinno 2015).

We characterized the strength of vegetation TN:TP total stoichiometric homeostasis (H_T) using

the homeostasis coefficient (Eq. 2). Equation 2a is a log-linear model where y is the live-AGB

TN:TP molar ratio, x is the TN:TP molar ratio of the resource, c is a constant that determines the

intercept and $1/H_T$ is the slope of the model between the log transformed ratios x and y and

represents the strength of homeostatic maintenance of nutrient ratios within the consumer

biomass. Reorganizing Eq. 2a to solve for $\frac{1}{H_T}$ we get Eq. 2b.

$$\log(y) = \log(c) + \frac{\log(x)}{H_T} \qquad \text{Eq. 2a}$$

$$\frac{1}{H_T} = \frac{\log(y) - \log(c)}{\log(x)} \qquad \text{Eq. 2b}$$

Hypothetically, values of $1/H_T$ range between zero and one, with values approaching one

indicating a non-homeostatic relationship and values approaching zero signifying consumer-

resource homeostasis. The degree of homeostasis was defined consistent with prior studies

(Makino et al. 2003; Persson et al. 2010; Feijoó et al. 2014) as the following:

1. $0.00 < 1/H_T < 0.25$: Homeostatic

2. $0.25 < 1/H_T < 0.50$: Weakly-Homeostatic

3. $0.50 < 1/H_T < 0.75$: Weakly Non-Homeostatic

4. $0.75 < 1/H_T < 1.00$: Non-Homeostatic

The stoichiometric homeostasis coefficient is an integrated metric to relate the ambient

environment's stoichiometry to that of the vegetation biomass. Moreover H_T can be used to

determine if the observed variation in the biota stoichiometry is the result of physiological

adjustments or reflects species-specific turnover (Elser et al. 2010). To assess the homeostatic

relationship of the vegetation relative to the ambient environment, N:P ratios of resource pools

were compared to the N:P ratios of AGB for each vegetation community. The communities'

uptake pathways differ between SAV and EAV - SAV primarily assimilate nutrients through the

water column while EAV obtain nutrients from the soils via soil solution (Richardson and

Marshall 1986; Reddy et al. 1999; Dierberg et al. 2002). Therefore, we compared AGB N:P

values of both EAV and SAV to surface water and soil N:P values (denoted as x in Eq. 2). Due

to offset sampling times, soil and AGB data were paired only when samples were collected

within six months of each other for each site. Site-specific surface water mean TN:TP values

during this six-month timeframe were paired with AGB values. Biota TN:TP molar ratios were

identified as each respective vegetation type live-AGB TN:TP molar ratio (y in Eq. 2). We used

the Theil-Sen single median linear model ('mblm' package; Komsta 2013) to evaluate biota and

resource TN:TP molar ratios to estimate c and $1/H_T$ in Eq. 2.

To evaluate the overall homeostatic versus non-homeostatic status among vegetation within a

FW, we evaluated the slope of the mblm linear models for each FW to determine if the slope was

significantly different from 0.50 using the One-Sample Wilcoxon rank signed test. This test is

similar to the t-test slope comparison method identified by Zar (2010) for parametric linear

models. The 0.50 threshold was selected as it is the criterion that differentiates homeostasis from

non-homeostasis (see above).

We also characterized the degree of stoichiometric homeostasis with respect to differential

resorption similar to the approach for stoichiometric homeostasis (H_T) above. Resorption

efficiency homeostasis (H_{RE}) was evaluated for EAV to assess whether RE changes with respect

to the ambient environment. We used mblm to evaluate the relative loss of nutrients after

resorption (y_{RE}) relative to resource TN:TP (x) where the slope of linear model is $1/H_{RE}$. The

relative loss of nutrients after resorption (y_{RE}) or the senescence efficiency is represented by Eq.

3. In the supplementary material we develop the framework to evaluate how resorption

contributes to homeostasis, the resulting relationship is given in Eq. 4 which is very similar to

Eq. 2. The subtraction of one on the left-hand side in Eq. 4 originates from the fact that this

equation states the vegetation's effort into homeostasis via resorption, where zero would yield an

effort to maintain full homeostasis and negative one would mean that there is no homeostasis

with respect to resorption (Supplemental Material). The regression constant c is different than in

Eq. 2, but is in both cases irrelevant to the calculation of stoichiometric homeostasis.

$$y_{RE} = \frac{(1 - RE_{TP})}{(1 - RE_{TN})} \qquad \text{Eq. 3}$$

$$\frac{1}{H_{RE}} - 1 = \frac{\log(y_{RE}) - \log(c)}{\log(x)} \qquad \text{Eq. 4}$$

235   In the supplemental material, we have developed a framework which can relate the two major

236   pathways for homeostasis (resorption and uptake) with each other. Total homeostasis (H_T; Eq. 2)

237   and the vegetation's effort to maintain homeostasis can then be partitioned into contributions

238   from resorption (Eq. 4) and uptake:

$$\left(1 - \frac{1}{H_{RE}}\right) + \left(1 - \frac{1}{H_U}\right) = \left(1 - \frac{1}{H_T}\right) \qquad \text{Eq. 5}$$

All statistical operations were performed with R© (Ver 3.1.2, R Foundation for Statistical

Computing, Vienna Austria), unless otherwise stated all statistical operations were performed

using the base R library. The critical level of significance was set at α=0.05.

Results

Surface water DOC, TN concentrations, DOC:TP and TN:TP molar ratios were overall higher in

FW 3 (SAV) while surface water TP concentrations and DOC:TN molar ratios were higher in

FW 1 (EAV) (Table 1). Soil organic matter, as indicated by loss-on-ignition, was greater in FW1

while soil Ca concentrations were greatest in FW3 soils indicating that FW1 soil is more organic

(i.e. peat) and FW3 more mineral (i.e. marl). In addition to qualitative differences between soil

composition, FW1 soils were generally more enriched with TP and TN. Among carbon and

nutrients, soil TP had the highest coefficient of variation between and within sites and FWs with

an overall coefficient of variation of 48%. Overall, FW 1 exhibited the highest coefficient of

variation for soil TP with 40% spatial and temporal variability (Table 1).

Tissue Stoichiometry

Live AGB TN : TP molar ratios were greatest in FW 3 (Table 1 and Fig. 2). Vegetation from

both FWs spanned their respective potential N and P-limiting conditions with most values above

the P limits identified for EAV (TN:TP>16) and SAV (TN:TP>27) communities (Table 1 and

Fig. 2). Furthermore, AGB TC:TN values were greater in FW 1, ranging from 19.3 to 82.8

whereas FW 3 ranged from 7.5 to 23.7 suggesting the potential for N-limitation in FW 1 (Fig. 2).

AGB TN:TP values were significantly different along the FW 1 transect (χ^2=19.4, df=4, ρ<0.01)

characterized by strong nutrient decline along the FW from inflow to outflow. Several values

below the TN:TP value of 14 (indicating N limiting conditions) were observed, most of which

occurred at the Inflow and Mid regions of the FWs and all values above the P-limiting criteria

occurred in the Outflow region. Meanwhile, nutrient limiting conditions were not significantly

different along the FW 3 flow path (χ^2=5.6, df=2, ρ=0.06) with the majority of the TN:TP values

greater than 27 indicating potential P-limitation, and four samples less than 27 occurring in the

Inflow and Mid regions of FW.

Stoichiometric Homeostasis

Changes in AGB stoichiometry occurred concurrent with changes in both surface water and soil

resource stoichiometry pools (Fig. 3). Log-log models comparing vegetation N:P to resource N:P

resulted in slopes significantly different from zero (Table 2) with varying levels of fit. The model

with the lowest R^2 was the SAV (FW 3) N:P to surface water N:P (Table 2). The relationship

between the vegetation and the ambient environment depends on which resource pool was

considered and thereby where each vegetation type falls on the H_T continuum. For instance,

EAV is weakly homeostatic if the water column is considered to be the major nutrient source and

non-homeostatic if nutrients are assumed to be taken up from the soil (Fig. 4 and Table 2). The

homeostasis coefficients, $1/H_T$, when considering soil as a potential resource pool, were

significantly greater than 0.5 for EAV (V=153870, $\rho<0.01$) and SAV (V=14455, $\rho<0.01$)

indicating that vegetation is non-homeostatic with respect to soil TN:TP (Fig. 4). In fact, the

homeostasis coefficient $1/H_T$ for EAV sourcing nutrient from soil is greater than 1. Meanwhile,

$1/H_T$ values when considering surface water as a potential resource pool were significantly less

than 0.5 for EAV (V=53539, $\rho<0.01$) and not significantly different for SAV (V=15452, $\rho=0.36$)

suggesting EAV and SAV with respect to surface water is weakly homeostatic to weakly non-

homeostatic, respectively (Fig. 4).

Resorption Efficiency

A total of 38 plant tissue samples were collected during the vegetation sampling events where

both live and senesced AGB were available from EAV (FW1). Cattail comprised most of the

species collected (N=36) with one sample each from lily and sawgrass. The estimated mean

RE_{TP} value for cattails was 74 ± 2 % (N=36), with the individual RE_{TP} values for water lily and

sawgrass being 70 % and 77 %, respectively (Table 3). Along FW 1, cattail RE_{TP} was not

significantly correlated with fractional distance downstream along the resource gradient (r_s

=0.21, ρ=0.22) or statistically different between sites (χ^2= 3.9, df=2, ρ=0.14). The estimated

mean RE_{TN} value for cattails was 32 ± 3 % (N=36), with RE_{TN} values for just sawgrass observed

to be 2.3% (Table 3). Much like RE_{TP}, cattail RE_{TN} was not significantly correlated with

fractional distance downstream (r_s = -0.04, ρ=0.84) or statistically different between sites (χ^2=

2.9, df=2, ρ=0.23). Both TP and TN RE did not significantly differ between nutrient limitation

categories as indicated by tissue TN:TP molar ratios (χ^2= 0.74, df=2, ρ=0.70 and χ^2= 0.38, df=2,

ρ=0.83, respectively).

Partitioned Contributors to Stoichiometric Homeostasis

Senescence efficiency ranged from 9% to 76% for P and 44% to 98% for N in cattail along FW

1. Therefore, the relative stoichiometric loss of nutrient after resorption (y_{RE}; Eq. 4) ranged from

0.16 to 0.73 for cattail along FW 1. Generally, the relative stoichiometric loss of nutrient after

resorption did not vary relative to the resource availability. When considering soil as a potential

resource pool relative to stoichiometric resorption, the slope was significantly different from zero

(V=33519, ρ<0.05) with a value of 0.24 indicating weakly homeostatic. Additionally, when

considering surface water as a potential resource pool the slope of the model was significantly

different from zero (V=21674, ρ<0.05) with a value of 0.05, suggesting a homeostatic

relationship.

Discussion

All organisms are faced with the challenges of acquiring sufficient energy and nutrients, with

most organisms dealing with a mixture of these resources in the environment that is imbalanced

relative to their requirements. Resource constraints and imbalances can alter individual species

strategies, population dynamics, inter-specific competition for resources, and even influence key

ecosystem processes (Frost et al. 2005). Ultimately, stoichiometric homeostasis is largely

controlled by a set of key physiological processes that regulate uptake, incorporation and

eventual release of nutrients (Sterner and Elser 2002). These processes must contend with what

is available in the ambient environment and can manifest as stoichiometric mismatches and lead

to nutritional imbalances. We used the concept of stoichiometric homeostasis to evaluate the

degree to which plants in a wetland ecosystem respond to variations in nutrient supply. The

dominant features of the variable nutrient supply in our system were both a decreasing P load

(i.e. uptake during transport) and a diminishing resource gradient (i.e. soil nutrient concentration)

with distance from inflow to outflow of two treatment wetland FWs.

Neither of the two types of vegetation (EAV and SAV) in the two studied FWs evaluated here

displayed clear homeostatic behavior, in that N:P stoichiometry of AGB remained independent

of the N:P stoichiometry of the resources (i.e. soil or water column nutrient pools). Considering

the water column as the resource supply, both EAV and SAV displayed weakly homeostatic to

weakly non-homeostatic behavior, respectively. However, if soil is considered as the resource

pool, EAV and SAV are considered weakly to strongly non-homeostatic, respectively. Generally,

EAV use nutrients from the soil while SAV obtains nutrients from the water column.

Early in the development of stoichiometric homeostasis theory it was hypothesized that

photoautotrophs (i.e. algae and plants) were considered to have very weak stoichiometric

homeostasis ($0.25 < 1/H < 0.50$), suggesting that they do not strongly regulate biomass

stoichiometry relative to the ambient environment due to luxury uptake of nutrients (Sterner and

Elser 2002; Reddy and DeLaune 2008; Elser et al. 2010). Since development of the concept,

however, several studies have suggested that plants can range from weakly homeostatic to non-

homeostatic (1/H >0.50) (Sterner et al. 1998; Elser et al. 2000; McGroddy et al. 2004; Güsewell

2004). Moreover, Demers et al. (2007) suggested that the relationship between external nutrient

concentrations (i.e. soil and water column) and foliar tissue nutrient concentrations in aquatic

macrophytes indicates a weakly non-homeostatic to non-homeostatic relationship similar to other

terrestrial and algae species. This conclusion was confirmed in our study from treatment

wetlands of south Florida which received high inputs of P as agricultural runoff.

A very important aspect of stoichiometric homeostasis is the proper pairing of resource to

organism along a trophic chain as this affects the efficiency of trophic transfer of. Theoretically,

$1/H$ values range from zero to one. However, values can exceed one, and in this study, a

homeostasis value greater than one was observed when comparing soil and EAV N:P. This is

manifested by a disproportionate increase in AGB N:P with increasing soil N:P. In a review of

stoichiometric homeostasis, Persson et al. (2010) reported several studies with $1/H$ values

greater than one. As suggested by Smalley et al. (2003), it is possible that non-inhibitory light

levels leading to high C-fixation and greater biosynthetic need for N and P to use the additional

C could lead to $1/H_{N:P} > 1$. EAV, and especially cattails, are generally not growth-limited by

light due to its tall structure above the water and broad leaves. Therefore, it is possible that EAV

can respond by disproportionally directing N use towards building other cellular machinery to

optimize nutrient use and functionality. An alternative mechanism may be that, in this P-limited

system, P acquisition becomes increasingly difficult as N:P increases because P is stored in

increasingly inaccessible forms within the resource pool.

It is possible that both vegetation types (EAV and SAV) take up nutrients from soil and water.

For EAV, homeostatic behavior appears stronger if the nutrient resource is assumed to come

from the soil rather than water column (Fig. 4). By the same token, homeostasis for SAV is

stronger if nutrients are assumed to be supplied from the water column (Fig. 3), although the

difference between the homeostasis coefficients for SAVs is decidedly lower compared to EAVs

(Fig 6). This may in turn partially be explained by lower variability in N:P ratios in EAV

compared to SAV, which in turn suggests less flexible N:P ratios and thus stronger homeostasis

in EAV.

As expected, stoichiometric mismatches were observed within our study system where resource

(i.e. soil or water column) N:P were greater than that of the AGB N:P (Table 1). Both soil and

water resource pools differed between FWs with water having a greater variability than that of

soil. This is notable as a strong soil P gradient is apparent along the FW (Osborne et al, *In Prep*),

potentially indicating changes in P-availability and the physiological limitation to use the

existing P-pool. This limitation has been noted in prior studies where uptake is lower in

treatments with higher available P (Shaver and Melillo 1984; Craft et al. 1995). Variability and

differences in the relative ratio of resources to consumer indicate that vegetation types access

available nutrient pools differently. In lake ecosystems, Rattray et al. (1991) determined that

SAV take up nutrients from the water column. However, other studies have demonstrated that,

while SAV are involved in active uptake of nutrient from the water column, it also modifies the

physical environment through photosynthesis and respiration causing the precipitation of P to

mineral bound P compounds (Dierberg et al. 2002; Reddy and DeLaune 2008). Submerged

macrophyte systems invoke P mineralization by inadvertently manipulating C (via dissolved

inorganic C) and pH through respiration and photosynthesis. In FW 3, pH values ranged from 6.9

to 10.4 (Julian et al. 2019) and, combined with a high mineral content environment, can lead to

coprecipitation of P with $CaCO_3$ (Boyer and Wheeler 1989)(Table 1). Much like their emergent

relatives, rooted SAV can also obtain nutrients from the soil but to a lesser degree (Brix 2003;

Dhote and Dixit 2009).   Stoichiometric mismatch in producer-consumer (i.e. detritus-

invertebrate) systems, typically identify nutritional deficiency in food sources (Filipiak and

Weiner 2014, 2017; Filipiak 2016). In this case stoichiometric mismatch is indicative less of

nutritional deficiency but more of limitation in nutrient utilization and availability as well as the

potential for plant-environment nutrient feedbacks discussed above. Regardless of the

stoichiometric mismatch observed, we still observe predominantly non-homeostatic responses by

vegetation across N to P-limiting conditions for both FWs despite being predominant P.

Generally, plant species respond to changes in nutrient available nutrients through the storage

and recycling of nutrients, ultimately shifting absolute and relative tissue nutrient concentrations

in a way suggesting a non-homeostatic relationship with their ambient environment (Güsewell

and Koerselman 2002; Güsewell et al. 2003; Rejmánková et al. 2008). The resorption of

nutrients from senescing tissues is often considered a strategy in nutrient-limiting conditions to

conserve scarce nutrients (Aerts and Chapin 1999; Rejmánková et al. 2008). Nutrient resorption

efficiency of EAV observed in this study (Table 3) (Table 1) is consistent with values observed

for aquatic macrophytes by studies elsewhere (Miao 2004; Rejmánková 2005). Rejmánková

(2005) evaluated resorption of macrophyte species from several wetlands across a broad spatial

distribution and observed that cattail achieved a maximum RE_{TP} of approximately 80% while the

study average was 70%. Rejmánková (2005) also observed that RE_{TP} was significantly greater in

P-limiting conditions. Meanwhile, Miao (2004) evaluated RE_{TP} of two aquatic macrophytes in a

mesocosm setting with two-nutrient treatment regimens broadly defined as enriched (high

nutrient availability) and unenriched (low nutrient availability). Miao (2004) observed RE_{TP}

ranging from 69 to 82% for sawgrass and 61 to 71% for cattail between treatments with no

treatment effect, indicating that ,despite available nutrient conditions, resorption remains

relatively high. Miao (2004) also observed RE_{TN} values ranging from 33 to 45% between cattail

and sawgrass species with no significant differences between treatment or species. In our study,

RE_{TP} or RE_{TN} was not significantly different along the flow way despite potential shifts in

relative N or P-limitation as indicated by foliar TN:TP values (Fig. 2) and nutrient stoichiometry

of other compartments (Julian et al. 2019). Mean RE_{TP} values observed at the inflow region,

where P resources are especially high, were slightly lower with more variability than middle and

outflow regions (Fig. 5), potentially suggesting some adaptation. We note that leaching of

senesced litter may overestimate active resorption calculated here and in previous studies.

In addition to nutrient uptake, nutrient resorption itself can also contribute to overall

stoichiometric homeostasis. In this study, we developed a concept based on first order dynamics

similar to Menge et al. (2009) of how nutrient resorption can be incorporated into the

stoichiometric homeostasis framework (Supplemental Material). By using equilibrium

assumptions, we were able to distinguish contributions of uptake versus resorption to

homeostasis (Fig. 6). In essence, we defined the effort put into homeostasis by the plant that

determines changes in stoichiometry from differential uptake and from differential resorption.

The definition of homeostasis is such that a value of zero indicates full homeostasis while a value

of one suggests non-homeostasis. Effort towards homeostasis is therefore the complement of a

homeostasis to one (Supplementary Information). The resulting calculations suggest that

nutrient resorption contributes approximately 11% and 15% to stoichiometric homeostasis in

EAV, when water and soil, respectively, are considered as the endmembers of nutrient supply

(Fig. 6). In Fig.67 we indicated a positive effort towards maintaining homeostasis via an arrow

down, with the length of the arrow indicating the magnitude of the effort. Due to the limitation of

the data collected, we were unable to determine resorption homeostasis for SAV. Despite this

limitation, given the resorption dynamics observed in EAV, it is possible that resorption

homeostasis by SAV would be similar, contributing slightly to overall total homeostasis.

Despite some shortcomings of this study, the data allow us to evaluate homeostasic behavior in

sub-tropical treatment wetland systems. As noted above, it is not clear to what degree the

vegetation uses water versus soil as the main nutrient resource, generally it is assumed that EAV

mine nutrients from the soil (via soil porewater) and SAV take up nutrients from the water

column and modify the chemical environment to facilitate abiotic mineralization. It is possible

that one nutrient is preferably obtained from the water column, and the other from the soil. Given

the higher N:P in the soil relative to the water column, it would be opportune for plants to

acquire N from the soil, while obtaining P from the water column. However, this inference may

be complicated by the fact that vegetation is specifically mining soil P, depleting soil P storages

and therefore increasing the N:P ratio in the soil. Given our data, it is not clear to what degree N

or P are accessible for plant uptake. Along the FW, the proportion of inorganic (rapidly taken up)

versus total nutrients varies in both the soil and water column (UF-WBL 2017). In the soil, the

majority of nutrients could be in a mineral form (predominately Ca-P minerals) and therefore not

as accessible to plants. Therefore, our calculation of homeostasis implicitly assumes that N and P

were mineralized according to their total nutrient ratio. While not studied here, soil microbial

interactions within the soil relative to ambient pool of nutrients could also influence site-specific

nutrient dynamics along the FW (Wright and Reddy 2001; Corstanje et al. 2007). Inorganic

nutrient addition experiments targeting soil and water column specifically, and perhaps at

various levels and ratios, would further create a more nuanced picture with respect to

homeostasis and preferential uptake pathways. Additionally, evaluation of SAV resorption and

resorption specific to other plant parts including plant rhizomes (i.e. plant roots) would also fill

knowledge gaps and allow better understanding of homeostasis due to resource partitioning

within wetland macrophytes. Nevertheless, treatment wetlands are useful systems that can be

used to study stoichiometric homeostasis as they present a wide nutrient gradient with large

variability in nutrient loading from inflow to the outflow.

Conclusions

This study suggests that vegetation within treatment wetlands maintains a relatively constant

uptake efficiencies for nitrogen and phosphorus as indicated by the non-homeostatic relationship

between foliar and active reservoirs.   Thus, tissue nutrient concentrations and nutrient

concentrations of the ambient environment scale with each other to some degree. In this study,

we developed a framework to link nutrient resorption to the homeostasis concept and found that

resorption efforts can alleviate nutrient imbalances in the resource pools, although the resource

pool (water or soil) cannot be clearly identified. However, the strongly non-homeostatic behavior

of EAV with respect to soil resources requires further attention and we speculate that a lowering

of the N:P ratio in the resource pool will lead to storage of P in increasingly plant-inaccessible

compounds.

Acknowledgements

We would like to thank SFWMD and UF Wetland Biogeochemistry Laboratory staff members

for providing the data used in this analysis. We would also like to thank Mark Brenner, Sue

Newman, Odi Villapando, Tom James, Jim Elser and the anonymous peer reviewer(s) and

editor(s) for their efforts and constructive review of this manuscript.

Conflict of Interest Statement

The authors declare that they have no conflict of interest.

Funding

Authors' Contributions

PJ performed data analyses including necessary calculations and statistical analyses and wrote

the manuscript. SG developed the expanded homeostasis framework, contributed to data

interpretation and writing. RKB and TZO were involved with data collection and writing. ALW

assisted with writing. JK was involved in data collection. All authors read and approved the final

manuscript.

References

Aerts R (1996) Nutrient Resorption from Senescing Leaves of Perennials: Are there General
Patterns? Journal of Ecology 84:597–608. doi: 10.2307/2261481

Aerts R, Chapin FS (1999) The mineral nutrition of wild plants revisited: a re-evaluation of
processes and patterns. Advances in ecological research 30:1–67

Boyer MLH, Wheeler BD (1989) Vegetation Patterns in Spring-Fed Calcareous Fens: Calcite
Precipitation and Constraints on Fertility. Journal of Ecology 77:597–609. doi:
10.2307/2260772

Brix H (2003) Plants used in constructed wetlands and their functions. In: Dias V, Vymazel J
(eds) 1st International Seminar on the use of Aquatic Macrophytes for Wastewater
Treatment in Constructed Wetlands. Lisbon, Portugal

Bruland GL, Osborne TZ, Reddy KR, et al (2007) Recent Changes in Soil Total Phosphorus in
the Everglades: Water Conservation Area 3. Environmental Monitoring and Assessment
129:379–395. doi: 10.1007/s10661-006-9371-x

Chapin III FS, Schulze E-D, Mooney HA (1990) The Ecology and Economics of Storage in
Plants. Annual Review of Ecology and Systematics 21:423–447

Chen H, Ivanoff D, Pietro K (2015) Long-term phosphorus removal in the Everglades
stormwater treatment areas of South Florida in the United States. Ecological Engineering
79:158–168. doi: 10.1016/j.ecoleng.2014.12.012

Chimney MJ, Pietro KC (2006) Decomposition of macrophyte litter in a subtropical constructed
wetland in south Florida (USA). Ecological Engineering 27:301–321. doi:
10.1016/j.ecoleng.2006.05.016

Corstanje R, Reddy KR, Prenger JP, et al (2007) Soil microbial eco-physiological response to
nutrient enrichment in a sub-tropical wetland. Ecological Indicators 7:277–289. doi:
10.1016/j.ecolind.2006.02.002

Craft CB, Vymazal J, Richardson CJ (1995) Response of everglades plant communities to
nitrogen and phosphorus additions. Wetlands 15:258–271. doi: 10.1007/BF03160706

De Deyn GB, Cornelissen JHC, Bardgett RD (2008) Plant functional traits and soil carbon
sequestration in contrasting biomes. Ecology Letters 11:516–531. doi: 10.1111/j.1461-
0248.2008.01164.x

DeBusk TA, Kharbanda M, Jackson SD, et al (2011) Water, vegetation and sediment gradients in
submerged aquatic vegetation mesocosms used for low-level phosphorus removal.
Science of The Total Environment 409:5046–5056. doi: 10.1016/j.scitotenv.2011.08.038

Demers JD, Driscoll CT, Fahey TJ, Yavitt JB (2007) Mercury cycling in litter and soil in
different forest types in the adirondack region, new york, usa. Ecological Applications
17:1341–1351. doi: 10.1890/06-1697.1

Dhote S, Dixit S (2009) Water quality improvement through macrophytes—a review.
Environmental Monitoring and Assessment 152:149–153. doi: 10.1007/s10661-008-
0303-9

Dierberg FE, DeBusk TA, Jackson SD, et al (2002) Submerged aquatic vegetation-based
treatment wetlands for removing phosphorus from agricultural runoff: response to
hydraulic and nutrient loading. Water research 36:1409–1422

Dinno A (2015) Dunn's test of multiple comparisons using rank sums. CRAN R-Project

Duarte CM (1992) Nutrient concentration of aquatic plants: Patterns across species. Limnol
Oceanogr 37:882–889. doi: 10.4319/lo.1992.37.4.0882

Elser JJ, Fagan WF, Denno RF, et al (2000) Nutritional constraints in terrestrial and freshwater
food webs. Nature 408:578

Elser JJ, Fagan WF, Kerkhoff AJ, et al (2010) Biological stoichiometry of plant production:
metabolism, scaling and ecological response to global change. New Phytologist 186:593–
608. doi: 10.1111/j.1469-8137.2010.03214.x

Feijoó C, Leggieri L, Ocón C, et al (2014) Stoichiometric homeostasis in the food web of a
chronically nutrient-rich stream. Freshwater Science 33:820–831. doi: 10.1086/677056

Fernández-Aláez M, Fernández-Aláez C, Bécares E (1999) Nutrient content in macrophytes in
Spanish shallow lakes. Hydrobiologia 408:317–326

Filipiak M (2016) Pollen Stoichiometry May Influence Detrital Terrestrial and Aquatic Food
Webs. Front Ecol Evol 4:. doi: 10.3389/fevo.2016.00138

Filipiak M, Weiner J (2014) How to Make a Beetle Out of Wood: Multi-Elemental
Stoichiometry of Wood Decay, Xylophagy and Fungivory. PLOS ONE 9:e115104. doi:
10.1371/journal.pone.0115104

Filipiak M, Weiner J (2017) Nutritional dynamics during the development of xylophagous
beetles related to changes in the stoichiometry of 11 elements: Stoichiometry of
xylophage development. Physiological Entomology 42:73–84. doi: 10.1111/phen.12168

Frost PC, Evans-White MA, Finkel ZV, et al (2005) Are you what you eat? Physiological
constraints on organismal stoichiometry in an elementally imbalanced world. Oikos
109:18–28. doi: 10.1111/j.0030-1299.2005.14049.x

Gu Q, Zamin TJ, Grogan P (2017) Stoichiometric homeostasis: a test to predict tundra vascular
plant species and community-level responses to climate change. Arctic Science 1–14.
doi: 10.1139/as-2016-0032

Güsewell S (2004) N□: P ratios in terrestrial plants: variation and functional significance. New
Phytologist 164:243–266. doi: 10.1111/j.1469-8137.2004.01192.x

Güsewell S, Koerselman W (2002) Variation in nitrogen and phosphorus concentrations of
wetland plants. Perspectives in Plant Ecology, Evolution and Systematics 5:37–61. doi:
10.1078/1433-8319-0000022

Güsewell S, Koerselman W, Verhoeven JTA (2003) Biomass n:p ratios as indicators of nutrient
limitation for plant populations in wetlands. Ecological Applications 13:372–384. doi:
10.1890/1051-0761(2003)013[0372:BNRAIO]2.0.CO;2

Julian P, Gerber S, Bhomia RK, et al (2019) Evaluation of nutrient stoichiometric relationships
among ecosystem compartments of a subtropical treatment wetland. Do we have
"Redfield wetlands"? Ecol Process 8:20. doi: 10.1186/s13717-019-0172-x

Julian P, Gerber S, Wright AL, et al (2017) Carbon pool trends and dynamics within a
subtropical peatland during long-term restoration. Ecol Process 6:43–57. doi:
10.1186/s13717-017-0110-8

Juston J, DeBusk TA (2006) Phosphorus mass load and outflow concentration relationships in
stormwater treatment areas for Everglades restoration. Ecological Engineering 26:206–
223. doi: 10.1016/j.ecoleng.2005.09.011

Kadlec RH, Wallace SD (2009) Treatment wetlands. CRC Press, Boca Raton, FL

Killingbeck KT (1996) Nutrients in Senesced Leaves: Keys to the Search for Potential
Resorption and Resorption Proficiency. Ecology 77:1716–1727. doi: 10.2307/2265777

King J, Zweig C (2016) P Flux Vegetation Assessments-Submerged Aquatic Vegetation (SAV)
Biomass Sampling and Analysis September 2016. South Florida Water Management
District, West Palm Beach, FL

Koerselman W, Meuleman AFM (1996) The Vegetation N:P Ratio: a New Tool to Detect the
Nature of Nutrient Limitation. The Journal of Applied Ecology 33:1441. doi:
10.2307/2404783

Komsta L (2013) Median-Based Linear Models. CRAN R-Project

Makino W, Cotner JB, Sterner RW, Elser JJ (2003) Are Bacteria More Like Plants or Animals?
Growth Rate and Resource Dependence of Bacterial C: N: P Stoichiometry. Functional
Ecology 17:121–130

McGroddy ME, Daufresne T, Hedin LO (2004) Scaling of C:N:P stoichiometry in forests
worldwide: implications of terrestrial redfield-type ratios. Ecology 85:2390–2401. doi:
10.1890/03-0351

Menge DNL, Pacala SW, Hedin LO (2009) Emergence and Maintenance of Nutrient Limitation
over Multiple Timescales in Terrestrial Ecosystems. The American Naturalist 173:164–
175. doi: 10.1086/595749

Miao S (2004) Rhizome growth and nutrient resorption: mechanisms underlying the replacement
of two clonal species in Florida Everglades. Aquatic Botany 78:55–66. doi:
10.1016/j.aquabot.2003.09.001

Miao SL, Zou CB (2012) Effects of inundation on growth and nutrient allocation of six major macrophytes in the Florida Everglades. Ecological Engineering 42:10–18. doi: 10.1016/j.ecoleng.2012.01.009

Newman S, Osborne TZ, Hagerthey SE, et al (2017) Drivers of landscape evolution: multiple regimes and their influence on carbon sequestration in a sub-tropical peatland. Ecol Monogr 87:578–599. doi: 10.1002/ecm.1269

Osborne TZ, Bruland GL, Newman S, et al (2011) Spatial distributions and eco-partitioning of soil biogeochemical properties in the Everglades National Park. Environmental Monitoring and Assessment 183:395–408. doi: 10.1007/s10661-011-1928-7

Peng H, Chen Y, Yan Z, Han W (2016) Stage-dependent stoichiometric homeostasis and responses of nutrient resorption in Amaranthus mangostanus to nitrogen and phosphorus addition. Scientific Reports 6:. doi: 10.1038/srep37219

Persson J, Fink P, Goto A, et al (2010) To be or not to be what you eat: regulation of stoichiometric homeostasis among autotrophs and heterotrophs. Oikos 119:741–751. doi: 10.1111/j.1600-0706.2009.18545.x

Rattray MR, Howard-Williams C, Brown JMA (1991) Sediment and water as sources of nitrogen and phosphorus for submerged rooted aquatic macrophytes. Aquatic Botany 40:225–237. doi: 10.1016/0304-3770(91)90060-I

Reddy KR, DeLaune RD (2008) Biogeochemistry of wetlands: science and applications. CRC Press, Boca Raton, FL

Reddy KR, Kadlec RH, Flaig E, Gale PM (1999) Phosphorus Retention in Streams and Wetlands: A Review. Critical Reviews in Environmental Science and Technology 29:83–146. doi: 10.1080/10643389991259182

Redfield AC (1958) The biological control of chemical factors in the environment. American Scientist 46:230A–221

Reed SC, Townsend AR, Davidson EA, Cleveland CC (2012) Stoichiometric patterns in foliar nutrient resorption across multiple scales. New Phytologist 196:173–180. doi: 10.1111/j.1469-8137.2012.04249.x

Rejmánková E (2005) Nutrient resorption in wetland macrophytes: comparison across several regions of different nutrient status. New Phytologist 167:471–482. doi: 10.1111/j.1469-8137.2005.01449.x

Rejmánková E, Macek P, Epps K (2008) Wetland ecosystem changes after three years of phosphorus addition. Wetlands 28:914–927. doi: 10.1672/07-150.1

Rejmánková E, Snyder JM (2008) Emergent macrophytes in phosphorus limited marshes: do phosphorus usage strategies change after nutrient addition? Plant Soil 313:141–153. doi: 10.1007/s11104-008-9687-0

632   Richardson CJ, Marshall PE (1986) Processes Controlling Movement, Storage, and Export of
633   Phosphorus in a Fen Peatland. Ecological Monographs 56:280–302. doi:
634   10.2307/1942548

635   Rivas-Ubach A, Sardans J, Pérez-Trujillo M, et al (2012) Strong relationship between elemental
636   stoichiometry and metabolome in plants. PNAS 109:4181–4186. doi:
637   10.1073/pnas.1116092109

638   Sardans J, Rivas-Ubach A, Peñuelas J (2012) The C:N:P stoichiometry of organisms and
639   ecosystems in a changing world: A review and perspectives. Perspectives in Plant
640   Ecology, Evolution and Systematics 14:33–47. doi: 10.1016/j.ppees.2011.08.002

641   Shaver GR, Melillo JM (1984) Nutrient Budgets of Marsh Plants: Efficiency Concepts and
642   Relation to Availability. Ecology 65:1491–1510. doi: 10.2307/1939129

643   Sistla SA, Appling AP, Lewandowska AM, et al (2015) Stoichiometric flexibility in response to
644   fertilization along gradients of environmental and organismal nutrient richness. Oikos
645   124:949–959. doi: 10.1111/oik.02385

646   Sistla SA, Schimel JP (2012) Stoichiometric flexibility as a regulator of carbon and nutrient
647   cycling in terrestrial ecosystems under change. The New Phytologist 196:68–78

648   Small GE, Helton AM, Kazanci C (2009) Can consumer stoichiometric regulation control
649   nutrient spiraling in streams? Journal of the North American Benthological Society
650   28:747–765. doi: 10.1899/08-099.1

651   Smalley GW, Coats DW, Stoecker DK (2003) Feeding in the mixotrophic dinoflagellate
652   Ceratium furca is influenced by intracellular nutrient concentrations. Marine Ecology
653   Progress Series 262:137–151. doi: 10.3354/meps262137

654   South Florida Water Management District (2013) Restoration Strategies Regional Water Quality
655   Plan: Science Plan for the Everglades Stormwater Treatment Areas. South Florida Water
656   Management District, West Palm Beach, FL

657   Sterner, Clasen, Lampert, Weisse (1998) Carbon:phosphorus stoichiometry and food chain
658   production. Ecology Letters 1:146–150. doi: 10.1046/j.1461-0248.1998.00030.x

659   Sterner RW, Elser JJ (2002) Ecological Stoichiometry: The Biology of Elements from Molecules
660   to the Biosphere. Princeton University Press

661   UF-WBL (2017) Evaluation of Soil Biogeochemical Properties Influencing Phosphorus Flux in
662   the Everglades Stormwater Treatment areas: 2016-2017 Annual Report. University of
663   Florida, Gainesville, FL

664   Van Heerwaarden LM, Toet S, Aerts R (2003) Current measures of nutrient resorption efficiency
665   lead to a substantial underestimation of real resorption efficiency: facts and solutions.
666   Oikos 101:664–669. doi: 10.1034/j.1600-0706.2003.12351.x

Villapando O, King J (2018) Appendix 5C-3: Evaluation of Phosphorus Sources, Forms, Flux,
and Transformation Processes in the Stormwater Treatment Areas. In: 2018 South Florida
Environmental Report. South Florida Water Management District, West Palm Beach, FL

Wright AL, Reddy KR (2001) Phosphorus loading effects on extracellular enzyme activity in
Everglades wetland soils. Soil Science Society of America Journal 65:588–595

Yu Q, Elser JJ, He N, et al (2011) Stoichiometric homeostasis of vascular plants in the Inner
Mongolia grassland. Oecologia 166:1–10. doi: 10.1007/s00442-010-1902-z

Zar JH (2010) Biostatistical Analysis, 15th edn. Prentice Hall

Figures and Tables

Figure 1. Surface water, soil and vegetation monitoring locations within Everglades Stormwater
Treatment Area-2 cells 1 (right) and 3 (left). Cell 1 is predominately emergent vegetation and
Cell 3 is predominately submerged aquatic vegetation. Operationally these cells are identified as
flow-way 1 and 3, respectively.

Figure 2. Distribution of plant foliar stoichiometric relationships within flow-way (FW) 1 and
FW 3 of Stormwater Treatment Area-2. Left and middle panels are histogram representations of
each respective stoichiometric relationship. The right panel is a cumulative distribution of
stoichiometric relationship for each FW.

Figure 3. Resource (Top: Soil and Bottom: Surface Water) and vegetation nitrogen to
phosphorus molar ratio (N:P) with Theil-Sen single median linear model $\pm$ 95% confidence
interval for each vegetation type. Note both axes are log-scale.

Figure 4. Homeostasis coefficient ($1/H_T$) and 95% confidence interval with respect to soil and
surface water resources for emergent aquatic vegetation and submerged aquatic vegetation (EAV
and SAV, respectively) vegetation communities and flow-way. Dashed line indicates the
homeostasis/non-homeostasis threshold (0.5) and the theoretical maximum $1 / H_T$ (i.e. 1.0).

Figure 5. Strip plot of (A) total phosphorus resorption efficiency (RE_{TP}) and (B) total nitrogen
resorption efficiency (RE_{TN}) of cattail within Stormwater Treatment Area-2 flow-way 1. Raw
data are represented by grey points, while mean $\pm$ standard error are represented by black points.

Figure 6. Median total homeostasis coefficient ($1/H_T$) for emergent aquatic vegetation (EAV;
FW1) with effort associated with maintaining total (H_T; red arrows), resorptive (H_{RE}; blue
arrows) and uptake (H_u; black arrows). A downward arrow indicates the effort into maintaining
homeostasis, representing the 1-1/H expression in Eq. 5.

Table 1. Summary statistics for variables and matrices used in this study of samples collected along the flow-way (FW) 1 and 3 transect within Stormwater Treatment Area-2. Summary statistics include mean, standard error (SE), sample size (N), range, and coefficient of variation (CV). Matrices include surface water, recently accreted soil and live aboveground biomass of sampled vegetation Note: Dissolved Organic Carbon = DOC; Total Phosphorus = TP; Total Nitrogen = TN; Total Carbon - TC; Loss-on-ignition = LOI; Calcium = Ca.

		Flow-way 1			Flow-way 3		
Matrix	Parameter (Units)	Mean ± SE (N)	Range	CV	Mean ± SE (N)	Range	CV
Surface Water	DOC (mM)	2.23 ± 0.05 (109)	1.26 – 3.31	23	2.57 ± 0.02 (177)	1.54 – 3.33	10
	TP (mM)	0.0021 ± 0.0002 (104)	0.0003 – 0.012	116	0.0011 ± 0.00008 (169)	0.0002 – 0.0094	91
	TN (mM)	0.12 ± 0.003 (107)	0.06 – 0.22	26	0.15 ± 0.001 (177)	0.11 – 0.30	14
	DOC:TP (molar)	2,649 ± 220 (95)	216 – 9,532	81	3,753 ± 187 (169)	290 – 1,461	65
	DOC:TN (molar)	18.9 ± 0.21 (101)	12.8 – 24.0	11	16.8 ± 0.12 (177)	9.30 – 19.1	10
	TN:TP (molar)	137.6 ± 11.3 (93)	15.5 – 410.3	80	216.0 ± 9.89 (169)	31.2 – 788.7	90
Soil	LOI (%)	79.5 ± 2.2 (27)	18.6 – 91.0	14	54.7 ± 4.7 (25)	20.3 – 83.9	43
	TC (mM kg^{-1})	36,051 ± 963 (27)	24,145 – 41,962	14	29,423 ± 1,895 (25)	15,320 – 40,797	32
	TP (mM kg^{-1})	27.6 ± 2.1 (27)	11.1 – 43.8	40	15.8 ± 1.05 (25)	11.6 – 29.0	33
	TN (mM kg^{-1})	2,037 ± 52 (27)	1,421 – 2,413	13	1,451 ± 110.7 (25)	368.3 – 2,213	38
	TC:TP (molar)	1,669 ± 190 (27)	671 – 3,511	59	2,104 ± 192.2 (25)	557.1 – 3,520	46
	TC:TN (molar)	17.7 ± 0.3 (27)	14.9 – 21.1	7	20.9 ± 0.4 (25)	17.9 – 24.0	9
	TN:TP (molar)	92.2 ± 9.7 (27)	39.5 – 192.4	55	104.7 ±10.6 (25)	23.5 – 187.3	51
	Ca (mM kg^{-1})	1,737 ± 206 (27)	649 – 4,465	62	3,987 ± 472.0 (25)	978.7 – 7,637	59
Vegetation	TC (mM kg^{-1})	35,347 ± 152 (39)	32,305 – 38,174	3	22,583 ± 711.3 (33)	15,486 – 29,932	18
	TP (mM kg^{-1})	43.8 ± 3.4 (39)	11.6 – 97.4	49	26.3 ± 5.9 (33)	6.78 – 109.1	72
	TN (mM kg^{-1})	672.5 ± 34.9 (39)	435.5 – 1,756	32	1,606 ± 137.7 (33)	671.1 – 3,476.9	49
	TC:TP (molar)	1,055 ± 96.4 (39)	361.5 – 3,110	57	845.3 ± 116.3 (33)	237.4 – 2,486.7	79
	TC:TN (molar)	56.0 ± 2.0 (39)	19.3 – 82.8	22	16.1 ± 0.84 (33)	7.50 – 23.7	30
	TN:TP (molar)	18.2 ± 1.2 (39)	7.67 – 37.6	41	48.1 ± 4.81 (33)	21.3 – 120.4	57

Table 2. Overall log-log median based model statistics of vegetation nitrogen to phosphorus (N:P) versus soil and surface water resource pools between emergent and submerged aquatic vegetation (EAV and SAV, respectively) communities within Stormwater Treatment Area 2 Flow-way 1 and Flow-way 3, respectively.

Resource	Community	Slope $(1/H_T)$	Intercept (c-value)	R^2	Median Absolute Deviation	V-statistic	df	ρ-value
Soil	EAV	1.62	-3.70	0.61	1.50	168032	37	<0.01
	SAV	0.66	0.84	0.72	0.47	19947	22	<0.01
Water Column	EAV	0.46	0.70	0.60	0.34	116683	37	<0.01
	SAV	0.51	0.93	0.17	0.94	23162	26	<0.01

Table 3. Resorption efficiency summary statistics for emergent aquatic vegetation species collected within Stormwater Treatment Area 2, FW 1. Total Phosphorus Resorption efficiency (RE_{TP}) calculated with equation 1, using both living and senescent living aboveground biomass. Summary statistics include mean, standard error, range, coefficient of variation and sample size (N).

Species	Total Phosphorus				Total Nitrogen			
	Mean ± SE (%)	Range (%)	Coefficient of Variation	N	Mean ± SE (%)	Range (%)	Coefficient of Variation	N
Typha domingensis Pers. (cattail)	74 ± 2	24 – 91	20.1	36	33 ± 2	1.8 - 56	39.8	29
Nymphaea odorata Aiton (water lily)	70	---	---	1	---	---	---	---
Cladium jamaicense Crantz (sawgrass)	77	---	---	1	2.3	---	---	1