Title: Subtle perturbations of the maize methylome reveal genes and transposons silenced by

DNA methylation

Authors: Sarah N. Anderson[1], Greg Zynda[2], Jawon Song[2], Zhaoxue Han[3], Matthew Vaughn[2],

Qing Li[4]*, Nathan M. Springer[1]*

*Co-corresponding authors

[1] Department of Plant and Microbial Biology; University of Minnesota; St. Paul, MN, 55108, USA

[2] Texas Advanced Computing Center, University of Texas; Austin TX 78758, USA

[3] State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University,

Yangling 712100, Shaanxi, China

[4] National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University,

Wuhan, Hubei, 430070, China

Corresponding authors:

Nathan M. Springer

140 Gortner Laboratory

1479 Gortner Ave.

St. Paul, MN, 55108

612-624-6241

springer@umn.edu

Qing Li

National Key Laboratory of Crop Genetic Improvement

Huazhong Agricultural University

No.1 Shizishan Street, Hongshan District

Wuhan, Hubei, 430070, China

qingli@mail.hzau.edu.cn

34   ABSTRACT

DNA methylation is a chromatin modification that can provide epigenetic regulation of gene and

transposon expression. Plants utilize several pathways to establish and maintain DNA

methylation in specific sequence contexts. The chromomethylase (CMT) genes maintain CHG

(where H = A, C or T) methylation. The RNA-directed DNA methylation (RdDM) pathway is

important for CHH methylation. Transcriptome analysis was performed in a collection of *Zea*

mays lines carrying mutant alleles for CMT or RdDM-associated genes. While the majority of

the transcriptome was not affected, we identified sets of genes and transposon families

sensitive to context-specific decreases in DNA methylation in mutant lines. Many of the genes

that are up-regulated in CMT mutant lines have high levels of CHG methylation, while genes

that are differentially expressed in RdDM mutants are enriched for having nearby mCHH

islands, providing evidence that context-specific DNA methylation directly regulates expression

of a small number of genes. The analysis of a diverse set of inbred lines revealed that many

genes regulated by CMTs exhibit natural variation for DNA methylation and gene expression.

Transposon families with differential expression in the mutant genotypes show few defining

features, though several families up-regulated in RdDM mutants show enriched expression in

endosperm, highlighting the importance for this pathway during reproduction. Taken together,

our findings suggest that while the number of genes and transposon families whose expression

is reproducibly affected by mild perturbations in context-specific methylation is small, there are

distinct patterns for loci impacted by RdDM and CMT mutants.

55   INTRODUCTION

The epigenome describes the potential for additional heritable information that can be passed

on through mitosis or meiosis (Hofmeister *et al.* 2017). DNA methylation is one molecular

mechanism that can provide epigenetic information. There is interest in the potential for cryptic

information present in genomes that is normally silenced by epigenetic mechanisms but could

be activated through epigenetic changes without requiring any genetic change.

Much is known about the mechanisms that control DNA methylation and the functional roles of

DNA methylation in regulating transposon and gene expression in the model plant *Arabidopsis*

thaliana. However, our knowledge of the regulating mechanisms and function of DNA

methylation is much more limited in crop plants. Evidence in rice and maize suggests that major

perturbations of DNA methylation disrupt development and the seeds/plants are not viable

(Yamauchi *et al.* 2014; Hu *et al.* 2014; Li *et al.* 2014). Forward genetic screens for factors

involved in epigenetic phenomena such as paramutation (Dorweiler *et al.* 2000; Hollick *et al.*

2005; Alleman *et al.* 2006; Hale *et al.* 2007; Jr *et al.* 2009) or transgene silencing (McGinnis *et*

al. 2006) have identified several genes that are associated with DNA methylation or chromatin

in maize (Hollick 2017). In addition, reverse genetic approaches have been utilized in attempts

to document the function of putative methyltransferase genes or other genes associated with

DNA methylation (Papa *et al.* 2001; Makarevitch *et al.* 2007; Li *et al.* 2014). To date, these

mutants have provided partial reductions in DNA methylation in specific sequence contexts but

no mutants with drastic reductions to genomic DNA methylation have been recovered in maize.

Surveys of natural variation for DNA methylation among diverse lines of maize have revealed

many examples of differentially methylated regions (DMRs) (Eichten *et al.* 2011, 2013; Regulski

et al. 2013; Li *et al.* 2015a). A subset of the genes located near DMRs exhibit a negative

correlation between DNA methylation and gene expression (Eichten *et al.* 2013; Li *et al.* 2015a).

This is primarily found at genes that have CG or CHG methylation in regions surrounding the

transcriptional start site (TSS) and show qualitative (on/off) expression variation among

genotypes (Li *et al.* 2015a). This suggests the potential for cryptic information in the maize

genome that is epigenetically silenced in some lines but can be active due to epigenetic

changes in other genotypes.

Several maize mutant lines with subtle perturbations of genomic DNA methylation have been

previously identified (Li *et al.* 2014). The mutants include *mop1* and *mop3*, two mutants

recovered in screens for factors required for maintenance of the paramutated state at the B'

locus (Dorweiler *et al.* 2000). The *Mop1* gene encodes an RNA-dependent RNA polymerase

related to RDR2 in *Arabidopsis* (Alleman *et al.* 2006) while *Mop3* is predicted to encode the

second largest subunit of RNA Pol IV (Sloan *et al.* 2014), which is allelic to *rmr6 (Erhard et al.*

2009). Mutant alleles for the two chromomethylase genes present in the maize genome, *Zmet2*

and *Zmet5* also influence context-specific DNA methylation patterns (Papa *et al.* 2001;

Makarevitch *et al.* 2007; Li *et al.* 2014). These genes are likely paralogs resulting from a whole

genome duplication event and are orthologous to *CMT3* from *Arabidopsis thaliana*. Previous

research has found that *mop1* and *mop3* genotypes have lost CHH methylation at many

genomic regions with elevated CHH, and there are changes in CG and CHG at these sites as

well (Li *et al.* 2014). However, as these types of regions are quite rare in the maize genome

these mutants have minimal effects on genome-wide levels of CG and CHG methylation. The

zmet2-m1 mutant and, to a lesser extent, the *zmet5-m1* mutant, result in reduction of CHG

methylation. These mutants also cause reductions of CWA methylation (where W is A or T) in

genomic regions with low, but detectable, CWA methylation (Li *et al.* 2014; Gouil and

Baulcombe 2016). Attempts to recover double mutants for *Zmet2/Zmet5* were unsuccessful,

suggesting at least partially redundant function for these paralogous genes.

Mutants for *mop1*, *mop3*, *zmet2* and *zmet5* are viable with relatively few major phenotypic

changes (Dorweiler *et al.* 2000; Papa *et al.* 2001). The *mop1* and/or *mop3* mutations have been

shown to play important roles in the regulation of specific maize loci (Dorweiler *et al.* 2000;

Alleman *et al.* 2006; Sloan *et al.* 2014), transgenes (McGinnis *et al.* 2006) or transposable

elements (Lisch *et al.* 2002; Woodhouse *et al.* 2006). Microarray profiling of gene expression

has revealed evidence for altered expression of small sets of genes in studies of *mop1*

(Madzima et al. 2014) and *zmet2 (Makarevitch et al. 2007)*. There is also evidence from an

RNAseq experiment for altered regulation of transposable element expression in apical

meristem tissue (Jia *et al.* 2009). Transcriptome analysis of *rmr6*, which is allelic to *mop3*,

provided evidence for a potential role in stress response (Forestan *et al.* 2016). The *rmr6*

mutation appears to increase the proportion of the genome that is transcribed but has subtle

effects at most loci with relatively few genes with significant changes in expression level

(Forestan *et al.* 2017). However, there have not been comprehensive studies on the overlap of

genes or transposons that are sensitive to mutations in different CMT or RdDM genes in maize.

Each of the mutant backgrounds used for this study has subtle effects on genomic methylation

levels and can produce viable plants. There are several phenotypic abnormalities observed in

mop1 and *mop3* stocks (Dorweiler *et al.* 2000; Barber *et al.* 2012; Sloan *et al.* 2014) although

the penetrance in multiple backgrounds has not been well characterized. We sought to

determine if the subtle changes in DNA methylation in these mutants would reveal genes or

transposons that are sensitive to these shifts in DNA methylation or chromatin. A limited number

of genes and transposon families exhibit altered expression in these genotypes. A subset of

these genes have high levels of DNA methylation in wild-type that are reduced in the mutant

genotypes. Many of these genes exhibit natural variation for DNA methylation and gene

expression. This provides evidence that the natural variation at these genes is due to epigenetic

rather than genetic variation and highlights cryptic information present in the maize genome that

could be accessible through alterations to the epigenome.

135   MATERIALS AND METHODS

Biological materials*: All the mutant and wild type samples used in this study are listed in Table

S1 and SRA accession numbers are listed for each dataset. Tissue for RNA and DNA isolations

was collected from three biological replicates. Plants were grown in standard greenhouse

conditions for 20 days to reach the V3 stage. The 2^(nd) and 3^(rd) leaves were collected individually

for each seedling. The 2^(nd) leaf was used to isolate DNA for genotyping, and the 3^(rd) leaf was used

for RNA isolation and sequencing. For each biological replicate, 4-6 seedlings were pooled.

Library preparation and sequencing*: Total RNA was isolated using the TRIZol reagent

following the manufacturer's protocol. RNA was quantified using RiboGreen and 3 µg total RNA

was used to construct libraries using TruSeq strand-specific kit (Illumina) following

manufacturer's suggestions. The final library was quantified using PicoGreen and twelve

libraries were pooled per Illumina lane. Library quality was checked using Agilent Bioanalyzer.

Sequencing was performed on HiSeq2500 using 2 x 50 bp mode.

Gene expression analysis*: Trim_glore was used to trim low-quality base from the 3' end of the

reads, as well as to remove adapters. Reads that passed quality control were mapped to B73

version 4 genome (Jiao *et al.* 2017) using Tophat2 (Kim *et al.* 2013), allowing at most 1

mismatch (-N 1) and the expected inner distance between mate pairs of 200 bp (-r 200). Reads

that are properly paired and uniquely mapped were filtered out using samtools (-f 0x0002 –q

50). HTSeq (Anders *et al.* 2015) was used to summarize the number of reads mapped to each

V4 gene model with the union mode, generating a matrix of count values for each gene in each

genotype.

Raw read counts were input into DEseq2 (Love *et al.* 2014) to perform differential expression

analysis. Pair-wise comparisons were made between each mutant and the appropriate wild

type. Genes with a FDR value of < 0.05 and $\log_2$(FoldChange) > 1 were called differentially

expressed genes. Detailed analysis was restricted to genes with consistent DE calls in at least

two mutant contrasts in the same pathway (RdDM or CMT). Genes were considered expressed

if at least 3 replicates in the libraries described had an RPM (reads per million) value > 1.

TE expression analysis: B73v4 (Jiao *et al.* 2017) TE annotation was modified to remove

helitrons and the file was resolved using RTrackLayer in R so that each base of the genome

was assigned to only a single TE. Exon regions were masked from the TE file using Bedtools

(Quinlan and Hall 2010) subtract. Gene annotations were added to this modified TE annotation

file, and mapped reads were assigned to features using HTSeq (Anders *et al.* 2015). A custom

script was used to read through the HTSeq sam output, assigning unique-mapping reads to

individual TE elements and multi-mapped reads to TE families if mapped positions hit only a

single TE family. Unique and multi-mapped reads were combined for per-family expression

counts, and RPM values were calculated by normalizing to the number of gene reads plus TE

family reads in each library. All reads mapped to gene annotations plus TE annotations were

excluded from the TE expression analysis. Differentially expressed TE families were determined

using DESeq2 (Love *et al.* 2014) using a log_2(FoldChange) cutoff of 1 and FDR adjusted p-value

cutoff of 0.05. Detailed analysis was restricted to TE families with consistent DE calls in at least

two mutant contrasts in the same pathway (RdDM or CMT).

WGBS data analysis: The WGBS datasets used in this study are detailed in Table S2 and SRA

accession numbers for each sample are provided. One µg DNA was used to prepare libraries

for whole-genome bisulfite sequencing using the KAPA library preparation kit. DNA was

sheared to a peak between 200-250 bp. End repair was performed to make blunt-ended

fragments, followed by adding base A to the 3' end, and adapter ligation. Size selection was

performed to enrich library with a size between 250-450 bp. Bisulfite conversion was then

carried out using Zymo DNA methylation lightning kit according to user's manual. Finally, library

was enriched using PCR amplification. Library quality was checked using the Agilent

Bioanalyzer. Library quantification was performed with qPCR before sequencing. Sequencing

was performed on HiSeq2000 with paired end 100 cycles.

Analysis was performed as previously described (Li *et al.* 2015a; Song *et al.* 2016). Read quality

was checked with FASTQC, adapters and low-quality bases at the 3' end of each read were

trimmed using Trim_glore. The high quality reads were mapped to B73 V4 genome (Jiao *et al.*

2017) using BSMAP (Xi and Li 2009) allowing at most 5 mismatches. Only properly paired

reads with unique mappings were kept and used for calling DNA methylation. Methylation calls

were performed using the methratio.py script from BSMAP. Finally, DNA methylation in each

context (CG, CHG, CHH) was summarized for each 100-bp non-overlapping tile of the 10 maize

chromosomes.

DMR calling: DMRs were called using previously described criteria (Li *et al.* 2015a). Briefly,

each 100-bp tile with > 6 CG/CHG sites, > 2X coverage and > 60% difference for CG/CHG were

compared. CHH DMRs were called using the same coverage and site number criteria, but with

a requirement for <5% CHH in one genotype and >25% CHH in another genotype, reflecting the

low level of CHH methylation in the maize genome.

mCHH Islands: High CHH bins were called genome-wide by requiring CHH methylation over

25% with at least 10 informative counts per bin. Genes and TEs were considered associated

with a mCHH island if when at least one high methylation bin was identified within the gene or in

the 2 kb region surrounding the gene.

Data availability: All data used in this study are deposited at the NCBI Sequence Read Archive

(SRA). Accession numbers for all libraries are listed in Table S1.

                                    RESULTS

215   **Alteration of gene and transposon expression in maize mutants with perturbed**

216   **methylomes:**

RNAseq was used to perform transcriptome profiling for several maize lines carrying mutations

in genes encoding CMT (this study) or RdDM components (Gent *et al.* 2014; Li *et al.* 2015b).

Together these factors are expected to be responsible for the majority of CHG and CHH

methylation in the maize genome. For CMT genes, three biological replicates of seedling leaf

tissue were profiled for mutations in two different genes, with multiple alleles utilized for one of

the genes (Table S1). In addition, for the RdDM genes we analyzed seedling leaf tissue for

mop1 and *mop3* (Li *et al.* 2015b), along with immature ear tissue for *mop1* (Gent *et al.* 2014).

The genetic background, read number and accession information for each sample is provided in

Table S1.

The expression of individual genes was estimated from the RNAseq data for each sample.

Differentially expressed (DE) genes in each mutant line (relative to the appropriate control) were

identified using DESeq2 followed by a requirement for a minimum of 2 fold-change and an FDR

value of less than 0.05 (Table S2). The observed differences in gene expression in the mutant

lines could be direct effects of the mutation on expression, indirect effects caused by direct

targets, or could be the result of introgressions of linked loci that contain cis-regulatory variation.

The number of genes in each 2 Mb bin with differential expression was assessed throughout the

genome (Figure S1). For mutations that were identified in one background and then

backcrossed into another background (*zmet2-m1, zmet2-m2, zmet5, mop1*), there were often a

cluster of DE genes surrounding the locus of the mutation itself. These regions often included

similar numbers of up- and down-regulated genes. For the other mutation (*mop3*) that was not

backcrossed into another genetic background, there is less evidence for expression changes at

linked genes (Figure S1). Based on these results we omitted DE genes located within 40 Mb of

the mutation in subsequent analyses.

A principle component analysis was performed using all DE genes to cluster samples used in

this study (Figure 1). This reveals that the major sources of differences are tissue and genetic

background, as demonstrated, respectively, by the relatively larger differences for *mop1-ear* and

mop3 which was not introgressed into B73 (Figure 1A). When comparing only leaf libraries in

the B73 background, few genotypes were substantially different from the wild-type controls,

suggesting limited changes to transcript levels induced by each mutation (Figure 1B). The

number of differentially expressed genes in each mutant genotype was highly variable (Figure

1C). In most cases the homozygous mutant individuals exhibit more up-regulated genes than

down-regulated genes, which is compatible with the concept that the CMT and RdDM genes

normally provide silencing activities.

Transposable elements (TEs) comprise a large portion of the maize genome, and typically have

high levels of CG and CHG methylation, with CHH methylation peaks at the edges of some TE

families. There are two classes of TEs, Class I (retrotransposons) and Class II (DNA

transposons), which transpose either through a copy-and-paste mechanism requiring an RNA

intermediate (retrotransposons) or through a cut-and-paste mechanism (DNA transposons)

(Wicker *et al.* 2007). Within each class are several orders divided into superfamilies,

distinguished by structural and protein-coding features. Families within each superfamily are

defined by sequence identity, and each family can contain any number of individual TE

elements (Jiao *et al.* 2017). Individual TE elements are defined at a single location within a

genome and are associated with a family, superfamily and class. We sought to document how

minor perturbations to the methylome impacted expression of TEs. Due to the highly repetitive

nature of TE sequences, we assessed per-family levels of expression by mapping RNA-seq

reads to the genome, reporting up to 20 best hits for each read using Tophat2. Per-family read

counts were determined by summing unique mapping reads (to a specific element) and multi-

mapping reads that align to only a single TE family. Overall, the total portion of RNAseq reads

that map to TE families is not significantly higher in the mutants than in wild-type plants

suggesting a lack of genome-wide activation of TEs in these mutants (Figure S2). In order to

assess expression of individual TE families in each genotype, per-family expression was

normalized by dividing the family counts by the total number of reads in the library assigned to

either TE families or genes, generating an RPM estimate. Using this approach we were able to

detect expression of 1,694 TE families in at least one of the genotypes used for this study. A

relatively small number of DE TE families ($\log_2$FC > 1, FDR < 0.05) were identified in each

mutant (Table S3; Figure 1D). Consistent with the role of DNA methylation in silencing TEs,

more families were identified as up-regulated rather than down-regulated in mutants compared

with WT controls. However, the majority of the TE families expressed in these libraries do not

exhibit significant changes in expression level in CMT or RdDM mutants in maize.

There is a significant overlap in the number of genes and TEs that exhibit consistent changes in

gene expression in at least two samples of CMT mutants (*zmet2-m1/zmet2-m2/zmet5*) or RdDM

mutants (*mop1/mop3*) (Figure S3). In order to understand the reproducible effects of these

pathways on expression, we focused our analyses on the set of 237 genes and 104 TE families

that exhibit consistent up- or down-regulation in multiple CMT or RdDM mutants. Hierarchical

clustering of the expression level of these genes or TE families in all samples reveals evidence

for many consistent changes in expression within a pathway (Figure 1E-F). Within CMT

mutants, many genes show shared expression changes between *zmet2-m1* and *zmet2-m2*,

though there is a smaller subset of genes primarily shared between *zmet2-m2* and *zmet5*

(Figure 1E). Although both *zmet2-m1* and *zmet2-m2* are predicted to encode loss of function

alleles, there are some genetic differences in the behavior of these alleles (Papa *et al.* 2001;

Makarevitch *et al.* 2007; Li *et al.* 2014). The *zmet2-m1* mutation exhibits partial dominance that

may reflect dominant negative action of the protein that could be produced from this allele (Papa

et al. 2001). Plants that are homozygous for *zmet2-m1* have the greatest loss of CHG

methylation and this could result from influence of the *ZMET2* protein product on functional

ZMET5 protein. The *mop1* and *mop3* seedling leaf samples have a number of examples of

consistent up-regulation but fewer examples of consistent down-regulation, consistent with the

greater number of up-regulated than down-regulated genes in RdDM mutants in general.

Some genes that are up-regulated in CMT mutants exhibit high CHG methylation levels:

The differential expression observed in each mutant background could result from direct

changes in DNA methylation or chromatin at these loci or could result from indirect effects due

to secondary effects from genes that are direct targets. Given that many of these mutants are

expected to affect DNA methylation levels, we might expect that wild-type plants would contain

high levels of DNA methylation for genes that exhibit increased expression in the mutants. The

context-specific DNA methylation profiles were assessed in wild-type B73 for genes that were

up- or down- regulated compared with all expressed genes (Figure 2A). Genes that are

differentially expressed in RdDM mutants exhibit slightly higher levels of CHH methylation

upstream of the transcription start sites relative to all expressed genes. The genes that are

down-regulated in CMT mutants do not show unusual patterns of DNA methylation. In contrast,

genes that are up-regulated in the CMT mutants exhibit distinct patterns of CG and CHG

methylation within gene bodies relative to other expressed genes (Figure 2A). Among the 112

genes up-regulated in CMT mutants, approximately half have high (>50%) and half have low

(<20%) methylation in the CG and CHG contexts (Figure 2B-C). In contrast, only ~4% of all

expressed genes have high CHG methylation in the same region. While a small number of

genes with high methylation overlap annotated TEs, most of the genes in this subset do not,

suggesting that this genic methylation is not solely due to nearby TEs. In wild-type samples,

genes with high CHG methylation in the gene body also have high CG methylation (Figure 2D).

However in *zmet2-m1* mutants, CHG methylation for these genes is reduced, with few examples

of a corresponding reduction in CG methylation (Figure 2D-E). As in the examples

(*Zm00001d045627* and *Zm00001d021982*) shown in Figure 2F, these genes have high levels of

CG and CHG methylation in wild-type B73, and the reduction in CHG methylation in *zmet2-m1*

mutants is associated with increased expression, suggesting that CMT-dependent silencing of

these genes depends on CHG but not CG methylation.

Genes that are up-regulated in RdDM mutants are enriched for being near mCHH islands:

A large number of maize genes (~60%) have been associated with the presence of a region of

elevated CHH in the promoter region, termed a mCHH island (Gent *et al.* 2013; Li *et al.* 2015b).

Genes with mCHH islands are enriched for high expression and the mCHH island often occurs

at the edge of the TE nearest these genes (Li *et al.* 2015b). These mCHH islands may form

important boundaries that could protect TE heterochromatin from the influence of genes (Li *et*

al. 2015b) and may also be important for long-distance interactions (Rowley *et al.* 2017). The

methylation within these mCHH islands requires *mop1* and *mop3* (Li *et al. 2014, 2015b)*. We

sought to determine if the genes that exhibit altered expression in *mop1* and *mop3* are enriched

for the presence of mCHH islands. Genes that are up- or down-regulated in the RdDM mutants

are enriched for the presence of mCHH islands, but this is only significant for the up-regulated

genes with 87.8% having a mCHH island within 2kb of the gene, compared to 64.6% of all

expressed genes (Figure 2G). The fact that both RdDM up- and down-regulated genes are often

near mCHH islands could be due to the fact that the mCHH island may provide long-range

interactions (Rowley *et al.* 2017) that could have either positive or negative influences on gene

expression.

In some cases, the mCHH island itself may result in transcriptional regulation. Work in

Arabidopsis has noted a positive feedback loop involving DNA methylation levels and

expression of the demethylase enzyme *ROS1* such that reduced levels of DNA methylation

result in lower *Ros1* expression but increased methylation is associated with elevated *Ros1*

expression (Williams *et al.* 2015). Reduced expression of maize DNA glycosylases has also

been observed in several transcriptome datasets of maize RdDM mutants (Williams *et al.* 2015;

Erhard *et al.* 2015). We find that one maize gene with sequence homology to *Ros1*,

Zm00001d038302, showed significantly reduced expression in the *mop1* and *mop3* mutants

and has a strong mCHH island in several inbred lines (Figure S4). This provides evidence to

support a requirement for RdDM and CHH methylation in the proper control of this gene in

maize.

Genes regulated by CMT are enriched for natural DMRs:

The genes that are sensitive to mutations in CMT or RdDM components may reflect examples

of natural variation for epigenetic regulation. Indeed, an earlier study has found that many of the

genes influenced by *Zmet2* exhibit variable expression patterns in different maize inbreds

(Makarevitch *et al.* 2007). We used WGBS data from B73 and 17 other diverse maize inbreds to

document natural variation for DNA methylation among maize inbreds. Differentially Methylated

Regions (DMRs) were identified in all three contexts (CG, CHG, and CHH) between B73 and

the other inbreds. More than 200,000 DMRs were called in the CG and CHG contexts, with over

50,000 DMRs in the CHH context. Each maize gene was classified based on whether there was

a DMR within 200 bp of the transcription start site for each of the three sequence contexts. A

relatively small portion (~3-7%) of maize genes have CG, CHG or CHH DMRs near the

promoter (Table S2). We proceeded to assess whether naturally variable DMRs were more

prevalent near genes that exhibit altered expression in CMT or RdDM mutants (Figure 3A).

Genes that are up- or down-regulated in CMT mutants exhibit a significant enrichment for CG

and CHG DMRs in their promoter regions. RdDM up-regulated genes are significantly enriched

for having CHH and CHG DMRs and also show an enrichment (though not significant) for CG

DMRs near the promoter (Figure 3A).

This suggests that many of the genes with altered expression in CMT or RdDM mutations may

have pre-existing natural variation for DNA methylation that would affect expression levels in

maize populations. RNAseq data from leaf tissue for ten of the inbred lines with WGBS data

was utilized to determine whether there was a significant association (p.value < 0.05, pearson

correlation) between context-specific methylation level at the DMR and gene expression levels.

We found that nearly 50% of the genes that exhibit altered expression in CMT or RdDM mutants

that are located near DMRs had natural variation for gene expression levels that was

significantly associated with DNA methylation levels. Two examples of CMT up-regulated genes

that exhibit significantly correlated expression and CHG methylation at the bin overlapping the

TSS among diverse lines are shown in Figure 3B-C. In wild-type maize inbred lines we see two

classes with respect to expression level and CHG methylation levels at the DMR near the TSS.

In one group of lines, including B73, the DMR is highly methylated and the gene is

transcriptionally silent. In the other group of genotypes (and in B73 *zmet2-m1* mutant lines -

blue dots) the DMR has low methylation and the gene is expressed. Although we were

interested in performing a similar analysis for natural variation in TE methylation and expression

we were not able to assess this due to the highly polymorphic nature of TEs among different

maize lines and the lack of de novo assemblies for other genotypes.

Properties of TEs with altered expression:

There are 104 TE families with altered expression in mutants that perturb RdDM or CMT

components in maize. These included 32 families up- and 7 families down-regulated in at least

two contrasts of RdDM mutants, and 48 families up- and 24 families down-regulated in at least

two contrasts of CMT mutants (Figure 1D). There are examples of both class I (specifically Long

Terminal Repeat or LTR) and class II (specifically Terminal Inverted Repeat or TIR) TE families

that exhibit altered expression in both RdDM and CMT mutants (Figure 4A). Most families with

varied expression were small (< 10 members), consistent with the genome-wide distribution

(Figure 4B). The relative age of the LTR families that have altered expression was assessed to

determine if they were particularly young or old families (Figure 4C). The relative ages of LTR

transposons can be approximated by comparing the sequence similarity of the two LTR

sequences. Since LTR sequences must be identical upon initial integration, a higher LTR

similarity denotes younger TE insertions. LTR families that are up-regulated in RdDM mutants

are enriched for younger LTR elements when compared with the distribution of ages present

genome-wide. We also tested the mean GC content of TEs within families to test whether

families depleted in cytosines are more susceptible to subtle perturbations in methylation, as is

the case for the *ONSEN* family in *Arabidopsis* (Cavrak *et al.* 2014). TE families up-regulated in

RdDM mutants do have a slightly lower GC content on average, though it is not clear if this

change alone is sufficient to cause the expression changes (Figure 4D).

We sought to further document the properties of these TE families through analysis of their

expression in nearly 100 developmental tissues or stages of B73. During typical development,

approximately 3,400 TE families are expressed in at least one tissue or stage. There are 5 TE

families up-regulated in RdDM mutants and 18 TE families up-regulated in CMT mutants that

are not expressed in any tissue or developmental stage assessed. The other families of TEs

that are up-regulated in RdDM mutants (25 families) or CMT mutants (27 families) were

assessed to determine if they exhibit distinct patterns of expression. Interestingly, approximately

one third of the TE families that are up-regulated in RdDM mutants show higher expression in

the endosperm than other tissues (Figure 5A). In contrast, the TE families up-regulated in CMT

mutants do not show any evidence for higher expression in a particular tissue type. The

enrichment for endosperm expression in TE families up-regulated in RdDM mutants does not

extend to genes up-regulated in the mutants and cannot be simply attributed to lower

expression of the *Mop1* and *Mop3* genes in these tissues (Figure 5). This result highlights the

potential for some TEs to escape RdDM-based silencing in endosperm, where dynamic

changes to DNA methylation may reinforce TE silencing in the embryo (Martínez and Slotkin

2012; Gehring 2013; Wang *et al.* 2015; Dong *et al.* 2017). Meanwhile, both genes and TE

families susceptible to mis-regulation in CMT mutants are less often expressed across

development, consistent with the greater developmental stability of CHG methylation over CHH

methylation (Kawakatsu *et al.* 2016, 2017; Narsai *et al.* 2017; Bouyer *et al.* 2017).

Evidence for locus-specific and coordinated changes in expression of TEs:

While per-family analysis is useful in capturing additional expression dynamics of repetitive

transposable elements, the expression of individual elements can be influenced by a variety of

location-specific attributes such as methylation levels and proximity to genes as well as family-

level attributes such as binding motifs and nucleotide content. We were interested in

documenting the relative behavior of different elements within the same family to understand

whether the changes in expression of TEs were occurring in an element-specific or family-wide

manner. Coordinate changes in expression could indicate the importance of RdDM or CMT for

family-wide regulation while element-specific changes could reflect influences at particular loci.

A set of TE families with <10 elements that had altered expression and for which at least 50% of

the reads could be uniquely assigned to specific element were identified and used for analysis

of coordinate versus locus-specific expression (Tables S3, S4). The unique mapping reads for

these families were used to evaluate element-specific expression. Half of the testable TE

families had expression of a single member of the family indicating locus-specific changes

(examples in Figure 6A, C). In the other half of the TE families there was evidence for

expression changes for multiple elements of the same family (Figure 6B, D). This suggests at

least some level of coordinate regulation of multiple members of the family by CMT or RdDM

pathways. However, even in examples of coordinate expression a single element accounted for

the vast majority of unique reads mapping to the family. Examples of both locus-specific and

coordinate changes in expression for both CMT and RdDM mutants were found but we were not

able to assess enough families to determine if there was any enrichment for the type of

regulation for these two silencing pathways.

453   DISCUSSION

Maize has been a model system for the discovery of several epigenetic phenomena such as

imprinting (Kermicle 1970; Kermicle and Alleman 1990), paramutation (Brink 1956; Chandler

2007; Hollick 2017) and transposon silencing (Chandler and Walbot 1986; Chomet *et al.* 1987).

An unresolved question is whether epigenetic regulation plays important roles in quantitative

trait variation beyond handful of well characterized loci. Our ability to document the full role for

epigenetic regulation and DNA methylation has been limited by our inability to recover plants

with major reductions in the level of DNA methylation (Li *et al.* 2014). Forward genetic screens

have uncovered a number of components of the RNA-directed DNA methylation (RdDM)

machinery as playing critical roles in maintenance of silenced paramutant states (Dorweiler *et*

al. 2000; Jr *et al.* 2009) or transgene silencing (McGinnis *et al.* 2006). These mutants have

substantial effects on CHH methylation in maize but have minimal effect on genome-wide levels

of CG or CHG methylation (Li *et al.* 2014). Reverse-genetic analyses have identified loss-of-

function alleles for a number of other genes predicted to play important roles in DNA

methylation but the only single mutants with significant effects on genome-wide DNA

methylation are the CMT genes of maize, *Zmet2* and *Zmet5 (Li et al. 2014)*. In this study we

have documented how these subtle perturbations of the maize methylome affect the

transcriptome in order to find genes subject to epigenetic regulation.

The effects of mutations in RdDM or CMT genes in maize are quite limited. Our evidence

suggests that there is little effect on the overall transcriptome of these plants. This might be

expected given the limited effect on overall plant phenotype for each of these mutations. A

recent study found that *rmr6* (allelic to *mop3*) mutants exhibited transcription changes from a

larger portion of the genome but much of this was associated with increased transcriptional

'noise' at lowly expressed regions (Forestan *et al.* 2017). However, there are sets of genes with

clear changes in expression in each of the mutant lines in our study. Similarly, a directed

analysis of differential expression in *rmr6* found a smaller set of genes with significant changes

(Forestan *et al.* 2017). While relatively few genes exhibit major changes in expression there are

a significant number of genes that exhibit similar expression changes in multiple RdDM or CMT

mutants. These findings are compatible with the concept that there are a small number of genes

in the maize genome that have epigenetic regulation that is solely dependent upon RdDM or

CMT mediated regulation. It is likely that a much larger number of genes are redundantly

regulated by the RdDM and CMT pathway along with MET1 mediated CG methylation.

Many of the genes that are up-regulated in CMT mutants exhibit high levels of CHG

methylation. The CMT mutants reduce this methylation and allow for increased expression.

Previous studies noted that the genes sensitive to *zmet2-m1* mutations varied in different maize

inbreds (Makarevitch *et al.* 2007). This prompted us to investigate whether the genes that are

up-regulated in CMT mutants might exhibit natural variation for DNA methylation levels. Many of

the genes that are up-regulated in CMT mutants have CHG DMRs nearby and many of these

exhibit variable levels of expression among maize genotypes that is negatively correlated with

CHG methylation levels. This suggests epiallelic diversity for targets of CMT-mediated gene

silencing. If these changes in expression lead to phenotypic variation, plant breeders are likely

able to select for preferred epigenetic states. However, it would also be possible to introduce

novel epigenetic variation through reductions of CHG methylation.

DNA methylation is often considered to play a primary role in maintaining genome integrity by

silencing transposable elements. Indeed, there are clear examples of release of transposon

silencing in mutants affecting DNA methylation in *Arabidopsis* (Miura *et al.* 2001; Mirouze *et al.*

2009) and maize (Lisch *et al.* 2002; Jia *et al.* 2009). However, generating a complete

understanding of transposon expression is complicated by the highly repetitive nature of

transposable elements. In order to survey expression using RNAseq most researchers focus on

unique mapping reads to ensure that expression is accurately attributed to the proper genomic

locus. In this study we elected to primarily focus on TE families rather than individual elements

and we utilized an approach that allowed for the combined use of unique and multiple-mapping

reads to assess TE family expression. We did find evidence that a number of TE families

require RdDM and/or CMT for silencing. There were few distinguishing features about these

TEs relative to others making it unclear why the silencing of these families was easily released

in these mutants. For several families we were able to document evidence for specific release of

silencing of a single member of the family while in other cases we found that multiple members

of the same family were reactivated.

This study defines a set of genes and TE families that are regulated by DNA methylation. The

silencing of these genes and TEs relies solely upon RdDM or CMT based epigenetic regulation.

These loci provide important insights into the mechanisms that allow for epigenetic regulation

and the natural variation for epigenetic regulation in maize.

520   ACKNOWLEDGEMENTS

521   We thank Peter Hermanson and Jonathan Giesler for technical assistance and Jonathan Gent

522   for helpful feedback. Funding for this work was grants from USDA-NIFA2016-67013-24747 to

523   N.M.S. and the National Science Foundation IOS-1237931 to N.M.S. and M.W.V. Q.L. was

524   supported by The National Key Research and Development Program of China (2016YFD0101003).

REFERENCES

Alleman M., Sidorenko L., McGinnis K., Seshadri V., Dorweiler J. E., *et al.*, 2006 An RNA-dependent RNA polymerase is required for paramutation in maize. Nature 442: 295–298.

Anders S., Pyl P. T., Huber W., 2015 HTSeq–a Python framework to work with high-throughput sequencing data. Bioinformatics 31: 166–169.

Barber W. T., Zhang W., Win H., Varala K. K., Dorweiler J. E., *et al.*, 2012 Repeat associated small RNAs vary among parents and following hybridization in maize. Proc. Natl. Acad. Sci. U. S. A. 109: 10444–10449.

Bouyer D., Kramdi A., Kassam M., Heese M., Schnittger A., *et al.*, 2017 DNA methylation dynamics during early plant life. Genome Biol. 18: 179.

Brink R. A., 1956 A Genetic Change Associated with the R Locus in Maize Which Is Directed and Potentially Reversible. Genetics 41: 872–889.

Cavrak V. V., Lettner N., Jamge S., Kosarewicz A., Bayer L. M., *et al.*, 2014 How a retrotransposon exploits the plant's heat stress response for its activation. PLoS Genet. 10: e1004115.

Chandler V. L., Walbot V., 1986 DNA modification of a maize transposable element correlates with loss of activity. Proc. Natl. Acad. Sci. U. S. A. 83: 1767–1771.

Chandler V. L., 2007 Paramutation: from maize to mice. Cell 128: 641–645.

Chomet P. S., Wessler S., Dellaporta S. L., 1987 Inactivation of the maize transposable element Activator (Ac) is associated with its DNA modification. EMBO J. 6: 295–302.

Dong X., Zhang M., Chen J., Peng L., Zhang N., *et al.*, 2017 Dynamic and Antagonistic Allele-Specific Epigenetic Modifications Controlling the Expression of Imprinted Genes in Maize Endosperm. Mol. Plant 10: 442–455.

Dorweiler J. E., Carey C. C., Kubo K. M., Hollick J. B., Kermicle J. L., *et al.*, 2000 Mediator of Paramutation1 is Required for Establishment and Maintenance of Paramutation at Multiple Maize Loci. Plant Cell 12: 2101–2118.

Eichten S. R., Swanson-Wagner R. A., Schnable J. C., Waters A. J., Hermanson P. J., *et al.*,

2011 Heritable epigenetic variation among maize inbreds. PLoS Genet. 7: e1002372.

Eichten S. R., Briskine R., Song J., Li Q., Swanson-Wagner R., *et al.*, 2013 Epigenetic and

genetic influences on DNA methylation variation in maize populations. Plant Cell 25: 2783–

2797.

Erhard K. F. Jr, Stonaker J. L., Parkinson S. E., Lim J. P., Hale C. J., *et al.*, 2009 RNA

polymerase IV functions in paramutation in Zea mays. Science 323: 1201–1205.

Erhard K. F. Jr, Talbot J.-E. R. B., Deans N. C., McClish A. E., Hollick J. B., 2015 Nascent

transcription affected by RNA polymerase IV in Zea mays. Genetics 199: 1107–1125.

Forestan C., Aiese Cigliano R., Farinati S., Lunardon A., Sanseverino W., *et al.*, 2016 Stress-

induced and epigenetic-mediated maize transcriptome regulation study by means of

transcriptome reannotation and differential expression analysis. Sci. Rep. 6: 30446.

Forestan C., Farinati S., Aiese Cigliano R., Lunardon A., Sanseverino W., *et al.*, 2017 Maize

RNA PolIV affects the expression of genes with nearby TE insertions and has a genome-wide

repressive impact on transcription. BMC Plant Biol. 17: 161.

Gehring M., 2013 Genomic imprinting: insights from plants. Annu. Rev. Genet. 47: 187–208.

Gent J. I., Ellis N. A., Guo L., Harkess A. E., Yao Y., *et al.*, 2013 CHH islands: de novo DNA

methylation in near-gene chromatin regulation in maize. Genome Res. 23: 628–637.

Gent J. I., Madzima T. F., Bader R., Kent M. R., Zhang X., *et al.*, 2014 Accessible DNA and

relative depletion of H3K9me2 at maize loci undergoing RNA-directed DNA methylation. Plant

Cell 26: 4903–4917.

Gouil Q., Baulcombe D. C., 2016 DNA Methylation Signatures of the Plant

Chromomethyltransferases. PLoS Genet. 12: e1006526.

Hale C. J., Stonaker J. L., Gross S. M., Hollick J. B., 2007 A novel Snf2 protein maintains trans-

generational regulatory states established by paramutation in maize. PLoS Biol. 5: 2156–2165.

Hofmeister B. T., Lee K., Rohr N. A., Hall D. W., Schmitz R. J., 2017 Stable inheritance of DNA

methylation allows creation of epigenotype maps and the study of epiallele inheritance patterns

in the absence of genetic variation. Genome Biol. 18: 155.

Hollick J. B., Kermicle J. L., Parkinson S. E., 2005 Rmr6 maintains meiotic inheritance of

paramutant states in Zea mays. Genetics 171: 725–740.

Hollick J. B., 2017 Paramutation and related phenomena in diverse species. Nature

reviews.Genetics 18: 5–23.

Hu L., Li N., Xu C., Zhong S., Lin X., *et al.*, 2014 Mutation of a major CG methylase in rice

causes genome-wide hypomethylation, dysregulated genome expression, and seedling lethality.

Proc. Natl. Acad. Sci. U. S. A. 111: 10642–10647.

Jia Y., Lisch D. R., Ohtsu K., Scanlon M. J., Nettleton D., *et al.*, 2009 Loss of RNA-dependent

RNA polymerase 2 (RDR2) function causes widespread and unexpected changes in the

expression of transposons, genes, and 24-nt small RNAs. PLoS Genet. 5: e1000737.

Jiao Y., Peluso P., Shi J., Liang T., Stitzer M. C., *et al.*, 2017 Improved maize reference genome

with single-molecule technologies. Nature 546: 524–527.

Jr K. F. E., Stonaker J. L., Parkinson S. E., Lim J. P., Hale C. J., *et al.*, 2009 RNA polymerase

IV functions in paramutation in Zea mays. Science 323: 1201–1205.

Kawakatsu T., Stuart T., Valdes M., Breakfield N., Schmitz R. J., *et al.*, 2016 Unique cell-type-

specific patterns of DNA methylation in the root meristem. Nature plants 2: 16058.

Kawakatsu T., Nery J. R., Castanon R., Ecker J. R., 2017 Dynamic DNA methylation

reconfiguration during seed development and germination. Genome Biol. 18: 171.

Kermicle J. L., 1970 Dependence of the R-Mottled Aleurone Phenotype in Maize on Mode of

Sexual Transmission. Genetics 66: 69–85.

Kermicle J. L., Alleman M., 1990 Gametic imprinting in maize in relation to the angiosperm life

cycle. Development (Cambridge, England).Supplement: 9–14.

Kim D., Pertea G., Trapnell C., Pimentel H., Kelley R., *et al.*, 2013 TopHat2: accurate alignment

of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. 14:

R36–2013–14–4–r36.

Li Q., Eichten S. R., Hermanson P. J., Zaunbrecher V. M., Song J., *et al.*, 2014 Genetic

perturbation of the maize methylome. Plant Cell 26: 4602–4616.

Li Q., Song J., West P. T., Zynda G., Eichten S. R., *et al.*, 2015a Examining the causes and

consequences of context-specific differential DNA methylation in maize. Plant Physiol. 168:

1262–1274.

Li Q., Gent J. I., Zynda G., Song J., Makarevitch I., *et al.*, 2015b RNA-directed DNA methylation

enforces boundaries between heterochromatin and euchromatin in the maize genome. Proc.

Natl. Acad. Sci. U. S. A. 112: 14728–14733.

Lisch D., Carey C. C., Dorweiler J. E., Chandler V. L., 2002 A mutation that prevents

paramutation in maize also reverses Mutator transposon methylation and silencing. Proc. Natl.

Acad. Sci. U. S. A. 99: 6130–6135.

Love M. I., Huber W., Anders S., 2014 Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 15: 550.

Madzima T. F., Huang J., McGinnis K. M., 2014 Chromatin structure and gene expression

changes associated with loss of MOP1 activity in Zea mays. Epigenetics 9: 1047–1059.

Makarevitch I., Stupar R. M., Iniguez A. L., Haun W. J., Barbazuk W. B., *et al.*, 2007 Natural

Variation for Alleles Under Epigenetic Control by the Maize Chromomethylase Zmet2. Genetics

177: 749–760.

Martínez G., Slotkin R. K., 2012 Developmental relaxation of transposable element silencing in

plants: functional or byproduct? Curr. Opin. Plant Biol. 15: 496–502.

McGinnis K. M., Springer C., Lin Y., Carey C. C., Chandler V., 2006 Transcriptionally silenced

transgenes in maize are activated by three mutations defective in paramutation. Genetics 173:

1637–1647.

Mirouze M., Reinders J., Bucher E., Nishimura T., Schneeberger K., *et al.*, 2009 Selective

epigenetic control of retrotransposition in Arabidopsis. Nature 461: 427–430.

Miura A., Yonebayashi S., Watanabe K., Toyama T., Shimada H., *et al.*, 2001 Mobilization of

transposons by a mutation abolishing full DNA methylation in Arabidopsis. Nature 411: 212–

214.

Narsai R., Gouil Q., Secco D., Srivastava A., Karpievitch Y. V., *et al.*, 2017 Extensive

transcriptomic and epigenomic remodelling occurs during Arabidopsis thaliana germination.

Genome Biol. 18: 172.

Papa C. M., Springer N. M., Muszynski M. G., Meeley R., Kaeppler S. M., 2001 Maize

chromomethylase Zea methyltransferase2 is required for CpNpG methylation. Plant Cell 13:

1919–1928.

Quinlan A. R., Hall I. M., 2010 BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26: 841–842.

Regulski M., Lu Z., Kendall J., Donoghue M. T., Reinders J., *et al.*, 2013 The maize methylome

influences mRNA splice sites and reveals widespread paramutation-like switches guided by

small RNA. Genome Res. 23: 1651–1662.

Rowley M. J., Rothi M. H., Böhmdorfer G., Kuciński J., Wierzbicki A. T., 2017 Long-range

control of gene expression via RNA-directed DNA methylation. PLoS Genet. 13: e1006749.

Sloan A. E., Sidorenko L., McGinnis K. M., 2014 Diverse gene-silencing mechanisms with

distinct requirements for RNA polymerase subunits in Zea mays. Genetics 198: 1031–1042.

Song J., Zynda G., Beck S., Springer N. M., Vaughn M. W., 2016 Bisulfite Sequence Analyses

Using CyVerse Discovery Environment: From Mapping to DMRs. In: *Current Protocols in Plant*

Biology, John Wiley & Sons, Inc.

Wang P., Xia H., Zhang Y., Zhao S., Zhao C., *et al.*, 2015 Genome-wide high-resolution

mapping of DNA methylation identifies epigenetic variation across embryo and endosperm in

Maize (Zea may). BMC Genomics 16: 21.

Wicker T., Sabot F., Hua-Van A., Bennetzen J. L., Capy P., *et al.*, 2007 A unified classification

system for eukaryotic transposable elements. Nature reviews.Genetics 8: 973–982.

Williams B. P., Pignatta D., Henikoff S., Gehring M., 2015 Methylation-sensitive expression of a

DNA demethylase gene serves as an epigenetic rheostat. PLoS Genet. 11: e1005142.

Woodhouse M. R., Freeling M., Lisch D., 2006 The mop1 (mediator of paramutation1) mutant

progressively reactivates one of the two genes encoded by the MuDR transposon in maize.

Genetics 172: 579–592.

Xi Y., Li W., 2009 BSMAP: whole genome bisulfite sequence MAPping program. BMC

Bioinformatics 10: 232.

Yamauchi T., Johzuka-Hisatomi Y., Terada R., Nakamura I., Iida S., 2014 The MET1b gene

encoding a maintenance DNA methyltransferase is indispensable for normal development in

rice. Plant Mol. Biol. 85: 219–232.

Figure 1: Summary of differentially expressed genes in mild methylation mutants. A-B: A

principal component analysis (PCA) was performed using log₂(RPM+1) expression values for

genes that are DE in at least one mutant line relative to the appropriate control. (A) The full set

of samples used for this study were assessed and we found that samples in other genetic

backgrounds (*mop3* and *Mop3*) or tissues (*mop1* ear and *Mop1* ear) have the highest level of

variation. (B) A second PCA was performed using only samples in the B73 genetic background

assessed in leaf tissue. WT samples are denoted with triangles and mutants with circles. C-D:

The number of up- (red) and down-regulated (blue) genes (C) and TEs (D) is shown for each

mutant relative to the appropriate wild-type control. The percent of DE genes or TE families that

are up-regulated is marked above each bar, and the number of genes or TEs with consistent

changes in two or more mutants from the same pathway are labeled. E-F: Each of the genes (E)

or TE families (F) that are DE in at least two CMT or RdDM mutants were used to perform

hierarchical clustering using the Euclidean method and the $\log_2$ of the fold-change relative to

wild-type is visualized with a heat map.

Figure 2: Methylation profiles of DE genes. A. The metaprofile of DNA methylation levels in wild-

type B73 seedling leaf tissue was assessed for different sets of DE genes. The DNA

methylation levels surrounding the TSS and TTS were plotted for all expressed genes and

genes DE in CMT (blue) and RdDM (orange) mutants. The three panels show the levels of CG,

CHG, and CHH methylation, with the y-axis showing DNA methylation levels. Error bars

represent standard error. B-C Histogram of the number of genes with different methylation

levels in the first 400 bp downstream of the TSS in the CG (B) and CHG (C) contexts, showing a

bimodal distribution of methylation values. D. Wild-type methylation levels in the CG and CHG

contexts are correlated. *zmet2-m1* mutant methylation data is shown for those genes with high

(>50%) CHG methylation in wild-type (orange dots), showing a loss of CHG but not CG

methylation in the mutant. E. Histogram of the difference between mutant and WT methylation

in the CG (red) and CHG (blue) contexts for those genes up-regulated in CMT mutants that

have WT CHG methylation >50%. F. IGV view of two up-regulated genes in CMT mutants:

Zm00001d045627 and *Zm00001d021982,* which have high CG and CHG methylation in WT

and reduced CHG methylation near the TSS (yellow box) in *zmet2-m1* mutants. G. The

proportion of genes within 2 kb of mCHH islands (mCHHi) is shown for different sets of genes.

The black bar shows the proportion of all genes with a mCHH island while the other bars show

the proportion of genes with altered expression in specific mutant backgrounds that have mCHH

islands, and * denotes significantly higher proportion than expected relative to all genes (p-val <

0.01, chi squared test).

Figure 3. Natural variation for methylation level. A. DMRs in diverse genotypes located within

200 bp the TSS for all genes and for genes with differential expression in either CMT or RdDM

mutants. Significant enrichment compared with expressed set is denoted with * (p-value < 0.01,

chi squared test). B-C. Examples of genes up-regulated in CMT that have negatively correlated

expression and CHG methylation at the bin overlapping the TSS. Data points show values for

B73 (red), *zmet2-m1* mutants (blue), and 9 diverse genotypes (B97, CML322, HP301, IL14,

Mo17, Oh43, P39, Tx303, and W22, black).

Figure 4. Attributes of TE families with altered expression in methylation mutants. A. TE super

family membership for TE families genome-wide and with varied expression in mutants, where

RST = SINE, RIT = LINE, RIL = LINE-L1-like, DTX = TIR-unclassified, DTT = TIR-Tc1/Mariner,

DTH = TIR-PIF/Harbinger, DTC = CACTA, RLX = LTR-unclassified, RLG = LTR-Gypsy, RLC =

LTR-Copia, DTM = TIR-Mu, and DTA = TIR-hAT. B. Size distribution for TE families genome-

wide and with varied expression in mutants, where small: 2-9 members, medium: 10-99

members, and large: >= 100 members. C. Boxplot of the average LTR similarity per-family for

LTR TE families genome-wide, along with those LTR families with differential expression in

methylation mutants. D. Boxplot of the average GC content per-family for TE families genome-

wide and with expression changes in the mutant. * denotes significant deviation from the mean

for all TE families (p-value < 0.01, t-test).

Figure 5. Developmental expression of TE families (A) and genes (B) up-regulated in RdDM and

CMT mutants, along with the typical expression of genes mutated in this study (C), where rows

show TE families or genes and columns show RNA-seq libraries. Developmental samples are

grouped by tissue type, with seed samples split into two clusters based on relative contribution

of endosperm: endosperm & late seed (12+ days after pollination) and embryo & early seed (up

to 10 days after pollination). For full list of tissue assignments and RNA-seq library accession

numbers, see Table S1. Approximately one third of TE families up-regulated in RdDM mutants

have higher expression in endosperm than other tissues across development, a pattern not

observed for genes. In contrast, many more TE families and genes up-regulated in CMT

mutants are never or lowly expressed during typical development.

Figure 6: TE families up-regulated in methylation mutants can have expression of a single

element or multiple elements. A-B Show two LTR families up-regulated in RdDM mutants, and

C-D show two LTR families up-regulated in CMT mutants. All families have both unique (red)

and multi-mapped (blue) reads. The proportion of unique-mapping reads assigned to each

element is shown for a representative library. Unique-mapping reads showed confident

expression of only a single member of a family (A and C) or coordinated expression of more

than one member of a family (B and D).

751   SUPPLEMENTAL MATERIALS

Supplemental Figures S1-S4: attached as separate document

Table S1: Complete list of datasets used in this study.

Table S2: Expression values, differential expression calls, list assignments, and DMR calls for

all genes.

Table S3: Expression values, differential expression calls, list assignments, and descriptions for

all TE families.

Table S4: Unique mapping read counts and descriptors for each unique transposable element.