TITLE

Endospores and other lysis-resistant bacteria comprise a widely shared core community

within the human microbiota

AUTHORS

Sean M. Kearney[1,2,3], Sean M. Gibbons[1,2,3], Mathilde Poyet[1,2,3], Thomas Gurry[1,2,3],

Kevin Bullock[2], Jessica R. Allegretti[4,5], Clary B. Clish[2], Eric J. Alm[1,2,3,*]

[1] Department of Biological Engineering, Massachusetts Institute of Technology,

Cambridge, MA, U.S.A.

[2] The Broad Institute, Cambridge, MA, U.S.A.

[3] The Center for Microbiome Informatics and Therapeutics, Cambridge, MA, U.S.A.

[4] Division of Gastroenterology, Brigham and Women's Hospital, Boston, MA, USA

[5] Harvard Medical School, Boston, MA, USA

* Corresponding author: ejalm@mit.edu

16 ABSTRACT

Endospore-formers in the human microbiota are well adapted for host-to-host

transmission, and an emerging consensus points to their role in determining health and

disease states in the gut. The human gut, more than any other environment,

encourages the maintenance of endospore formation, with nearly 60% of genera in the

microbiome carrying genes attributed to this trait. However, there has been limited work

on the ecological role of endospores and other stress-resistant cellular states in the

human gut. Here, we applied a culture-independent protocol to enrich for endospores

and other stress-resistant cells in human feces, and provide evidence that organisms

found in resistant states are adapted for cross-host dissemination. Furthermore,

organisms with resistant cellular states respond to shared environmental signals that

likely mediate colonization of recently disturbed human guts.

30 INTRODUCTION

To date, there is limited work investigating the relevance of stress-resistant cellular

states in the propagation, survival, and function of organisms in the mammalian

gastrointestinal (GI) tract. The gut is the only known environment with such a

considerable abundance of organisms that form endospores, considered the most

stress-resistant of all cell-types[1]. Anaerobic, endospore-forming Firmicutes are

numerically dominant members of the GI tract of most animal species[2,3]. Within this

group of organisms, the presence of genes for endospore formation suggests that

growth in the GI tract favors the maintenance of this large gene repertoire[2]. The

apparent utility of these genes is to allow organisms to enter metabolically dormant

states that aid in survival and transmission to new hosts. Passage through the GI tract

is likely to trigger sporulation[4,5], but the mechanisms by which this process occurs and

the signals that induce sporulation here are mostly unknown, even for well-studied

pathogens like *Clostridioides difficile*. Resistant cellular states like endospores appear to

be adaptive in the mammalian gut environment. It is likely that other non-endospore-

forming taxa have evolved analogous resistance strategies for passing between hosts.

Many endospore-forming organisms in the human gut are in the class Clostridia, the

most well-studied of which are the pathogens *C. difficile* and *Clostridium perfringens*[6,7].

However, Clostridia also includes abundant organisms not known to form endospores,

like *Faecalibacterium prausnitzii*[8] and *Roseburia intestinalis*[9]. For *C. difficile*, the role of

sporulation is central to disease etiology[10], particularly in patients who experience

recurrence. Sporulation and rising levels of antibiotic resistance allow *C. difficile* to

persist in the face of antibiotic assault, ensuring that it remains in the environment to

rapidly re-colonize its host.

Among Clostridia that do not cause disease, multiple strains of endospore-forming

organisms have the capacity to induce T regulatory cells and associated anti-

inflammatory cytokines[11,12] involved in sensitivity to, for example, peanut antigen[13].

These organisms have recently been shown to provide pathogen resistance in neonatal

mice[14]. Similarly, endospore-forming commensals of the murine GI tract have a central

role in mediating the induction of a Th17-type T helper cell response[15–17]. Many

Clostridia also produce butyrate as an end-product of metabolism, which can regulate

how immune cells interact with gut commensal bacteria[18–22]. This group of organisms

also boosts production of serotonin by enterochromaffin cells in the intestine, crucial for

motility in the gut[23]. An ecological understanding of sporulation and induction of other

resistant states could be informative for how these phenotypes interact with host

immunity. For example, such an understanding could inform whether inflammation acts

positively or negatively on endospore formation, or whether endospores themselves

have immunomodulatory effects.

Environmental stress resistance protects cells faced with unfavorable conditions. The

signals triggering resistance are likely quite varied. Even for well-studied endospore-

forming bacteria, inducing sporulation *in vitro* can be difficult, and across strains of one

species, signals that induce sporulation in one strain may be insufficient to induce

sporulation in others[24]. Further, even organisms that abundantly form endospores in

their native environment may not do so under conditions permitting vegetative growth.

For instance, *Paenibacillus larvae* in honeybees will only form endospores *in vitro* under

idiosyncratic conditions designed to mimic the host environment[25]. Similarly, certain

strains of *Clostridium perfringens* rarely form endospores *in vitro* unless exposed to a

specific set of environmental stressors[24]. The discrepancy in phenotype of organisms in

their native environments compared to *in vitro* argues for culture-free approaches to

investigate such phenotypes *in situ*. Enriching for stress-resistant cells in environmental

samples provides a means to uncover the actual context in which these states form.

Here, we investigate which organisms are present as endospores or as other resistant cell types in the human gastrointestinal tract. We modified previously described methods to enrich fecal samples for endospores and obtain paired bulk community and resistant fraction 16S rDNA sequence data for 24 healthy individuals and one individual across 24 days. We consistently enriched for putatively endospore-forming taxa in all samples, as well as other taxa, predominantly from the Actinobacteria phylum, that show high levels of lysis resistance. We compared resistant OTUs (rOTUs) and non-resistant OTUs (nOTUs) to identify ecological characteristics differing between these groups. Using a database of rOTUs, we find consistent signals for these organisms in their responses to a variety of disturbances across multiple independent data sets. Overall, we show a tight association between the ecological role of these resistant organisms and their distribution within and across human hosts.

RESULTS AND DISCUSSION

Sequencing resistant fraction reveals resistant taxa present in human feces

We modified a culture-independent method[26] to generate paired bulk community and resistant fraction 16S rDNA amplicon data from human feces (Figure 1A). Aggregating the data across our cohort, we see expansion of classes with known endospore-formers in the resistant fraction: Clostridia, Erysipelotrichia, and Bacilli (Figure 1B). We also see

depletion of classes lacking endospore-formers (Bacteroides, Betaproteobacteria,

Verrucomicrobia, Gammaproteobacteria).

Organisms in the class Actinobacteria were enriched in the resistant fraction, but lack

genes considered essential for endospore formation. Although exospore formation is

well documented in some families of Actinobacteria (e.g. Actinomycetaceae and

Streptomycetaceae), these families have only modest representation in our data. We

see high-level resistance primarily from *Bifidobacterium* and *Collinsella*, whose

representative genomes lack orthologs for genes thought to be essential to exospore

formation.

We suspect that high level resistance in the Actinobacteria is mediated primarily by

resistance to lysozyme conferred by cell wall structures common to Actinobacteria[27].

Lysozyme is one of the most common and important defense mechanisms used by

neutrophils, monocytes, macrophages, and epithelial cells[28,29]. It is abundant in human

milk, a source of *Bifidobacterium* species transferred to breast-feeding infants, and in

saliva and mucus, where it serves an antibacterial role[30]. Attempts to deplete

Actinobacteria with achromopeptidase, which has previously been shown to break down

Actinobacterial peptidoglycan, had variable efficacy across samples (data not shown).

Thus, factors other than cell wall structure may contribute to Actinobacteria resistance.

To quantify the extent of lysis resistance, we calculated the proportion of normalized

reads for each 100% OTU (unique sequence) in the resistant fraction to the sum of its

reads in the bulk community and the resistant fraction. We then obtain a finite quantity

even for organisms not observed in one of the paired samples. When the proportion

exceeds 0.5 we call an OTU enriched in the resistant fraction (Figure 1A). An OTU is

enriched in at least one of the samples in which it is present is considered a resistant

OTU (rOTU), and non-resistant (nOTU) otherwise. Using the above definitions, all of the

rOTUs are either Firmicutes or Actinobacteria (Figure 1C). In fact, when grouping OTUs

at the genus level, the top two most enriched genera (*Bifidobacterium*, *Collinsella*) are

both Actinobacteria.

Resistant fractions consist of a few dominant and many rare OTUs

In order to investigate ecological properties of the resistant cell fraction, we first

examined the community structure of resistant fractions and compared these to their

bulk community counterparts. Using alpha diversity metrics, it is clear that resistant

fractions are significantly less diverse than their bulk community counterparts (Figure

2A). However, this difference is driven largely by differences in evenness between the

communities, as enrichments are significantly less even. The OTU richness between

resistant fractions and bulk community samples tend to be no different. Thus, in

general, resistant fractions are dominated by a few highly abundant OTUs and many

OTUs with just a single count.

rOTUs are more shared than nOTUs among individuals

Resistant fraction OTUs are shared between individuals more often than bulk

community OTUs (Figure 2B). By comparing the ratio of OTUs found in a single person

to OTUs found in multiple individuals, we find a significantly higher proportion of OTUs

shared in the resistant fraction than in the bulk community (Fisher Exact Test, odds ratio

= 1.68, p-value = 1.5e-45, Figure 2B). Furthermore, the majority of OTUs found in the

resistant fractions are not observed in their bulk community counterparts. We interpret

these results to mean that the resistant fractions comprise a seed bank of generally rare

OTUs that are widespread within our cohort.

To further test the hypothesis that resistant states contribute to prevalence, we

examined the frequency with which rOTUs were found among the bulk communities

across individuals compared to nOTUs (Supplementary Figure 1). First, nOTUs, which

are never enriched in the resistant fraction, are significantly less likely to be shared

among multiple individuals than rOTUs (Mann Whitney U Test comparing the

distribution of the number of individuals sharing each rOTU to the number of individuals

sharing each nOTU, p-value = 4.8e-8). We again see this result by calculating the

correlation between the frequency of resistance (the number of times an organism is

enriched in the resistant fraction divided by the number of times it is observed) and

sharedness (number of individuals an OTU is observed in divided by the total number of

individuals), giving a weak, but positive and highly significant correlation (Spearman rho,

correlation = 0.12, p-value = 7.3e-25; Kendall tau, correlation = 0.10, p-value = 1.97e-

24). This result suggests that organisms that do not form resistant states are less likely

to be found across multiple individuals than those that do. In fact, rOTUs tend to be less

dominant members of the community (median rOTUs = 1 count, median nOTUs = 2

counts, Mann-Whitney U Test, p-value = 3.5e-13). This result indicates that organisms

found in resistant states generally have lower overall abundance. We suggest that

organisms with resistant states sacrifice abundance within an individual for higher rates

of dispersal across individuals.

Representation of organisms in resistant fractions is heterogeneous across and within

individuals

A tradeoff between resistance and population size within a host suggests that there may

be strong positive selection for such resistance phenotypes. Variation in this trait among

related organisms could be indicative of selection. In order to visualize how much of a

population is present in a resistant state within a given sample, we scaled 16S rDNA

abundance data using V4 16S rDNA qPCR-based estimates of community size and

defined the resistant fraction as the ratio of these scaled reads for each OTU. We plot

this quantity on a phylogeny representing 99% OTUs (clustered at 99% nucleotide

identity) present in at least 8 individuals and up to 24 individuals (Figure 3). First, we

note the high variability in the resistant fraction within and across taxa (the average

variation is over 50-fold within each taxon). For one Roseburia 99% OTU in particular,

this quantity varies over 3 orders of magnitude, suggesting this OTU contains

organisms present in a resistant state in some individuals, but not in others.

Furthermore, within a person, OTUs with the same genus classification can be

discordant in their degree of resistance. In the individual time series, for example, one

Ruminococcus 100% OTU is almost always enriched, and another is never enriched

(Supplementary Figure 4). The closest matching genomes to these two organisms show

differences in sporulation gene content, with the resistant *Ruminococcus* sharing 48/58

core sporulation genes[31], and the non-resistant only 41/58 (Supplementary Figure 5 and

Supplementary Table 5). We also see that spore maturation proteins *spmA* and *spmB*

vary in their presence in genomes of genera with variable enrichment phenotypes.

These genes are involved in spore cortex dehydration and heat resistance in *B. subtilis*

and *C. perfringens*, so their loss might contribute to differences in the recovery of

resistant cells in this work.

Formation of resistant cells itself might be selected on in this system. There is evidence

that the sporulation phenotype is evolving in mammalian guts, as several gut isolates of

B. subitilis lack genes that negatively regulate sporulation compared to their laboratory

counterparts[34]. Knowing which organisms can form resistant cells in a community does

not provide complete information about which organisms do. Formation of resistant

states *in vivo* seems to be highly context dependent. We also note that loss of a single

gene (i.e. *spo0A*) in *C. difficile* is sufficient for loss of sporulation, such that retaining

endospore formation requires strong purifying selection.

rOTUs share signals for growth within an individual

Previous evidence has shown that bile acids contribute to outgrowth of *C. difficile*

endospores *in vivo*[35]. As a first pass to measure bile acid concentration in the gut, we

tracked a *Bilophila* OTU in the time series. *Bilophila* are known to use taurine derived

from taurocholic acid, a primary bile acid, as an electron acceptor for sulfite reduction[36].

Among rOTUs within the bulk community time series, the average Spearman correlation

to *Bilophila* is 0.110 compared to 0.015 for nOTUs (Mann Whitney U Test p-value =

3.17e-14). The abundance distribution for *Bilophila* provides information about the

abundance of rOTUs. As, taurocholate is a known germinant for several endospore-

forming species[2], *Bilophila* abundance might act as a proxy for taurocholate

concentration or the capacity for sulfite reduction, which requires reducing anaerobic

conditions.

More broadly, the consistent correlations between rOTUs and *Bilophila* implied that

rOTUs would in general be more correlated to each other. By hierarchically clustering

the Spearman correlation profiles of all OTUs present for at least half of the time series

(Figure 4A), we find that the rOTUs mostly fall into one of two clusters, with nOTUs in

the other cluster (Fisher exact test, p-value = 3.13e-25). The average correlation

between rOTUs in the time series to each other is 0.156 compared to 0.014 for nOTUs

to each other (Wilcox rank sum test, p-value = 0): nearly all of the strong correlated

behavior in this person is within this group of organisms (Figure 4A). We interpret this

result to mean that the dynamic behavior of rOTUs is strongly coupled: these OTUs

respond coherently to environmental signals, which are likely linked to bile acids given

their correlations to *Bilophila*.

rOTUs link growth to fatty acid metabolism

To address whether bile-related signals relate to the dynamics of rOTUs in the time
series, we conducted untargeted metabolomics with standards for fatty acid
metabolism. We then calculated the Spearman correlations between the median
abundance profile of OTUs in the rOTU-dominated correlation cluster and metabolites
for which we had standard markers. This cluster tends to correlate positively with long-
chain saturated fatty acids, and negatively with long-chain polyunsaturated fatty acids
and, notably, taurocholate (Supplementary Table 3). We suspect that taurocholate
metabolism by members of this group drives down the concentration of taurocholate in
stool. Additionally, saturated fatty acid concentration in the stool measures fatty acids
escaping absorption in the small intestine. This process would be negatively impacted
by microbial metabolism of taurocholate, as it more efficiently emulsifies saturated fats
than glycine-conjugated primary bile acids[36]. Fecal concentrations of taurocholate
reflect secretion of unmetabolized taurocholate, which should increase if taurocholate
metabolism by the gut microbiota decreases.

Resistant cells lose resistance in response to physiological bile acid concentrations

As a more direct test of the coupling of rOTUs to bile acid concentration, we dosed
ethanol-treated feces (to kill vegetative cells without the additional harshness of the
resistant fraction DNA enrichment protocol) with increasing concentrations of bovine bile

in aqueous solution. We then measured the depletion of OTUs from the endospore-

enrichment using 16S rDNA sequencing (Figure 4B). When correcting for biomass via

qPCR, nearly 20% of OTUs identified in the resistant fraction apparently germinated in

response to bile acids (log-link quasipoisson generalized linear model, p-value < 0.05,

Supplementary Table 4). The true fraction of resistant cells that lose resistance in

response to bile acids is likely higher, as many endospores require an activation step

(i.e. heating at 80°C or treatment with lysozyme as for *C. difficile*[37]) before they will

respond to germinants.

Notably, most ethanol-resistant OTUs began to show a germination-like response at

0.5% bile (Figure 4B), which is near the concentrations found in the human small

intestine[,38]. Although Clostridia and other putative endospore-formers make up the

majority of organisms that lose resistance in response to bile acids, genera in the

Actinobacteria and other resistant cells also show this response when approaching

physiological concentrations. These conserved responses suggest that the same cues

can mediate loss of resistance in distantly related organisms, similar to the conserved

resuscitation response of dormant bacteria to peptidoglycan[39].

rOTUs exhibit shared dynamics in diverse contexts

Correlated behavior, increased prevalence, and shared signals for growth among

rOTUs indicated that these organisms might exhibit a global response during

disturbances of various kinds. To test this hypothesis, we made a sequence database of

rOTUs within our cohort, and used this database to identify putative rOTUs in other

datasets (Figure 5A). We expected that increased prevalence and shared signals for

growth would lead to enhanced colonization of the developing infant gut microbiota[40].

The lysozyme-resistant members of the Actinobacteria and Bacillales dominate the

infant gut microbiota for most of the first 80 days of life and do not equilibrate until the

infant starts a full adult diet (Figure 5B). Early colonization by these rOTUs connects a

resistant state to development of the infant gut microbiome. Here, lysozyme resistance

might be essential for semi-selective transmission of *Bifidobacterium,* as human breast

milk is rich in lysozyme, potentially lysing non-resistant cells[41]. Others have shown

endospore-formers negatively associate with vertical transmission from mother to

infant[42], but other environmentally resistant states as in the Actinobacteria may be

important for vertical transmission.

Depletion of endospore-forming clades is common during infection with *C. difficile.* We

predicted a strong signal for rOTUs in individuals infected with *C. difficile*, due to its

sporulation requirement for transmission[10]. We find a significant depletion of rOTUs

dependent on *C. difficile* infection status (Figure 5C), with a serial depletion of rOTUs

from healthy to first time diagnosis to recurrent patients[43]. Because of this depletion in

rOTUs, we expected that fecal microbiota transplant (FMT) might transfer relatively

more rOTUs than other OTUs[44]. Indeed, among OTUs shared with donors, 90% of

rOTUs increase in abundance following FMT, compared to 77% for the rest of the

community (Fisher exact test, p-value = 0.008) (Figure 5D).

We suspected that rOTUs are a particularly malleable component of the microbiota. To

test this hypothesis, we measured the turnover of rOTUs in the time series of an

otherwise healthy male individual who was infected by Salmonella[47]. New rOTUs almost

completely replaced old rOTUs following this perturbation. By contrast, fewer OTUs

from the rest of the community were lost and gained. This result holds both when

examining the number of OTUs replaced (Fisher exact test, p-value = 6e-12) as well as

the change in abundance of these OTUs (Figure 5E). We see again that rOTUs exhibit

coherent responses to changes in the gut environment, most pronounced in systems

with dramatic perturbations. Colonization of newly vacant niches favors rOTUs, likely

transmitted in an endospore or other resistant state to germinate in an environment

replete with nutrients (including untransformed bile acids). In the absence of a fully

functioning microbiota, rOTUs appear to fill open niches more readily than nOTUs.

CONCLUSION

Gut bacteria in the resistant fraction were more shared across individuals and showed

more correlated dynamics compared to non-resistant organisms. Resistant taxa show

greater turnover following large-scale disturbance events, as in the case of *C. difficile*

and *Salmonella* infection, which suggests that many of these organisms are sensitive to

environmental fluctuations and respond to stress by entering into a dormant, seed-like

state. Environmental sensitivity and high turnover rates of resistant taxa provide an

opportunity to manipulate the composition of the human gut microbiota through targeted

perturbations and replacements. Because of the therapeutic relevance of Clostridia

endospores[11–14,23], determining the exact conditions that permit their replacement may

be of high value for future microbiota-based therapeutics. Here, we found that the

growth of many resistant organisms was associated with dietary fatty acids. If this result

extends to more individuals, one can imagine a therapeutic strategy coupling dietary

changes with introduced resistant cells to enable robust colonization and engraftment.

MATERIALS AND METHODS

See supplemental information for materials and methods

DATA AVAILABILITY STATEMENT

All amplicon sequencing data generated in this study have been can be accessed on

the US National Center for Biotechnology Information SRA database under BioProject

PRJNA389431. Metabolomics data and WGS metagenomic data will be made available

on request.

CODE AVAILABILITY STATEMENT

All custom scripts generated in R to analyze the data in this paper will be made

available through GitHub (https://github.com/microbetrainer/Spores). The sequence

processing pipeline is currently available online (http://amplicon-sequencing-

pipeline.readthedocs.io/en/latest/)

356 **Acknowledgments**

357 We thank Fatima Hussain and Mathieu Groussin for extensive discussion and

358 experimental advice. We thank the MIT BioMicro Center for sequencing service.

360 Funding was provided by the Broad Institute BN10 Training Grants. SM Kearney was

361 funded by an NSF Graduate Research Fellowship.

Figure 1. Resistant fraction sequencing of human fecal bacteria. (A) Overview of resistant cell enrichment and 16S rDNA sequencing protocol. Resistant fraction samples are treated with a series of physical, enzymatic, and chemical lysis steps to deplete vegetative cells. DNA from bulk community and resistant fraction samples are extracted via a mechanical lysis protocol, and 16S rDNA libraries prepared. Communities are analyzed to determine the change in abundance of each OTU in the resistant fraction relative to the bulk community. (right) Phase contrast images of bulk community and resistant fraction – phase bright cells are endospores. Endospores stain green when heat fixed with malachite green, vegetative cells appear red from safranin counter stain. (B) Representative results of 16S rDNA profile for bulk community and endospore-enriched samples. Reads from each OTU are summed across 24 individuals to give a meta-bulk and meta-endospore community. Phylogenetic classes in black text increase with resistant fraction; gray text classes decrease with resistant fraction. (C) Distribution of resistant fraction proportion across phyla aggregated across individuals filtered to remove OTUs with single counts in a sample (for visualization purposes). Colors represent phyla. OTUs with a resistant fraction proportion of 0 are absent from

381   the resistant fraction; OTUs with a resistant fraction proportion of 1 are absent from the
382   bulk community and only found in the resistant fraction.
383

Figure 2. Resistant fraction OTUs are more shared across individuals than bulk community OTUs. (A) Alpha diversity metrics measured for the bulk community (x-axis) and resistant fraction enrichments (y-axis). P-values are for the test of differences between alpha-diversity metric distributions using paired Wilcoxon Rank Sum Test. (B) The number of OTUs found in only a single individual in the bulk communities and endospore enrichments are indicated in orange. The number of OTUs found in more than one individual are indicated through varying shades of purple. The odds ratio for the ratio of shared OTUs to unique OTUs for the endospore enrichments compared to the bulk communities is 1.69.

396
397
Figure 3. Taxa show heterogeneous patterns of resistant cell fractions across
individuals. Phylogenetic placement of the fraction of resistant organisms for taxa
present within at least 8 individuals estimated by the ratio of counts scaled by qPCR-
estimates of biomass in the resistant fractions and bulk communities. Tree branch
colors represent the degree to which a taxonomic group was enriched in the resistant
fraction with pink branches never enriched and blue and green branches enriched at
least once. Classes within each phylum are shown with a colored bar. Arrows indicate
OTUs showing the maximum (black) and minimum (gray) within-OTU variability in
enrichment scores.

Figure 4. Common signals govern resistant state exit and growth in the GI tract.
(A) Correlation matrix of OTU time series data within a single individual's bulk
community hierarchically clustered (Ward's D) by Spearman correlation profile to other
OTUs over time; bars at the tips of the Ward hierarchical cluster represent rOTUs
(blue), nOTUs (pink), *Bilophila* (green). Strength of the correlation varies from red
(perfectly anti-correlated) to blue (perfectly correlated). (B) Abundance of OTUs in the
resistant fraction as a function of bile acid exposure for nine phylogenetically distant
OTUs.
418

419
Figure 5. Resistant OTUs show disproportionate turnover in diverse contexts. (A)
Overview of approach for identifying resistant-cell forming OTUs in 16S rDNA
sequencing datasets. rOTU database sequences are matched to sequences in other
datasets, and then patterns within those datasets among the identified rOTUs are
determined. (B) Fraction of rOTUs present during microbial colonization of an infant gut
annotated with major diet and health perturbations. rOTUs encompass both putative
endospore-forming organisms and those not known to form endospores, but which
possess a resistant state (Actinobacteria and non-endospore-forming Firmicutes) (C)
Fraction of rOTUs present as a function of *C. difficile* infection status (fCDI = first time *C.*
difficile diagnosis, rCDI = at least 3 episodes of *C. difficile* infection following initial
treatment) (D) Fraction of rOTUs and all other OTUs (non-resistant OTUs) transferred
from donors to recipients by fecal microbiota transplant. (E) Time series of rOTUs (top)
and all other (non-resistant) OTUs (bottom) from a human male infected with

Salmonella, with OTUs significantly more abundant pre-infection (dark gray) and
significantly more abundant post-infection (light gray).

MATERIALS AND METHODS

Contact for reagent and resource sharing

Further information may be obtained from the Lead Contact Eric J. Alm (Email: ejalm@mit.edu; address: Massachussetts Institute of Technology Cambridge, MA, 02139, USA)

Experimental Model and Subject Details

Human Subjects

Human subject enrollment and sample collection was approved by the Institutional Review Board of the Massachusetts Institute of Technology (IRB Approval Number: 1510271631). Informed consent was obtained from all subjects. 12 male and 12 female healthy human subjects (age range 21-65) with no history of antibiotic use in the last six months were enrolled in the study. In total, 24 fecal samples were collected from these individuals and an additional 24 fecal samples were collected from one male individual (age 24) over 24 days for culturing and DNA isolation.

Bacterial Cell Culture Conditions

Identities and culture purity of bacterial species were confirmed by Sanger sequencing

of the 16S rRNA genes followed by comparison to sequences in public databases.

Bacteria were grown in their respective media for all experiments except where

otherwise detailed. For routine culture, frozen glycerol stocks of each strain are first

streaked onto agar supportive of growth before inoculation of single colonies into liquid

medium. All bacteria were grown to log-phase at 37° C under anaerobic conditions

(atmosphere 85% N_2, 5% H_2, 10% CO_2), except when preparing endospores.

Method Details

Fecal Sample Processing and Storage

Fecal samples were collected and processed in a biosafety cabinet within 30 minutes of

defecation. Samples (5 g) were suspended in 20 mL of 1% sodium hexametaphosphate

solution (a flocculant) in order to bring biomass into solution as described previously[26].

Fecal samples were bump vortexed with glass beads to homogenize, and centrifuged at

50 x g for 5 min at room temperature to sediment particulate matter and beads.

Triplicate aliquots of 1 mL of the supernatant liquid were transferred into cryovials and

stored at -80° C until processing. For the time series, samples were collected at

approximately 24-hour intervals to reduce variation from uneven sampling frequency.

Resistant Fraction Enrichment from Fecal Samples

We modified a previously published method[26] for endospore sequencing to increase

throughput and decrease signal from contaminating, non-endospore forming organisms.

Fecal samples previously frozen at -80° C were thawed at 4° C prior to use, and 500 µL

was aliquoted for resistant fraction, while the remaining 500 µL was saved for bulk

community DNA extraction. Samples were centrifuged at 4° C and 10,000 x g for 5

minutes, washed and then resuspended in 1 mL Tris-EDTA pH 7.6. Samples were

heated at 65° C for 30 minutes with shaking at 100 rpm and then cooled on ice for 5

minutes. Lysozyme (10 mg/mL) was added to a final concentration of 2 mg/mL and the

samples were incubated at 37° C for 30 minutes with shaking at 100 rpm. At 30

minutes, 50 uL Proteinase K (>600 mAU/ml) (Qiagen) was added and the samples

incubated for an additional 30 minutes at 37° C. Next, 200 uL 6% SDS, 0.3 N NaOH

solution was added and the samples incubated for 1 hour at room temperature with

shaking at 100 rpm. Samples were then centrifuged at 10,000 rpm for 30 minutes. At

this step, a pellet containing resistant endospores should be visible or slightly visible in

the sample, and the pellet is washed three times at 10,000 x g with 1 mL chilled sterile

ddH2O. The pellet is then resuspended in 100 uL ddH2O, and treated with 2 uL DNAse

I (Ambion) to remove residual contaminating DNA with incubation at 37° C for 30 min.

The DNAse is killed by addition of 10 µL Proteinase K (Qiagen) and incubation at 50° C

for 15 minutes, followed by incubation at 70° C for 10 minutes to inactivate Proteinase

K. At this step, microscopic examination of samples is used to confirm the presence of

phase-bright (or phase-dark) spores. The sample is then ready for downstream

extraction and sequencing.

Extraction of Nucleic Acids

We extracted DNA from both the original sample suspended in sodium hexametaphosphate and the output of the resistant fraction. Both the original sample and the resistant fraction were extracted with MoBio PowerSoil Isolation Kit (MoBio Laboratories, Inc.) with three 10 minute bead-beating steps followed by sequential collection of ⅓ of the solution to enhance recovery of endospore DNA as shown previously[46]. DNA was extracted from bacterial pure cultures, fecal enrichment cultures, and endospores using the same protocol as for fecal samples in order to achieve consistency between the methods. DNA from bacterial colonies for 16S rDNA Sanger sequencing confirmation or qPCR was obtained by homogenizing colonies in alkaline polyethylene glycol buffer as described previously [47].

16S rDNA Library Preparation and Sequencing

Libraries for paired-end Illumina sequencing were constructed using a two-step 16S rRNA PCR amplicon approach as described previously with minor modifications [48]. The first-step primers (PE16S_V4_U515_F, 5′ ACACG ACGCT CTTCC GATCT YRYRG TGCCA GCMGC CGCGG TAA-3′; PE16S_V4_E786_R, 5′-CGGCA TTCCT GCTGA ACCGC TCTTC CGATC TGGAC TACHV GGGTW TCTAA T 3′) contain primers U515F and E786R targeting the V4 region of the 16S rRNA gene, as described previously [48]. Additionally, a complexity region in the forward primer (5′-YRYR-3′) was added to help the image-processing software used to detect distinct clusters during Illumina next-

generation sequencing. A second-step priming site is also present in both the forward

(5′-ACACG ACGCT CTTCC GATCT-3′) and reverse (5′-CGGCA TTCCT GCTGA

ACCGC TCTTC CGATC T-3′) first-step primers. The second-step primers incorporate

the Illumina adapter sequences and a 9-bp barcode for library recognition (PE-III-PCR-

F, 5′-AATGA TACGG CGACC ACCGA GATCT ACACT CTTTC CCTAC ACGAC

GCTCT TCCGA TCT 3′; PE-III-PCR-001-096, 5′-CAAGC AGAAG ACGGC ATACG

AGATN NNNNN NNNCG GTCTC GGCAT TCCTG CTGAA CCGCT CTTCC GATCT 3′,

where N indicates the presence of a unique barcode.

Real-time qPCR before the first-step PCR was done to ensure uniform amplification and

avoid overcycling all templates. Both real-time and first-step PCRs were done similarly

to the manufacturer's protocol for Phusion polymerase (New England BioLabs, Ipswich,

MA). For qPCR, reactions were assembled into 20 µL reaction volumes containing the

following: DNA-free H_2O, 8.9 µL, HF buffer, 4 µL, dNTPs 0.4 µL, PE16S_V4_U515_F (3

µM), 2 µL, PE16S_V4_E786_R (3 µM) 2 µL, BSA (20 mg/mL), 0.5 µL, EvaGreen (20X),

1 µL, Phusion, 0.2 µL, and template DNA, 1 µL. Reactions were cycled for 40 cycles

with the following conditions: 98° C for 2 min (initial denaturation), 40 cycles of 98 C for

30 s (denaturation), 52° C for 30 s (annealing), and 72° C for 30s (extension). Samples

were diluted based on qPCR amplification to the level of the most dilute sample, and

amplified to the maximum number of cycles needed for PCR amplification of the most

dilute sample. For first step PCR, reactions were scaled (EvaGreen dye excluded, water

increased) and divided into three 25-µl replicate reactions during both first- and second-

step cycling reactions and cleaned after the first-and second-step using Agencourt

AMPure XP-PCR purification (Beckman Coulter, Brea, CA) according to manufacturer

instructions. Second-step PCR contained the following: DNA-free H_2O, 10.65 µL, HF

buffer, 5 µL, dNTPs 0.5 µL, PE-III-PCR-F (3 µM), 3.3 µL, PE-III-PCR-XXX (3 µM) 3.3

µL, Phusion, 0.25 µL, and first-step PCR DNA, 2 µL. Reactions were cycled for 10

cycles with the following conditions: 98° C for 30 s (initial denaturation), 10 cycles of 98°

C for 30 s (denaturation), 83° C for 30 s (annealing), and 72° C for 30s (extension).

Following second-step clean-up, product quality was verified by DNA gel

electrophoresis and sample DNA concentrations determined using Quant-iT PicoGreen

dsDNA Assay Kit (Thermo Fisher Scientific). The libraries were multiplexed together

and sequenced using the paired-end with 250-bp paired end reads approach on the

MiSeq Illumina sequencing machine at the BioMicro Center (Massachusetts Institute of

Technology, Cambridge, MA).

qPCR

For testing of the resistant fraction protocol, qPCR was carried out as described in the

16S rDNA Library Preparation and Sequencing section. Total bacterial abundance

was quantified using the same primers. For quantification of Firmicutes and

Actinobacteria, primer sequences were obtained from[49]. Primers were used at the same

concentrations as 16S primers, and annealing temperatures were adjusted to the

appropriate temperature for the corresponding primer pairs.

16S rDNA Sequence Data Processing and Quality Control

Paired-end reads were joined with PEAR[50] using default settings. Sequence data files

were processed using a custom amplicon sequencing pipeline. Sequences at least 252

bp long after quality filtering (quality trim Q25) were retained. After quality filtering, the

complexity region between the adapters and the primer along with the primer sequence,

was removed. All sequences not matching the first 15 bases of the primer were

removed. Sequences were trimmed to 252 base pairs and clustered at 100% identity

with UCLUST. OTUs were classified using RDP [51,52]. Sequences that had fewer than 10

reads across individuals were excluded from downstream analysis. The resulting count

tables were used as input for analysis within R.

Identifying High Confidence Endospore-Forming & Resistant OTUs

We developed a workflow for identifying organisms showing increased abundance in

the resistant fraction relative to the bulk community. We use as input an OTU table with

normalized counts obtained using cumulative sum scaling as implemented in the

package metagenomSeq in R, which improves estimation of true differences between

samples that would be lost in scaling to the total number of reads due to differences in

read depth[53]. Because of biases due to PCR amplification and sequencing and within

OTU-heterogeneity, we avoided making strong quantitative claims about the proportions

of spores formed by any given OTU. Instead, we examined the frequency with which an

OTU was enriched in the endospore-enriched fraction outside of Poisson noise within

individuals. More precisely, we used the counts of an OTU in the bulk community and in

the resistant fraction to estimate the Poisson variance – if the difference between counts

was within a pre-specified (we used 2 standard deviations) window of the estimated

variance, we excluded calculations on such OTUs within a sample. Next, because there

were several OTUs found in the resistant fraction that were absent from the bulk

community, we excluded calculations involving these OTUs, which would fail to provide

a finite estimate of the level of resistant fraction and would have apparently deflated

prevalence in the bulk community samples. For the remaining OTUs, an enriched OTU

was one that had more counts in the resistant fraction than in the bulk community and

an unenriched OTU was one that had fewer counts in the resistant fraction than in the

bulk community. The level of enrichment was calculated as the ratio of counts in the

resistant fraction to the sum of counts in the bulk community and in the resistant

fraction.

To compile a list of high-confidence resistant fraction-enriched organisms, we took a

similar strategy as before, but also included OTUs which had 0 counts in the bulk

community but non-zero counts in the resistant fraction. The OTUs increased in

abundance in the resistant fraction compared to the bulk community in more than half of

the samples present (excluding singletons) were included in this list.

Prevalence and abundance comparisons

To compare prevalence among OTUs in the resistant fraction and the bulk community,

we counted the number of times an OTU had non-zero abundance in one of our

subjects in either the resistant fraction or the bulk community. We used the Mann-

Whitney U test both to check for differences in read depth across the fractions and to

compare the centers of these distributions. In order to compare prevalence of OTUs

categorized as enriched to those categorized as unenriched (again excluding

singletons), we used the Mann-Whitney U test on the prevalence estimated for these

OTUs within the bulk community across the categories.

To determine the effect of abundance on prevalence estimations as above, we ran

binomial logistic regression with log-transformed normalized counts within the bulk

community as the predictor variable and endospore-enrichment status (rOTU or nOTU)

as the binary response variable. A significantly positive slope indicates that endospore-

enriched organisms are more likely than unenriched to derive from abundant organisms

in the bulk community.

Genomic Spore Gene Content

Protein sequences in *Bacillus subtilis subtilis 168* from genes identified as shared

among all spore-forming Bacilli and Clostridia[31] were downloaded from UniProt

(http://www.uniprot.org/) to make a spore gene database. All genomes as of August

2016 from 9 genera of the Clostridia in containing OTUs that were both significantly

enriched at times in the resistant fraction and significantly unenriched were downloaded

from NCBI. A standard tblastn approach was used to identify homologues in the

downloaded genomes with the corresponding genes in the spore gene database. After

identifying presence/absence of spore genes, genome spore gene profiles were

hierarchically clustered using UPGMA on the binary distance (Jaccard) matrix.

Metabolite profiling

Metabolites were measured using liquid chromatography tandem mass spectrometry

(LC-MS) method operated on a Nexera X2 U-HPLC (Shimadzu Scientific Instruments;

Marlborough, MA) coupled to a Q Exactive hybrid quadrupole orbitrap mass

spectrometer (Thermo Fisher Scientific; Waltham, MA) methods. Stool samples

(200mg/mL in 1% sodium hexametaphosphate) were homogenized using a TissueLyser

II (Qiagen). Stool homogenates (30 µL) were extracted using 90 µL of methanol

containing PGE2-d4 as an internal standard (Cayman Chemical Co.; Ann Arbor, MI)

and centrifuged (10 min, 10,000 x g, 4°C). The supernatants (2 µL) were injected onto a

150 x 2.1 mm ACQUITY UPLC BEH C18 column (Waters; Milford, MA). The column

was eluted isocratically at a flow rate: 450µL/min with 20% mobile phase A (0.1% formic

acid in water) for 3 minutes followed by a linear gradient to 100% mobile phase B

(acetonitrile with 0.1% formic acid) over 12 minutes. MS analyses were carried out

using electrospray ionization in the negative ion mode using full scan analysis

over m/z 70-850 at 70,000 resolution and 3 Hz data acquisition rate. Additional MS

settings were: ion spray voltage, -3.5 kV; capillary temperature, 320°C; probe heater

temperature, 300 °C; sheath gas, 45; auxiliary gas, 10; and S-lens RF level 60. Raw

data were processed using TraceFinder 3.3 (Thermo Fisher Scientific; Waltham, MA)

and Progenesis QI (Nonlinear Dynamics; Newcastle upon Tyne, UK) software for

detection and integration of LC-MS peaks.

Bile germination tests

Treatment of fecal samples with ethanol has previously been shown to allow culture-

based recovery of endospore-forming organisms[2]. To this end, fresh fecal samples

were homogenized in 50% ethanol (250 mg/mL), incubated for 1 hour under aerobic

conditions with shaking at 100 rpm, and washed three times (5 min, 10,000 x g) with

sterile water to remove residual ethanol. Serial dilutions from 1e-4-10% (w/v) bile bovine

oxgall (Sigma) were prepared in sterile water and 2.5 mL ethanol-treated fecal

suspension mixed in triplicate with 2.5 mL each of these bile solutions. Samples were

incubated under aerobic conditions for 2 hours at 37° C with 200 rpm shaking, and then

transferred to -80° C prior to resistant fraction extraction and 16S rDNA library

preparation.

Bile germination analysis

We transformed 16S rDNA sequencing counts generated by the bile germination tests

again using the cumulative sum-scaling transformation[53]. Under the assumption that

cells in the resistant fraction can only decrease or remain the same during treatment,

we searched for negative relationships between bile acid concentration and abundance

that would indicate and OTU had germinated. To identify significant negative

relationships, we first fit a generalized linear model (GLM) with a log-link quasi-Poisson

distribution to the normalized counts of OTUs present in the control sample with bile

acid concentration as the predictor variable. We then identified the OTU with the

strongest positive trend in the data (that with the highest positive slope and lowest p-

value). We assume that OTUs increase due only to compositional effects (that is, this

OTU has not germinated but its abundance apparently increases due to loss of other

OTUs), and we use the slope estimated from the fit of this model to detrend the other

dose-response data so as to constrain the abundance of this apparently increasing OTU

to be constant. We do so by dividing counts of all OTUs by exp(slope*bile acid

concentration), which is also a measure of the depletion of the endospore-enrichment

biomass. From this detrended dose-response data, we again fit a quasipoisson GLM

and identify putatively germinating OTUs as those having a significant ($p < 0.05$)

negative slope.

Analysis of Infant Gut Time Series

SRA files containing 16S rDNA Sequences were downloaded from Genbank under

accession no. SRA012472)[40]. Sequences were generated using a Roche 454

pyrosequencer. In order to simplify analysis of the dataset, these sequences were again

processed using the protocol outlined for processing of the original dataset in this paper.

However, sequences were quality trimmed using Q20 to 230 base pairs, and the

retained sequences were used to call 100% OTUs. OTUs were assigned taxonomies

using RDP and 100% OTUs were collapsed into taxonomic names. As very few

sequences matched between datasets when using uclust, these taxonomic names were

instead used to identify organisms as potential resistant cell-formers based on the

correspondence to the RDP-assigned taxonomic names of high confidence resistant

cell-formers identified previously. While this approach loses information given the noted

heterogeneity in resistance phenotypes even among closely related strains, the original

sequences themselves are still proxies for having this phenotype, and so the results of

such analysis must be interpreted keeping this observation in mind.

The relative abundance of organisms identified in the infant gut time series as putative

resistant-cell formers were summed, and the dynamics of this resistant cell-forming

population in the infant gut was visualized over time.

Analysis of 16S rDNA sequence files from first time and recurrent *C. difficile*

infection

The open reference 97% OTU table including RDP taxonomic annotations from

Allegretti et al 2016 was used for this analysis[43]. OTU IDs were mapped using uclust to

the corresponding genus level OTUs identified as rOTUs from this study. Patients were

grouped either as healthy, first-time *C. difficile* infection (fCDI), or recurrent *C. diffiicle*

infection (rCDI), and the fraction of rOTUs was calculated by summing their relative

abundances within each patient. A Mann Whitney U test was used to determine whether

there were significant differences in the total relative abundance of rOTUs across

groups with a Bonferroni multiple hypothesis test correction.

Analysis of 16S rDNA sequence files from fecal microbiota transplant in relapsing

***C. difficile* infection**

This dataset was obtained from Youngster et al, 2014[44]. To simplify analysis, an existing

closed-reference GreenGenes 97% OTU table generated by the original authors was

used. Closed-reference OTU IDs were mapped back to GreenGenes reference

sequences, and sequences were assigned to the resistant cell-former database

sequences again using uclust as for the adult time series.

Unique pre-FMT, post-FMT, and donor samples were separated in the dataset. We

again identified organisms that had significantly different relative abundance (Benjamini-

Hochberg adjusted Mann-Whitney U test $p < 0.05$) across the groups for our analysis.

We again obtained four categories of OTUs: nonresistant and resistant cell-formers that

were elevated in the donor and the post-FMT samples relative to the pre-FMT samples.

We used the Fisher exact test on the contingency table containing the number of OTUs

in each of the previously mentioned categories to identify whether OTU engraftment

from the donor was different across the groupings.

Analysis of 16S rDNA sequence files in adult time series pre- and post-

Salmonella Infection

Illumina HiSeq sequencing files containing 16S rDNA sequences from the stool of a

healthy adult male[45] were downloaded and processed as described for the original

dataset in this paper, except that sequences were trimmed to 101 base pairs as

described previously before calling 100% OTUs due to the use of shorter read

sequencing technology. Sequences were assigned to the resistant cell-former database

sequences using uclust constrained with the parameters: --id 99 –usersort –libonly, in

order that sequences from this dataset would be assigned only to resistant cell-formers.

In order to assess the presence of differential turnover between resistant and non-

resistant cell formers in this dataset, we identified organisms that had significantly

different relative abundance (Benjamini-Hochberg adjusted Mann-Whitney U test $p <$

0.05) before Salmonella infection starting at day 151 (days 0-150) and after the end of

infection at day 159 (days 160-252). We partitioned these OTUs into four sets for our

analysis: non-resistant and resistant cell formers whose median abundance was higher

post-infection and those whose median abundance was lower post-infection. We used

the Fisher exact test on the contingency table containing the number of OTUs in each of

the previously mentioned categories to identify whether the OTU turnover was different

across the groupings.

Supplementary Methods

Preparation of Endospores of *Bacillus subtilis*

Endospores of *B. subtilis* for testing resistant fraction protocols were prepared as

described previously[54]. *B. subtilis subtilis* strain AG174 (kindly provided by Alan

Grossman, MIT) was streaked onto nutrient agar and grown overnight. A single colony

was inoculated into 25 mL Difco Sporulation Medium (DSM), and grown at 37° C until

mid-log phase (OD600 = 0.5), and diluted 1:10 into 250 mL of pre-warmed 37° C DSM

in a 2 L flask. Cells were grown for 48 hrs at 37° C and 150 rpm under aerobic

conditions. When 90% of cells were free phase-bright spores, cultures were centrifuged

at 10,000 x g and washed twice with 200 mL 4 C sterile distilled water. The pellet was

resuspended in 200 mL cold distilled water and left at 4° C overnight, before repeating

the washing procedure. No further purification steps were used as only spores were

observed with microscopic examination. Spore viability and counts were confirmed by

germination and growth after 24 hours on nutrient agar at 37° C.

Fluorometric Quantification of Dipicolinic Acid Content

Fecal samples were thawed at 4° C and 10 mg (50 µL) suspended in 0.9 ml sodium

acetate buffer and 0.1 ml aluminium chloride ($AlCl_3$, 0.5 M) as described previously[50].

Samples were microwaved in microcentrifuge tubes for 20 s to disrupt endospores.

After cooling on ice for 10 minutes, samples were centrifuged at 10,000 x g for 10 min,

and 100 µL of the sample supernatant mixed with 100 µL terbium chloride solution

($TbCl_3$, 30 µM) in clear-bottom black 96-well microtiter plates. Fluorescence was

immediately measured using a plate reader (Synergy H1) with the following settings:

time-resolved fluorescence (delay 50 µs, interval 1200 µs) at an excitation

wavelength of 272 nm, emission wavelength of 545 nm, and 10 endpoint readings per

sample at 30°C. The number of spores in the samples was determined using standard

addition method with endospores of *B. subtilis* or dipicolinic acid. Endospore content

was expressed as equivalents of *B. subtilis*.

Fecal spore-former culture library & isolate revival

Thousands of isolates were collected from 12 healthy donor fecal samples (manuscript in preparation), and isolates from a single donor from the cohort in this study was used for all isolate experiments. Fecal material was diluted 1:15 in anaerobic PBS + 0.1% L-cysteine and homogenized in an anaerobic chamber (Coy Anaerobic Systems, 20% CO_2; 5% H_2; 75% N_2). 500 ul aliquots of the homogenized fecal material were stored in 50% glycerol at -80 C for continued culturing. Samples were treated with equivolume amounts of 100% ethanol for 1 hour, and serial dilutions were plated on Gifu Anaerobic Medium (GAM) or Brain Heart Infusion, Supplemented (BHIS) (citation). Individual colonies were selected after 24 hours of growth under anaerobic conditions at 37° C, and re-streaked onto fresh agar plates. Single colonies were grown in liquid medium, and archived in 50% glycerol at -80° C. Aliquots of liquid culture were taken for 16S rDNA Sanger Sequencing by first amplifying with the 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'-TACGGYTACCTTGTTACGACTT-3') primers, and then submitted with the U515F (5'-GTGCCAGCMGCCGCGGTAA-3') internal primer. Sequences were assigned taxonomy using the RDP database [51,52].

Testing Resistance of Actinobacteria

For revival, frozen glycerol stocks of bacteria were streaked in an anaerobic chamber onto pre-reduced (24 h) BHIS or GAM agar medium and incubated overnight under

anaerobic conditions at 37° C. For testing Actinobacteria isolates for lysis resistance,

plates with revived organisms were incubated for 4 weeks under anaerobic conditions at

room temperature to induce nutrient starvation. Plates were scraped of colonies and

colonies resuspended in 1% sodium hexametaphosphate and subjected to spore

enrichment as described previously. Results from these experiments in Supplementary

Figure 3A.

SUPPLEMENTARY RESULTS AND DISCUSSION

Quantification of fecal-resident endospores

Although endospore-forming organisms are dominant in the gastrointestinal tract, the

abundance of endospores has not been well quantified. Using a unique marker of

endospores, dipicolinic acid, others have estimated a range in human fecal samples

between 1e06 and 1e09 endospores per gram wet weight of stool[56], with a geometric

mean of 1e08. With the same protocol, we estimate the endospore burden at 1.5e07-

4.0e09 (95% CI) endospores per gram, which represents 0.03-0.7% (95% CI)

endospores per cell estimated by qPCR (Supplementary Figure 2).

Using modifications on previously described methods[26], we depleted vegetative cells in

human fecal samples to enrich for resistant cells (Fig 1A). By qPCR on this resistant

fraction, our estimate of the resistant cell fraction is slightly higher than that given by

dipicolinic acid quantification: 2e08-7e09 (95% CI) resistant cells per gram

(Supplementary Figure 2A), giving 0.2%-27% resistant cells per cell (Supplementary

Figure 2B). Culture-based studies of germinable endospores widen these estimates and

generally deplete them finding 1e05-1e09 germinable endospores per gram of stool,

giving a wide range of 0.3-140% endospores per cell. We assume that most fecal

samples contain at least 1e07 (and at most 1e12) endospores, which is roughly 100

times the infectious dose for endospores of *C. difficile*.

Most Clostridial OTUs in the resistant fraction show genetic capacity for sporulation

Many prominent gut commensals, including members of the genera *Faecalibacterium*

apparently form endospores *in vivo* and have most of the genes to do so

(Supplementary Table 1, 5, and Supplementary Figure 6), but have not been shown to

form endospores *in vitro*[57]. We believe that for *Faecalibacterium* species as for

endospores of *P. larvae* and *C. perfringens*, *in vivo* conditions drive endospore

formation, and these conditions are not readily replicated *in vitro*. Despite being an

extensively studied, numerically abundant, prevalent, and disease-relevant commensal,

endospore formation has not been demonstrated for this organism.

Clostridial OTUs in general are more shared, whether present as rOTUs or nOTUs

When restricting our analysis of prevalence to only the Clostridia, there is no difference

in sharing between Clostridia rOTUs and nOTUs (Mann Whitney U Test, p-value =

0.9952). The majority of the sharing signal is phylogenetic – when comparing

prevalence of non-Clostridia nOTUs to Clostridia nOTUs and rOTUs combined, the

difference in prevalence is very pronounced (Mann Whitney U Test, p-value = 3.0e-24).

Even Clostridia OTUs not identified as rOTUs may still contain organisms with the

potential to enter resistant states, but may not necessarily be doing so in all individuals

or at all times. Alternatively, other conserved phylogenetic traits within the Clostridia

could contribute to the widespread sharing of these organisms, but because sporulation

is required for transmission of *C. difficile*[10], it seems possible that resistance is more

broadly required for efficient transmission of the Clostridia in general.

When assessing differential sharing of rOTUs and nOTUs determined from the time

series within the cross-section, we again find that the sharing is primarily associated

with the Clostridia, as there is no significant difference in sharing among time series-

defined Clostridia nOTUs and rOTUs (Mann Whitney U Test p-value = 0.10). In part,

this result arises because there is a high degree of overlap between the OTUs defined

as rOTUs and nOTUs in the time series and in the cross-section. Although infrequent,

there are 27 inconsistent classifications (rOTUs become nOTUs and vice versa) and

268 consistent classifications between the time series and the cross-sectional data. Of

the inconsistencies, all are Firmicutes, highlighting that endospore formation is likely

driven by host- or strain-specific variability. However, it is clear that organisms linked to

endospore formation are more prevalent across unrelated hosts.

If we compare the abundance of rOTUs and nOTUs from the Clostridia, we still find that

11/24 individuals have Clostridia rOTUs at a significantly lower abundance than nOTUs

(none have significantly higher rOTU than nOTU abundance). In general, when

combining the independent tests of differences between the abundance of Clostridia

nOTUs and rOTUs, we find a significant tendency for Clostridia rOTUs to be at lower

abundance (Fisher's Method, $X^2_{df=48}$ = 289.6 p-value = 0, 0, all OTUs, Clostridia OTUs).

We wanted to determine if Clostridia drove a differential persistence signal in the time

series. Comparing the persistence of only Clostridia nOTUs and rOTUs, we find that

Clostridia nOTUs are significantly more persistent than rOTUs defined within the time

series (Wilcoxon rank sum test, p-value = 2.5e-05), and for those Clostridia nOTUs and

rOTUs shared with the cross-sectional data (Wilcoxon rank sum test, p-value = 2.4e-5).

This result suggests a tradeoff in forming resistant cells and within-host persistence in

this person. Only those Clostridia that are enriched in the resistant state have reduced

within-host persistence. Similarly, the Actinobacteria (all of which are rOTUs) are also

more ephemeral than nOTUs (p-value = 2.2e-5), suggesting a shared life history

strategy in their resistance phenotype.

Organisms in Clostridium cluster XI (including *C. sordelli*, *C. scindens,* and *C.difficile*)

link their growth with the concentration of bile acids in the colon. In the phylogenetically

diverse group of organisms encompassed by Clostridium cluster IV (including the

genera *Clostridium, Eubacterium, Ruminococcus, Coprococcus, Dorea, Lachnospira,*

Roseburia and *Butyrivibrio*), many are linked to the production of secondary bile acids,

and several also exhibit preferential growth in the mucus layer[20,58]. The enrichment for

some of these organisms among rOTUs may then derive from initiation of vegetative

growth in the small intestine through bile acid-mediate germination, growth in the

mucosa, limited growth in the lumen, and enhanced shedding of endospores into the

lumen relative to vegetative cells – driving the observation of reduced abundance and

persistence. These pieces of evidence suggested that the dynamics of rOTUs may be in

part governed by bile acids and shared niche space, and as such, we expect these

organisms to show similar dynamic behavior in other contexts.

Differences in abundance drive differences in persistence

The difference in persistence between rOTUs and nOTUs is largely attributable to a

difference in relative abundance between these groups of organisms, with rOTUs

having significantly lower median abundance than nOTUs in the time series (Mann

Whitney U test, p-value = 3.8e-5). We recall that rOTUs across people tended to have

lower median abundance than nOTUs. By comparing the abundance distributions of

rOTUs and nOTUs within individuals, we find that this result holds for 16/24 individuals

sampled (Mann Whitney U Test, p < 0.05 for each test). Combining the results from

these tests (Fisher's Method, $X^2_{df=48}$ = 412.0, p-value = 0) suggests that overall,

resistant OTUs are less abundant than non-resistant OTUs. We propose that the

reduced abundance of rOTUs accompanies reduced persistence within a host

associated with lysis resistance.

Common microbiota is more likely to source from rOTUs than nOTUs

Comparing the fractions of rOTUs and nOTUs present in each individual to assess

sharing still reveals a tendency for increased sharing across individuals (Wilcoxon rank

sum test, p-value = 4.3e-03, Supplementary Figure 3), meaning that individuals are

more likely to source their microbiota from rOTUs than nOTUs. Similarly, comparing the

frequency of appearance and disappearance of rOTUs and nOTUs in the time series

gives the same observation of reduced persistence (Mann Whitney U Test, p-value =

2.7e-13). Together, these results imply that resistance states enable increased

prevalence at the cost of within-host abundance. Because of the heterogeneity in

endospore formation within species groupings, this result weakens when grouping

OTUs at the 99% level (100% OTUs used previously), but is still significant (p-value =

8.6e-29, mean ρ = 0.083 (rOTUs), 0.047 (nOTUs)).

Metabolomic correlations to Bilophila

While there were no significant correlations between the *Bilophila* OTU and any

individual bile acid in feces measured with untargeted metabolomics, it had a significant

positive correlation (unadjusted p-value = 2.1e-06, Spearman ρ = 0.84) to a lipid marker

of milk fat consumption, pentadecanoic acid[59]. Milk fat consumption, and not

consumption of other kinds of fat, has previously been shown to lead to expansions of

Bilophila in mouse models, which were correlated with increased taurocholate

concentration in the gall bladder[36]. Thus, this particular OTU-metabolite relationship

serves as an independent verification that experimentally validated relationships play

out *in vivo.*

Introduced endospore-formers compete with endogenous endospore-formers

To further understand the dynamics of the endospore-forming fraction of the microbiota, the subject of the time series began taking tablets once daily (3e5 CFU/tablet) of a probiotic prepared as endospores (Miyarisan *Clostridium butyricum* Tablets) on day 6 of the time course. On day 8, an OTU corresponding to *C. butyricum* became detectable in both the bulk community and in the endospore fraction. These observations lead to two conclusions: (1) the endospores are capable of outgrowth *in vivo* in the presence of a complex community, and (2) the organisms are forming endospores *in vivo*, as the levels present in stool exceed the dose in the tablet. *C. butyricum* administered as endospores has previously been shown to prevent antibiotic-associated diarrhea in children (Seki et al., 2003), as well as to ameliorate DSS-induced colitis in mice through induction of IL-10 producing macrophages[60], suggesting that it grows *in vivo*, and is sufficient to block colonization by pathogenic endospore-formers such as *C. difficile*. Given these properties, we wanted to examine the dynamic behavior of this organism *in vivo*, and relate it to the activity of endogenous endospore-formers.

When examining relationships of introduced *C. butyricum* to metabolites in the stool, we found a strong anti-correlation to pentadecanoic acid (unadjusted p-value = 6.3e-05, ρ = -0.72). Recall that *Bilophila* strongly correlated with pentadecanoic acid, and that most other endospore-formers fell into the correlation cluster containing *Bilophila*. By contrast, this OTU, which had been introduced as an endospore, exhibited the opposite

behavior. We believe that this result may be a signal of competitive exclusion: because

endogenous endospore-forming commensals occupy the dominant niche, tethering their

growth to shared signals, the introduced endospore-formers are restricted to a smaller,

potentially orthogonal niche. Further work remains to determine the signals that mediate

colonization of introduced endospores.

EXTENDED DISCUSSION

Anaerobic endospore-forming organisms dominate the mammalian gastrointestinal tract

as no other studied environment. Why does this environment, compared to most others,

favor the growth of organisms with a metabolically costly and risky strategy? While it

seems obvious that this trait should increase transmission, and here we provide

evidence for this effect through our observation of increased prevalence, it is not

obvious that increased transmissibility is cost-free. We propose that a tradeoff exists in

maintaining a sporulation phenotype: endospore formation incurs diminished vegetative

growth, resulting in lower population sizes and thus decreased persistence within a host

due to a greater probability of extinction within. Because sporulation and presumably

other resistance traits seem to permit long-term survival outside of a host, this stability

provides a means for maintaining these resistant-cell formers within host populations

over time. By contrast, other strict anaerobic organisms in the gastrointestinal tract by

definition exhibit the opposite strategy: instead of investing in endospore formation, they

invest in larger population sizes that help maintain themselves within a given host.

1008

If other organisms adapted to a gut environment did not have a strategy for

transmission across hosts, we would only see endospore-formers across people. Of

course, this is not the case, so how is it that these organisms transmit? There are a

number of possibilities, but persister states may allow non-endospore-forming

organisms to enter a metabolically inert state upon exit from the gastrointestinal tract.

Toxin-antitoxin systems, associated with persistence in *E. coli*, are overrepresented in

Bacteroidetes, Alpha- and Gammaproteobacteria[61], and Bacteroidetes are among the

most metabolically inactive cells in human fecal samples[62]. However, we also observe

that among the non-Clostridia nOTUs, abundance is higher within hosts, suggesting that

simply by maintaining high population levels, these organisms increase their ability to

transmit to new hosts. When host populations are high density, gut residents do not

require long-term environmental survival; the probability of encountering a new host will

be high. However, when host populations are low density, long-term environmental

survival becomes a necessary strategy. Over longer evolutionary timescales,

endospore-formers should be selected for in hosts with low population densities.

1024

Resistant cells appeared to be primed to enter uncolonized gastrointestinal tracts. There

may be a few reasons for this observation. Environmental resistance enables them to

survive for long periods of time without diminished viability. Specific molecular indicators

of a suitable host, such as bile acids, may permit germination and subsequent

outgrowth under the right host conditions. Outgrowth seems to require strict anaerobic

conditions and low reduction potentials as evidenced by the entrainment of rOTUs to

the dynamic behavior of the strictly anaerobic sulfite-reducer, *Bilophila*. Similarly, growth

within the gastrointestinal tract permits sporulation for a number of these organisms as

well, giving them an exit strategy if host conditions are sub-optimal.

1034

We demonstrate that even though resistant cells themselves appear to be lowly

abundant in human fecal material (on the order of 0.1-10% total biomass), sharing of

rOTUs and the Clostridia in particular is favored across unrelated individuals. The

increased sharing comes at the cost of vegetative cell abundance and persistence

within a host. Because of heterogeneity in the resistance phenotype even within strains

of the same species, few OTUs consistently form resistant cells at high levels across all

individuals in our dataset. This heterogeneity drives a signal where organisms forming

these resistant cells within any given person may not be doing so in other individuals,

yet when considering endospore-forming potential as a whole (i.e. through its

phylogenetic conservation within the Clostridia), OTUs with organisms that potentially

can form endospores are more likely than others to be found among many unrelated

individuals. Within a person, the organisms we observe forming resistant cells exhibit

distinct correlated behavior coupled to the dynamic behavior of select non-endospore-

forming taxa that likely respond to shared environmental signals.

1049

1050

1051

Supplementary Figure 1. Resistant OTUs are more prevalent across individuals in

their bulk communities. The prevalence of OTUs across 24 individuals in our cohort

as a function of fraction of times the OTU was enriched in the resistant fraction relative

to the bulk community. The area of a point corresponds to the number of observations

of OTUs at a given fractional enrichment, fractional sharing pair. Pink points correspond

to nOTUs (never enriched in the endospore fraction) and blue points correspond to

rOTUs (those ever enriched in the endospore fraction). The histogram on top of the plot

relates the density of observations at each fractional enrichment level; the histogram on

the side of the plot relates the density of observations for nOTUs (pink) and rOTUs

(blue) across the sharing gradient.

1063

Supplementary Figure 2. Resistant cells generally comprise fewer than 2% of

cells in the gut microbiota (A) Resistant cell abundance sorted by fraction of total cells

estimated using qPCR on the resistant fraction quantified with a DNA standard curve of

a mock community consisting of cultured isolates of *E. coli*, *L. reuteri*, *V. parvula*, *B.*

fragilis, and *C. difficile*. (B) Endospore relative abundance as the fraction of total cells

estimated using qPCR, sorted from highest to lowest.

Isolate Tested

of *B. subtilis* endospores added to sample

Supplementary Figure 3. Cultured Actinobacteria isolated from the human gut are

significantly depleted when enriching for resistant cells, but endospores are

reliably recovered. (A) Fraction of DNA remaining after resistant cell enrichment is

applied to cultured Actinobacteria. Fractions are estimated by qPCR for four human gut-

derived isolates from the phylum Actinobacteria calculated by comparing the threshold

cycle (Ct) for amplification of DNA using the V4 16S rDNA primers used for library

assembly. Ct is converted to DNA concentration using a standard curve of DNA

dilutions for each isolate. Error bars are the standard error of the mean. (B) Estimate of

the recovery of endospores of *B. subtilis* added at different amounts into independent

fecal samples (individual points) and treated with standard DNA extraction (blue

diamonds) or resistant fraction (red squares). Endospore recovery is estimated using

qPCR with primers targeting *aprE* (*aprE* F: 5'-TTTACGATGGCGTTCAGCAAC-3', *aprE*

R: 5'-GGAAGTGCCTTCATTTCCGGCT-3')[63]. The line y = x indicates stoichiometric

recovery, and the horizontal dashed line shows the limit of detection of the primer set.

1085

Supplementary Figure 4. Time series of two distinct _Ruminococcus_ OTUs show

discordance in resistant fraction tendency. (A) Bulk community and resistant fraction

time series for a _Ruminococcus_ OTU with corresponding V4 16S region matching at

100% sequence identity to _Ruminococcus lactaris_ ATCC 29176. (B) Bulk community

and resistant fraction time series for a _Ruminococcus_ OTU with corresponding V4 16S

region matching at 97% sequence identity to _Ruminococcus faecis_ strain Eg2.

1092

1093

1094

Supplementary Figure 5. Variably endospore-enriched taxa lack several genes shared between previously identified endospore-forming Bacilli and Clostridia. Sporulation genes (y-axis) considered conserved between all endospore-forming *Bacilli* and *Clostridia*[31] were mapped to genomes of genera (using tblastn with an E-value cutoff of 1e-5) containing OTUs observed as significantly increased and decreased in the resistant fraction across individuals measured by Mann Whitney U test (FDR adjusted p-value < 0.10). Genomes on the x-axis are all genomes corresponding to the named genera downloaded from the NCBI database as of August 2017. Color bar on the x-axis corresponds to genera. Genomes are clustered using UPGMA hierarchical clustering on presence-absence profiles of sporulation gene content; genes profiles

1105   across genomes are clustered using the same procedure. Anaerotruncus colihomininis

1106   (gray) is a known endospore-former that lacks several putatively conserved genes.

1107 **Supplementary Table 3. Significant (unadjusted p < 0.05) Spearman Correlations**

1108 **between metabolites and rOTU cluster.**

Metabolite Name	m/z	Spearman Rho	Unadjusted p-value	FDR q-value
sebacate	201.1119	0.74	5.64E-05	3.89E-03
arachidonate	303.2321	-0.593	2.72E-03	3.13E-02
docosapentaenoate	329.2476	-0.582	3.39E-03	3.13E-02
eicosapentaenoate	301.2164	-0.58	3.51E-03	3.13E-02
nonadecanoate	297.279	-0.578	3.63E-03	3.13E-02
pentadecanoate	241.2162	0.591	2.81E-03	3.13E-02
3-oxooctadecanoate	297.2427	0.619	1.58E-03	3.13E-02
docosahexaenoate	327.232	-0.565	4.63E-03	3.55E-02
hydrocinnamate	149.0592	0.558	5.25E-03	3.62E-02
undecanedionate	215.1276	0.617	1.67E-03	3.62E-02
taurocholate	514.2825	-0.549	6.23E-03	3.91E-02
crustecdysone	525.3053	-0.608	2.37E-02	1.09E-01
adrenate	331.2633	-0.474	2.04E-02	1.09E-01
10-nonadecanoate	295.2633	-0.473	2.07E-02	1.09E-01
dimethylurate	195.0511	0.469	2.23E-02	1.09E-01
eicosatrienoate	305.2477	-0.438	3.33E-02	1.44E-01
arachidate	311.2947	-0.424	3.99E-02	1.62E-01
urobilin	593.3324	0.417	4.35E-02	1.67E-01

1109

1110

REFERENCES

1. Filippidou, S. *et al.* Under-detection of endospore-forming Firmicutes in metagenomic data. *Comput. Struct. Biotechnol. J.* **13,** 299–306 (2015).

2. Browne, H. P. *et al.* Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation. *Nature* **533,** 543–546 (2016).

3. Ley, R. E. *et al.* Evolution of Mammals and Their Gut Microbes. *Science* **320,** 1647–1651 (2008).

4. Angert, E. R. & Losick, R. M. Propagation by sporulation in the guinea pig symbiont Metabacterium polyspora. *Proc. Natl. Acad. Sci.* **95,** 10218–10223 (1998).

5. Flint, J. F., Drzymalski, D., Montgomery, W. L., Southam, G. & Angert, E. R. Nocturnal Production of Endospores in Natural Populations of Epulopiscium-Like Surgeonfish Symbionts. *J. Bacteriol.* **187,** 7460–7470 (2005).

6. Alexander, C. J., Citron, D. M., Brazier, J. S. & Goldstein, E. J. Identification and antimicrobial resistance patterns of clinical isolates of Clostridium clostridioforme, Clostridium innocuum, and Clostridium ramosum compared with those of clinical isolates of Clostridium perfringens. *J. Clin. Microbiol.* **33,** 3209–3215 (1995).

7. Paredes-Sabja, D., Torres, J. A., Setlow, P. & Sarker, M. R. Clostridium perfringens Spore Germination: Characterization of Germinants and Their Receptors. *J. Bacteriol.* **190,** 1190–1201 (2008).

8. Sokol, H. *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci.* **105,** 16731–16736 (2008).

9. Png, C. W. *et al.* Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. *Am. J. Gastroenterol.* **105,** 2420–2428 (2010).

10. Deakin, L. J. *et al.* The *Clostridium difficile spo0A* Gene Is a Persistence and Transmission Factor. *Infect. Immun.* **80,** 2704–2711 (2012).

11. Atarashi, K. *et al.* Induction of Colonic Regulatory T Cells by Indigenous Clostridium Species. *Science* **331,** 337–341 (2011).

12. Atarashi, K. *et al.* Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature* **500,** 232–236 (2013).

13. Stefka, A. T. *et al.* Commensal bacteria protect against food allergen sensitization. *Proc. Natl. Acad. Sci.* **111,** 13145–13150 (2014).

14. Kim, Y.-G. *et al.* Neonatal acquisition of Clostridia species protects against colonization by bacterial pathogens. *Science* **356,** 315 (2017).

15. Ivanov, I. I. *et al.* Induction of Intestinal Th17 Cells by Segmented Filamentous Bacteria. *Cell* **139,** 485–498 (2009).

16. Kuwahara, T. *et al.* The Lifestyle of the Segmented Filamentous Bacterium: A Non-Culturable Gut-Associated Immunostimulating Microbe Inferred by Whole-Genome Sequencing. *DNA Res.* **18,** 291–303 (2011).

17. Sczesnak, A. *et al.* The Genome of Th17 Cell-Inducing Segmented Filamentous Bacteria Reveals Extensive Auxotrophy and Adaptations to the Intestinal Environment. *Cell Host Microbe* **10,** 260–272 (2011).

18. Smith, P. M. *et al.* The Microbial Metabolites, Short-Chain Fatty Acids, Regulate Colonic Treg Cell Homeostasis. *Science* **341,** 569–573 (2013).

19. Furusawa, Y. *et al.* Commensal microbe-derived butyrate induces the differentiation

of colonic regulatory T cells. *Nature* **504,** 446–450 (2013).

20. Van den Abbeele, P. *et al.* Butyrate-producing Clostridium cluster XIVa species

specifically colonize mucins in an in vitro gut model. *ISME J.* **7,** 949–961 (2013).

21. Eeckhaut, V. *et al.* Butyrate production in phylogenetically diverse Firmicutes

isolated from the chicken caecum: Butyrate-producing bacteria from the chicken

caecum. *Microb. Biotechnol.* **4,** 503–512 (2011).

22. Louis, P., Young, P., Holtrop, G. & Flint, H. J. Diversity of human colonic butyrate-

producing bacteria revealed by analysis of the butyryl-CoA:acetate CoA-transferase

gene. *Environ. Microbiol.* **12,** 304–314 (2010).

23. Yano, J. M. *et al.* Indigenous Bacteria from the Gut Microbiota Regulate Host

Serotonin Biosynthesis. *Cell* **161,** 264–276 (2015).

24. Kaplan, I. & Williams, J. W. Spore Formation among the Anaerobic Bacteria: I. The

Formation of Spores by Clostridium sporogenes in Nutrient Agar Media. *J. Bacteriol.*

42, 265 (1941).

25. Dingman, D. W. & Stahly, D. P. Medium promoting sporulation of Bacillus larvae and

metabolism of medium components. *Appl. Environ. Microbiol.* **46,** 860–869 (1983).

26. Wunderlin, T., Junier, T., Roussel-Delif, L., Jeanneret, N. & Junier, P. Endospore-

enriched sequencing approach reveals unprecedented diversity of *Firmicutes* in

sediments: Endospore-forming enrichment. *Environ. Microbiol. Rep.* **6,** 631–639

(2014).

27. Sekar, R. *et al.* An Improved Protocol for Quantification of Freshwater Actinobacteria

by Fluorescence In Situ Hybridization. *Appl. Environ. Microbiol.* **69,** 2928–2935

(2003).

28. Fahlgren, A., Hammarström, S., Danielsson, Å. & HAMMARSTRÖM, M.-L.

Increased expression of antimicrobial peptides and lysozyme in colonic epithelial

cells of patients with ulcerative colitis. *Clin. Exp. Immunol.* **131,** 90–101 (2003).

29. Keshav, S., Chung, P., Milon, G. & Gordon, S. Lysozyme is an inducible marker of

macrophage activation in murine tissues as demonstrated by in situ hybridization. *J.*

Exp. Med. **174,** 1049 (1991).

30. Gueimonde, M., Laitinen, K., Salminen, S. & Isolauri, E. Breast milk: a source of

bifidobacteria for infant gut development and maturation? *Neonatology* **92,** 64–66

(2007).

31. Galperin, M. Y. *et al.* Genomic determinants of sporulation in *Bacilli* and *Clostridia*:

towards the minimal set of sporulation-specific genes: Distribution of sporulation

genes in *Bacilli* and *Clostridia*. *Environ. Microbiol.* **14,** 2870–2890 (2012).

32. Popham, D. L., Illades-Aguiar, B. & Setlow, P. The Bacillus subtilis dacB gene,

encoding penicillin-binding protein 5*, is part of a three-gene operon required for

proper spore cortex synthesis and spore core dehydration. *J. Bacteriol.* **177,** 4721–

4729 (1995).

33. Orsburn, B., Sucre, K., Popham, D. L. & Melville, S. B. The SpmA/B and DacF

proteins of Clostridium perfringens play important roles in spore heat resistance.

FEMS Microbiol. Lett. **291,** 188–194 (2009).

34. Serra, C. R., Earl, A. M., Barbosa, T. M., Kolter, R. & Henriques, A. O. Sporulation

during Growth in a Gut Isolate of Bacillus subtilis. *J. Bacteriol.* **196,** 4184–4196

(2014).

35. Francis, M. B., Allen, C. A., Shrestha, R. & Sorg, J. A. Bile Acid Recognition by the

Clostridium difficile Germinant Receptor, CspC, Is Important for Establishing

Infection. *PLoS Pathog.* **9,** e1003356 (2013).

36. Devkota, S. *et al.* Dietary-fat-induced taurocholic acid promotes pathobiont

expansion and colitis in Il10–/– mice. *Nature* (2012). doi:10.1038/nature11225

37. Sorg, J. A. & Sonenshein, A. L. Inhibiting the Initiation of Clostridium difficile Spore

Germination using Analogs of Chenodeoxycholic Acid, a Bile Acid. *J. Bacteriol.* **192,**

4983–4990 (2010).

38. Ceuppens, S. *et al.* Survival and Germination of Bacillus cereus Spores without

Outgrowth or Enterotoxin Production during In Vitro Simulation of Gastrointestinal

Transit. *Appl. Environ. Microbiol.* **78,** 7698–7705 (2012).

39. Shah, I. M., Laaberki, M.-H., Popham, D. L. & Dworkin, J. A eukaryotic-like Ser/Thr

kinase signals bacteria to exit dormancy in response to peptidoglycan fragments.

Cell **135,** 486–496 (2008).

40. Koenig, J. E. *et al.* Succession of microbial consortia in the developing infant gut

microbiome. *Proc. Natl. Acad. Sci.* **108,** 4578–4585 (2011).

41. Chandan, R. C., Shahani, K. M. & Holly, R. G. Lysozyme Content of Human Milk.

Nature **204,** 76 (1964).

42. Nayfach, S., Rodriguez-Mueller, B., Garud, N. & Pollard, K. S. An integrated

metagenomics pipeline for strain profiling reveals novel patterns of bacterial

transmission and biogeography. *Genome Res.* **26,** 1612–1625 (2016).

43. Allegretti, J. R. *et al.* Recurrent *Clostridium difficile* infection associates with distinct

bile acid and microbiome profiles. *Aliment. Pharmacol. Ther.* **43,** 1142–1153 (2016).

44. Youngster, I. *et al.* Fecal Microbiota Transplant for Relapsing Clostridium difficile

Infection Using a Frozen Inoculum From Unrelated Donors: A Randomized, Open-

Label, Controlled Pilot Study. *Clin. Infect. Dis.* **58,** 1515–1522 (2014).

45. David, L. A. *et al.* Host lifestyle affects human microbiota on daily timescales.

Genome Biol **15,** R89 (2014).

46. Bueche, M. *et al.* Quantification of Endospore-Forming Firmicutes by Quantitative

PCR with the Functional Gene spo0A. *Appl. Environ. Microbiol.* **79,** 5302–5312

(2013).

47. Chomczynski, P. & Rymaszewski, M. Alkaline polyethylene glycol-based method for

direct PCR from bacteria, eukaryotic tissue samples, and whole blood.

Biotechniques **40,** 454 (2006).

48. Preheim, S. P., Perrotta, A. R., Martin-Platero, A. M., Gupta, A. & Alm, E. J.

Distribution-Based Clustering: Using Ecology To Refine the Operational Taxonomic

Unit. *Appl. Environ. Microbiol.* **79,** 6593–6603 (2013).

49. Fierer, N., Jackson, J. A., Vilgalys, R. & Jackson, R. B. Assessment of soil microbial

community structure by use of taxon-specific quantitative PCR assays. *Appl.*

Environ. Microbiol. **71,** 4117–4120 (2005).

50. Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate Illumina

Paired-End reAd mergeR. *Bioinformatics* **30,** 614–620 (2014).

51. Maidak, B. L. *et al.* The ribosomal database project (RDP). *Nucleic Acids Res.* **24,**

82–85 (1996).

52. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for

rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl.*

Environ. Microbiol. **73,** 5261–5267 (2007).

53. Paulson, J. N., Stine, O. C., Bravo, H. C. & Pop, M. Differential abundance analysis

for microbial marker-gene surveys. *Nat. Methods* **10,** 1200–1202 (2013).

54. Nicholson, W. & Setlow, P. Purification of spores on step gradients of Renografin. in

Molecular Biological Methods for Bacillus 391–450 (John Wiley, 1990).

55. Fichtel, J., Köster, J., Rullkötter, J. & Sass, H. Spore dipicolinic acid contents used

for estimating the number of endospores in sediments: Determination of endospores

in sediments. *FEMS Microbiol. Ecol.* **61,** 522–532 (2007).

56. Khanna, S. *et al.* A Novel Microbiome Therapeutic Increases Gut Microbial Diversity

and Prevents Recurrent *Clostridium difficile* Infection. *J. Infect. Dis.* **214,** 173–181

(2016).

57. Lopez-Siles, M. *et al.* Cultured Representatives of Two Major Phylogroups of Human

Colonic Faecalibacterium prausnitzii Can Utilize Pectin, Uronic Acids, and Host-

Derived Substrates for Growth. *Appl. Environ. Microbiol.* **78,** 420–428 (2012).

58. Lopetuso, L. R., Scaldaferri, F., Petito, V. & Gasbarrini, A. Commensal Clostridia:

leading players in the maintenance of gut homeostasis. *Gut Pathog.* **5,** 1 (2013).

59. Smedman, A. E., Gustafsson, I.-B., Berglund, L. G. & Vessby, B. O. Pentadecanoic

acid in serum as a marker for intake of milk fat: relations between intake of milk fat

and metabolic risk factors. *Am. J. Clin. Nutr.* **69,** 22–29 (1999).

60. Hayashi, A. *et al.* A Single Strain of Clostridium butyricum Induces Intestinal IL-10-

Producing Macrophages to Suppress Acute Experimental Colitis in Mice. *Cell Host*

Microbe **13,** 711–722 (2013).

61. Makarova, K. S., Wolf, Y. I. & Koonin, E. V. Comprehensive comparative-genomic

analysis of Type 2 toxin-antitoxin systems and related mobile stress response

systems in prokaryotes. *Biol. Direct* **4,** 19 (2009).

62. Maurice, C. F., Haiser, H. J. & Turnbaugh, P. J. Xenobiotics Shape the Physiology

and Gene Expression of the Active Human Gut Microbiome. *Cell* **152,** 39–50 (2013).

63. Sadeghi, A., Mortazavi, S. A., Bahrami, A. R. & Sadeghi, B. Design of multiplex PCR

for simultaneous detection of rope‐forming Bacillus strains in Iranian bread dough.

J. Sci. Food Agric. **92,** 2652–2656 (2012).

1278