Transgenic mouse lines expressing the 3xFLAG-dCas9 protein for enChIP

analysis

Toshitsugu Fujita, Fusako Kitaura, Asami Oji, Naoki Tanigawa, Miyuki Yuno, Masahito Ikawa,

Ichiro Taniuchi, and Hodaka Fujii*

Toshitsugu Fujita, Hodaka Fujii

Department of Biochemistry and Genome Biology, Hirosaki University Graduate School of

Medicine, Zaifu-cho 5, Hirosaki, Aomori 036-8562, Japan

Toshitsugu Fujita, Fusako Kitaura, Naoki Tanigawa, Miyuki Yuno, Hodaka Fujii

Chromatin Biochemistry Research Group, Combined Program on Microbiology and Immunology,

Research Institute for Microbial Diseases, Osaka University, Yamadaoka 3-1, Suita, Osaka

565-0871, Japan

Asami Oji, Masahito Ikawa

Genome Information Research Center and Department of Experimental Genome Research,

Research Institute for Microbial Diseases, Osaka University, Yamadaoka 3-1, Suita, Osaka

565-0871, Japan

Ichiro Taniuchi

Laboratory for Transcriptional Regulation, RIKEN Center for Integrative Medical Sciences,

1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa 230-0045, Japan

Short title: Transgenic mice expressing 3xFLAG-dCas9

Keywords: enChIP, dCas9, ChIP, chromatin immunoprecipitation, transgenic mouse

*Corresponding author

E-mail: hodaka@hirosaki-u.ac.jp

Department of Biochemistry and Genome Biology, Hirosaki University Graduate School of

Medicine, Zaifu-cho 5, Hirosaki, Aomori, 036-8562 Japan

Abstract

We developed the engineered DNA-binding molecule-mediated chromatin immunoprecipitation

(enChIP) technology to isolate specific genomic regions while retaining their molecular

interactions. In enChIP, the locus of interest is tagged with an engineered DNA-binding molecule,

such as a modified form of the clustered regularly interspaced short palindromic repeats

(CRISPR) system containing a guide RNA (gRNA) and a catalytically inactive form of Cas9

(dCas9). The locus is then affinity-purified to enable identification of associated molecules. In

this study, we generated transgenic mice expressing 3xFLAG-tagged *Streptococcus pyogenes*

dCas9 (3xFLAG-dCas9) and retrovirally transduced gRNA into primary CD4$^+$ T cells from these

mice for enChIP. Using this approach, we achieved high yields of enChIP at the targeted genomic

region. Our novel transgenic mouse lines provide a valuable tool for enChIP analysis in primary

mouse cells.

Introduction

Identification of molecules associated with a genomic region of interest *in vivo* is an essential step in understanding the regulatory mechanisms underlying that region's functions. To this end, we previously developed engineered DNA-binding molecule-mediated chromatin immunoprecipitation (enChIP) technology to isolate genomic regions of interest along with their interacting molecules (Fujita *et al.* 2013; Fujita & Fujii 2013). In enChIP, the locus of interest is tagged with engineered DNA-binding molecules such as transcription activator-like (TAL) proteins (Bogdanove & Voytas 2011) or a variant of the clustered regularly interspaced short palindromic repeats (CRISPR) system (Harrison *et al.* 2014; Wright *et al.* 2016) containing a guide RNA (gRNA) and a catalytically inactive form of Cas9 (dCas9). The tagged locus is then affinity-purified to enable identification of associated molecules. Locus-tagging can be achieved in cells by expressing engineered DNA-binding molecules (Fujita *et al.* 2013; Fujita & Fujii 2013, 2014b; Fujita *et al.* 2015; Fujita *et al.* 2016a; Fujita *et al.* 2017b), or *in vitro* using recombinant or synthetic engineered DNA-binding molecules (Fujita & Fujii 2014a; Fujita *et al.* 2016b; Fujita *et al.* 2017a) Combination of enChIP with mass spectrometry (MS), RNA sequencing, and next-generation sequencing (NGS) enables identification of proteins (Fujita *et al.* 2013; Fujita & Fujii 2013, 2014b), RNAs (Fujita *et al.* 2015), and other genomic regions (Fujita *et al.* 2017a; Fujita *et al.* 2017b) that interact with specific loci of interest in a non-biased manner.

To perform locus-tagging in primary cells, it is necessary to express both dCas9 and gRNA by transduction or other methods. However, the low transduction efficiency of some cell lineages results in a low percentage of cells expressing both dCas9 and gRNA. To resolve this technical issue, we generated transgenic (Tg) mouse lines expressing 3xFLAG-tagged *Streptococcus*

69 *pyogenes* dCas9 (3xFLAG-dCas9), either constitutively or inducibly. To facilitate their use in

70 various experimental contexts, expression of the tagged dCas9 and/or a reporter green fluorescent

71 protein (GFP) can be flexibly induced or abolished by Cre- or FLPe-mediated site-specific

72 recombination events that delete expression-modulating cassettes. We anticipate that these novel

73 Tg mouse lines will serve as a powerful tool for efficient enChIP analysis in primary cells.

Results and Discussion

Generation of Tg mouse lines expressing 3xFLAG-dCas9

To facilitate enChIP analysis using primary mouse cells, we generated two Tg mouse lines expressing 3xFLAG-dCas9 (Fig. 1A, B). One line, 3xFLAG-dCas9-IRES-EGFP, harbors 3xFLAG-dCas9 and IRES-EGFP in the *Rosa26* locus (Fig. 1A). In the other line, 3xFLAG-dCas9 and IRES-EGFP are present at the *Rosa26* locus, but expression of 3xFLAG-dCas9 can be induced by Cre-mediated deletion of the STOP cassette (along with the neo^r gene), and EGFP expression can be disrupted by FRT-mediated deletion of the IRES-EGFP cassette (Fig. 1B (i)). 3xFLAG-dCas9/CTV (neo$^+$/ GFP$^+$) mice were crossed with CAG-Cre mice (Matsumura *et al.* 2004) to delete the STOP cassette and neo^r gene, yielding 3xFLAG-dCas9/CTV (GFP$^+$) mice (Fig. 1B (ii)). 3xFLAG-dCas9/CTV (GFP$^+$) mice can be further crossed with CAG-FLPe mice (Schaft *et al.* 2001) to delete the IRES-EGFP cassette, yielding 3xFLAG-dCas9/CTV mice (Fig. 1B (iii)). Targeted integration of transgenes was confirmed by PCR (Fig. 1C). All mice were viable and fertile with normal litter sizes and did not exhibit any morphological abnormalities. Expression of EGFP was observed throughout the body, including thymocytes and splenocytes (Fig. 2A), and 3xFLAG-dCas9 was detected in NE of thymocytes (Fig. 2B).

enChIP analysis using primary CD4$^+$ T cells

Next, we performed enChIP analysis using primary cells from the Tg mice. For these experiments, CD4$^+$ T cells were purified from 3xFLAG-dCas9-IRES-EGFP mice and 3xFLAG-dCas9/CTV (GFP$^+$) mice and activated with anti-CD3 and anti-CD28 Abs. The activated cells were

transduced with a retroviral vector expressing gRNA targeting the c-*myc* promoter (m-c-*myc*

gRNA #1/pSIR-hCD2) or a negative control vector (pSIR-hCD2), and 2 days later, hCD2$^+$ cells

were isolated and expanded in media containing IL-2. Cells were fixed with formaldehyde and

subjected to enChIP analysis using anti-FLAG Ab. Yields of enChIP were monitored by real-time

PCR. As shown in Figure 3, efficient enrichment of the c-*myc* promoter region, but not an

irrelevant locus (*Gapdh*), was detected in samples expressing gRNA targeting the c-*myc* locus.

By contrast, no enrichment was observed for samples in the absence of gRNA. The yields of

enChIP were comparable between 3xFLAG-dCas9-IRES-EGFP mice and 3xFLAG-dCas9/CTV

(GFP$^+$) mice. These results demonstrate that primary cells from these Tg mice can be used for

enChIP analysis.

Conclusions

We generated Tg mouse lines expressing 3xFLAG-tagged dCas9 and retrovirally transduced

gRNA targeting a genomic locus into primary CD4$^+$ T cells from these mice. Using this approach,

high yields of enChIP could be achieved. Thus, these Tg mouse lines represent a useful tool for

enChIP analysis in primary mouse cells. In addition, the Tg mouse strains generated in this study

could be used for CRISPR interference (CRISPRi) experiments (Qi *et al.* 2013) using primary

mouse cells.

Experimental procedures

Plasmids

To construct pCAG1.2-PM, a modified pCAGGS plasmid (Niwa *et al.* 1991) was digested with

*Sac*I. Two oligonucleotides, *Mlu*I–*Pme*I oligo-S (27551) and *Mlu*I–*Pme*I oligo-A (27552), were

annealed, phosphorylated, and inserted into the digested plasmid, yielding two plasmids,

pCAG1.2-PM (*Pme*I–*Mlu*I) and pCAG1.2-MP (*Mlu*I–*Pme*I), distinguished by the orientations of

the oligonucleotides. To construct 3xFLAG-dCas9/pCAG1.2-PM, pCAG1.2-PM was digested

with *Eco*RV and *Not*I. 3xFLAG-dCas9/pMXs-puro (Addgene #51240) was digested with *Pac*I,

blunted, and further digested with *Not*I. The vector backbone and the coding sequence of

3xFLAG-dCas9 were purified by agarose gel electrophoresis and ligated. To construct

3xFLAG-dCas9-IG/pCAG1.2-PM, 3xFLAG-dCas9/pCAG1.2-PM and 3xFLAG-dCas9/pMXs-IG

(Addgene #51258) were digested with *Not*I and *Sal*I, respectively. After blunting, the plasmids

were further digested with *Asc*I. The vector backbone and the coding sequence of

3xFLAG-dCas9 were purified by agarose gel electrophoresis and ligated. To construct

3xFLAG-dCas9-IG/pSKII-ROSA, pSKII-ROSA26arm0.5kb-zeo was digested with *Bam*HI and

*Sal*I, and 3xFLAG-dCas9-IG/pCAG1.2-PM was digested with *Mlu*I and *Pac*I. After blunting, the

vector backbone and the coding sequence of 3xFLAG-dCas9 were purified by agarose gel

electrophoresis and ligated.

To construct 3xFLAG-dCas9/CTV, CTV (Addgene #15912) was digested with *Asc*I, and

3xFLAG-dCas9/pMXs-puro was digested with *Pac*I and *Not*I. After blunting, the vector

backbone and the coding sequence of 3xFLAG-dCas9 were purified by agarose gel

electrophoresis and ligated.

To construct a gBlock targeting the mouse c-*myc* promoter, the oligonucleotides mcMYC

promoter 1 sense (27822) and mcMYC promoter 1 antisense (27823) were annealed,

phosphorylated, and inserted into gRNA cloning vector *Bbs*I ver. 1, which was digested with

*Bbs*I. To construct m-c-*myc* gRNA #1/pSIR, the gBlock was excised with *Xho*I and *Hin*dIII and

inserted into *Xho*I/*Hin*dIII-cleaved pSIR (Clontech). To construct a retroviral vector for

expression of gRNA against the mouse c-*myc* promoter (m-c-*myc* gRNA #1/pSIR-hCD2),

pSIR-hCD2 (Addgene #51143) and m-c-*myc* gRNA #1/pSIR were digested with *San*DI and

*Hin*dIII. The vector backbone and insert were purified by agarose gel electrophoresis and ligated.

Oligonucleotides used for construction of plasmids are shown in Table 1.

Mice

Embryonic stem (ES) cells (EGR-101) (Fujihara *et al.* 2013) were transfected with linearized

3xFLAG-dCas9-IG/pSKII-ROSA, as described previously (Oji *et al.* 2016). ES cells retaining the

transgene in the *Rosa26* locus were injected into blastocysts (ICR × ICR) to generate chimeric

mice. The chimeric mice were crossed with B6D2F1 mice to generate heterozygous

3xFLAG-dCas9-IRES-EGFP mice (strain name:

B6;D2-Gt(ROSA)26Sor<em1(3×FLAG-dCas9-EGFP)Osb>).

B6JN/1 ES cells (Moriyama *et al.* 2014), derived from a hybrid between C57BL/6J and

C57BL/6N, were electroporated with 3xFLAG-dCas9/CTV plasmid linearized with *Asi*SI. To

generate chimeric mice, ES cells containing the *neo^r* gene, transgene, and *GFP* gene in the

Rosa26 locus were injected into blastocysts (Balb/c × Balb/c) by the animal facility group at

RIKEN IMS. The chimeric mice were crossed with C57BL/6 mice to generate heterozygous

3xFLAG-dCas9/CTV (neo$^+$/GFP$^+$) mice (strain name:

C57BL/6-Gt(ROSA)26Sor<tm1(CAG-3xFLAG-dCas9-EGFP)Hfuj>). The 3xFLAG-dCas9/CTV

(neo$^+$/GFP$^+$) mice were crossed with CAG-Cre mice (Matsumura *et al.* 2004) to generate

3xFLAG-dCas9/CTV (GFP$^+$) mice (strain name:

C57BL/6-Gt(ROSA)26Sor<tm1.1(CAG-3xFLAG-dCas9-EGFP)Hfuj>). The

3xFLAG-dCas9/CTV (GFP$^+$) mice can be crossed with CAG-FLPe mice (Schaft *et al.* 2001) to

generate 3xFLAG-dCas9/CTV mice (strain name:

C57BL/6-Gt(ROSA)26Sor<tm1.2(CAG-3xFLAG-dCas9)Hfuj>).

All animal experiments were approved by the Institutional Animal Care and Use Committee at

Research Institute for Microbial Diseases, Osaka University.

Genotyping

For genotyping, genomic DNA was extracted and subjected to PCR with KOD FX (Toyobo).

PCR conditions were as follows: heating at 94°C for 2 min; followed by 38 cycles of 98°C for 10

s, 62°C for 30 s, and 68°C for 4 min. Primers used for genotyping PCR are shown in Table 1.

Cell staining and flow cytometry

Thymi and spleens were isolated from euthanized mice and used to prepare single cells. For

surface staining, cells were stained for 30 minutes at 4°C with fluorochrome-conjugated

antibodies (Abs): fluorescein isothiocyanate (FITC)-conjugated mouse CD4 (130-102-541,

Miltenyi) and phycoerythrin (PE)-conjugated human CD2 (hCD2) (347597, BD Bioscience).

Flow cytometric analysis was performed on a FACSCalibur (BD Biosciences) and analyzed with

the FlowJo software (TreeStar).

Immunoblot analysis

Nuclear extracts (NE) were prepared with NE-PER Nuclear and Cytoplasmic Extraction Reagents

(Thermo Fisher Scientific). Aliquots of NE (10 µg) were subjected to immunoblot analysis with

anti-FLAG M2 Ab (F1804, Sigma-Aldrich), as described previously (Fujita & Fujii 2013).

Transduction of gRNA and isolation of transduced cells

Transduction of pSIR-hCD2 or m-c-*myc* gRNA #1/pSIR-hCD2 into primary CD4[+] T cells was

performed as described previously (Naoe *et al.* 2007). Briefly, CD4[+] T cells were purified from

spleens of the Tg mice using the Mouse CD4[+] T cell isolation kit (Miltenyi, 130-104-454).

Purified CD4[+] T cells were activated with anti-CD3 Ab (3 µg/ml, 145-2C11, 553057, BD

Pharmingen) and anti-CD28 Ab (3 µg/ml, 37.51, 553295, BD Pharmingen). pSIR-hCD2 or

m-c-*myc* gRNA #1/pSIR-hCD2 was transfected into Plat-E cells (Morita *et al.* 2000) along with

pPAM3 (Miller & Buttimore 1986) using Lipofectamine 3000 (Invitrogen) to produce retroviral

particles. Activated CD4[+] T cells were transduced with the retroviral particles by the spin

infection method (Naoe *et al.* 2007). After culturing for 2 days in RPMI complete media

containing mouse IL-2 (20 ng/ml, 402-ML, R & D Systems), hCD2[+] cells were purified using

human CD2 MicroBeads (130-091-114, Miltenyi) and used for enChIP analysis.

enChIP real-time PCR

enChIP real-time PCR was performed as described (Fujita & Fujii 2013). Primers used in the

analysis are shown in Table 1.

Acknowledgments

We thank S. Muroi for genotyping ES cells, T. Ishikura for injection of ES cells into blastocysts,

and K. Rajewsky for providing a plasmid (Addgene plasmid # 15912).

Funding

This work was supported by the Takeda Science Foundation (T.F.), Grant-in-Aid for Scientific

Research (C) (#15K06895) (T.F.), and Grant-in-Aid for Scientific Research (B) (#15H04329) (T.F.,

H.F.), 'Transcription Cycle' (#15H01354) (H.F.) from the Ministry of Education, Culture, Sports,

Science and Technology of Japan.

Abbreviations: enChIP, engineered DNA-binding molecule-mediated chromatin

immunoprecipitation; CRISPR, clustered regularly interspaced short palindromic repeats; dCas9,

catalytically inactive form of Cas9; gRNA, guide RNA; GFP, green fluorescent protein; MS,

mass spectrometry; NGS, next-generation sequencing; Tg, transgenic.

Conflicts of interests

T.F. and H.F. have patents on enChIP ("Method for isolating specific genomic region using

molecule binding specifically to endogenous DNA sequence"; patent number: Japan 5,954,808;

patent application number: WO2014/125668). T.F. and H.F. are founders of Epigeneron, LLC.

Availability of data and materials

All data generated or analyzed during the current study are included in this published article. Tg

mice generated in this study can be obtained from RIKEN BioResource Center.

Authors' contributions

H.F. designed and performed experiments (design and construction of transgenes, flow

cytometric analysis, immunoblot analysis, transduction of gRNA retroviruses), wrote the

manuscript, and directed and supervised the study. T.F. and M.Y. performed enChIP analyses.

N.T. constructed the retrovirus vector expressing gRNA targeting the c-*myc* locus, and performed

transduction of gRNA retroviruses and enChIP analysis. F.K. and M.Y. maintained the mouse

colony. F.K. screened ES cells and genotyped mice. A.O. and M.I. performed CRISPR-mediated

knock-in of 3xFLAG-dCas9-IRES-EGFP transgenes. I.T. generated ES cells harboring the

3xFLAG-dCas9/CTV transgene, chimeric mice, and knock-in mice.

References

Bogdanove, A.J. & Voytas, D.F. (2011) TAL effectors: customizable proteins for DNA targeting.

Science **333**, 1843-1846.

Fujihara, Y., Kaseda, K., Inoue, N., Ikawa, M. & Okabe, M. (2013) Production of mouse pups

from germline transmission-failed knockout chimeras. *Transgenic Res.* **22**, 195-200.

Fujita, T., Asano, Y., Ohtsuka, J., Takada, Y., Saito, K., Ohki, R. & Fujii, H. (2013)

Identification of telomere-associated molecules by engineered DNA-binding molecule-mediated

chromatin immunoprecipitation (enChIP). *Sci. Rep.* **3**, 3171.

Fujita, T. & Fujii, H. (2013) Efficient isolation of specific genomic regions and identification of

associated proteins by engineered DNA-binding molecule-mediated chromatin

immunoprecipitation (enChIP) using CRISPR. *Biochem. Biophys. Res. Commun.* **439**, 132-136.

Fujita, T. & Fujii, H. (2014a) Efficient isolation of specific genomic regions retaining molecular

interactions by the iChIP system using recombinant exogenous DNA-binding proteins. *BMC Mol.*

Biol. **15**, 26.

Fujita, T. & Fujii, H. (2014b) Identification of proteins interacting with genomic regions of

interest in vivo using engineered DNA-binding molecule-mediated chromatin

immunoprecipitation (enChIP). *Bio-Protoc.* **4**, e1124.

Fujita, T., Kitaura, F., Yuno, M., Suzuki, Y., Sugano, S. & Fujii, H. (2017a) Locus-specific ChIP

combined with NGS analysis reveals genomic regulatory regions that physically interact with the

Pax5 promoter in a chicken B cell line. *DNA Res.* **24**, 537-548.

Fujita, T., Yuno, M. & Fujii, H. (2016a) Allele-specific locus binding and genome editing by

CRISPR at the p16INK4a locus. *Sci. Rep.* **6**, 30485.

Fujita, T., Yuno, M. & Fujii, H. (2016b) Efficient sequence-specific isolation of DNA fragments

and chromatin by in vitro enChIP technology using recombinant CRISPR ribonucleoproteins.

Genes Cells **21**, 370-377.

Fujita, T., Yuno, M., Okuzaki, D., Ohki, R. & Fujii, H. (2015) Identification of non-coding RNAs

associated with telomeres using a combination of enChIP and RNA sequencing. *PLoS One* **10**,

e0123387.

Fujita, T., Yuno, M., Suzuki, Y., Sugano, S. & Fujii, H. (2017b) Identification of physical

interactions between genomic regions by enChIP-Seq. *Genes Cells* **22**, 506-520.

Harrison, M.M., Jenkins, B.V., O'Connor-Giles, K.M. & Wildonger, J. (2014) A CRISPR view of

development. *Genes Dev.* **28**, 1859-1872.

Matsumura, H., Hasuwa, H., Inoue, N., Ikawa, M. & Okabe, M. (2004) Lineage-specific cell

disruption in living mice by Cre-mediated expression of diphtheria toxin A chain. *Biochem.*

Biophys. Res. Commun. **321**, 275-279.

Miller, A.D. & Buttimore, C. (1986) Redesign of retrovirus packaging cell lines to avoid

recombination leading to helper virus production. *Mol Cell Biol.* **6**, 2895-2902.

Morita, S., Kojima, T. & Kitamura, T. (2000) Plat-E: and efficient and stable system for transient

packaging of retroviruses. *Gene Ther.* **7**, 1063-1066.

Moriyama, S., Takahashi, N., Green, J., Hori, S., Kubo, M., Cyster, J. & Okada, T. (2014)

Sphingosine-1-phosphate receptor 2 is critical for follicular helper T cell retention in germinal

centers. *J. Exp. Med.* **211**, 1297-1305.

Naoe, Y., Setoguchi, R., Akiyama, K., Muroi, S., Kuroda, M., Hatam, F., Littman, D. & Taniuchi,

I. (2007) Repression of interleukin-4 in T helper type 1 cells by Runx/Cbf beta binding to the Il4

silencer. *J. Exp. Med.* **204**, 1749-1755.

Niwa, H., Yamamura, K. & Miyazaki, J. (1991) Efficient selection for high-expression

transfectants with a novel eukaryotic vector. *Gene* **108**, 193-199.

Oji, A., Noda, T., Fujihara, Y., Miyata, H., Kim, Y., Muto, M., Nozawa, K., Matsumura, T.,

Isotani, A. & Ikawa, M. (2016) CRISPR/Cas9 mediated genome editing in ES cells and its

application for chimeric analysis in mice. *Sci. Rep.* **6**, 31666.

Qi, L.S., Larson, M.H., Gilbert, L.A., Doudna, J.A., Weissman, J.S., Arkin, A.P. & Lim, W.A.

(2013) Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene

expression. *Cell* **152**, 1173-1183.

Schaft, J., Ashery-Padan, R., van der Hoeven, F., Gruss, P. & Stewart, A. (2001) Efficient FLP

recombination in mouse ES cells and oocytes. *Genesis* **31**, 6-10.

Wright, A.V., Nuñez, J.K. & Doudna, J.A. (2016) Biology and Applications of CRISPR Systems:

Harnessing Nature's Toolbox for Genome Engineering. *Cell* **164**, 29-44.

Figure legends

Fig. 1. Generation of Tg mouse lines expressing 3xFLAG-dCas9. (**A**) Scheme of the targeted

locus of 3xFLAG-dCas9-IRES-EGFP. (**B**) Scheme of the targeted loci of (i)

3xFLAG-dCas9/CTV (neo$^+$/GFP$^+$); (ii) 3xFLAG-dCas9/CTV (GFP$^+$); and (iii)

3xFLAG-dCas9/CTV. (**C**) Genotyping PCR of Tg mouse lines.

Fig. 2. Expression of 3xFLAG-dCas9. (**A**) Expression of GFP in thymocytes and splenocytes

from Tg mice. Fluorescence in the FL-1 channel (GFP) is shown for C57BL/6 mice (WT: black)

and Tg mice (Tg: green). Numbers represent mean fluorescence intensities (MFI) in the FL-1

channel. (**B**) Expression of 3xFLAG-dCas9 in thymocytes from Tg mice. Expression of

3xFLAG-dCas9 was detected by immunoblot analysis with anti-FLAG Ab. Coomassie Brilliant

Blue (CBB) staining is shown as a protein loading control.

Fig. 3. enChIP analysis of CD4$^+$ T cells from 3xFLAG-dCas9 Tg mice. (**A**) Scheme of enChIP

analysis using Tg mouse lines. (**B**) Positions of primer sets used for enChIP real-time PCR. (**C**)

Yields of enChIP analysis for the target site. Error bars represent differences between duplicate

analyses. N.D.: not detected.

TABLE

Table 1. Oligodeoxyribonucleotides used in this study

Number	Name	Sequence (5' → 3')	Experiments
27551	*Mlu*I–*Pme*I oligo-S	cacgcgtgtttaaacgagct	Construction of pCAG1.2-PM and pCAG1.2-MP
27552	*Mlu*I–*Pme*I oligo-A	cgtttaaacacgcgtgagct	Construction of pCAG1.2-PM and pCAG1.2-MP
27699	ROSA-5' TF2	ctcagagagcctcggctagg	Genotyping of all Tg mouse lines
27644	Cas9-R	cagggcgataagattaccaaac	Genotyping of 3xFLAG-dCas9-IRES-EGFP
26013	oIMR0013	cttgggtggagaggctattc	Genotyping of 3xFLAG-dCas9/CTV (neo$^+$/GFP$^+$)
28110	Cas9-R 27643	gtaggctttctgcctccttcact	Genotyping of 3xFLAG-dCas9/CTV (neo$^+$/GFP$^+$) and 3xFLAG-dCas9/CTV (GFP$^+$)
27822	mcMYC promoter 1 sense	caccgcgcccgggacgtgcgtgacg	Construction of gRNA targeting c-*myc* promoter: sense strand
27823	mcMYC promoter 1 antisense	aaaccgtcacgcacgtcccgggcgc	Construction of gRNA targeting c-*myc* promoter: antisense strand
28157	mcMYC-0.62k-F	tatacgtggcagtgagttgctga	enChIP real-time PCR (c-*myc*)
28158	mcMYC-0.44k-R	ctctaaggctggggaaaacagaa	enChIP real-time PCR (c-*myc*)
26574	mGAPDH-prom-F	gggttcctataaatacggactgc	enChIP real-time PCR (*Gapdh*)
26575	mGAPDH-prom-R	agcatccctagacccgtacagt	enChIP real-time PCR (*Gapdh*)

Figure 1

A

3xFLAG-dCas9-IRES-EGFP

B

(i) 3xFLAG-dCas9/CTV (neo+/GFP+)

(ii) 3xFLAG-dCas9/CTV (GFP+)

(iii) 3xFLAG-dCas9/CTV

C

(a) ROSA-5' TF2 (27699) x Cas9-R (27644)

(b) ROSA-5' TF2 (27699) x oIMR0013 (26013)

(c) ROSA-5' TF2 (27699) x Cas9-R 27643 (28110)

lanes

1, 3, 6: C57BL/6

2:      3xFLAG-dCas9-IRES-EGFP

4, 7:   3xFLAG-dCas9/CTV (neo+/GFP+)

5, 8:   3xFLAG-dCas9/CTV (GFP+)